



LJMU Research Online

Allen, L, Reeve, R, Nousek-McGregor, A, Villacampa, J and MacLeod, R

Are orchid bees useful indicators of the impacts of human disturbance?

<http://researchonline.ljmu.ac.uk/id/eprint/10703/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Allen, L, Reeve, R, Nousek-McGregor, A, Villacampa, J and MacLeod, R (2019) Are orchid bees useful indicators of the impacts of human disturbance? Ecological Indicators, 103. pp. 745-755. ISSN 1470-160X

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 **Are orchid bees useful indicators of the impacts of**
2 **human disturbance?**

3 Authors: Laura Allen ^{a, b}, Richard Reeve ^{a, b}, Anna Nousek-McGregor ^c, Jaime Villacampa,
4 Ross MacLeod ^a.

5 ^a Institute of Biodiversity, Animal Health and Comparative Medicine, College of Medical,
6 Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

7 ^b Boyd Orr Centre for Population and Ecosystem Health, University of Glasgow, Glasgow,
8 UK

9 ^c School of Life Sciences, University of Glasgow, Glasgow, UK

10 Corresponding author: Laura Allen, email: laurajallen@hotmail.com

11

12 **Abstract**

13 Biodiversity and ecosystem functions are threatened by human disturbance, and tropical
14 forests are one the most vulnerable habitats. Monitoring the impacts of disturbance and
15 the success of conservation projects is crucial, and to do this effectively it is important to
16 identify suitable measures that are sensitive to ecosystem disturbance. Orchid bees
17 (Euglossini) are a specialist group with mutualistic relationships with many plant species
18 and can fly long distances, making them important pollinators of widely dispersed plant
19 species. A loss of specialist pollinators such as these could have severe consequences for
20 the plants that rely on their services. We therefore aimed to answer the following
21 question: are orchid bees useful indicators of the impacts of human disturbance? If so,
22 what measures of orchid bee diversity are most sensitive? And do orchid bees provide any
23 indication of changes in pollination services along a disturbance gradient? Orchid bees
24 were collected from 18 sites across a gradient of disturbance in a tropical forest region in
25 southeast Peru. Alpha diversity across the gradient was compared using Hills numbers.
26 Beta diversity was assessed using community composition, species contributions to beta
27 diversity, beta diversity partitioning and novel measures of redundancy and
28 representativeness. The potential pollination services available at each site were
29 measured using artificial flowers and counts of pollinator visits. Alpha diversity of orchid
30 bees showed low sensitivity to disturbance. Beta diversity measures were more
31 informative, with disturbed sites found to be highly redundant in the ecosystem compared
32 to the less disturbed sites. However, the most sensitive measure across the gradient was
33 abundance - there was a significant decrease in the number of bees caught as disturbance
34 increased, with likely consequences for pollination services. These results suggest that
35 orchid bees may be useful indicators of the impacts of human disturbance, but alpha
36 diversity is a poor metric for this purpose. In order to understand how human disturbance
37 is affecting biodiversity, multiple diversity indices should be considered, and in the case of
38 orchid bees, redundancy and abundance could be useful for detecting sensitive responses
39 to forest disturbance.

40 **Keywords**

41 Biodiversity indices, indicator, Euglossini, disturbance, agriculture, rainforest.

43 **1.1 Introduction**

44 Covering only a small percentage of the world's surface, tropical forest harbours over 50%
45 of terrestrial biodiversity. Yet this cover is decreasing globally, with forest being cut down
46 for timber and conversion to agriculture (FAO, 2016), with severe consequences for
47 biodiversity (Alroy, 2017; Gibson et al., 2011) and ecosystem functions (DeFries et al.,
48 2004). It is therefore of the utmost importance that we have effective tools for detecting
49 changes in biodiversity and ecosystem function in response to anthropogenic disturbance
50 (Feest et al., 2010; Hill et al., 2016; Kremen, 2005; Pimm and Raven, 2000). Indicator
51 groups have been widely used for this purpose (Cleary, 2004; Krug et al., 2017; Schulze et
52 al., 2004), however choosing suitable indicator groups can be challenging (Broszeit et al.,
53 2017; Fleishman and Murphy, 2009; Gao et al., 2015). Ideally they should be efficient to
54 survey, show a predictable, sensitive response to environmental change, correlate well
55 with overall biodiversity responses and play an important role in the ecosystem (Brown,
56 1997; Hilty and Merenlender, 2000).

57 Bees are the most important group of pollinators (Bawa, 1990) but have been declining
58 globally (Allen-Wardell et al., 1998; Potts et al., 2010), with land conversion and habitat
59 loss the leading causes (Winfrey et al., 2009). Orchid bees (Hymenoptera: Apidae:
60 Euglossini) are found throughout the Neotropics and are one of the best studied groups of
61 bees in the region. They exhibit many of the recommended features that could make them
62 suitable indicators of disturbance impacts (Brown 1997; Favila & Halffter 1997; Gardner et
63 al. 2008a; Goodsell, Underwood & Chapman 2009), including being widespread, common,
64 cost-effective, easy to sample using standardised methods (Pearson and Dressler, 1985)
65 and having well developed taxonomic literature (eg. Dressler 1982a; Roubik 2004; Nemésio
66 & Silveira 2007a). They have close associations with plant species and play important
67 ecological roles as pollinators, able to access flowers that are unavailable to many other
68 insects (Dressler, 1982a), due to their long tongues and also the large body size of some
69 species. They can fly long distances, making them valuable long-distance pollinators
70 (Janzen, 1971) for many widely-spaced plant species and many orchids are entirely
71 dependent on orchid bees for their pollination (Dressler, 1982a). Another key feature of a
72 good indicator is that they should display a clear, graded response to environmental
73 change. There is some evidence that orchid bees display graded responses to human
74 disturbance, but the results have been mixed and part of our motivation is to study this in

75 greater detail. Orchid bees can persist in heavily degraded ecosystems in some cases,
76 including forest fragments (Storck-Tonon et al., 2013) and farmland (Otero and Sandino,
77 2003; Sandino, 2004), although this may depend on the proximity to intact forest (Briggs et
78 al., 2013). At the same time, there is concern that orchid bees are sensitive to forest loss
79 (Roubik and Hanson, 2004) and some species are thought to be at risk of extinction
80 (Nemesio, 2013), which would have major consequences for the plants they pollinate. The
81 loss of old-growth forest and canopy disruption is likely to result in a loss of orchid bee
82 resources (Hietz et al., 2006; Nöske et al., 2008), such as orchids, other epiphytes and
83 specific tree saps (Dressler, 1982a; Roubik and Hanson, 2004), with probable negative
84 consequences for the orchid bee species that depend on them. Together these features
85 suggest that orchid bees could be an excellent candidate for use as an indicator of faunal
86 and floral responses to tropical forest disturbance.

87 So far, studies on orchid bee responses to forest disturbance have primarily focused on the
88 effects of forest fragmentation, with mixed patterns identified. The abundance and
89 diversity of euglossine bees have been suggested to decline with decreased forest
90 fragment size (Brosi, 2009; Brosi et al., 2008), but other studies found no effect of
91 fragmentation on abundance or richness (Storck-Tonon et al., 2013). A few studies have
92 compared agricultural land to intact forest but these have mostly been focused on less
93 diverse Central American habitats and generally only compared two or three land uses.
94 Briggs *et al.* (2013) found that polyculture could sustain orchid bee communities similar to
95 forest habitats in composition, but abundance was higher in the polyculture than in either
96 forest or monoculture, although abundance did decline with increasing distance from the
97 forest. Abundance and richness of orchid bees have been found to decrease in some
98 agricultural matrix habitats compared to forests (Aguiar et al., 2015), although others
99 have been found to support similar communities to forest (Rosa et al., 2015) and some
100 found higher capture rates in agricultural areas than in forest (Otero and Sandino, 2003;
101 Sandino, 2004). However, studies of forest disturbance gradients are relatively uncommon,
102 and none have covered the response of orchid bees across a continuous, multi-level
103 gradient from monoculture agriculture to intact forest. There are several studies on the
104 mutualistic relationships between orchid bees and orchids (Dressler 1967; Ackerman 1983),
105 but again there is a lack of studies relating orchid bee diversity to overall pollination
106 services available in the ecosystem.

107 As well as the challenge of selecting suitable indicator groups, another difficulty is
108 deciding how to quantify biodiversity. There are many indices available to measure the
109 alpha, beta and gamma components of diversity. Alpha diversity is usually the main focus
110 of diversity studies, and species richness the most commonly used measure because it is
111 intuitive, simple and often sensitive. However, species richness lends a lot of weight to
112 rare species, while other indices such as Shannon and Simpson diversity place more weight
113 on the more dominant species in the community. Recent discussions on diversity
114 partitioning have concluded that diversity profiles (a set of multiple diversity measures
115 that differ in the relative weight given to rare and common species, and are particularly
116 powerful as a graphical tool) based on Hill numbers (Hill, 1973) could help avoid a narrow
117 focus on a single result (Chao et al., 2012; Jost, 2006). Beta diversity is also often of
118 interest in ecological studies but has an even wider array of indices to choose from than
119 alpha diversity (Anderson et al., 2011; Tuomisto, 2010). Beta diversity measures differ in
120 their approaches to quantifying similarity or dissimilarity between sites, and the
121 importance of species abundance differences and the relative importance of species
122 turnover or richness differences (Socolar et al., 2016). Beta diversity can also be placed
123 into a unified framework for partitioning diversity effectively using diversity profiles
124 (Reeve et al., 2016). The large number of diversity measures available have been
125 developed to fit a broad range of questions and few perform consistently well under
126 different conditions, which means that focusing on a single index may risk overlooking
127 important ecological patterns (Santini et al., 2017). We are therefore proponents of using
128 multiple measures of alpha and beta diversity to maximise our understanding of
129 biodiversity responses to anthropogenic disturbance.

130 In this study, we investigated how orchid bee communities respond to habitat conversion
131 across a gradient of human disturbance, from minimally disturbed tropical forest to
132 banana plantations. We aimed to study how alpha and beta diversity changed across this
133 disturbance gradient and aimed to investigate which measures were most sensitive for
134 detecting orchid bee responses to disturbance. Identifying the aspects of the orchid bee
135 community most sensitive to disturbance can inform which metrics to use when applying
136 orchid bees as indicators of environmental change. Identifying which components of the
137 community respond to disturbance may also help us better understand what they are
138 responding to and pave the way for further investigation into the mechanisms involved.

139 We assessed alpha diversity using Hill numbers and used several approaches to quantify
140 beta diversity, including two recently developed measures, redundancy and
141 representativeness (Reeve et al., 2016). Finally, we studied the activity of general
142 pollinators across the gradient using artificial flowers. We aimed to identify if there was
143 any change in pollinator activity across the disturbance gradient, and if orchid bee
144 diversity provided any indication of the potential general pollination services available in
145 the ecosystem (Engel and Irwin, 2003). We conducted our study in the highly biodiverse
146 Manu Biosphere Reserve, Peru, where there has been limited work on the orchid bee
147 communities, and none to our knowledge on their responses to disturbance.

148 **1.2 Materials and methods**

149 **1.2.1 Study area**

150 This study was based around the Manu Learning Centre (-12.789882, -71.391753, 470 m
151 above sea level), a research station run by the Crees Foundation, in the cultural zone of
152 the Manu Biosphere Reserve in southeast Peru, a UNESCO World Heritage Site. This zone
153 contains a mixture of protected areas of lowland tropical forest interspersed with areas of
154 high human impact, including logging and agriculture, and is intended as a buffer for Manu
155 National Park. Eighteen sites were chosen to represent a gradient of human disturbance
156 from banana monoculture to minimally disturbed tropical forest. A stratified site selection
157 approach ensured sampling was not dominated by a single disturbance history, with three
158 sites chosen for each of the six of major land uses in the local area, covering a gradient of
159 human disturbance (supplementary materials Figure S.1). These were ranked from highest
160 to lowest disturbance intensity based on land use (Beck et al., 2002; Eggleton et al.,
161 2002), with 1 being the most disturbed and 6 the least disturbed (details in Table 1). For
162 each land use type, one of the replicates was named 'A', 'B' or 'C', semi-randomly, taking
163 into consideration suitable site groupings for access and analysis. We sampled all 'A' sites
164 in the first week, then the 'B' and 'C' sites in the following weeks, so that one of each
165 habitat types would be sampled concurrently.

166 At each site, data were collected on the vegetation structure to assess how this changed
167 across the disturbance gradient. Three vegetation points were randomly selected within
168 each site, and a 25m² plot marked out at each. Within this plot, canopy cover was

169 quantified using a quadrat held above the sampler's head and the quadrat used to
170 estimate the percentage of canopy cover at five points within the circle. Understory
171 vegetation density was estimated at the four corners of the sampling plot using the
172 modified Braun-Blanquet scale as described in Hurst and Allen (2007). We counted the
173 number of trees with a diameter >5cm at breast height and measured the diameters of the
174 three largest trees within each vegetation plot, and measured leaf litter depth at 16
175 random points within the plot. These followed the protocols for vegetation assessment
176 used by Whitworth et al., (2016). Weather data were collected at the research station,
177 with temperature, rainfall and humidity data collected daily at 7am, manually checking a
178 thermometer, humidity meter and rain gauge at the weather station located at the
179 research centre each morning. Elevation data for each site were obtained from Google
180 Earth Pro 7.3.2.5491, and distance to the main river, the Alto Madre de Dios, was
181 measured in QGIS 2.18.7 (QGIS Development Team, 2017), measuring the straight line
182 distance from the centre of each site to the nearest edge of the river using the 'measure
183 line' tool and manually comparing the distances to the edges of the river to find the
184 shortest.

185

186

Table 1 - Disturbance gradient description

Disturbance Rank	Disturbance type	Description
1	Banana plantation	Primarily banana monocultures, but occasionally including a few other crops; generally open habitat with low vegetation.
2	Agroforestry	Banana plantations interspersed with native tree species, providing more shade and habitat complexity.
3	Disturbed secondary forest	Uncultivated but heavily disturbed forest lying in between plantations, with more canopy cover and thick understory vegetation.
4	Cleared forest regenerating	Forest once cleared for agriculture and grazing, now regenerating under protection for over 30 years to form a closed canopy with thick understory vegetation.
5	Mixed history forest	Selectively logged forest with small patches cleared for cultivation, regenerating under protection for over 30 years. The canopy is well defined and the understory less dense, but large trees are absent.
6	Minimally disturbed primary forest	Primary forest protected for over 30 years, with little evidence of previous disturbance. The canopy is high and well defined, large trees are present and the understory is more open.

187

188 Sites of the different disturbance types were interspersed in space as much as possible.
 189 The potentially confounding effects of large landscape and climatic differences were
 190 minimised by selecting sites within a small area (20 km²), while ensuring sites of the same
 191 disturbance type were far enough apart (>500 m) to avoid sampling pseudo-replication
 192 (Ramage et al., 2013). To minimise spatial effects on the results we kept groups of sites of
 193 different disturbance levels as close together as possible given other constraints and the

194 locations of these habitats. We used linear models to validate the results of our
195 correlation tests and to check for any significant effect of these other environmental
196 variables, including them as explanatory variables along with disturbance if they had any
197 significant effect on the response.

198 This project was conducted in two parts, the primary study looking at changes in the
199 orchid bee community along the gradient and then a second study exploring potential
200 pollination services across the same sites. We cover the methods and results of the main
201 study first, followed by the pollination study.

202

203 **1.2.2 Orchid bee diversity**

204 **1.2.2.1 Data collection**

205 Orchid bees were sampled in the morning between 09:00-12:00 and in the afternoon
206 between 12:30-15:30 (± 15 mins). Each site was sampled for two morning and two
207 afternoon sessions (a total of 12 hours per site) and these four sampling sessions were
208 pooled to form a single sample for each site (a total of 18 samples). To reduce potential
209 biases from weather or other potential temporal sampling biases, two sites were sampled
210 simultaneously - one more disturbed (rank 1-3) and one less (rank 4-6), and on each day
211 different disturbance types were sampled in the mornings and afternoons.

212 At each site, two sampling stations were set up 50 m apart to reduce any bias from a
213 single within-site location choice. At each of these stations, eight balls of cotton were
214 hung from branches at a height of 1.5 m, with 2 m between cotton balls. Each cotton ball
215 was baited with two drops of one of the following eight attractants: wintergreen oil,
216 methyl salicylate, eucalyptus oil, eucalyptol, vanillin (3 tsp vanillin dissolved in 50 ml 96%
217 ethanol), benzyl acetate, clove oil, eugenol. The bait stations were monitored over the
218 three-hour sampling period, and orchid bees attracted to the baits were caught using hand
219 nets then killed and preserved in 70% ethanol. Orchid bees attracted to the survey area
220 that did not settle at a specific bait but came within 1m of a bait and flitted between
221 baits, were also captured. Multiple researchers conducted the hand netting, three per site
222 each day. This included trained researchers experienced with capturing insects with nets,
223 and less experienced volunteers. To reduce biases from experience, we ensured that the

224 teams surveying the different sites were as balanced as possible, with less experienced
225 people paired with more experienced people and the teams rotated between the sites.
226 Some of the potential issues associated with capture rates are explored in the discussion.

227 Preserved orchid bees were identified in Cusco, using a stereo microscope, published keys,
228 checklists and descriptions (Bonilla-Gomez and Nates-Parra, 1992; Dressler, 1978, 1979,
229 1982b, 1982c, 1982d, 1984; Faria and Melo, 2007; Hinojosa-Díaz and Engel, 2011, 2012,
230 2014; Kimsey, 1979, 1982; Melo, 2014; Moure, 1965; Nemésio, 2011, 2009; Nemésio and
231 Silveira, 2007b; Niemack et al., 2012; Roubik, 2004; Roubik and Hanson, 2004). For as
232 many species as possible, identification was verified by consulting the collections at the
233 Department of Entomology at the Universidad de San Antonio Abad de Cusco, and at the
234 Museo de Historia Natural, Universidad Nacional Mayor de San Marcos, Lima. Reference
235 specimens were deposited in both these collections in September 2016.

236 **1.2.2.2 Analysis**

237 A principal component analysis (PCA) of the vegetation structure data across the gradient
238 was performed using the vegan package v.2.4-6 in R (Oksanen et al., 2018), and we used a
239 Spearman rank correlation test (Spearman, 1904) to check the correlation between
240 disturbance rank and the first principal component of the vegetation PCA.

241 To test if alpha diversity varied with human disturbance, orchid bee alpha diversity was
242 calculated for each site using Hill numbers, a family of diversity measures parameterised
243 by a viewpoint parameter, q , with increasing emphasis on dominant species as the value of
244 q increases (Hill, 1973; Reeve et al., 2016). We calculated diversity at $q = 0, 1, 2$ and ∞ ,
245 as these correspond to the following commonly used diversity measures: species richness,
246 Shannon entropy, Simpson diversity and Berger Parker diversity respectively, so our
247 measures can be easily compared to previous studies.

248 All analyses were conducted using R version 3.3.2 (R Core Team, 2017) in RStudio version
249 1.0.136 (RStudio Team, 2016). Observed alpha diversity was calculated using the package
250 rdiversity v.0.4.3 (Mitchell and Reeve, 2016). Estimates of species richness, Shannon
251 entropy and Simpson diversity were calculated using the package iNEXT v.2.0.14 (Hsieh et
252 al., 2016) to rarefy or extrapolate estimates to a standard sample size of 50 individuals at

253 each site (Chao et al., 2014; Colwell et al., 2012). This sample size represented
254 approximately double the smallest sample size, the maximum that can be reliably
255 extrapolated at $q = 0$ with high confidence (Hsieh et al., 2016). This provided
256 approximately 90% sample coverage for all sites (supplementary materials Figure S.3). The
257 diversity estimates at each value of q were bootstrapped 1000 times using the iNEXT
258 function in R, to calculate a confidence interval around our mean estimates. This allowed
259 us to test for consistency in the direction of change in the diversity estimates while
260 accounting for the uncertainty in the diversity estimates. We did this by extracting the
261 1000 raw estimates of the diversity of each site, and then calculating the proportion of
262 times a less disturbed site was more diverse than a more disturbed site. All disturbance
263 type pairs were compared, and a Spearman rank correlation test (Spearman, 1904) was
264 used to test the correlation between the proportion of times that the less disturbed site
265 was the more diverse of the pair and the difference in disturbance rank between the sites.

266 To understand how beta diversity was affected by human disturbance, we quantified the
267 change between sites along the gradient using several methods, including community
268 composition, beta diversity partitioning and two recently developed beta diversity
269 measures, redundancy and representativeness (Reeve et al., 2016). The total beta
270 diversity of the gradient, measured as the total variance of the community matrix, was
271 calculated using the `beta.div.comp` function in the R package `adespatial` v.0.1-1 (Dray et
272 al., 2018) along with the partitioning of the total beta diversity into nestedness and
273 turnover. A Hellinger transformation of the species x site abundance matrix was used, as
274 the Hellinger distance provides a good compromise between linearity and resolution and
275 has been found to correlate better with ‘true’ distances in simulations than many
276 alternatives (Legendre and Gallagher, 2001). Beta diversity decomposition was then
277 calculated for this matrix using the quantitative form of Sorensen’s dissimilarity
278 coefficient, in order to account for differences in relative abundances as well as species
279 identity (Legendre and De Cáceres, 2013). A redundancy analysis (RDA) was used to
280 examine the change in species composition across the gradient, applied to the Hellinger
281 transformed community matrix and constrained by disturbance rank, elevation and
282 distance to the river, using the `vegan` package in R (Oksanen et al., 2018). The use of this
283 transformation overcomes many of the issues associated with raw Euclidean distances,
284 including many zeros and large differences in abundances, which may lend

285 disproportionate weight to rare species (Borcard et al., 2011; Legendre and Gallagher,
286 2001). Species contributions to beta diversity (SCBD index), which is the relative degree of
287 variation in the abundance of individual species across the study gradient (Legendre and
288 De Cáceres, 2013), were calculated with the beta.div function in the adespatial package,
289 using the Hellinger dissimilarity coefficient (Legendre and De Cáceres, 2013). The
290 abundances of the species with the highest SCBD values were compared across the
291 gradient, with some low SCDB species included for contrast.

292 The redundancy of the communities at each site was calculated using the redundancy (ρ)
293 measure, available in the package rdiversity (Mitchell and Reeve, 2016). This is a measure
294 of beta diversity that represents the extent to which the diversity of the overall
295 metacommunity (the diversity of the whole gradient in this case) would be preserved if a
296 single community or site was lost (Reeve et al., 2016). We also calculated the
297 representativeness ($\bar{\rho}$) of the sites, which is a measure of how well a single site represents
298 the overall metacommunity. Representativeness considers how much of the
299 metacommunity diversity a site holds (i.e. the redundancy) relative to the size of the
300 community at that site, providing a correction for the different sample sizes across the
301 gradient. Both redundancy and representativeness were calculated at $q = 1$, as this
302 provides an intermediate level of conservatism and is a key value of q due to its
303 correspondence to many measures of beta diversity through relative entropy and K-L
304 divergence (Reeve et al., 2016).

305 We used disturbance as a continuous explanatory variable since this allowed us to consider
306 land-use on a continuous spectrum of disturbance intensity, with the possibility of other
307 land-uses falling at intermediate intensities. We could have used an ordered discrete
308 variable, which would have been better able to detect a signal in the presence of
309 unevenness in disturbance differences, but this would have required more data to fit. Our
310 simpler approach is supported by the strong correlation of our continuous disturbance rank
311 with vegetation structure data. Nonetheless, because we could not be sure of the exact
312 difference in disturbance between each level, where possible we used a Spearman rank
313 correlation test to assess the patterns of diversity along this gradient. This approach
314 makes no assumptions regarding the shape of the relationship between the variables, only
315 that the pattern is monotonic along the gradient, and is therefore a conservative

316 approach for these analyses. Spearman rank correlation tests were used to test for
317 correlations of disturbance rank with abundance, observed alpha diversity at $q = 0, 1, 2$
318 and ∞ , estimated alpha diversity at $q = 0, 1$ and 2 , and representativeness and
319 redundancy at $q = 1$. In addition to checking each value of q independently, we tested if
320 the overall pattern of alpha diversity change across the disturbance gradient was
321 significant at $\alpha = 0.05$. To overcome the non-independence of the q values from one
322 another, we used a permutation test to calculate the combined p value for the correlation
323 between disturbance rank and observed alpha diversity at $q = 0, 0.5, 1, 2$ and ∞ and
324 estimated alpha diversity at $q = 0, 0.5, 1$ and 2 ($q = \infty$ could not be included because the
325 estimates all converged at 1 , so there was no variation; we included $q = 0.5$ because q is
326 on a logarithmic scale, so this prevents biased weighting of rare species when calculating
327 the overall pattern of alpha diversity across multiple values of q). The permutation test
328 involved randomly re-labelling the study sites, and then re-calculating the significance of
329 the correlation between disturbance rank (now randomly re-assigned) and alpha diversity
330 at each value of q (Good, 2000). These permuted p -values for each value of q were
331 combined to determine the significance of the overall alpha diversity change across the
332 gradient (Fisher, 1925). This was repeated for 10000 permutations, and the permuted p -
333 values compared to those obtained with the original data, to determine the probability
334 that the observed correlation of diversity could have occurred by chance. To check that
335 the patterns of change in redundancy and representativeness across the gradient were not
336 specific to our chosen value of $q = 1$, we also used a permutation test to obtain the
337 combined p value for the correlation of disturbance rank with redundancy and
338 representativeness calculated at $q = 0, 0.5, 1, 2$ and ∞ .

339 As well as testing for the effect of disturbance on diversity and abundance, the potential
340 effects of altitude, rainfall and distance to river were investigated using linear models, to
341 confirm whether the effects of disturbance identified with the correlation tests remained
342 significant when accounting for these variables. The diversity and abundance response
343 variables were log-transformed and modelled with a gaussian distribution. The model
344 summaries and residuals were inspected to evaluate model fit. None of the additional
345 environmental variables resulted in a significant improvement compared the model that
346 included only disturbance rank. Full details of the models tested can be found in Table S.2
347 of the supplementary materials. In the case of the relationship between abundance and

348 disturbance rank, visual inspection of the data prompted us to also test a quadratic
349 polynomial, but the AIC values indicated that a linear relationship was a better fit ($\Delta AIC =$
350 1.42). Throughout the results section, the statistics reported are the results of the
351 Spearman rank correlation test unless stated otherwise.

352 **1.2.3 Pollination services**

353 **1.2.3.1 Data collection**

354 The potential for the provision of pollination services (from any pollinators) across the
355 gradient was tested using artificial flowers filled with sugar solution (Internicola et al.,
356 2007; Real, 1981). The flowers were 5 cm in diameter and constructed from thin craft
357 foam with a central well containing 1.5 ml of the sugar solution (1:1 sugar and water).
358 Red, blue and yellow flowers were used, with five of each colour on an array that was
359 suspended at a height of 1.3 m at the survey site. Two flower arrays (30 flowers in total)
360 were used at each site, separated by approximately 30 m, and each monitored by a
361 member of the research team. All insects (of any Order) that arrived at the flower array
362 were counted as potential pollination events, apart from individuals that simply moved
363 from one location on the array to another. The flowers were monitored for three periods
364 of 45 minutes separated by 15-minute intervals during which the flowers were covered.
365 This was repeated for four mornings, between 08:45 and 13:00, at each of the 18 sites.

366 **1.2.3.2 Analysis**

367 Spearman rank correlation tests were used to test the correlation between the number of
368 potential pollination events (flower visits) with disturbance rank and with observed orchid
369 bee species richness.


370 **1.3 Results**

371 **1.3.1 Orchid bee diversity**

372 **1.3.1.1 Vegetation structure**

373 The first component of the vegetation structure PCA (PC1) explained 33% of the variation
374 in vegetation structure, with higher rankings (lower disturbance) correlated positively with

375 canopy cover, tree count and diameter, and negatively with leaf litter depth and
 376 understory herb abundance (Figure S.2). There was a strong correlation between
 377 vegetation structure (PC1) and disturbance rank, supporting the disturbance ranking used
 378 to represent human disturbance intensity along the gradient ($p < 0.00001$, $\rho = 0.86$;
 379 Figure 1). The results of all correlation tests are provided in Table S.1 in the
 380 supplementary materials.




381

382 Figure 1 - Correlation between vegetation structure and disturbance rank ($p < 0.001$, $\rho = 0.86$). Disturbance
 383 rank runs from the most disturbed (rank 1) to the least disturbed habitat (rank 6). The first axis of the principal
 384 component analysis of vegetation structure (PC1) was positively correlated with canopy cover, tree count and
 385 diameter, and negatively with leaf litter depth and understory herb abundance.


386 1.3.1.2 Alpha diversity

387 We collected 1783 individuals of 31 species of orchid bee. Overall observed alpha diversity
 388 decreased across the disturbance gradient (permuted combined p value for $q = 0, 0.5, 1, 2$
 389 and ∞ : $p = 0.025$; Figure 2). This was mostly driven by the patterns of alpha diversity at
 390 low values of q , with observed species richness ($q = 0$) and observed Shannon diversity ($q =$
 391 1) both decreasing with increased disturbance across the gradient ($\rho = 0.57$, $p = 0.01$;
 392 $\rho = 0.51$, $p = 0.03$). There was no significant change in alpha diversity across the
 393 gradient at higher values of q . However, when estimated diversity was examined,

394 correcting for sample size ($n = 50$), there were no significant differences in estimated
 395 diversity across the disturbance gradient, either overall (permuted combined p value for q
 396 $= 0, 0.5, 1$ and 2 : $p = 0.330$) or for any individual value of q , due to the wide confidence
 397 intervals around the mean estimates (Figure 3 and Figure S.4).



398
 399 Figure 2 – Orchid bee observed alpha diversity across the disturbance gradient. Increasing values of q indicate
 400 increased emphasis on dominant species ($q = 0 =$ species richness, $1 =$ Shannon diversity, $2 =$ Simpson
 401 diversity, $\infty =$ Berger Parker diversity). Disturbance rank is shown by a colour gradient, dark red for the most
 402 disturbed sites, dark blue for the least disturbed. Overall diversity was significantly higher in less disturbed sites
 403 (permuted combined p value = 0.025), and at $q = 0$ ($\rho = 0.57$, $p = 0.01$) and $q = 1$ ($\rho = 0.51$, $p = 0.03$), but
 404 not significantly different at or above $q = 2$ ($\rho = 0.45$, $p = 0.06$) Because q is on a log scale, the broken axis
 405 and dashed lines indicate inferred values as diversity was calculated only for the values at either side of the
 406 break (0 and ∞); along the solid line, q was calculated at intervals of 0.1.




407
 408 Figure 3 – The correlation between diversity estimates and disturbance ranking. Using the iNEXT package in R,
 409 we generated 1000 estimates for the diversity of each site. We then compared all possible habitat pairs to
 410 calculate the proportion of times that the less disturbed habitat was estimated to have a higher diversity than
 411 the more disturbed habitat, shown on the y axis (p^qD). This was calculated for diversity at a) $q = 0$ ($p = 0.33$; $\rho = 0.27$),
 412 b) $q = 1$ ($p = 0.37$; $\rho = 0.25$), and c) $q = 2$ ($p = 0.42$, $\rho = 0.22$); $n = 50$. The difference in disturbance
 413 ranking between the sites is shown along the x-axis – high values mean the sites are further apart along the
 414 disturbance gradient.

415
 416

417 1.3.1.3 Abundance

418 Abundance of orchid bees declined across the disturbance gradient ($\rho = 0.63$, $p = 0.005$),
 419 with less than a quarter of the number of bees found in the most disturbed habitat
 420 compared to the best of the less disturbed forest sites (Figure 4).




421 Figure 4 – Change in orchid bee abundance across the disturbance gradient. Abundance is the number of
 422 orchid bees captured at each site, plotted against the disturbance rank of the site ($\rho = 0.63$, $p = 0.005$).
 423 Disturbance rank runs from 1 (most disturbed) to 6 (least disturbed).
 424

425 1.3.1.4 Beta diversity


426 Although there was some change in the number of orchid bee species found across the
 427 disturbance gradient, beta diversity was driven primarily by a turnover of species, rather
 428 than a change in richness. The results of the beta diversity partitioning were a total beta
 429 diversity of 0.18 across the gradient (a total beta diversity of 1 would indicate that the
 430 sites had completely distinct communities; Legendre & De Cáceres 2013), with 77% of this
 431 due to species turnover and 22% due to differences in richness (nestedness).

432 The composition of the orchid bee community changed across the gradient, with different
 433 communities found in the more and less disturbed sites, as demonstrated by a separation
 434 along the RDA1 axis (Figure 5), which captured 37% of the variation in community
 435 composition. The Species Contributions to Beta Diversity (SCBD) index identified the

436 species that changed most in abundance along the gradient (Figure 6). *Euglossa chalybeata*
 437 and *Euglossa orellana* appear to be forest specialists that are lost as forest disturbance
 438 increases, whereas *Euglossa despecta* appears to favour intermediate levels of
 439 disturbance.



440 Figure 5 – RDA of orchid bee community composition across the disturbance gradient, constrained by
 441 disturbance rank, elevation and distance to the river. Disturbance rank is represented by a colour gradient from
 442 dark red (most disturbed) to dark blue (least disturbed), with the rank of each site also shown numerically.
 443




444
445
446
447
448

Figure 6 – Differences in abundance across the gradient of the species that contribute most to beta diversity. A total of 31 species were found. The grey bars (a-e) show the top five contributors to beta diversity, in order of contribution. Two examples of low contributors to beta diversity are shown in white (n and o) for comparison. Disturbance rank runs from 1-6, with 1 representing the most disturbed sites.

449
450
451
452
453
454
455


The less disturbed sites hold communities that are less redundant (Figure 7a; at $q = 1$: $p = 0.02$, $\rho = -0.55$; permuted combined p value for $q = 0, 0.5, 1, 2$ and ∞ : $p = 0.016$) and more representative of the overall metacommunity (Figure 7b; at $q = 1$: $p = 0.04$, $\rho = 0.4922$; permuted combined p value for $q = 0, 0.5, 1, 2$ and ∞ : $p = 0.024$) than the more disturbed sites. This suggests that the overall diversity of the region (all sites across the gradient) would suffer a greater loss should the community of one of the less disturbed sites be lost, than if the community of a highly disturbed site was lost.



456
 457 Figure 7 – Change in the a) redundancy (ρ) of sites ($p = 0.02$, $\rho = -0.55$) and b) representativeness ($\bar{\rho}$) of
 458 sites ($p = 0.04$, $\rho = 0.49$) across the disturbance gradient at $q = 1$. The disturbance rank runs from 1 (most
 459 disturbed) to 6 (least disturbed).

460 1.3.2 Pollination services

461 There was no relationship between the number of visits from potential pollinator insects
 462 and disturbance rank (Figure 8a; $\rho = 0.1160$, $p = 0.6467$) or orchid bee diversity (Figure
 463 8b; $\rho = -0.19$, $p > 0.4$ at $q = 0, 1$ and 2). The artificial flowers received similar numbers
 464 of visits across the disturbance gradient, with slightly more visits occurring in lower
 465 disturbance sites but with a large amount of variance within each disturbance level. We
 466 found no evidence that orchid bee diversity (specialised pollinators) indicated any trend in
 467 the activity of pollinators in general.



468
469 Figure 8 – Potential pollination events. Number of insect visits to artificial flower arrays correlated with (a)
470 disturbance rank ($\rho = 0.12$, $p = 0.65$) and (b) observed orchid bee species richness ($q = 0$; $\rho = -0.20$, $p =$
471 0.44). Disturbance ranking runs from 1 (most disturbed) to 6 (least disturbed).

472 1.4 Discussion

473 Orchid bees declined in abundance as disturbance increased, and community composition
474 changed across the gradient, suggesting that orchid bees can be useful as indicators of the
475 impacts of human disturbance. Redundancy of the orchid bee community at a site
476 increased with increased disturbance and representativeness of the sites decreased. No
477 changes in species richness or higher order alpha diversity could be detected after
478 controlling for sample sizes, suggesting that, unlike many other groups, alpha diversity of
479 orchid bees is unlikely to be a strong indicator of human disturbance impacts. We found no
480 change in pollinator visits in response to disturbance nor any correlation between
481 pollinator visits and orchid bee species richness.

482 Previous studies that suggested orchid bees show little response to human disturbance
483 focused on different disturbance types and fewer sites with less replication than covered
484 in our study (Nemésio and Silveira, 2006; Rasmussen, 2009). The first of these studies
485 sampled a similar number of bees as in our study over a longer period in only six sites,
486 focusing on the effect of distance to the forest edge. They found no change in observed
487 species richness, and erratic differences in abundance, uncorrelated with distance to
488 edge, but they did find some change in community composition, though many species were

489 shared across sites (Nemésio and Silveira, 2006). The second study considered alpha
490 diversity at $q = 0, 1$ and 2 as well as abundance and community similarity between sites
491 and found no significant relationship between any of these with disturbance level
492 (Rasmussen, 2009). However, only three sites were compared, one of each disturbance
493 type: primary forest, a reforested site and one heavily disturbed forest. Like these studies,
494 we also found no significant change in alpha diversity after controlling for sample size.
495 However, it should be noted that this lack of a significant response appeared to be
496 because there were many more bees caught in the less disturbed sites, consistent with the
497 marked decrease in orchid bee abundance with increased disturbance we detected. When
498 comparing observed alpha diversity, without controlling for the number of bees caught,
499 there was a decrease in species richness and Shannon diversity as disturbance increased.
500 Since standardised sampling effort was used there is no reason to suspect the observed
501 differences were caused by any bias in the sampling design, and so this difference should
502 not be dismissed. That there are fewer bees in the disturbed sites means that the
503 confidence interval around the estimated true number of species present is large, however
504 if sampling time was extended to catch the same number of bees as found in the least
505 disturbed sites, it would be unsurprising if there were still fewer species (supplementary
506 materials Figure S.4). In any case, the fact that during a set time period, fewer individuals
507 of fewer species are likely to visit plants in disturbed habitats is an ecologically
508 meaningful result, as it is likely to impact plant pollination, even if similar total numbers
509 of species could eventually be detected over a longer time.

510 This kind of study is prone to problems with biases arising from study design and the
511 assistance of volunteers. As well as training, and pairing experienced staff with
512 inexperienced samplers, we randomised the location of the teams to avoid systematic
513 biases. The high turnover of volunteers allowed us to keep this strategy constant over the
514 sampling period. Capture success was high, and although there were some escapes, many
515 of these individuals could be seen leaving and returning to the bait, allowing a second
516 capture attempt. We did not record miss rates but there was no noticeable difference
517 observed between genera or species identifiable in flight, although this would be valuable
518 to investigate in future studies. Another known issue in studies like this is the potential for
519 differences in the attractive radius of the baits between sites to affect capture rates,
520 because differences in vegetation structure and shade between the sites could affect the

521 evaporation rates and how much the baits might be carried on the wind. Further research
522 is needed to fully address this (Nemésio, 2012), but we followed standard practice from
523 previous studies in using standardised bait protocols across all sites, including regularly
524 refreshing the bait to avoid evaporation issues. Since capture rate was higher in the less
525 open forest, there was also no evidence for sampling issues related to scent dispersal
526 being inhibited by vegetation. Due to this study region consisting of a matrix of different
527 habitat types, it was unavoidable that some sampling sites would not be far from other
528 disturbance types. We set up the bait stations near the middle of a habitat type to ensure
529 that the majority of bees were likely to be attracted from within that site, with a
530 minimum of 100m to the neighbouring habitat. It is plausible that some bees may have
531 been attracted from neighbouring habitats, yet this should theoretically weaken our power
532 to detect a correlation between the orchid bee community and habitat type, lending
533 confidence to the patterns that we have detected, as discussed in Brosi (2009).
534 Furthermore, for ecological purposes we care about the bees that visit a site to provide
535 pollination services, so it is still relevant if some bees visit from neighbouring habitats, as
536 we are interested in the disturbance impacts on orchid bees in the context of this matrix
537 landscape.

538 Regarding the value of orchid bees as indicators, in this study we have shown that changes
539 in the orchid bee population are strongly correlated with intensity of forest disturbance,
540 and identified which measures are most useful for detecting these responses. The results
541 of the vegetation surveys showed a loss of canopy cover and number of large trees as
542 disturbance increased (Figure S.2). This may also result in a change in microclimate and a
543 loss of epiphytes. A loss of resources such as nectar and nesting habitats may have
544 contributed to the decline in orchid bee abundance, and a change in the plant species
545 composition could have influenced the change in orchid bee species along the gradient. A
546 more detailed exploration of the mechanisms behind the orchid bee responses are
547 important in order to understand what exactly they are indicative of: is it a change in
548 vegetation structure (shown to be closely correlated with disturbance in our study), a
549 decline in overall biodiversity, a loss of important nesting habitats, a loss of epiphytes or
550 particular flower species? These are important points for further investigation, as there is
551 a general lack of information about the relationship between ecological indicators and
552 indicandum (Gao et al., 2015). It would also be worth expanding on research assessing how

553 far these disturbance effects penetrate into forest edges (Nemesio and Silveira, 2006),
554 considering the different indices used in this study and comparing different land uses and
555 regions. For basic assessments, orchid bee abundance is a simple measure that can be
556 monitored by conservation managers without the need for much analytical expertise or
557 even species identification. Although this should not be used in isolation, it could serve as
558 a useful warning flag of negative impacts of disturbance. The Hill numbers diversity
559 measures in the rdiversity package require more technical skills but allow efficient
560 calculation of a broad suite of indices, including alpha and beta diversity (redundancy and
561 representativeness) at multiple values of q , using a single input of the species x site data
562 matrix and only simple code modifications to calculate multiple indices.

563 Species richness has been recognised as a poor index for detecting the effect of
564 disturbance on some other taxonomic groups, and it has been suggested that community
565 composition may be more sensitive (Stork et al., 2017). For example, Samejima *et al.*
566 (2004) found that stingless bees (Meliponini) showed a change in community composition in
567 response to human disturbance. We also found that community composition changed
568 across the disturbance gradient, but community composition can be expected to change
569 for many reasons, including non-disturbance related changes in vegetation and other
570 habitat features. Due to the small spatial scale of this study and the history of the area,
571 we expect that the habitats of the study sites would have been very similar prior to the
572 anthropogenic disturbance; this assumption is less likely to hold true across larger spatial
573 scales, making it harder to link changes in species composition to habitat disturbance
574 across space. However, as a potential indicator, identifying shifts in orchid bee community
575 composition could be useful for monitoring a site over time, to provide an indication of
576 whether the site is being disturbed to a degree that is negatively impacting the ecosystem
577 (Santini et al., 2017).

578 In ecological terms, reduced orchid bee abundance in highly disturbed habitats is of
579 concern, as it is indicative of a potential cascade effect resulting from the loss of forest
580 canopy, along with habitat complexity and epiphytic diversity (Barthlott et al., 2001). This
581 could result in a loss of specialist pollinators that are crucial for the persistence of many
582 plant species, which may reduce the resilience of the remaining degraded forest, as the
583 ecological networks have been weakened. However, the services provided by orchid bees
584 may be partially maintained by a well-connected patchwork of habitats including high

585 quality forest, as the dispersal distances of this group (Janzen, 1971) enables them to
586 make opportunistic visits to degraded habitats when resources are available. This way,
587 they may be able to provide pollination services to habitats that possibly lack the
588 resources to support viable orchid bee populations independently, but this will require a
589 substantial area of intact forest to be maintained nearby.

590 We found that the redundancy (ρ) of the sites increased with increased disturbance.
591 Redundancy is therefore likely to be a useful measure of disturbance impacts, because it
592 considers both the species present and their abundance, so gives a more complete picture
593 of how the sites differ. From these results, we can see that the impact of losing a
594 minimally disturbed site would more severely impact the overall diversity of the region
595 than the loss of a highly disturbed site. Similarly, when we considered the
596 representativeness ($\bar{\rho}$) of the sites at $q = 1$, we found that in each of the less disturbed
597 sites (ranks 4-6) about 80% of the overall biodiversity of the study area could be found, on
598 average, whereas the more disturbed sites (ranks 1-3) only held an average of about 60%.

599 Pollination potential showed no correlation with orchid bee abundance or alpha diversity
600 (Figure 8; supplementary materials Table S.1). This is probably because visitors to the
601 flower arrays were mostly sweat bees (family Halictidae) and these visited the flowers in
602 high abundance. Other visitors included flies, wasps, butterflies, a few orchid bees, and
603 even a hummingbird on one occasion. An important caveat in interpreting this experiment
604 was that we were only able to measure the potential opportunities for pollination (the
605 number of visits the flower received), and were not able to account for the fact that
606 insects differ widely in their effectiveness as pollinators (Ivey et al., 2003; King et al.,
607 2013; Primack and Silander, 1975; Ramsey, 1988; Schemske and Horvitz, 1984). Turnover
608 of species between sites also means that higher bee diversity than expected is likely to be
609 required to deliver pollination services over large spatial scales (Winfree et al., 2018).
610 Halictid bees are considered valuable pollinators and visit many different plant species
611 (Lindsey, 1984), although they do not show such distinctive host adaptations as the orchid
612 bees. Orchid bees do pollinate a wide range of plants; however, they are particularly
613 important due to their many species-specific relationships and cannot be easily substituted
614 by more generalist pollinators. Further research into the pollination networks of orchid
615 bees and the plants they visit would be helpful to understand how a loss or change in the

616 orchid bee community might impact the plant community. Exploring functional differences
617 (e.g. tongue length and body size) between the orchid bee species affected by disturbance
618 would also add valuable information on the contributions of specific orchid bee species as
619 specialist pollinators.

620 **1.5 Conclusion**

621 Orchid bees show a clear negative response to human disturbance across a tropical forest-
622 agricultural gradient. They are also efficient to sample and play a key role in pollination
623 services. We therefore suggest they can be a useful addition to the indicator groups
624 available for studying the impacts of forest loss on biodiversity and ecosystem functioning.
625 In choosing whether to use orchid bees in future biodiversity studies it should be noted
626 that it is important to consider more than one indicator group in any assessment (Hilty
627 and Merenlender, 2000; Lawton et al., 1998). When orchid bees are selected as a suitable
628 indicator group, our results show that abundance, redundancy (ρ) and representativeness
629 ($\bar{\rho}$) provide the most sensitive measures for detecting the response of orchid bees to
630 human disturbance. In order to understand the response of biodiversity to human
631 disturbance, it is essential to consider the response measures carefully, as a measure that
632 works well for one group may not always be the best for another, and often multiple
633 indices are necessary.

634

635 **1.6 Acknowledgements**

636 Collection permits were provided by the Peruvian National Forestry and Wildlife Service,
637 SERFOR (N° 0233-2016-SERFOR-DGGSPFFS and N° 041-2014-SERFOR-DGGSPFFS). Permission
638 was also obtained from the landowners whose properties were used as sample sites. Claus
639 Rasmussen and Jeanne Robinson provided helpful advice on collection, specimen
640 preparation and identification. Diana Silva Dávila and Evelyn Sanchez Sandoval helped with
641 access to collections and processing our deposited specimens at the Museo de Historia
642 Natural, Universidad Nacional Mayor de San Marcos. Araseli Elme Tumpe and Edgar
643 Montesinos helped with identification and preparation of specimens. Erick Yabar and
644 Abdhel Bustamante provided advice and access to facilities at the Universidad Nacional de
645 San Antonio Abad del Cusco. Staff and volunteers of the Crees Foundation provided
646 valuable assistance with fieldwork.

647 **1.7 Funding**

648 This work was funded by the University of Glasgow, the Crees Foundation and the T & J
649 Meyer Foundation, with additional support from the Darwin Initiative Sustainable Manu
650 project and BBSRC grant BB/P004202/1.

651 **1.8 Competing interests**

652 The authors have no competing interests to declare.

653 **1.9 Figures**

654 Figure 2 to be printed in colour.

655 **1.10 Data Access**

656 The data have been made publicly available and can be accessed from the University of
657 Glasgow Enlighten repository: <http://dx.doi.org/10.5525/gla.researchdata.589>

658

1.11 References

659

- 660 Ackerman, J.D., 1983. Specificity and mutual dependency of the orchid-euglossine bee
661 interaction. *Biol. J. Linn. Soc.* 20, 301-314.
- 662 Aguiar, W.M., Sofia, S.H., Melo, G., Gaglianone, M., 2015. Changes in Orchid Bee
663 Communities Across Forest- Agroecosystem Boundaries in Brazilian Atlantic Forest
664 Landscapes. *Environ. Entomol.* 1-7. doi:10.1093/ee/nvv130
- 665 Allen-Wardell, A.G., Bernhardt, P., Bitner, R., Burquez, A., Cane, J., Cox, P.A., Dalton,
666 V., Feinsinger, P., Ingram, M., Jones, C.E., Kennedy, K., Kevan, P., Koopowitz, H.,
667 Medellin, R., Medellin-Morales, S., Nabhan, G.P., Pavlik, B., Tepedino, V., Torchio, P.,
668 1998. The Potential Consequences of Pollinator Declines on the Conservation of
669 Biodiversity and Stability of Food Crop Yields. *Conserv. Biol.* 12, 8-17.
- 670 Alroy, J., 2017. Effects of habitat disturbance on tropical forest biodiversity. *Proc. Natl.*
671 *Acad. Sci.* 114, 6056-6061. doi:10.1073/pnas.1611855114
- 672 Anderson, M.J., Crist, T.O., Chase, J.M., Vellend, M., Inouye, B.D., Freestone, A.L.,
673 Sanders, N.J., Cornell, H.V., Comita, L.S., Davies, K.F., Harrison, S.P., Kraft, N.J.B.,
674 Stegen, J.C., Swenson, N.G., 2011. Navigating the multiple meanings of beta diversity:
675 A roadmap for the practicing ecologist. *Ecol. Lett.* 14, 19-28. doi:10.1111/j.1461-
676 0248.2010.01552.x
- 677 Barthlott, W., Schmit-Neuerbug, V., Nieder, J., Engwald, S., 2001. Diversity and
678 Abundance of Vascular Epiphytes: A Comparison of Secondary Vegetation and Primary
679 Montane Rain Forest in the Venezuelan Andes. *Plant Ecol.* 152, 145-156.
- 680 Bawa, K., 1990. Plant-Pollinator Interactions in Tropical Rain Forests. *Annu. Rev. Ecol.*
681 *Syst.* 21, 399-422.
- 682 Beck, J., Schulze, C.H., Linsenmair, K.E., Fiedler, K., 2002. From forest to farmland:
683 diversity of geometrid moths along two habitat gradients on Borneo. *J. Trop. Ecol.* 18,
684 33-51. doi:10.1017/S026646740200202X
- 685 Bonilla-Gomez, M.A., Nates-Parra, G., 1992. Abejas Euglosinas de Colombia (Hymenoptera:
686 Apidae) 1. Claves Ilustradas. *Caldasia* 17, 149-172.
- 687 Borcard, D., Gillet, F., Legendre, P., 2011. *Numerical Ecology with R.* Springer Science and
688 Business Media, New York Dordrecht London Heidelberg. doi:10.1007/978-1-4419-7976-
689 6
- 690 Briggs, A.H.M., Perfecto, I., Brosi, B.J., 2013. The Role of the Agricultural Matrix: Coffee
691 Management and Euglossine Bee (Hymenoptera: Apidae: Euglossini) Communities in
692 Southern Mexico. *Environ. Entomol.* 42, 1210-1217.
- 693 Brosi, B.J., 2009. The effects of forest fragmentation on Euglossine bee communities
694 (Hymenoptera: Apidae: Euglossini). *Biol. Conserv.* 142, 414-423.
695 doi:10.1016/j.biocon.2008.11.003

- 696 Brosi, B.J., Daily, G.C., Shih, T.M., Oviedo, F., Durán, G., 2008. The effects of forest
697 fragmentation on bee communities in tropical countryside. *J. Appl. Ecol.* 45, 773-783.
698 doi:10.1111/j.1365-2664.2007.01412.x
- 699 Broszeit, S., Beaumont, N.J., Uyarra, M.C., Heiskanen, A.S., Frost, M., Somerfield, P.J.,
700 Rossberg, A.G., Teixeira, H., Austen, M.C., 2017. What can indicators of good
701 environmental status tell us about ecosystem services?: Reducing efforts and increasing
702 cost-effectiveness by reapplying biodiversity indicator data. *Ecol. Indic.* 81, 409-442.
703 doi:10.1016/j.ecolind.2017.05.057
- 704 Brown, K.S., 1997. Diversity, disturbance, and sustainable use of Neotropical forests:
705 insects as indicators for conservation monitoring. *J. Insect Conserv.* 1, 25-42.
- 706 Chao, A., Chiu, C., Hsieh, T.C., 2012. Proposing a resolution to debates on diversity
707 partitioning. *Ecology* 93, 2037-2051.
- 708 Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E., Colwell, R.K., 2014. Rarefaction and
709 extrapolation with Hill numbers: a framework for sampling and estimation in species
710 diversity studies. *Ecol. Monogr.* 84, 45-67.
- 711 Cleary, D.F.R., 2004. Assessing the use of butterflies as indicators of logging in Borneo at
712 three taxonomic levels. *J. Econ. Entomol.* 97, 429-435. doi:10.1603/0022-0493-
713 97.2.429
- 714 Colwell, R.K., Chao, A., Gotelli, N.J., Lin, S.Y., Mao, C.X., Chazdon, R.L., Longino, J.T.,
715 2012. Models and estimators linking individual-based and sample-based rarefaction,
716 extrapolation and comparison of assemblages. *J. Plant Ecol.* 5, 3-21.
717 doi:10.1093/jpe/rtr044
- 718 DeFries, R.S., Foley, J.A., Asner, G.P., 2004. Land-use choice: balancing human needs and
719 ecosystem function. *Front. Ecol. Environ.* 2, 249-257. doi:10.1890/1540-
720 9295(2004)002[0249:LCBHNA]2.0.CO;2
- 721 Dray, S., Blanchet, G., Borcard, D., Clappe, S., Guenard, G., Jombart, T., Larocque, G.,
722 Legendre, P., Madi, N., Wagner, H.H., 2018. *adespatial: Multivariate Multiscale Spatial*
723 *Analysis.*
- 724 Dressler, R.L., 1967. Why do euglossine bees visit orchid flowers? *Atas do Simpósio sobre a*
725 *Biota Amaz.* 5, 171-180.
- 726 Dressler, R.L., 1978. An infrageneric classification of *Euglossa*, with notes on some
727 features of special taxonomic importance (Hymenoptera; Apidae). *Rev. Biol. Trop.* 26,
728 187-198.
- 729 Dressler, R.L., 1979. *Eulaema bombiformis*, *E. meriana* and Mullerian Mimicry in Related
730 Species (Hymenoptera: Apidae). *Biotropica* 11, 144-151.
- 731 Dressler, R.L., 1982a. Biology of the Orchid Bees (*Euglossini*). *Annu. Rev. Ecol. Syst.* 13,
732 373-394. doi:10.1146/annurev.es.13.110182.002105
- 733 Dressler, R.L., 1982b. New species of *Euglossa* II (Hymenoptera: Apidae). *Rev. Biol. Trop.*
734 30, 121-129.

- 735 Dressler, R.L., 1982c. New species of *Euglossa* III. The *bursigera* species group. *Rev. Biol.*
736 *Trop.* 30, 31-40.
- 737 Dressler, R.L., 1982d. New species of *Euglossa* IV. The *cordata* and *purpurea* species groups
738 (Hymenoptera: Apidae). *Rev. Biol. Trop.* 30, 141-152.
- 739 Dressler, R.L., 1984. *Euglossine* Bees of the Tambopata Reserved Zone, Madre de Dios,
740 Peru. *Rev. Peru. Entomol.* 27, 75-79.
- 741 Eggleton, P., Bignell, D.E., Hauser, S., Dibog, L., Norgrove, L., Madong, B., 2002. Termite
742 diversity across an anthropogenic disturbance gradient in the humid forest zone of West
743 Africa. *Agric. Ecosyst. Environ.* 90, 189-202. doi:10.1016/S0167-8809(01)00206-7
- 744 Engel, E.C., Irwin, R.E., 2003. Linking pollinator visitation rate and pollen receipt. *Am. J.*
745 *Bot.* 90, 1612-1618. doi:10.3732/ajb.90.11.1612
- 746 Faria, L., Melo, G., 2007. Species of *Euglossa* (*Glossura*) in the Brazilian Atlantic forest,
747 with taxonomic notes on *Euglossa stellfeldi* Moure (Hymenoptera, Apidae, *Euglossina*).
748 *Rev. Bras. Entomol.* 51, 275-284. doi:10.1590/S0085-56262007000300004
- 749 Favila, M. and Halffter, G., 1997. The use of indicator groups for measuring biodiversity as
750 related to community structure and function. *Acta Zool. Mex.* 72, 1-25.
- 751 Feest, A., Aldred, T.D., Jedamzik, K., 2010. Biodiversity quality: A paradigm for
752 biodiversity. *Ecol. Indic.* 10, 1077-1082. doi:10.1016/j.ecolind.2010.04.002
- 753 Fisher, R., 1925. *Statistical methods for research workers*, Biological monographs and
754 manuals. doi:10.1056/NEJMc061160
- 755 Fleishman, E., Murphy, D.D., 2009. A realistic assessment of the indicator potential of
756 butterflies and other charismatic taxonomic groups. *Conserv. Biol.* 23, 1109-1116.
757 doi:10.1111/j.1523-1739.2009.01246.x
- 758 Food and Agriculture Organisation of the United Nations, 2016. *Global Forest Resources*
759 *Assessment 2015*. 2nd Edition, Rome (available at: [http://www.fao.org/3/a-](http://www.fao.org/3/a-i4793e.pdf)
760 [i4793e.pdf](http://www.fao.org/3/a-i4793e.pdf))
- 761 Gao, T., Nielsen, A.B., Hedblom, M., 2015. Reviewing the strength of evidence of
762 biodiversity indicators for forest ecosystems in Europe. *Ecol. Indic.* 57, 420-434.
763 doi:10.1016/j.ecolind.2015.05.028
- 764 Gardner, T., Barlow, J., Araujo, I.S., Avila-Pires, T.C., Bonaldo, A., Costa, J., Esposito,
765 M.C., Ferreira, L. V, Hawes, J., Hernandez, M.I.M., Hoogmoed, M.S., Leite, R.N., Lo-
766 Man-Hung, N.F., Malcolm, J.R., Martins, M.B., Mestre, L. a M., Miranda-Santos, R.,
767 Overal, W.L., Parry, L., Peters, S.L., Ribeiro-Junior, M.A., da Silva, M.N.F., da Silva
768 Motta, C., Peres, C. A, 2008. The cost-effectiveness of biodiversity surveys in tropical
769 forests. *Ecol. Lett.* 11, 139-50. doi:10.1111/j.1461-0248.2007.01133.x
- 770 Gibson, L., Lee, T.M., Koh, L.P., Brook, B.W., Gardner, T.A., Barlow, J., Peres, C. A,
771 Bradshaw, C.J.A., Laurance, W.F., Lovejoy, T.E., Sodhi, N.S., 2011. Primary forests are
772 irreplaceable for sustaining tropical biodiversity. *Nature* 478, 378-81.
773 doi:10.1038/nature10425

- 774 Good, P., 2000. Permutation tests: a practical guide to resampling methods for testing
775 hypotheses, 2nd edition. Springer Series in Statistics. Springer Science and Business
776 Media., New York.
- 777 Goodsell, P.J., Underwood, A.J., Chapman, M.G., 2009. Evidence necessary for taxa to be
778 reliable indicators of environmental conditions or impacts. *Mar. Pollut. Bull.* 58, 323-
779 331. doi:10.1016/j.marpolbul.2008.10.011
- 780 Hietz, P., Buchberger, G., Winkler, M., 2006. Effect of forest disturbance on abundance
781 and distribution of epiphytic bromeliads and orchids. *Ecotropica* 12, 103-112.
- 782 Hill, M.O., 1973. Diversity and evenness: a unifying notation and its consequences. *Ecology*
783 54, 427-432.
- 784 Hill, S.L.L., Harfoot, M., Purvis, A., Purves, D.W., Collen, B., Newbold, T., Burgess, N.D.,
785 Mace, G.M., 2016. Reconciling Biodiversity Indicators to Guide Understanding and
786 Action. *Conserv. Lett.* 9, 405-412. doi:10.1111/conl.12291
- 787 Hilty, J., Merenlender, A., 2000. Faunal indicator taxa selection for monitoring ecosystem
788 health. *Biol. Conserv.* 92, 185-197. doi:10.1016/S0006-3207(99)00052-X
- 789 Hinojosa-Díaz, I. A., Engel, M.S., 2011. Revision of the orchid bee subgenus *Euglossella*
790 (Hymenoptera, Apidae), Part I, the *decorata* species group. *Zookeys* 140, 27-69.
791 doi:10.3897/zookeys.140.1923
- 792 Hinojosa-Díaz, I.A., Engel, M.S., 2012. *Alloglossura*, a new subgenus of the orchid bee
793 genus *Euglossa* (Hymenoptera: Apidae), with a review of the included species. *Sci. Pap.*
794 *Nat. Hist. Museum, Univ. Kansas* 45, 1-37.
- 795 Hinojosa-Díaz, I.A., Engel, M.S., 2014. Revision of the orchid bee subgenus *Euglossella*
796 (Hymenoptera: Apidae), Part II: The *viridis* and *mandibularis* species groups. *J.*
797 *Melitology* 36.
- 798 Hsieh, T.C., Ma, K.H., Chao, A., 2016. iNEXT: an R package for rarefaction and
799 extrapolation of species diversity (Hill numbers). *Methods Ecol. Evol.* 7, 1451-1456.
800 doi:10.1111/2041-210X.12613
- 801 Hurst, J.M., Allen, R.B., 2007. A permanent plot method for monitoring indigenous forests.
802 Landcare Research New Zealand, Canterbury.
- 803 Internicola, A.I., Page, P.A., Bernasconi, G., Gigord, L.D.B., 2007. Competition for
804 pollinator visitation between deceptive and rewarding artificial inflorescences: an
805 experimental test of the effects of floral colour similarity and spatial mingling. *Funct.*
806 *Ecol.* 21, 864-872. doi:10.1111/j.1365-2435.2007.01303.x
- 807 Ivey, C.T., Martinez, P., Wyatt, R., 2003. Variation in pollinator effectiveness in swamp
808 milkweed, *Asclepias incarnata* (Apocynaceae). *Am. J. Bot.* 90, 214-225.
809 doi:10.3732/ajb.90.2.214
- 810 Janzen, D.H., 1971. Euglossine Bees as Long-Distance Pollinators of Tropical Plants.
811 *Science* 171, 203-205.

- 812 Jost, L., 2006. Entropy and diversity. *Oikos* 113, 363-375. doi:10.1111/j.2006.0030-
813 1299.14714.x
- 814 Kimsey, L., 1979. An Illustrated Key to the Genus *Exaerete* with Descriptions of Male
815 Genitalia and Biology (Hymenoptera: Euglossini, Apidae). *J. Kansas Entomol. Soc.* 52,
816 735-746.
- 817 Kimsey, L.S., 1982. Systematics of Bees of the Genus *Eufriesea*. University of California
818 Press, Berkeley and Los Angeles.
- 819 King, C., Ballantyne, G., Willmer, P.G., 2013. Why flower visitation is a poor proxy for
820 pollination: Measuring single-visit pollen deposition, with implications for pollination
821 networks and conservation. *Methods Ecol. Evol.* 4, 811-818. doi:10.1111/2041-
822 210X.12074
- 823 Kremen, C., 2005. Managing ecosystem services: What do we need to know about their
824 ecology? *Ecol. Lett.* 8, 468-479. doi:10.1111/j.1461-0248.2005.00751.x
- 825 Krug, C.B., Schaepman, M.E., Shannon, L.J., Cavender-Bares, J., Cheung, W., McIntyre,
826 P.B., Metzger, J.P., Niinemets, Ü., Obura, D.O., Schmid, B., Strassburg, B.B., Van
827 Teeffelen, A.J., Weyl, O.L., Yasuhara, M., Leadley, P.W., 2017. Observations,
828 indicators and scenarios of biodiversity and ecosystem services change – a framework
829 to support policy and decision-making. *Curr. Opin. Environ. Sustain.* 29, 198-206.
830 doi:10.1016/j.cosust.2018.04.001
- 831 Lawton, J.H., Bignell, D.E., Bolton, B., Bloemers, G.F., Eggleton, P., Hammond, P.M.,
832 Hodda, M., Holt, R.D., Larsen, T.B., Mawdsley, N. A, Stork, N.E., Srivastava, D.S.,
833 Watt, A. D., 1998. Biodiversity inventories, indicator taxa and effects of habitat
834 modification in tropical forest. *Nature* 391, 72-76. doi:10.1038/34166
- 835 Legendre, P., De Cáceres, M., 2013. Beta diversity as the variance of community data:
836 Dissimilarity coefficients and partitioning. *Ecol. Lett.* 16, 951-963.
837 doi:10.1111/ele.12141
- 838 Legendre, P., Gallagher, E.D., 2001. Ecologically meaningful transformations for
839 ordination of species data. *Oecologia* 129, 271-280. doi:10.1007/s004420100716
- 840 Lindsey, A.H., 1984. Reproductive Biology of Apiacea. I. Floral Vistors to *Thaspium* and
841 *Zizia* and Their Importance in Pollination. *Am. J. Bot.* 71, 375-387.
- 842 Melo, G.A.R., 2014. Notes on the systematics of the orchid-bee genus *Eulaema*
843 (Hymenoptera, Apidae). *Rev. Bras. Entomol.* 58, 235-240.
- 844 Mitchell, S., Reeve, R., 2016. rdiversity: Measurement and Partitioning of Similarity-
845 Sensitive Biodiversity.
- 846 Moure, J.S., 1965. Some New Species of Euglossine Bees (Hymenoptera: Apidae). *J. Kansas*
847 *Entomol. Soc.* 38, 266-277.
- 848 Nemésio, A., 2009. Taxonomic notes on *Euglossa* (Glossuopoda) with a key to the known
849 species (Hymenoptera: Apidae: Euglossina). *Zootaxa* 56, 45-56.

- 850 Nemésio, A., 2011. The orchid-bee fauna (Hymenoptera: Apidae) of a forest remnant in
851 southern Bahia, Brazil, with new geographic records and an identification key to the
852 known species of the area 54, 47-54.
- 853 Nemésio, A., 2012. Methodological Concerns and Challenges in Ecological Studies with
854 Orchid Bees (Hymenoptera: Apidae: Euglossina). *Bioscience* 26, 2012.
- 855 Nemésio, A., 2013. Are orchid bees at risk? First comparative survey suggests declining
856 populations of forest-dependent species. *Brazilian J. Biol.* 73, 367-374.
- 857 Nemésio, A., Silveira, F. A, 2007a. Diversity and distribution of orchid bees (Hymenoptera:
858 Apidae) with a revised checklist of species. *Neotrop. Entomol.* 36, 874-888.
859 doi:10.1590/S1519-566X2007000600008
- 860 Nemésio, A., Silveira, F. A, 2007b. Orchid bee fauna (Hymenoptera: Apidae: Euglossina) of
861 Atlantic Forest fragments inside an urban area in southeastern Brazil. *Neotrop.*
862 *Entomol.* 36, 186-191.
- 863 Nemésio, A., Silveira, F. A, 2006. Edge effects on the orchid-bee fauna (Hymenoptera:
864 Apidae) at a large remnant of Atlantic Rain Forest in southeastern Brazil. *Neotrop.*
865 *Entomol.* 35, 313-323. doi:10.1590/S1519-566X2006000300004
- 866 Niemack, R.S., Bennett, D.J., Hinojosa-Diaz, I., Chaboo, C.S., 2012. A contribution to the
867 knowledge of the orchid bee fauna of the Los Amigos Biological Station, Madre de Dios,
868 Peru (Hymenoptera: Apidae: Euglossini). *Check List* 8, 215-217.
- 869 Nöske, N.M., Hilt, N., Werner, F.A., Brehm, G., Fiedler, K., Sipman, H.J.M., Gradstein,
870 S.R., 2008. Disturbance effects on diversity of epiphytes and moths in a montane forest
871 in Ecuador. *Basic Appl. Ecol.* 9, 4-12. doi:10.1016/j.baae.2007.06.014
- 872 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,
873 P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Henry, M., Stevens, H., Szoecs, E.,
874 Wagner, H., 2018. *vegan: Community Ecology Package*.
- 875 Otero, J.T., Sandino, J.C., 2003. Capture Rates of Male Euglossine Bees across a Human
876 Intervention Gradient, Chocó Region, Colombia. *Biotropica* 35, 520-529.
877 doi:10.1111/j.1744-7429.2003.tb00608.x
- 878 Pearson, D.L., Dressler, R.L., 1985. Two-year study of male orchid bee (Hymenoptera:
879 Apidae: Euglossini) attraction to chemical baits in lowland south-eastern Perú. *J. Trop.*
880 *Ecol.* 1, 37. doi:10.1017/S0266467400000067
- 881 Pimm, S.L., Raven, P., 2000. Extinction by numbers. *Nature* 403, 843-845.
882 doi:10.1038/35002708
- 883 Potts, S.G., Biesmeijer, J.C., Kremen, C., Neumann, P., Schweiger, O., Kunin, W.E., 2010.
884 Global pollinator declines: Trends, impacts and drivers. *Trends Ecol. Evol.* 25, 345-353.
885 doi:10.1016/j.tree.2010.01.007
- 886 Primack, R.B., Silander, J.A., 1975. Measuring the relative importance of different
887 pollinators to plants. *Nature* 255, 143-144. doi:10.1038/255143a0


- 888 QGIS Development Team, 2017. QGIS Geographic Information System. Open Source
889 Geospatial Foundation Project.
- 890 R Core Team, 2017. R: A language and environment for statistical computing.
- 891 Ramage, B.S., Sheil, D., Salim, H.M.W., Fletcher, C., Mustafa, N.Z.A, Luruthusamay, J.C.,
892 Harrison, R.D., Butod, E., Dzulkiply, A.D., Kassim, A.R., Potts, M.D., 2013.
893 Pseudoreplication in Tropical Forests and the Resulting Effects on Biodiversity
894 Conservation. *Conserv. Biol.* 27, 364-372. doi:10.1111/cobi.12004
- 895 Ramsey, M.W., 1988. Differences in pollinator effectiveness of birds and insects visiting
896 *Banksia menziesii* (Proteaceae). *Oecologia* 76, 119-124. doi:10.1007/BF00379609
- 897 Rasmussen, C., 2009. Diversity and abundance of orchid bees (Hymenoptera: Apidae,
898 Euglossini) in a tropical rainforest succession. *Neotrop. Entomol.* 38, 812-9.
- 899 Real, L.A., 1981. Uncertainty and Pollinator-Plant Interactions: The Foraging Behavior of
900 Bees and Wasps on Artificial Flowers. *Ecology* 62, 20-26.
- 901 Reeve, R., Leinster, T., Cobbold, C.A., Thompson, J., Brummitt, N., Mitchell, S.N.,
902 Matthews, L., 2016. How to partition diversity. arXiv 1404.6520, 1-9.
- 903 Rosa, J.F., Ramalho, M., Monteiro, D., Silva, M.D., 2015. Permeability of matrices of
904 agricultural crops to Euglossina bees (Hymenoptera, Apidae) in the Atlantic Rain Forest.
905 *Apidologie* 46, 691-702. doi:10.1007/s13592-015-0359-9
- 906 Roubik, D., Hanson, P., 2004. Orchid bees of tropical America: Biology and Field Guide.
907 InBio Press, Heredia, Costa Rica.
- 908 Roubik, D.W., 2004. Sibling Species of Glossura and Glossuropoda in the Amazon Region
909 (Hymenoptera: Apidae: Euglossini). *J. Kansas Entomol. Soc.* 77, 235-253.
910 doi:10.2317/0308.20.1
- 911 RStudio Team, 2016. RStudio.
- 912 Samejima, H., Marzuki, M., Nagamitsu, T., Nakasizuka, T., 2004. The effects of human
913 disturbance on a stingless bee community in a tropical rainforest. *Biol. Conserv.* 120,
914 577-587. doi:10.1016/j.biocon.2004.03.030
- 915 Sandino, J.C., 2004. Are there any agricultural effects on the capture rates of male
916 euglossine bees (Apidae: Euglossini)? *Rev. Biol. Trop.* 52, 115-118.
- 917 Santini, L., Belmaker, J., Costello, M.J., Pereira, H.M., Rossberg, A.G., Schipper, A.M.,
918 Ceaușu, S., Dornelas, M., Hilbers, J.P., Hortal, J., Huijbregts, M.A.J., Navarro, L.M.,
919 Schiffers, K.H., Visconti, P., Rondinini, C., 2017. Assessing the suitability of diversity
920 metrics to detect biodiversity change. *Biol. Conserv.* 213, 341-350.
921 doi:10.1016/j.biocon.2016.08.024
- 922 Schemske, D., Horvitz, C., 1984. Variation Among Floral Visitors in Pollination Ability: A
923 Precondition for Mutualism Specialization. *Science* 225, 519-521.
924 doi:10.1126/science.225.4661.519

- 925 Schulze, C.H., Waltert, M., Kessler, P.J.A, Pitopang, R., Shahabuddin, Veddeler, D.,
926 Mühlenberg, M., Gradstein, S.R., Leuschner, C., Steffan-Dewenter, I., Tschardtke, T.,
927 2004. Biodiversity indicator groups of tropical land-use systems: Comparing plants,
928 birds, and insects. *Ecol. Appl.* 14, 1321-1333. doi:10.1890/02-5409
- 929 Socolar, J.B., Gilroy, J.J., Kunin, W.E., Edwards, D.P., 2016. How Should Beta-Diversity
930 Inform Biodiversity Conservation? *Trends Ecol. Evol.* 31, 67-80.
931 doi:10.1016/j.tree.2015.11.005
- 932 Spearman, C., 1904. The Proof and Measurement of Association between Two Things. *Am.*
933 *J. Psychol.* 15, 72-101. doi:10.1093/pan/mpj017
- 934 Storck-Tonon, D., Morato, E., Melo, A., Oliveira, M., 2013. Orchid Bees of forest fragments
935 in Southwestern Amazonia. *Biota Neotrop.* 13, 133-141. doi:10.1590/S1676-
936 06032013000100015
- 937 Stork, N.E., Srivastava, D.S., Eggleton, P., Hodda, M., Lawson, G., Leakey, R.R.B., Watt,
938 A.D., 2017. Consistency of effects of tropical-forest disturbance on species composition
939 and richness relative to use of indicator taxa. *Conserv. Biol.* 00, 1-10.
940 doi:10.1111/cobi.12883
- 941 Tuomisto, H., 2010. A diversity of beta diversities: Straightening up a concept gone awry.
942 Part 2. Quantifying beta diversity and related phenomena. *Ecography* 33, 23-45.
943 doi:10.1111/j.1600-0587.2009.06148.x
- 944 Whitworth, A., Villacampa, J., Brown, A., Huarcaya, R.P., Downie, R., MacLeod, R., 2016.
945 Past human disturbance effects upon biodiversity are greatest in the canopy; A case
946 study on rainforest butterflies. *PLoS One* 11. doi:10.1371/journal.pone.0150520
- 947 Winfree, R., Aguilar, R., Vázquez, D., LeBuhn, G., Aizen, M., 2009. A Meta-Analysis of
948 Bees' Responses to Anthropogenic Disturbance. *Ecology* 8, 2068-2076.
- 949 Winfree, R., Reilly, J.R., Bartomeus, I., Cariveau, D.P., Williams, N.W., Gibbs, J., 2018.
950 Species turnover promotes the importance of bee diversity for crop pollination at
951 regional scales. *Science* 793, In press. doi:10.1126/SCIENCE.AAO2117

952

953

1.12 Supplementary materials



954

955


956

957

958

Figure S.1 – Map of our study sites in Manu, Peru. Colours represent disturbance gradient, dark red = 1. Most disturbed, dark blue = 6. Least disturbed. The map also shows: elevation contour lines; the Manu Learning Centre research station; the nearest local town, Salvación; the local road; the main river, Alto Madre de Dios, and its tributaries.

959



960

961 Figure S.2 – PCA of vegetation structure across the gradient, with sites colour coded by their disturbance level
 962 (dark blue least disturbed – dark red most disturbed).




Figure S.3 – Sample completeness at $q = 0$. The sample completeness for each site is shown in a separate panel. The labels above each panel describe the disturbance rank (1-6, 1 is most disturbed) and habitat type of each site (BA: Banana, AF: Agroforestry, SF: Disturbed secondary forest, CCR: Cleared regenerating forest, MXD: Mixed history regenerating forest and MIN: minimally disturbed primary forest), as well as which of the replicates it was (A, B or C). The solid black line shows the rarefied estimate of sample coverage, and the dashed line is the extrapolated estimate, with the circle indicating the sample size collected. The grey shading indicates the 95% confidence interval around the coverage estimate. Coverage is shown up to 100 individuals for easy comparison between sites, but some sites did exceed this number; estimated diversity was compared at $n = 50$.




Figure S.4 – Estimated richness at each site extrapolated to a sample size of 250 individuals. Solid line shows rarefied estimate, dashed line shows extrapolation, and filled circle indicates the collected sample size. Shaded area indicates 95% confidence interval with 1000 bootstraps.

Table S.1 – Results of the Spearman rank correlation tests.

The p and rho values of the results of correlations between variables 1 and 2 are shown for all tests, along with the 95% confidence interval for the rho values.

Variable 1	Variable 2	p	rho	rho 95% CI
Abundance	Disturbance rank	0.0051	0.6301	0.2534 - 0.8303
Observed diversity $q = 0$	Disturbance rank	0.0141	0.5672	0.1538 - 0.808
Observed diversity $q = 1$	Disturbance rank	0.0302	0.5110	0.0105 - 0.8576
Observed diversity $q = 2$	Disturbance rank	0.0581	0.4546	-0.0517 - 0.817
Observed diversity $q = \infty$	Disturbance rank	0.0801	0.4232	-0.0701 - 0.7895
Vegetation structure PC1	Disturbance rank	0.0000	0.8621	0.6124 - 0.9679
Estimated diversity $q = 0$ (mean)	Disturbance rank	0.4472	0.1912	-0.3512 - 0.6686
Estimated diversity $q = 1$ (mean)	Disturbance rank	0.3747	0.2226	-0.2824 - 0.622
Estimated diversity $q = 2$ (mean)	Disturbance rank	0.2300	0.2978	-0.1826 - 0.6794
$q = 0$ raw iNEXT diversity estimates (x1000)	Difference in disturbance rank	0.3318	0.2693	-0.2551 - 0.6685
$q = 1$ raw iNEXT diversity estimates (x1000)	Difference in disturbance rank	0.3782	0.2453	-0.295 - 0.6639
$q = 2$ raw iNEXT diversity estimates (x1000)	Difference in disturbance rank	0.4201	0.2250	-0.3067 - 0.6463
Redundancy (ρ) $q = 1$	Disturbance rank	0.0168	-0.5549	-0.7938 - -0.1434
Representativeness (ρ) $q = 1$	Disturbance rank	0.0380	0.4922	0.0075 - 0.8126
Pollinator visits	Disturbance rank	0.6467	0.1160	-0.3499 - 0.5397
Pollinator visits	Observed diversity $q = 0$	0.4380	-0.1950	-0.5965 - 0.2459
Pollinator visits	Observed diversity $q = 1$	0.4331	-0.1971	-0.6293 - 0.3067
Pollinator visits	Observed diversity $q = 2$	0.4429	-0.1930	-0.6426 - 0.3347

Table S.2 – Model selection.

Where the correlations tested in Table S.1 were significant, linear models were used to control for the effect of other environmental variables. The log likelihoods and degrees of freedom for all models tested are presented here.

Model	Log Likelihood (Max. Likelihood)	Degrees of freedom
log(Abundance)~ Disturbance rank	-16.6356	3
log(Abundance)~ Disturbance rank + Elevation	-16.2785	4
log(Abundance)~ Disturbance rank + Distance to river	-14.0386	5
log(Obs. Diversity q = 0) ~ Disturbance rank	-1.1849	3
log(Obs. Diversity q = 0) ~ Disturbance rank + Elevation	-1.0606	4
log(Obs. Diversity q = 0) ~ Disturbance rank + Distance to river	0.4615	4
log(Obs. Diversity q = 1) ~ Disturbance rank	-2.3260	3
log(Obs. Diversity q = 1) ~ Disturbance rank + Elevation	-2.1954	4
log(Obs. Diversity q = 1) ~ Disturbance rank + Distance to river	-1.3630	4
log(Obs. Diversity q = 2)~ Disturbance rank	-6.2357	3
log(Obs. Diversity q = 2) ~ Disturbance rank + Elevation	-5.9173	4
log(Obs. Diversity q = 2) ~ Disturbance rank + Distance to river	-5.6520	4
log(Obs. Diversity q = ∞) ~ Disturbance rank	-5.0875	3
log(Obs. Diversity q = ∞) ~ Disturbance rank + Elevation	-4.6954	4
log(Obs. Diversity q = ∞) ~ Disturbance rank + Distance to river	-4.8372	4
Vegetation structure PC1 ~ Disturbance rank	-6.4559	3
Vegetation structure PC1 ~ Disturbance rank + Elevation	-6.2303	4
Vegetation structure PC1 ~ Disturbance rank + Distance to river	-5.7609	4
log(Est. diversity q = 0) ~ Disturbance rank	6.70	3
log(Est. diversity q = 0) ~ Disturbance rank + Elevation	6.82	4
log(Est. diversity q = 0) ~ Disturbance rank + Distance to river	6.83	4
log(Est. diversity q = 1) ~ Disturbance rank	-2.15	3
log(Est. diversity q = 1) ~ Disturbance rank + Elevation	-1.86	4
log(Est. diversity q = 1) ~ Disturbance rank + Distance to river	-2.10	4
log(Est. diversity q = 2) ~ Disturbance rank	-5.87	3
log(Est. diversity q = 2) ~ Disturbance rank + Elevation	-5.50	4

Model	Log Likelihood (Max. Likelihood)	Degrees of freedom
log(Est. diversity $q = 2$) ~ Disturbance rank + Distance to river	-5.70	4
log(ρ Redundancy $q = 1$) ~ Disturbance rank	-13.3025	3
log(ρ Redundancy $q = 1$) ~ Disturbance rank + Elevation	-12.7122	4
log(ρ Redundancy $q = 1$) ~ Disturbance rank + Distance to river	-11.5194	4
log(ρ Representativeness $q = 1$) ~ Disturbance rank	6.1746	3
log(ρ Representativeness $q = 1$) ~ Disturbance rank + Elevation	6.1913	4
log(ρ Representativeness $q = 1$) ~ Disturbance rank + Distance to river	9.1013	4

Table S.3 – Results of the best fitting models.

The estimated coefficients and significance are presented for the explanatory variables found to produce the best fitting models identified in Table S.2.

Model	Variable	Est. coefficient	SE	t	p
log(Abundance) ~ Disturbance rank + Distance to River	Rank	0.317	0.082	3.871	0.002
	Distance to river	-0.001	0.000	-2.240	0.041
log(Obs. Diversity $q = 0$) ~ Disturbance rank	Rank	0.103	0.038	2.724	0.015
log(Obs. Diversity $q = 1$) ~ Disturbance rank	Rank	0.077	0.040	1.914	0.074
log(Obs. Diversity $q = 2$) ~ Disturbance rank	Rank	0.087	0.050	1.730	0.103
log(Obs. Diversity $q = \infty$) ~ Disturbance rank	Rank	0.091	0.047	1.946	0.069
Vegetation structure PC1 ~ Disturbance rank	Rank	0.492	0.051	9.703	0.000
log(Est. diversity $q = 0$) ~ Disturbance rank	Rank	0.020	0.024	0.813	0.428
log(Est. diversity $q = 1$) ~ Disturbance rank	Rank	0.031	0.040	0.772	0.451
log(Est. diversity $q = 2$) ~ Disturbance rank	Rank	0.055	0.049	1.112	0.282
log(ρ Redundancy $q = 1$) ~ Disturbance rank	Rank	-0.217	0.074	-2.920	0.010
log($\bar{\rho}$ Representativeness $q = 1$) ~ Disturbance rank + Distance to river	Rank	0.0711	0.0227	3.136	0.0068
	Distance to river	-0.0002	<0.0001	-2.401	0.0298

Table S.4 – Moran's I

Results of the tests for spatial autocorrelation performed on the residuals of the best fitting models (Table S.3), for all models where a significant correlation was found.

Model	Observed	Expected	SD	p	Observed - Expected
log(Abundance) ~ Disturbance rank + Distance to river	-0.1074	-0.0588	0.0916	0.5950	-0.0487
log(Obs. Diversity q = 0) ~ Disturbance rank	-0.1237	-0.0588	0.0940	0.4901	-0.0649
log(Obs. Diversity q = 1) ~ Disturbance rank	-0.1346	-0.0588	0.0879	0.3883	-0.0757
log(Obs. Diversity q = 2) ~ Disturbance rank	-0.0950	-0.0588	0.0879	0.6806	-0.0362
log(Obs. Diversity q = ∞) ~ Disturbance rank	-0.0788	-0.0588	0.0912	0.8270	-0.0199
Vegetation structure PC1 ~ Disturbance rank	-0.0176	-0.0588	0.0958	0.6669	0.0412
log(Est. diversity q = 0) ~ Disturbance rank	-0.0403	-0.0588	0.0920	0.8401	0.0186
log(Est. diversity q = 1) ~ Disturbance rank	-0.0492	-0.0588	0.0889	0.9140	0.0096
log(Est. diversity q = 2) ~ Disturbance rank	-0.0648	-0.0588	0.0883	0.9464	-0.0059
log(ρ Redundancy q = 1) ~ Disturbance rank	0.00095	-0.0588	0.0914	0.5130	0.0598
log(ρ Representativeness q = 1) ~ Disturbance rank	-0.1439	-0.0588	0.0924	0.3570	-0.0851