



The effect of high-fat diet on the morphological properties of the forelimb musculature in hypertrophic myostatin null mice

Article

Accepted Version

Elashry, M. I., Elashry, A., Glanske, K., Matsakas, A., Wenisch, S., Arnhold, S. and Patel, K. (2019) The effect of high-fat diet on the morphological properties of the forelimb musculature in hypertrophic myostatin null mice. *Journal of Anatomy*, 235 (4). pp. 825-835. ISSN 0021-8782 doi: <https://doi.org/10.1111/joa.13025> Available at <http://centaur.reading.ac.uk/83669/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/joa.13025>

Publisher: Wiley-Blackwell

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **The effect of high-fat diet on the morphological properties of the forelimb**
2 **musculature in hypertrophic myostatin null mice.**

3 Mohamed I. Elashry^{3, 4*}; Asmaa Eldaey^{3, 5}; Kristina Glenske⁵; Antonios Matsakas²;
4 Sabine Wenisch⁵; Stefan Arnhold⁴ and Ketan Patel¹

5 ¹School of Biological Sciences, Whiteknights, University of Reading, RG6 6UB, United
6 Kingdom

7 ²Molecular Physiology Laboratory, Centre for Atherothrombotic and Metabolic
8 Disease, Hull York Medical School, University of Hull

9 ³Anatomy and Embryology Department, Faculty of Veterinary Medicine, University of
10 Mansoura 35516, Egypt

11 ⁴Institute of Veterinary Anatomy, Histology and Embryology, Justus Liebig University
12 of Giessen 35392, Germany

13 ⁵Clinic of Small Animals, c/o Institute of Veterinary Anatomy, Histology and
14 Embryology, Justus Liebig University of Giessen 35392, Germany

15

16 ***Corresponding author:**

17 Mohamed Elashry

18 Institute of Veterinary Anatomy, Histology and Embryology,

19 Justus-Liebig-University of Giessen, Frankfurter Str. 98, 35392 Gießen

20 Tel. +49 641 99-38105

21 Email: mohammed.elashry@vetmed.uni-giessen.de

22

23 **Acknowledgements**

24 We gratefully thank University of Reading, Wellcome Trust (078649), UK, Engemann
25 research foundation, Faculty of Veterinary medicine, University of Mansoura, Egypt,
26 and University of Gießen, Germany for generous funding of this work.

27

28 **Abstract**

29 Obesity is a worldwide nutritional disorder affecting body performance including
30 skeletal muscle. Inhibition of myostatin not only increases the muscle mass but also it
31 reduces body fat accumulation. We examined the effect of high-fat diet on the
32 phenotypic properties of forelimb muscles from myostatin null mice. **Male wild-type and**
33 **myostatin null mice were fed on either normal diet or high-fat diet (45 % fat) for ten**
34 **weeks. *M. triceps brachii Caput longum*; *M. triceps brachii Caput laterale*; *M. triceps***
35 ***brachii Caput mediale*; *M. extensor carpi ulnaris* and *M. flexor carpi ulnaris* were**
36 processed for fiber type composition using immunohistochemistry and morphometric
37 analysis. Although the muscle mass revealed no change under high-fat diet, there were
38 morphometric alterations in the absence of myostatin. **We show that high-fat diet**
39 **reduces the cross-sectional area of the fast (IIB and IIX) fibers in *M. triceps brachii***
40 ***Caput longum* and *M. triceps brachii Caput laterale* of both genotypes. In contrast,**
41 **increases of fast fibers area were observed in both *M. extensor carpi ulnaris* of wild-**
42 **type and *M. flexor carpi ulnaris* of myostatin null. Meanwhile, a high-fat diet increases**
43 **the area of the fast IIA fibers in wild-type, myostatin null displays a muscle-dependent**
44 **alteration in the area of the same fiber type. The combined high-fat diet and myostatin**
45 **deletion shows no effect on the area of slow type I fibers. Despite, a high-fat diet causes**
46 **a reduction in the area of the peripheral IIB fibers in both genotypes, only myostatin**
47 **null shows an increase in the area of the central IIB fibers. We provide evidence that a**
48 **high-fat diet induces a muscle-dependent fast to slow myofiber shift in the absence of**
49 **myostatin. Taken together, the data suggest that the morphological alterations of**
50 **muscle fibers under combined high-fat diet and myostatin deletion reflect a functional**
51 **adaptation of the muscle to utilize the high energy intake.**

52 **Keywords:** skeletal muscle, high-fat diet, myostatin, myosin heavy chain, muscle fiber

53

54 **Running Headline/ Short title**

55 High-fat diet in myostatin null mice

56

57

58 **1. Introduction**

59 Obesity has become a challenging chronic nutritional threat affecting human health,
60 performance and quality of life worldwide (Apovian, 2016). Obesity originates in an
61 imbalance between energy consumption and energy expenditure (Lebrasseur, 2012).
62 As a metabolic disorder, obesity increases the risk of cardiovascular disease, type 2
63 diabetes, and cancer (Muio & Newgard, 2006). However, the impact of obesity on
64 skeletal muscle remains controversial, although obesity coincides with impaired
65 locomotor ability of skeletal muscle (reviewed by Tallis et al. 2018). Beyond the
66 essential contractile function, skeletal muscle is one of the main regulator for glucose
67 disposal and glycogen storage in human body and is crucial for minimizing the risk of
68 insulin resistance (Stump et al. 2006).

69 Skeletal muscle is an adapting tissue altering its physiological components under the
70 effect of either external or internal stimuli to overcome functional demands (Egan &
71 Zierath, 2013). Phenotypic evaluation of skeletal muscle is often based on profiling the
72 heterogeneous population of muscle fibers categorized on their speed of contraction
73 (underpinned by the expression of specific myosin heavy chain isoforms- MHC) or their
74 mode of metabolism (oxidative versus glycolytic). Generally, glycolytic muscles tend to
75 be fast twitching, fatigue susceptible and rely on anaerobic metabolism to produce
76 energy whereas oxidative muscles are slow contracting, fatigue resistant and generate
77 energy via aerobic metabolism (Zierath & Hawley, 2004; Matsakas & Patel, 2009). It
78 has been reported that obesity impairs muscle mass and function as in case of
79 sarcopenic obesity (Prado et al. 2012). The accumulation of fatty tissue can induce
80 skeletal muscle inflammation. Studies have shown an increase of macrophage and
81 inflammatory cytokines involving tumor necrosis factor- alpha (TNF α) and monocyte
82 chemoattractant protein-1 (MCP-1) were associated with obesity (Valerio et al. 2006;
83 Varma et al. 2009; Lumeng & Saltiel, 2011). At the molecular level, it has been reported
84 that obesity interferes with calcium signaling and 5-adenosine monophosphate-
85 activated protein kinase (AMPK) causing muscle fiber shift and alteration in the
86 contractile properties (Tallis et al. 2018). High-fat diet has been shown to elicit changes
87 in mouse skeletal muscle gene expression and increase MHC type I in the quadriceps
88 muscle (Lange et al. 2007; Wilde et al. 2008). Conversely, skeletal muscle composition
89 from obese humans and animals has shown an increased proportion of fast twitch at
90 the expense of slow twitch type I fibers (Matsakas & Patel, 2009). In addition, cytokines

91 and other molecules produced by adipocytes have been shown to impair the contractile
92 function of the muscle leading to muscle fiber atrophy, inflammation and insulin
93 resistance (Srikanthan et al. 2010; Pellegrinelli et al. 2015). Additionally, development
94 of obesity is associated with a reduction of mitochondrial contents in skeletal muscle
95 which results in impairment of fatty acid oxidation (Holloway et al. 2009).

96 Myostatin is a member of transforming growth factor- β (TGF- β) superfamily, acts as a
97 powerful negative regulator for skeletal muscle development (Sharma et al. 2001).
98 Myostatin interferes with the cell-cycle progression and inhibits myoblasts proliferation
99 (Thomas et al. 2000). Genetic deletion of myostatin results in significant increase of
100 muscle mass due to a combination of hyperplasia (an increase of muscle fiber number)
101 and hypertrophy (an increase of muscle fiber size) (McPherron et al. 1997).
102 Interestingly, as myostatin deletion reduced the body fat accumulation concomitantly,
103 it has become a suitable target for therapeutic investigations against obesity
104 (McPherron & Lee, 2002) as well as type 2 diabetes mellitus (Allen et al. 2011).
105 Moreover, myostatin inhibition alters the phenotype of white adipose tissue into brown
106 adipose tissue indicating a higher energy utilization and enhanced fatty acid oxidation
107 (Zhang et al. 2012).

108 The present study examined for the first time, the effect of high-fat diet (HFD) on the
109 myofiber morphology of forelimb musculature **in the myostatin null (*Mstn*^{-/-}) mice in
110 comparison to C57/BL6 wild-type (*Mstn*^{+/+}) counterpart**. After 10 weeks of HFD intake,
111 **five anatomically and phenotypically different forelimb muscles nominated as follow;**
112 ***M. triceps brachii* Caput longum (T.long); *M. triceps brachii* Caput laterale (T.lateral),**
113 ***M. triceps brachii* Caput mediale (T.medial); *M. extensor carpi ulnaris* (ECU) and *M.***
114 ***flexor carpi ulnaris* (FCU) were processed for immunohistochemistry using antibodies**
115 **against myosin heavy chain type IIB, IIA and type I isoforms. Evaluation of body mass,**
116 **wet muscle specific-weight, cross-sectional area (CSA) and fiber type composition**
117 **were analyzed. Our data demonstrates that although the muscle mass revealed no**
118 **change under HFD, there were a remarkable alteration in the absence of myostatin in**
119 **terms of CSA and phenotype of all muscle fibers types. Our detailed analysis revealed**
120 **that HFD reduced the CSA of the fast glycolytic (IIB and IIX) fibers in T.lateral and**
121 **T.long of both genotypes. In contrast, the CSA of the same fiber type was increased in**
122 **T.medial and ECU of *Mstn*^{+/+} as well as, T.medial and FCU muscles of *Mstn*^{-/-}. HFD**
123 **increases CSA of the fast oxidative IIA fibers in a muscle and genotype-dependent**

124 manner. We show that slow type I fibers of *Mstn*^{-/-} were less liable to the effect of HFD
125 intake. We found evidence of fiber size alteration based on their anatomical distribution
126 following combined myostatin deletion and HFD intake. We provide evidence that
127 myostatin deletion together with HFD induced a muscle-dependent fast to slow
128 myofiber shift. The data suggest that HFD modulates the phenotypic properties of the
129 myostatin null muscles to maximize energy utilization.

130

131 **2. Materials and Methods**

132 2. 1. Animals

133 **Males 4-5 months *Mstn*^{+/+} and *Mstn*^{-/-} mice** were raised in the biological unit, University
134 of Reading, United Kingdom. *Mstn*^{-/-} mice were donated by Dr. Lee (McPherron et al.
135 1997). All the standard procedures were approved by the Animal Care and Ethical
136 Review Committee (AWERB) and performed under a project license (number 7516)
137 from the United Kingdom Home Office under Animals (Scientific Procedures) Act 1986.
138 *Mstn*^{+/+} and *Mstn*^{-/-} mice were administered *ad libitum* either a standard chow (normal
139 diet, ND) or a high-fat diet (HFD), composed of 45 % fat, 20 % protein and 35%
140 carbohydrates (Special diet services (SDS) code: 824053). The animals were kept
141 under standard environmental conditions at 21 C and cyclic 12 hours light/dark groups
142 up to 10 weeks. After the completion of the experiment, the animals were humanely
143 euthanized using Schedule 1 procedure.

144

145 2. 2. Tissue collection

146 In order to evaluate whether HFD affects the phenotype of the muscles in the absence
147 of myostatin, five representative muscles from the upper and lower forelimb were
148 surgically removed as follow; M. triceps brachii Caput longum (T.long); M. triceps
149 brachii Caput laterale (T.lateral), M. triceps brachii Caput mediale (T.medial); M.
150 extensor carpi ulnaris (ECU) and M. flexor carpi ulnaris (FCU). The dissected muscles
151 were frozen using isopentane pre-cooled with liquid nitrogen and were embedded in
152 tissue tech OCT (Sakura, VWR) using ethanol precooled in dry ice. 10 µm thickness
153 transverse mid-belly cryosections from each muscle were placed on poly-L-lysine
154 coated slides (VWR, Germany) and left to dry at room temperature at least for one
155 hour before stored at -80C.

156 2. 3. Immunohistochemistry

157 Muscle sections were washed three times in PBS for 15 minutes, then permeabilized
158 in buffer solution composed of 20 mM Hepes (Biochrom, Germany), 300 mM sucrose
159 (Merck, Germany), 50 mM NaCl (Roth, Germany), 3 mM MgCl₂ (Roth, Germany) and
160 0.5% Triton-X100 (pH7, Roth, Germany) for 15 minutes at room temperature. The
161 muscle sections were preblocked in wash buffer containing 5% fetal calf serum (v/v)
162 (Gibco, Thermo Fisher Scientific) and 0.05 % Triton X-100 (v/v) in phosphate buffered
163 saline (PBS) for 30 minutes at room temperature. The myofiber type was investigated
164 based on the expression of certain myosin heavy chain protein. The myofiber type IIB,
165 IIA and type I were identified using BFF3 mouse IgM (1:1), A.474 mouse IgG (1:4) and
166 A4.840 mouse IgM (1:1) monoclonal primary antibodies (DSHB) respectively as
167 previously reported (Matsakas et al. 2009). Type IIB and IIA myofibers were
168 immunostained in the same muscle section. However, type I myofibers were identified
169 on serial sections. Muscle sections with no primary antibody added were used as a
170 negative control (Fig. 1a and b). All the primary antibodies were incubated with muscle
171 sections overnight at 4C. After five times washing steps for 10 minutes, the primary
172 antibodies were detected using Alexa Fluor 633 goat anti-mouse IgM (Molecular
173 Probes, A21046 diluted in wash buffer 1:200) for MHC I and MHC IIB and Alexa Fluor
174 488 Goat-anti-mouse IgG (Molecular probes, A11029 diluted in wash buffer 1:200)
175 secondary antibody. The muscle sections were incubated with the secondary
176 antibodies for 45-60 minutes at room temperature in the dark. The nuclei were
177 counterstain with 4', 6-Diamidino-2-Phenylindole, Dihydrochloride (DAPI, Thermo
178 Fisher scientific) and the slides were mounted using a fluorescent mounting medium
179 (Mowiol 4-88, Calbiochem).

180 2. 4. Imaging and myofiber quantifications

181 Immunostained muscle sections were morphologically evaluated and photographed
182 using a Zeiss Axioscop2 fluorescence microscope connected to a digital camera and
183 a computer operated with the Axiovision software (Zeiss). Briefly, uniform, transverse
184 and undamaged muscle sections were selected for photography. To evaluate the total
185 myofiber number and the percentage of each fiber type (IIB, IIA and, type I), the whole
186 muscle cross section was photographed in separate images and was reconstructed
187 using the photo merge tool in Photoshop CS6 (64 Bit). The total fiber number, number
188 of IIB, IIA and type I fibers were quantified via manual tracking of the immunostained

189 fibers. The quantification of type IIX fibers was performed via subtracting the number
190 of all immune positive fibers from the total fiber number per muscle section.
191 Measurement of the cross sectional area (CSA) for each fiber type was performed for
192 all muscles by using the measuring tool in the Axiovision software (Zeiss). Minimum of
193 250 fibers was measured for each fiber type. For some muscles, the available fibers
194 from certain type were insufficient. Therefore, all the available fibers from that particular
195 type were measured.

196 2. 5. Statistical analysis

197 In order to analyze the effect of genotype ($Mstn^{+/+}$ vs. $Mstn^{-/-}$) and high caloric
198 supplement (ND vs. HFD) on total muscle fiber number, measurement of CSA and the
199 percentage of type IIB, IIX, IIA and I fibers of T.lateral, T.long, T.medial, ECU and FCU
200 muscles, a two-way ANOVA was performed. Multiple comparisons and the variables
201 interactions were evaluated using Tukey's and Sidak's Post hoc test. Statistical
202 analysis was conducted by using Graph Pad Prism 6 software. All values are presented
203 as mean \pm SEM and $p < 0.05$ was considered to be significant.

204

205 3. Results

206 3. 1. Effect of high-fat diet on the total body mass and individual muscle weight

207 The present study examined the effect of high-fat diet on the phenotypic properties of
208 forelimb muscles in the myostatin null and wild type mice. As expected, body mass
209 was significantly higher ($p < 0.05$) in $Mstn^{-/-}$ ND compared to $Mstn^{+/+}$ ND mice. We found
210 that a 10-week high-fat diet regime resulted in significantly greater ($p < 0.01$) body mass
211 (g) independent of genotype (i.e. 20% and 14% increase in $Mstn^{+/+}$ and $Mstn^{-/-}$
212 respectively) (Table 1). Next, we evaluated the effect of HFD on specific muscle weight
213 (mg). No significant differences were detected in $Mstn^{+/+}$ and $Mstn^{-/-}$ muscles after HFD
214 compared to ND (Table 1).

215 3. 2. Evaluation of the muscle fiber number following a high-fat diet

216 We next determined the effect of HFD on muscle fiber number studied (Fig. 1a-f). As
217 expected, myostatin deletion induced muscle-dependent hyperplasia of 232 ± 12 %,
218 142 ± 2 %, 140 ± 2 %, 130 ± 8 % and 98 ± 1 % for T.lateral, FCU, T.long, ECU and T.medial

219 muscles respectively in comparison to *Mstn*^{+/+} kept on a ND. T.lateral, FCU and T.long
220 muscles of *Mstn*^{-/-} demonstrated the highest increases in myofiber number ($p < 0.0001$)
221 followed by ECU and T.medial muscles ($p < 0.01$) compared to matched muscles of
222 *Mstn*^{+/+} ND. However, HFD did not induce any significant changes in total muscle fiber
223 number in either genotype (Fig. 1g). These data indicate that neither myofiber
224 hyperplasia nor fiber splitting nor fiber loss occurs under HFD. These data also suggest
225 that muscle-specific differences in the amount of myofiber hyperplasia may be due to
226 differential levels of myostatin expression within individual muscle.

227 **3.3. The effect of high-fat diet on the CSA of muscle fiber type**

228 Evaluation of the muscle fiber CSA would indicate whether high-fat diet alters
229 myofiber type lead to either atrophy (a decrease of the muscle fiber size) or
230 hypertrophy (an increase of the muscle fiber size). Therefore, measurement of the CSA
231 of different fiber types IIA, IIX and IIB for each muscle of both genotypes was
232 performed. The analysis showed that *Mstn*^{+/+} and *Mstn*^{-/-} muscles responded differently
233 regarding the CSA of type IIB fiber following HFD. *Mstn*^{+/+}HFD displayed significant
234 reduction of IIB fiber CSA in T.long and FCU muscles ($p < 0.0001$ and $p < 0.001$).
235 Conversely, IIB fiber CSA was increased in T.medial and ECU muscles ($p < 0.0001$ and
236 $p < 0.01$) compared to the matched *Mstn*^{+/+} fed on a ND (Fig. 2c, f, i, and l). Similarly,
237 *Mstn*^{-/-}HFD showed reduction in CSA of IIB fiber in T.lateral and T.long muscles
238 ($p < 0.001$ and $p < 0.0001$), in contrast, there were increased CSA of IIB in T.medial and
239 FCU muscles ($p < 0.001$) compared to matched muscles of *Mstn*^{-/-}ND (Fig. 2a, c, f and
240 l). The data analysis for T.long and FCU muscles demonstrated significant interactions
241 ($p < 0.05$ and $p < 0.0001$) between the effects of genotype and the diet on the CSA of IIB
242 fibers. Thus, it might suggest that both genotypes responded differently under HFD
243 (Fig. 2c and l). The same trend was observed when we examined the CSA of type IIX
244 fibers. HFD resulted in significantly smaller CSA of type IIX fibers ($p < 0.001$ and
245 $p < 0.0001$) for T.lateral muscles of *Mstn*^{+/+} and T.long of *Mstn*^{-/-} compared to genotype-
246 matched mice on a ND. Interestingly, however both *Mstn*^{+/+} and *Mstn*^{-/-} showed
247 decreased CSA of type IIX of FCU muscle under HFD ($p < 0.001$ and 0.01) compared
248 to genotype-matched mice kept on ND (Fig. 2b, d, and m). Although ECU muscle of
249 *Mstn*^{+/+}HFD exhibited an elevation of the CSA of type IIX fibers ($p < 0.001$), such
250 increase was stronger in *Mstn*^{-/-}HFD involving both T.medial and ECU muscles ($p < 0.05$
251 and $p < 0.0001$) compared to the corresponding muscles from mice kept on a ND (Fig.

252 2g, and j). The data analysis revealed that four out of five muscles showed significant
253 interactions ($p < 0.001$, $p < 0.0001$, $p < 0.05$ and $p < 0.001$) between the effects of HFD and
254 the genotype on the CSA of type IIX fibers. The results suggest that myostatin deletion
255 affect the response of fast fibers (type IIB and IIX) under HFD.

256 Next, the CSA of MHC type IIA fibers referred to the fast oxidative fibers for all muscles
257 under ND and HFD was analyzed. Under HFD, an increased CSA of type IIA fibers
258 was detected in T.long and T.medial muscles of $Mstn^{+/+}$ ($p < 0.05$ and $p < 0.0001$)
259 compared to $Mstn^{+/+}$ ND (Fig. 2e and h). In contrast, the response of the $Mstn^{-/-}$ HFD
260 was slightly variable, meanwhile, CSA of type IIA fibers was decreased in T.long and
261 T.medial muscles ($p < 0.0001$ and 0.01), there was an increase in type IIA CSA in ECU
262 and FCU muscles ($p < 0.001$) compared to $Mstn^{-/-}$ ND muscles (Fig. 2e, h, k, and n).
263 The analysis demonstrated significant interactions between the effect of HFD and
264 genotype ($p < 0.0001$, $p < 0.001$, $p < 0.001$ and $p < 0.0001$) for T.long, T.medial, ECU and
265 FCU muscles. Due to the absence of the sufficient number of MHC type I referred to
266 the slow oxidative fibers in T.lateral, T.long, ECU and FCU muscles for CSA
267 measurement, only T.medial muscle could be analyzed. The data showed a significant
268 reduction in the CSA of type I fibers in the $Mstn^{+/+}$ under HFD ($p < 0.05$) compared to
269 $Mstn^{+/+}$ kept on a ND. However, the $Mstn^{-/-}$ demonstrated no difference in the CSA of
270 type I fiber of T.medial muscle following HFD period (Fig. 2o). These data were
271 accompanied with significant interactions ($p < 0.01$) between the effect of HFD and the
272 effect of genotype suggesting the influence of genotype on the slow fiber type under
273 HFD (Fig. 2o).

274 **3.4. Effect of high-fat diet on the CSA of myofibers based on their regional** 275 **distribution**

276 Given our data, we next examined whether high-fat consumption alters the myofiber
277 CSA based on their anatomical distribution. Therefore, we analyzed the CSA of MHC
278 type IIB, IIX and type IIA fibers from the peripheral (PF) and central (CT) regions within
279 the mid-belly of $Mstn^{+/+}$ and $Mstn^{-/-}$ muscles. Type I fibers were excluded from the
280 analysis due to insufficient numbers in some of the muscles under investigation. The
281 CSA of PFIIB fibers of $Mstn^{+/+}$ HFD were reduced in T.lateral, T.long and FCU muscles
282 ($p < 0.0001$, $p < 0.0001$ and $p < 0.01$) compared to PFIIB fibers of $Mstn^{+/+}$ ND (Fig. 3a, c
283 and k). Similarly, $Mstn^{-/-}$ HFD showed a reduction in the CSA of PF type IIB fibers of
284 T.lateral and T.long muscles ($p < 0.0001$ and 0.0001) compared to the matched region

285 of $Mstn^{-/-}$ ND (Fig. 3a and c). In contrast, PFIIB fibers of T.medial and FCU muscles of
286 $Mstn^{-/-}$ HFD demonstrated an increase of the CSA of the same fiber type ($p<0.05$ and
287 $p<0.0001$) compared to $Mstn^{-/-}$ ND (Fig. 3e and k). Furthermore, we quantified the CSA
288 of CTIIB fibers in each genotype following HFD in comparison to ND. Under HFD,
289 $Mstn^{+/+}$ showed an increase of CSA in the CTIIB fibers of T.lateral and T.medial
290 muscles ($p<0.0001$). However, T.long and FCU muscles of the same genotype
291 demonstrated a decrease in the CSA of CTIIB fibers ($p<0.0001$ and $p<0.01$) compared
292 to the matched muscle region of $Mstn^{+/+}$ ND (Fig. 3a, c, e, h and k). Similarly, significant
293 increases in the CSA of CTIIB fibers of T.lateral, ECU and FCU muscles of $Mstn^{-/-}$ HFD
294 ($p<0.0001$) compared to the corresponding muscle region of $Mstn^{-/-}$ ND (Fig. 3a, c, e,
295 h, and k). The analysis revealed, significant interactions in almost all muscles
296 ($p<0.0001$) between the effect of HFD, the effect of genotype and most importantly the
297 anatomical localization of the muscle fiber. Furthermore, under high-fat diet, a common
298 trend of loss in the CSA of PFIIB compared to the same muscle region on a ND could
299 be detected. Although, there were either increased or decreased CSA of CTIIB in
300 $Mstn^{+/+}$, at least three muscles of $Mstn^{-/-}$ showed an increase in the CSA of CTIIB fibers
301 following HFD.

302 Subsequently, we evaluated whether HFD alters the CSA of MHC type IIX fibers in
303 terms of the anatomical distribution (PF vs. CT) of the myofiber. Although, the majority
304 of $Mstn^{+/+}$ HFD displayed no change in the CSA of PFIIX fibers compared to the
305 matched region of $Mstn^{+/+}$ ND, only $Mstn^{-/-}$ HFD had larger CSA in PFIIX fibers of
306 T.medial and ECU muscles ($p<0.0001$ and $p<0.001$) compared to the matched region
307 of $Mstn^{-/-}$ ND (Fig. 3b, d, f, i and l). Regarding the CSA of CTIIX fibers measurement,
308 $Mstn^{+/+}$ HFD showed an elevation in the CSA of CTIIX fibers of T.medial and ECU
309 muscles ($p<0.0001$ and 0.01) in comparison to the matched region of $Mstn^{+/+}$ ND. In
310 contrast, both T.lateral and FCU muscles of $Mstn^{+/+}$ HFD showed a reduction in the
311 CSA of the CTIIX fibers ($p<0.0001$) compared $Mstn^{+/+}$ ND. Similarly, $Mstn^{-/-}$ HFD
312 displayed increase in the CTIIX fibers of T.long, T.medial and ECU muscles ($p<0.0001$)
313 compared to matched-muscle region of $Mstn^{-/-}$ ND (Fig. 3b, d, f, i and l). The data was
314 accompanied with significant interaction ($p<0.0001$) indicating the effect of genotype
315 under HFD on all muscles.

316 Furthermore, we examined the impact of HFD on the CSA of MHC type IIA fibers from
317 PF and CT regions of T.medial, ECU and FCU muscles of both genotypes following

318 the HFD course. The analysis revealed a significant increase in the CSA of PF and CT
319 IIA fibers of ECU ($p < 0.01$ and 0.01) and FCU muscles ($p < 0.001$ and 0.001) of $Mstn^{-/-}$
320 HFD compared to the matched region of $Mstn^{-/-}$ ND. However, T.medial muscle of $Mstn^{-/-}$
321 HFD showed a decrease in the CSA of only PFIIA fibers ($p < 0.0001$) compared to the
322 corresponding PFIIA region in $Mstn^{-/-}$ ND. Moreover, CTIIA fibers of $Mstn^{+/+}$ HFD muscle
323 showed either increase in T.medial or a decrease in ECU muscles compared to
324 matched-muscle region kept on ND (Fig. 3g, j, and m).

325 **3. 5. Morphological evaluation of the muscle phenotype following a high-fat diet**

326 Having shown that the muscles of both genotypes responded differently in terms of the
327 CSA of the muscle fibers. We next examined whether HFD affects the muscle
328 phenotypic profile. The morphological evaluation revealed marked increase of slow
329 MHC fibers compared to fast MHC in $Mstn^{+/+}$ and $Mstn^{-/-}$ under HFD compared to
330 matched genotype muscle kept on a ND (Fig. 4f). Thus, quantifications of the
331 percentage of MHC type I, IIA, IIX and IIB fibers in the mid-belly of each muscle were
332 analyzed. All the muscles except the T.lateral muscle displayed a common observation
333 of myofiber shift from fast to slow MHC isoform due to the HFD. T.long and FCU
334 muscles showed a reduction in the percentage of type IIB ($p < 0.001$ and $p < 0.01$) and
335 ($p < 0.05$ and 0.01) in $Mstn^{+/+}$ HFD and $Mstn^{-/-}$ HFD compared to matched genotype
336 muscle kept on a ND. The reduction of MHC type IIB fibers was accompanied by a
337 shift towards an increase of the percentage of MHC type IIX fiber as shown in T.long
338 ($p < 0.001$) and FCU muscles ($p < 0.01$ and 0.05) of the $Mstn^{+/+}$ HFD and $Mstn^{-/-}$ HFD
339 respectively when compared to the corresponding genotype kept on a ND (Fig. 4a and
340 b). Furthermore, the fast to slow myofiber shift was also observed in T.medial muscle
341 of $Mstn^{-/-}$ HFD via a reduction of the percentage of MHC type IIX ($p < 0.001$) at the
342 expense of an increase of MHC type IIA ($p < 0.05$) compared to $Mstn^{-/-}$ ND (Fig. 4c). This
343 finding was supported with significant interaction ($p < 0.0001$) indicative for the effect of
344 myostatin deletion under HFD. Alternatively, ECU m of $Mstn^{+/+}$ HFD displayed another
345 form of fast to slow MHC shift, as a reduction in the percentage of MHC type IIX fibers
346 ($p < 0.0001$) towards an increase in the percentage of MHC type IIA fibers ($p < 0.01$) was
347 detected compared to $Mstn^{+/+}$ ND. Interestingly, however, in ECU muscle of $Mstn^{-/-}$ HFD
348 only MHC type IIB into IIX fiber shift ($p < 0.01$) was detected in comparison to $Mstn^{-/-}$
349 ND. In addition, a significant interaction ($p < 0.0001$) between the effects of genotype
350 and HFD could be detected (Fig. 4d). In contrast, no significant change was observed

351 in the myofiber composition of T.lateral muscle for both $Mstn^{+/+}$ and $Mstn^{-/-}$ following
352 HFD course (Fig. 4e). These data not only provides strong evidence that a muscle-
353 dependent attempt of fast to slow MHC shift but also point out the effect of myostatin
354 deletion on the muscle response in the course of HFD intake.

355 **4. Discussion**

356 The aim of this study was to determine the effect of high-fat diet intake on the
357 morphological properties and fiber composition of five phenotypically different forelimb
358 muscle of myostatin null mice.

359 Our data revealed that after 10 weeks of HFD intake, both $Mstn^{+/+}$ HFD and $Mstn^{-/-}$ HFD
360 showed increased body mass with no increase in the muscle-specific weight compared
361 to genotype-matched littermates under ND. The higher body mass may be attributed
362 to accumulation of abdominal body fat as shown previously (Matsakas et al. 2015).
363 Similarly, a study on rats showed no change in muscle weight following 16 weeks of
364 high-fat diet intake (Campbell et al. 2015). However, another report found that an
365 increase in body mass and a reduction in fat deposition were present following
366 inhibition of myostatin and HFD in skeletal muscle but not in adipose tissue (Guo et al.
367 2009).

368 Although the deletion of myostatin results in an increase in muscle fiber number as
369 previously shown (McPherron & Lee, 1997; Elashry et al. 2009; Elashry et al. 2017),
370 HFD for 10 weeks showed no alteration in fiber number which excludes not only the
371 possibility of myofiber hyperplasia but also myofiber apoptosis/loss due to lipotoxicity.
372 In agreement with our results, a study demonstrated that HFD intake for up to 16 weeks
373 failed to induce autophagy or even apoptosis in soleus and plantaris muscles of rat
374 (Campbell et al. 2015).

375 Despite the absence of muscle mass changes under HFD, there were remarkable
376 alterations in myofiber CSA of $Mstn^{-/-}$ mice. We have observed a reduction of CSA of
377 fast type IIB fibers in (T.long, FCU) and (T.lateral, T.long) muscles for $Mstn^{+/+}$ and $Mstn^{-/-}$
378 $^{-/-}$ respectively. However, there was an increase in the CSA of the same fiber isoform
379 in T.medial and ECU muscles of $Mstn^{+/+}$ and T.medial and FCU muscles of $Mstn^{-/-}$
380 under HFD. There was a significant interaction between genotype and diet on the CSA
381 of IIB and IIX fibers. These data suggest that the phenotype of muscles affects their
382 response to HFD. In this respect, a study using mixed high-fat and sucrose diet in rats

383 to evaluate muscle performance under obesity, showed that despite the quadriceps
384 muscle was functionally attenuated, the slow twitch soleus muscle was resistant to
385 diet-based muscle alteration (Collins et al. 2017). Taken together, the alteration of CSA
386 of type IIB and IIX fibers under HFD seems to be relevant to the muscle phenotype as
387 well as, to its metabolic activity. Furthermore, simultaneous myostatin deletion and
388 HFD induce muscle- dependent adaptation in order to utilize the excess of energy
389 intake. The present results also revealed that both genotypes responded differently
390 under HFD regarding the CSA of MHC type IIA fibers.

391 Our data revealed a reduction in CSA of IIB fibers from PF region under HFD as shown
392 in T.lateral and T.long muscles of both genotypes. The data point out that PF fibers
393 are more prone to HFD-induced fiber size reduction compared to the corresponding
394 muscle region under ND. Consequently, there was an increase in the CSA of CTIIB
395 fibers in T.lateral, ECU and FCU muscles of *Mstn*^{-/-} under HFD as well as, in CTIIX
396 fibers as shown in T.long, T.medial and ECU muscles compared to matching muscle
397 region for ND. These data were accompanied by a significant interaction indicating the
398 different response of *Mstn*^{-/-} under HFD. Moreover, such regional alteration might be a
399 compensatory adaptation to restore the PF fiber loss. Indeed, the PFIIB and PFIIX
400 fibers are the major store of muscle protein plus their glycolytic nature which it could
401 point out towards an alteration of the muscle metabolism. In the same line, we have
402 shown previously that fast myofibers are more liable to age-related muscle loss
403 (Elashry et al. 2009) as well as, being more prone to atrophy after five weeks of dietary
404 restriction (Elashry et al. 2017). The results suggest a compensatory mechanism in
405 order to maximize energy utilization in the course of the HFD regime by either reducing
406 the size of the fast MHC fibers or increasing the size of the slow fiber isoform. In the
407 same line, it has been reported that following three weeks of HFD, the whole body
408 metabolism was directed toward lipid oxidation with fast to slow myofiber shift in order
409 to maintain muscle-dependent insulin sensitivity (Trajcevski et al. 2013). MHC type IIA
410 fibers increased in CSA for both PF and CTIIA fibers in ECU and FCU muscles of *Mstn*^{-/-}
411 ⁻HFD. The data revealed enhanced oxidative response in the absence of myostatin.
412 In agreement with our results, a study in cats reported increased CSA of the oxidative
413 fibers in the oxidative region of the muscle compared to the same fiber type in glycolytic
414 region suggesting that the fiber size adaptation is more likely to compensate functional
415 demands (Gonyea, 1979).

416 Despite four out of five muscles displaying fast to slow myofiber shift, the mechanism
417 of fiber type shift varied between the muscles in the absence of myostatin. Along the
418 same lines, it has been shown in rats that HFD intake promoted the oxidative
419 metabolism in the fast twitch Extensor digialis longus (EDL) muscle via upregulation
420 of mitochondrial uncoupling protein 3 and pyruvate dehydrogenase kinase 4 and porin
421 expression in conjunction with, fast to slow shift of MHC expression (Mizunoya et al.
422 2013). Similarly, it has been reported that the muscle with high levels of oxidative
423 muscle fibers showed enhanced oxidation and reduction of body fat accumulation
424 (Abou Mrad et al. 1992). On the other hand, the muscle oxidative capacity is affected
425 by the quality and the number of mitochondria (Chanséaume & Morio, 2009). Taken
426 together, we postulate that HFD may enhance the oxidative metabolism of the muscle
427 fiber perhaps by increasing the mitochondrial biogenesis leading to slow fiber type shift.
428 This is in line with our data regarding the reduction of the MHC IIB and IIX isoforms
429 CSA in combination with the increased size of MHC IIA isoform. In agreement with this
430 concept, similar report revealed changes in the oxidative metabolism in case of insulin
431 resistance due to suppression of mitochondrial biogenesis (Johannsen & Ravussin,
432 2009; Abdul-Ghani & DeFronzo, 2010). Similarly, a study in rats revealed that HFD
433 intake was associated with intramyocellular lipid (IMCL) that require a comparable
434 increase of mitochondrial contents to maintain proper oxidative capacity (Van den
435 Broek et al. 2010).

436 Our data revealed no change in the myofiber composition of the T.lateral muscle of
437 both genotypes following HFD treatment. This muscle is composed primarily of IIB
438 fibers and responds only by losing the size of the fast fibers rather than enhancing the
439 slow phenotype shift under HFD. In the same line, a study in mouse extensor digialis
440 longus m (EDL) showed that although HFD induced a fast to slow fiber type shift in
441 $Mstn^{+/+}$, $Mstn^{-/-}$ under HFD displayed no changes in terms of MHC quantification and
442 succinate dehydrogenase (SDH) analysis (Matsakas et al. 2015). It has been
443 documented that, following five weeks of HFD in mice, the fast twitch EDL
444 demonstrated neither increase in fatty acid content nor increase in mitochondrial
445 oxidation, in contrast, slow twitch soleus muscle displayed a reduced force production
446 and fast to slow shift in troponin C type (Ciapaite et al. 2015). A similar study revealed
447 that, although no change in the muscle mass following 12 weeks of HFD in mice, there
448 was reduction in the contractile properties, increase of mitochondrial oxidation and

449 increased percentage of type IIX MHC in the EDL muscle indicating a phenotypic
450 alteration (Eshima et al. 2017).

451

452 **5. Conclusion**

453 The present study elucidated the effect of HFD on the morphological properties of the
454 forelimb skeletal musculature in the myostatin null mice. Although the significant
455 increase in the body weight, no change in the muscle mass was detected after 10
456 weeks of HFD intake. We provide evidence that HFD induces a reduction in the CSA
457 of the fast MHCIIB and MHCIIX myofibers of myostatin null in muscle and genotype-
458 dependent manner. HFD increases CSA of oxidative type IIA fibers of T.long and
459 T.medial of *Mstn*^{+/+} and ECU and FCU muscles of the *Mstn*^{-/-}. Moreover, type I fibers
460 of *Mstn*^{-/-} was less liable to HFD intake. We show that the anatomically located
461 peripheral fibers are more prone to HFD-induced fiber atrophy compared to the central
462 fibers. We provide evidence that combined HFD together with myostatin absence
463 induced a muscle-dependent fast to slow myofiber shift. Thus, it can be assume that
464 HFD reverses the hypermuscular phenotype of the myostatin null mice to utilize the
465 high energy intake. Our work provides a platform for further investigations interms of
466 using myostatin inhibition as a therapeutic application for metabolic diseases including
467 obesity.

468 **Competing interest**

469 All the authors have declared no competing interests regarding the publication of this
470 article.

471 **Author contributions**

472 MIE, AE, KG and AM have contributed to data collection and analysis, designed the
473 experiments, conceived and prepared the manuscript draft. SA and SW have
474 contributed to the manuscript preparation and data interpretation. AM and KP has
475 revised, formulated and finalized the submitted manuscript.

476 **6. References**

477 **Abdul-Ghani MA, DeFronzo RA** (2010) Pathogenesis of insulin resistance in skeletal
478 muscle. *Journal of biomedicine & biotechnology* **2010**, 476279.

479 **Abou Mrad J, Yakubu F, Lin D, Peters JC, Atkinson JB, Hill JO** (1992) Skeletal
480 muscle composition in dietary obesity-susceptible and dietary obesity-resistant rats.
481 *The American journal of physiology* **262**, R684-8.

482 **Allen DL, Hittel DS, McPherron AC** (2011) Expression and function of myostatin in
483 obesity, diabetes, and exercise adaptation. *Medicine and science in sports and*
484 *exercise* **43**, 1828–1835.

485 **Apovian CM** (2016) Obesity. Definition, comorbidities, causes, and burden. *The*
486 *American journal of managed care* **22**, s176-85.

487 **Campbell TL, Mitchell AS, McMillan EM, et al.** (2015) High-fat feeding does not
488 induce an autophagic or apoptotic phenotype in female rat skeletal muscle.
489 *Experimental biology and medicine* (Maywood, N.J.) **240**, 657–668.

490 **Chanséaume E, Morio B** (2009) Potential mechanisms of muscle mitochondrial
491 dysfunction in aging and obesity and cellular consequences. *International journal of*
492 *molecular sciences* **10**, 306–324.

493 **Ciapaite J, van den Berg SA, Houten SM, Nicolay K, van Dijk KW, Jeneson JA**
494 (2015) Fiber-type-specific sensitivities and phenotypic adaptations to dietary fat
495 overload differentially impact fast- versus slow-twitch muscle contractile function in
496 C57BL/6J mice. *The Journal of nutritional biochemistry* **26**, 155–164.

497 **Collins KH, Hart DA, Smith IC, et al.** (2017) Acute and chronic changes in rat soleus
498 muscle after high-fat high-sucrose diet. *Physiological reports* **5**.

499 **Egan B, Zierath JR** (2013) Exercise metabolism and the molecular regulation of
500 skeletal muscle adaptation. *Cell metabolism* **17**, 162–184.

501 **Elashry MI, Matsakas A, Wenisch S, Arnhold S, Patel K** (2017) The effect of caloric
502 restriction on the forelimb skeletal muscle fibers of the hypertrophic myostatin null
503 mice. *Acta histochemica* **119**, 582–591.

504 **Elashry MI, Otto A, Matsakas A, El-Morsy SE, Patel K** (2009) Morphology and
505 myofiber composition of skeletal musculature of the forelimb in young and aged wild
506 type and myostatin null mice. *Rejuvenation research* **12**, 269–281.

507 **Eshima H, Tamura Y, Kakehi S, et al.** (2017) Long-term, but not short-term high-fat
508 diet induces fiber composition changes and impaired contractile force in mouse fast-
509 twitch skeletal muscle. *Physiological reports* **5**.

510 **Gonyea WJ** (1979) Fiber size distribution in the flexor carpi radialis muscle of the cat.
511 *The Anatomical record* **195**, 447–454.

512 **Guo T, Jou W, Chanturiya T, Portas J, Gavrilova O, McPherron AC** (2009)
513 Myostatin inhibition in muscle, but not adipose tissue, decreases fat mass and
514 improves insulin sensitivity. *PloS one* **4**, e4937.

515 **Holloway GP, Bonen A, Spriet LL** (2009) Regulation of skeletal muscle mitochondrial
516 fatty acid metabolism in lean and obese individuals. *The American journal of clinical*
517 *nutrition* **89**, 455S-62S.

518 **Johannsen DL, Ravussin E** (2009) The role of mitochondria in health and disease.
519 *Current opinion in pharmacology* **9**, 780–786.

520 **Lange P de, Moreno M, Silvestri E, Lombardi A, Goglia F, Lanni A** (2007) Fuel
521 economy in food-deprived skeletal muscle. Signaling pathways and regulatory
522 mechanisms. *FASEB journal: official publication of the Federation of American*
523 *Societies for Experimental Biology* **21**, 3431–3441.

524 **Lebrasseur NK** (2012) Building muscle, browning fat and preventing obesity by
525 inhibiting myostatin. *Diabetologia* **55**, 13–17.

526 **Lumeng CN, Saltiel AR** (2011) Inflammatory links between obesity and metabolic
527 disease. *The Journal of clinical investigation* **121**, 2111–2117.

528 **Matsakas A, Foster K, Otto A, et al.** (2009) Molecular, cellular and physiological
529 investigation of myostatin propeptide-mediated muscle growth in adult mice.
530 *Neuromuscular disorders: NMD* **19**, 489–499.

531 **Matsakas A, Patel K** (2009) Skeletal muscle fibre plasticity in response to selected
532 environmental and physiological stimuli. *Histology and histopathology* **24**, 611–629.

533 **Matsakas A, Prosdocimo DA, Mitchell R, et al.** (2015) Investigating mechanisms
534 underpinning the detrimental impact of a high-fat diet in the developing and adult
535 hypermuscular myostatin null mouse. *Skeletal muscle* **5**, 38.

536 **McPherron AC, Lawler AM, Lee SJ** (1997) Regulation of skeletal muscle mass in
537 mice by a new TGF-beta superfamily member. *Nature* **387**, 83–90.

538 **McPherron AC, Lee SJ** (1997) Double muscling in cattle due to mutations in the
539 myostatin gene. *Proceedings of the National Academy of Sciences of the United*
540 *States of America* **94**, 12457–12461.

541 **McPherron AC, Lee S-J** (2002) Suppression of body fat accumulation in myostatin-
542 deficient mice. *The Journal of clinical investigation* **109**, 595–601.

543 **Mizunoya W, Iwamoto Y, Shirouchi B, et al.** (2013) Dietary fat influences the
544 expression of contractile and metabolic genes in rat skeletal muscle. *PloS one* **8**,
545 e80152.

546 **Muoio DM, Newgard CB** (2006) Obesity-related derangements in metabolic
547 regulation. *Annual review of biochemistry* **75**, 367–401.

548 **Pellegrinelli V, Rouault C, Rodriguez-Cuenca S, et al.** (2015) Human Adipocytes
549 Induce Inflammation and Atrophy in Muscle Cells during Obesity. *Diabetes* **64**,
550 3121–3134.

551 **Prado CMM, Wells JCK, Smith SR, Stephan BCM, Siervo M** (2012) Sarcopenic
552 obesity. A Critical appraisal of the current evidence. *Clinical nutrition (Edinburgh,*
553 *Scotland)* **31**, 583–601.

554 **Sharma M, Langley B, Bass J, Kambadur R** (2001) Myostatin in muscle growth and
555 repair. *Exercise and sport sciences reviews* **29**, 155–158.

556 **Srikanthan P, Hevener AL, Karlamangla AS** (2010) Sarcopenia exacerbates
557 obesity-associated insulin resistance and dysglycemia. Findings from the National
558 Health and Nutrition Examination Survey III. *PloS one* **5**, e10805.

559 **Stump CS, Henriksen EJ, Wei Y, Sowers JR** (2006) The metabolic syndrome. Role
560 of skeletal muscle metabolism. *Annals of medicine* **38**, 389–402.

561 **Tallis J, James RS, Seebacher F** (2018) The effects of obesity on skeletal muscle
562 contractile function. *The Journal of experimental biology* **221**.

563 **Thomas M, Langley B, Berry C, et al.** (2000) Myostatin, a negative regulator of
564 muscle growth, functions by inhibiting myoblast proliferation. *The Journal of*
565 *biological chemistry* **275**, 40235–40243.

566 **Trajcevski KE, O'Neill HM, Wang DC, et al.** (2013) Enhanced lipid oxidation and
567 maintenance of muscle insulin sensitivity despite glucose intolerance in a diet-
568 induced obesity mouse model. *PloS one* **8**, e71747.

569 **Valerio A, Cardile A, Cozzi V, et al.** (2006) TNF-alpha downregulates eNOS
570 expression and mitochondrial biogenesis in fat and muscle of obese rodents. *The*
571 *Journal of clinical investigation* **116**, 2791–2798.

572 **Van den Broek NMA, Ciapaite J, Feyter HMML de, et al.** (2010) Increased
573 mitochondrial content rescues in vivo muscle oxidative capacity in long-term high-
574 fat-diet-fed rats. *FASEB journal: official publication of the Federation of American*
575 *Societies for Experimental Biology* **24**, 1354–1364.

576 **Varma V, Yao-Borengasser A, Rasouli N, et al.** (2009) Muscle inflammatory
577 response and insulin resistance. Synergistic interaction between macrophages and
578 fatty acids leads to impaired insulin action. *American journal of physiology.*
579 *Endocrinology and metabolism* **296**, E1300-10.

580 **Wilde J de, Mohren R, van den Berg S, et al.** (2008) Short-term high fat-feeding
581 results in morphological and metabolic adaptations in the skeletal muscle of
582 C57BL/6J mice. *Physiological genomics* **32**, 360–369.

583 **Zhang C, McFarlane C, Lokireddy S, et al.** (2012) Inhibition of myostatin protects
584 against diet-induced obesity by enhancing fatty acid oxidation and promoting a
585 brown adipose phenotype in mice. *Diabetologia* **55**, 183–193.

586 **Zierath JR, Hawley JA** (2004) Skeletal muscle fiber type. Influence on contractile and
587 metabolic properties. *PLoS biology* **2**, e348.

588

589

590

591

592

593

594

595

596

597 .

598

599

600

601

602

603

604

605

606

607

608 **Table legends**

609 **Table (1)** Effect of high-fat diet on body mass and individual muscle weight

610 Average body mass (g) and muscle wet weight (mg) of normal diet (ND) and high food
611 diet (HFD) Mstn^{+/+} and Mstn^{-/-} mice (N= 8 for each experimental group). Significant
612 increase in body mass of Mstn^{+/+} and Mstn^{-/-} following 10 weeks of HFD. No significant
613 differences were detected in Mstn^{+/+} and Mstn^{-/-} muscles after HFD compared to ND.
614 All values presented as mean± SEM. **=p<0.01 for the comparison within the same
615 genotype and Φ =p<0.05 for the comparison between different genotype kept on a ND.

616 **Figures legends**

617 **Fig. 1** Evaluation of the muscle fiber number following high-fat diet

618 (a-f) Representative double labelled immunofluorescent images of ECU muscle show
619 MHCIIb (red), MHCIIa (green) and MHCIIx (black) positive fibers of Mstn^{+/+} and Mstn^{-/-}
620 mice kept on ND and HFD. (g) Average total muscle fiber number of T.lateral, T.long,
621 T.medial, ECU and FCU muscles of ND and HFD Mstn^{+/+} and Mstn^{-/-} mice (N=8 for
622 each experimental group). The muscles of Mstn^{-/-} show various degrees of myofiber
623 number increase compared to Mstn^{+/+}. T.medial muscle displays no changes following
624 Mstn^{-/-}. The muscles of both Mstn^{+/+} and Mstn^{-/-} on HFD demonstrate no detectable
625 difference in the muscle fiber number compared to genotype-matched muscles on ND.
626 All values presented as mean± SEM. **=p<0.01 and ****=p<0.001. Scale bar in a, b=
627 20 μm and in c, d, e, f= 200 μm. DAPI used as a nuclear counter stain in a, b.

628 **Fig. 2** Effect of high-fat diet on CSA of muscle fiber type

629 (a-o) Average CSA measurements (μm²) of MHC type IIB, IIX, IIA and type I myofibers
630 of ND and HFD Mstn^{+/+} and Mstn^{-/-} mice (250 fibers per each type and group were
631 measured using the axiovision software). Average CSA of types IIB (a) and IIX (b)
632 myofibers of T.lateral muscle; type IIB (c), IIX (d) and IIA (e) myofibers of T.long
633 muscle; type IIB (f), IIX (g), IIA (h) and type I (o) myofibers of T.medial muscle; type IIB
634 (i), IIX (j), IIA (k) of ECU muscle and type IIB (l), IIX (m), IIA (n) myofibers of FCU
635 muscle. All values presented as mean± SEM. *=p<0.05, **=p<0.01, ***=p<0.001 and
636 ****=p<0.001.

637 **Fig. 3** Effect of high-fat diet on the CSA of myofibers based on their regional distribution

638 (a-m) Average CSA measurement of peripheral (PF) and central (CT) myofibers of ND
639 and HFD Mstn^{+/+} and Mstn^{-/-} mice (250 fibers per each type and group were measured
640 using the axiovision software). Average CSA measurements of type IIB (a) and IIX (b)

641 myofibers of T.lateral muscle; type IIB (c) and IIX (d) myofibers of T.long muscle; type
642 IIB (e), IIX, (f) and IIA (g) myofibers of T.medial muscle; type IIB (h), IIX (i) and IIA (j)
643 myofibers of ECU muscle and types IIB (k), IIX (l) and IIA (m) myofibers of FCU muscle.
644 All values presented as mean± SEM. *=p<0.05, **=p<0.01, ***=p<0.001 and
645 ****=p<0.001.

646 **Fig. 4** Morphological evaluation of the muscle phenotype following high-fat diet
647 (a-e) Average percentage of type IIB, IIX, IIA and I myofibers of ND and HFD Mstn^{+/+}
648 and Mstn^{-/-} mice (N= 8 for each experimental group). (a) T.long, (b) FCU, (c) T.medial,
649 (d) ECU and (e) T.lateral muscles. (f) Reconstructive double immunofluorescent
650 images for ECU muscle of Mstn^{+/+} and Mstn^{-/-} following ten weeks on a ND and HFD.
651 Immunofluorescence of the muscle mid-belly cryo-sections show muscle phenotypic
652 composition based on MHCIIIB (red), MHCIIIX (unstained) and MHCIIA (green)
653 isoforms. All values presented as mean± SEM. (IIB) *=p<0.05, **=p<0.01 and
654 ***=p<0.001; (IIX) σ=p<0.05, σ σ =p<0.01, σ σ σ =p<0.001 and σ σ σ σ =p<0.0001;
655 (IIA) θ=p<0.05; θ θ=p<0.01. Scale bar in f = 200 μm.

656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674

675

676

Weight	Mstn ^{+/+}		Mstn ^{-/-}	
	ND	HFD	ND	HFD
Total body weight	30.8±0.4	36.9±0.6**	36±0.9 ^Φ	40.7±0.3**
M. triceps brachii Caput laterale	3.5±0.2	3.3±0.28	7±0.2	7.52±0.6
M. triceps brachii Caput longum	127±0.7	138±6.5	223±7.6	242±3.8
M. triceps brachii Caput mediale	10±1	9.2±0.4	13.4±0.5	15.2±1.2
M. extensor carpi ulnaris	9.7±0.1	9.5±0.2	14±0.8	15.6±0.3
M. flexor carpi ulnaris	3.5±0.2	3.3±0.3	7±0.3	7.5±0.6

677

678 **Table (1)** Effect of high-fat diet on body mass and individual muscle weight

679

680

681

682

683

684

685

686


687

688

689

690

691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715



716
717

Fig. 1 Evaluation of the muscle fiber number following high-fat diet




Fig. 2 Effect of high-fat diet on CSA of muscle fiber type

718

719

720

721


722

723

724

725

726



728

729 **Fig. 3** Effect of high-fat diet on the CSA of myofibers based on their regional distribution

730

731

732

733


734

735

736

737

738



740

741 **Fig. 4** Morphological evaluation of the muscle phenotype following high-fat diet