

**St. Catherine University**  
**SOPHIA**

---

Sr. Seraphim Gibbons Undergraduate Symposium

2013 Sr. Seraphim Gibbons Undergraduate  
Research Symposium

---

## 3 Step Synthesis of Benzocaine

Zoe A. Peterson  
*St. Catherine University*

Jordyn Wentland  
*St. Catherine University*

Follow this and additional works at: [https://sophia.stkate.edu/undergraduate\\_research\\_symposium](https://sophia.stkate.edu/undergraduate_research_symposium)

---

Peterson, Zoe A. and Wentland, Jordyn, "3 Step Synthesis of Benzocaine" (2013). *Sr. Seraphim Gibbons Undergraduate Symposium*. 27.  
[https://sophia.stkate.edu/undergraduate\\_research\\_symposium/2013/Sciences/27](https://sophia.stkate.edu/undergraduate_research_symposium/2013/Sciences/27)

This Event is brought to you for free and open access by the Conferences and Events at SOPHIA. It has been accepted for inclusion in Sr. Seraphim Gibbons Undergraduate Symposium by an authorized administrator of SOPHIA. For more information, please contact [amshaw@stkate.edu](mailto:amshaw@stkate.edu).

Benzocaine is a local anesthetic from the ester family. The drug benzocaine is normally used as a topical pain reliever or as a common ingredient in cough drops. Benzocaine is used in multiple forms including lotion, gel, liquid, lozenges, and sprays. When Benzocaine is applied in any form it temporarily numbs or blocks the nerve endings, which leads to a decrease in the amount of pain. A three-step synthesis is used to create benzocaine. First p-aminobenzoic acid is produced from taking p-acetotoluidide and oxidizing it with magnesium sulfate. Then to create an amino acid, p-aminobenzoic acid is then heated with hydrochloric acid. To esterify the amino acid and produce benzocaine ethanol must be added in the last step of the synthesis. To make this experiment greener we used hydrochloric acid in the last step instead of sulfuric acid.