

University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Faculty Publications in Food Science and Technology

Food Science and Technology Department

2013

The protein structure determines the sensitizing capacity of Brazil nut 2S albumin (*Ber e1*) in a rat food allergy model

Jolanda HM Van Bilsen TNO, j.vanbilsen@tno.nl

Léon MJ Knippels TNO

André H. Penninks TNO

Willem F. Nieuwenhuizen

Harmen HJ De Jongh TNO

See next page for additional authors

Follow this and additional works at: http://digitalcommons.unl.edu/foodsciefacpub



Part of the Food Science Commons

Van Bilsen, Jolanda HM; Knippels, Léon MJ; Penninks, André H.; Nieuwenhuizen, Willem F.; De Jongh, Harmen HJ; and Koppelman, Stef J., "The protein structure determines the sensitizing capacity of Brazil nut 2S albumin (Ber e1) in a rat food allergy model" (2013). Faculty Publications in Food Science and Technology. 128. http://digitalcommons.unl.edu/foodsciefacpub/128

This Article is brought to you for free and open access by the Food Science and Technology Department at Digital Commons@University of Nebraska -Lincoln. It has been accepted for inclusion in Faculty Publications in Food Science and Technology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

| Authors Jolanda HM Van Bilsen, Léon MJ Knippels, André H. Penninks, Willem F. Nieuwenhuizen, Harmen HJ De Jongh, and Stef J. Koppelman |
|---|
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |



POSTER PRESENTATION

Open Access

Criteria used to categorise proteins as allergens for inclusion in allergenonline.org: a curated database for risk assessment

Richard Goodman^{1*}, Ronald van Ree², Stefan Vieths³, Fatima Ferreira⁴, Motohiro Ebisawa⁵, Hugh A Sampson⁶, Joseph Baumert⁷, John Wise⁷, Steve L Taylor⁷

From 5th International Symposium on Molecular Allergology (ISMA 2013) Vienna, Austria. 6-7 December 2013

Background

New proteins introduced in foods through genetic engineering or processing are often evaluated for potential risks of eliciting food allergy (Codex, 2003). The primary risk is the transfer of an allergen or a protein nearly identical to an allergen and capable of causing IgE mediated cross-reactions into a new food source. The AllergenOnline.org database was developed to provide a curated set of allergenic sequences for bioinformatics comparisons to identify proteins that should be tested further (e.g. serum IgE binding). Curation is necessary as the scientific literature and general sequence databases include thousands of proteins labeled as allergens that lack proof of allergenic activity.

Methods

In 2006 we developed guidelines for reviewing and classifying candidate proteins as "allergens", "putative allergens" and those with "insufficient evidence" of causing IgE mediated allergic reactions in humans. Airway, contact, venom, salivary and food allergens are included. Criteria were developed for judging allergic subjects; allergen sources; protein characteristics, sequences; allergenic activity and IgE binding. Candidate allergens and peer reviewed publications are identified from the NCBI Protein and PubMed databases using keywords for review by our panel. Data, evaluations and decisions are stored in an achieved data management system during an annual update process. Access and FASTA searches are free, anonymous and unrecorded.

Results

We evaluate: descriptions of allergic and control subjects (numbers and ages; types of reactions); symptom onset; route of exposure (contact, ingestion, inhalation or injection); diagnostic methods (history, skin prick test or other challenge, specific IgE, basophil activation). We consider allergen source information (taxonomy, tissue source, maturity or processing); evidence the protein is present in the test material; protein characterization and source (purified vs. extract, native vs. recombinant); sequence and methods (cDNA or protein sequence); molecular mass and glycosylation status. Specific IgE measurements should include standards, controls, details of critical reagents and scoring criteria. Tests should purified proteins as well as extracts to demonstrate relevance. Four examples are presented.

Conclusions

Data quality and evidence of the importance of individual allergens varies markedly while the rate of publication of poorly described allergens and sequences is rising rapid.

Authors' details

¹University of Nebraska-Lincoln, Food Allergy Research and Resource Program, Lincoln, USA. ²Academic Medical Center, Dept. of Experimental Immunology, Amsterdam, Netherlands. ³Paul-Ehrilich-Institut, Div. of Allergology, Langen, Germany. ⁴University of Salzburg, Christian Doppler Laboratory for Allergy Diagn, Salzburg, Austria. ⁵Sagamihara National Hospital, Clinical Research Center for Allergy and Rheumatol, Sagamihara, Japan. ⁶Icahn School of Medicine at Mount Sinai, Division of Allergy, New York, USA. ⁷University of Nebraska—Lincoln, Food Allergy Research and Resource Program, Lincoln, USA.

Full list of author information is available at the end of the article



¹University of Nebraska-Lincoln, Food Allergy Research and Resource Program, Lincoln, USA

Published: 17 March 2014

doi:10.1186/2045-7022-4-S2-P12

Cite this article as: Goodman *et al.*: Criteria used to categorise proteins as allergens for inclusion in allergenonline.org: a curated database for risk assessment. *Clinical and Translational Allergy* 2014 **4**(Suppl 2):P12.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit

