

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/197342>

Please be advised that this information was generated on 2019-06-02 and may be subject to change.

# Towards reconstructing the quantum effective action of gravity

Benjamin Knorr<sup>1,\*</sup> and Frank Saueressig<sup>1,†</sup>

<sup>1</sup>*Institute for Mathematics, Astrophysics and Particle Physics (IMAPP),  
Radboud University Nijmegen, Heyendaalseweg 135, 6525 AJ Nijmegen, The Netherlands*

Starting from a parameterisation of the quantum effective action for gravity we calculate correlation functions for observable quantities. The resulting templates allow to reverse-engineer the couplings describing the effective dynamics from the correlation functions. Applying this new formalism to the autocorrelation function of spatial volume fluctuations measured within the Causal Dynamical Triangulations program suggests that the corresponding quantum effective action consists of the Einstein-Hilbert action supplemented by a non-local interaction term. We expect that our matching-template formalism can be adapted to a wide range of quantum gravity programs allowing to bridge the gap between the fundamental formulation and observable low-energy physics.

## INTRODUCTION

A characteristic feature of quantum gravity research is its fragmentation into disjoint branches including, *e.g.*, string theory [1, 2], loop quantum gravity [3, 4], the Asymptotic Safety program [5–8], Causal Dynamical Triangulations (CDT) [9], Causal Set Theory [10, 11], Group Field Theory [12, 13] or non-local gravity theories [14–16]. Each approach formulates its own fundamental dynamics typically at the Planck scale. The complexity of these formulations makes it quite hard to derive physical consequences from the fundamental formulation. A canonical way towards addressing this problem would be the construction of the quantum effective action  $\Gamma$  which encodes the dynamics of a quantum theory taking all quantum fluctuations into account. In this way it stores the outcome of a large number of (scattering) processes in an economical way. Generally, finding the exact form of  $\Gamma$  is considered equivalent to solving the theory. Not surprisingly calculating the quantum effective action is a hard problem. While fundamental Lagrangians describing our known physical theories are local and often restricted to a small number of interaction terms,  $\Gamma$  generically contains all possible interactions permitted by the symmetries of the theory. Furthermore, quantum corrections related to massless particles like the graviton give rise to non-local terms [17].

In contrast, correlation functions built from fluctuations of physical quantities like volumes and curvatures are accessible even at the non-perturbative level [18–21]. In this work we explicitly demonstrate that this information allows to reconstruct (parts of) the underlying quantum effective action, thereby taking a first explicit step in such a reconstruction program. Starting from the two-point autocorrelation functions of three-volume fluctuations measured in Monte Carlo simulations within CDT [18–21] we determine several couplings appearing in  $\Gamma$ . Our analysis provides first-hand evidence for the presence of non-local terms which could affect the gravitational dynamics at cosmic scales.

## THE QUANTUM EFFECTIVE ACTION FOR GRAVITY

In the case of gravity, the quantum effective action may be built from the spacetime metric and its curvature tensors. The foliation structure, constituting an elementary building block in the CDT program [22, 23], suggests to write  $\Gamma$  using the Arnowitt-Deser-Misner (ADM)-formalism, reviewed *e.g.* in [24]. In this case the spacetime metric  $g_{\mu\nu}$  is decomposed into a lapse function, a shift vector and a metric  $\sigma_{ij}$  measuring distances on spatial slices  $\Sigma$  orthogonal to a normal vector  $n_\mu$ . Curvature tensors constructed from  $g_{\mu\nu}$  can be separated into terms containing the intrinsic and extrinsic curvatures defined with respect to the foliation. For example, the Gauss-Codazzi equations relate the Ricci scalar  $R$  constructed from  $g_{\mu\nu}$  to the intrinsic Ricci scalar on the spatial slice  ${}^{(3)}R$ , the extrinsic curvature  $K_{ij}$ , and its trace  $K = \sigma^{ij} K_{ij}$  via

$$R = {}^{(3)}R - K_{ij}K^{ij} + K^2, \quad (1)$$

up to a total derivative.

The local part of the quantum effective action can then be organised in terms of a derivative expansion. The lowest order terms coincide with the Einstein-Hilbert action

$$\Gamma^{\text{local}} = \frac{1}{16\pi G_N} \int d^4x \sqrt{g} [2\Lambda - R] + \dots \quad (2)$$

where  $G_N$  and  $\Lambda$  are Newton's constant and the cosmological constant. The dots represent terms containing four or more derivatives as  $\int d^4x \sqrt{g} R^2$ , or the infamous Goroff-Sagnotti counterterm [25–27]. In principle, there could be additional coupling constants for each of the terms appearing on the right-hand side of (1), which will not be resolved here.

The non-local part of  $\Gamma$  typically contains inverse powers of the Laplacian  $\square \equiv -g^{\mu\nu} D_\mu D_\nu$  acting on curvature tensors. At second order in the curvature this leads to

terms of the form<sup>1</sup>

$$\Gamma^{\text{non-local}} = -\frac{b^2}{96\pi G_N} \int d^4x \sqrt{g} \mathcal{R} \mathcal{F}(\square) \mathcal{R}, \quad (3)$$

where  $\mathcal{R}$  is linear in the curvature tensors and the functions  $\mathcal{F}(\square)$  are known as form factors.<sup>2</sup> In the following, we are interested in the special case where

$$\mathcal{R} = R + \frac{1}{2} {}^{(3)}R \quad \text{and} \quad \mathcal{F}(\square) = \square^{-2}. \quad (4)$$

This specific combination gives rise to mass-type contributions to two-point correlation functions. Notably, there are other non-local terms giving rise to similar mass terms and (4) should be seen as one particular representative for this class. The task at hand is then to derive the values of the parameters  $G_N, \Lambda, b, \dots$  in terms of the parameters defining the fundamental theory.<sup>3</sup> In general, the latter set will vary from theory to theory. For CDT they are given by the bare Newton's constant  $\kappa_0$  and the relative size of spatial and time-like lines encoded in  $\Delta$ .

## DERIVATION OF THE MATCHING TEMPLATE

CDT simulations have been performed for spatial slices  $\bar{\Sigma}$  possessing the topology of a 3-sphere [18] and recently also for toroidal geometry [19–21]. Quite remarkably, the resulting profiles for the expectation value of three-volumes  $V_3(\bar{\Sigma}, t)$  as a function of the Euclidean time parameter  $t \in [0, 1[$  agrees with spacetime metrics of the form

$$\bar{g}_{\mu\nu} = \text{diag}(1, a(t)^2 \bar{\sigma}_{ij}(x)), \quad (5)$$

where

$$\begin{aligned} \text{torus:} \quad & a(t) = 1, & \bar{\sigma}_{ij} &= \delta_{ij}, \\ \text{3-sphere:} \quad & a(t) = \sin(\pi t), & \bar{\sigma}_{ij}(x) &= \bar{\sigma}_{ij}^{S^3}(x). \end{aligned} \quad (6)$$

For concreteness, we will focus mainly on the toroidal case and only briefly comment on the analogous analysis for the spherical case. In the former case the measured volume profile is essentially flat. Requiring that  $\bar{\Sigma} = S^1 \times S^1 \times S^1$  is a solution to the equations of motion

then fixes  $\Lambda = 0$ . The higher-derivative and non-local terms in the quantum effective action do not contribute to the dynamics for this case.

Motivated by the existence of a well-defined background geometry, one can then study the autocorrelation of three-volume fluctuations around the background,

$$\mathfrak{V}_2(t', t) = \langle \delta V_3(t') \delta V_3(t) \rangle. \quad (7)$$

Based on the quantum effective action, the fluctuations in the spatial metric are defined in the standard way, setting

$$\sigma_{ij}(t, x) = \bar{\sigma}_{ij} + \delta\sigma_{ij}(t, x). \quad (8)$$

The fluctuations in the 3-volume can then be found by expanding  $V_3(\Sigma, t) \equiv \int d^3x \sqrt{\sigma}$  in powers of the fluctuations. To leading order,

$$\delta V_3(t) = \frac{1}{2} \int d^3x \sqrt{\bar{\sigma}} \bar{\sigma}^{ij} \delta\sigma_{ij} + \mathcal{O}(\delta\sigma^2). \quad (9)$$

Introducing the fluctuation field  $\hat{\sigma}(t, x) \equiv \bar{\sigma}^{ij} \delta\sigma_{ij}(t, x)$ , the correlator  $\mathfrak{V}_2$  is given by the integral over the propagator  $\mathfrak{G}_{\hat{\sigma}\hat{\sigma}}(t', x'; t, x) \equiv \langle \hat{\sigma}(t', x') \hat{\sigma}(t, x) \rangle$

$$\mathfrak{V}_2(t', t) = \frac{1}{4} \int d^3x' \sqrt{\bar{\sigma}} \int d^3x \sqrt{\bar{\sigma}} \langle \hat{\sigma}(t', x') \hat{\sigma}(t, x) \rangle. \quad (10)$$

The computation of the two-point function proceeds by expanding the quantum effective action to second order in  $\hat{\sigma}$ ,  $\Gamma^{\text{quad}} = \frac{1}{32\pi G} \int d^4x \sqrt{g} \hat{\sigma} \Gamma^{(2)} \hat{\sigma}$ . The propagator  $\mathfrak{G}_{\hat{\sigma}\hat{\sigma}}(t', x'; t, x)$  can then be expressed in terms of the eigenvalue spectrum  $\{\bar{\lambda}_n\}$  and normalised eigenfunctions  $\Phi_n(t, x)$  of the differential operator  $\Gamma^{(2)}$ ,

$$\mathfrak{G} = 16\pi G_N \sum_n \frac{1}{\bar{\lambda}_n} \Phi_n^*(t', x') \Phi_n(t, x). \quad (11)$$

For compact spaces and correlation functions involving fluctuations which are averaged over  $\bar{\Sigma}$  the construction of the two-point function can be simplified by the following observation. On compact spaces the eigenfunctions  $\Phi_n(t, x)$  can be expanded in a complete set of orthonormal functions  $\psi_k(x)$  defined on  $\bar{\Sigma}$ ,

$$\Phi_n(x, t) = \sum_k \phi_{n,k}(t) \psi_k(x). \quad (12)$$

The spatial integrals appearing in (10) then project the expansion (12) on the spatially constant mode  $\psi_0(x) \equiv (V_3(\bar{\Sigma}))^{-1/2}$ . Hence

$$\mathfrak{V}_2(t', t) = 4\pi G_N V_3(\bar{\Sigma}) \sum_n' \frac{1}{\bar{\lambda}_n} \phi_n^*(t') \phi_n(t), \quad (13)$$

where  $\{\bar{\lambda}_n\}$  is the eigenvalue spectrum of  $\Gamma^{(2)}$  restricted to constant spatial modes. The prime indicates that the

<sup>1</sup> For UV modifications of gravity including these types of form factors see *e.g.* [28, 29].

<sup>2</sup> The form factors  $\mathcal{F}(\square)$ , defined through the matrix elements  $\langle x | \mathcal{F}(\square) | y \rangle \equiv L(x - y)$  allow to write non-local terms  $\int d^4x \sqrt{g(x)} \int d^4y \sqrt{g(y)} R(x) L(x - y) R(y)$  into quasi-local form. Regularity of the non-local terms may require supplementing the operator appearing in the structure functions by non-trivial endomorphism terms built from the curvature [30]. We do not include them in the expressions (3), since their resolution depends on correlators which are beyond the scope of this work.

<sup>3</sup> For the implementation of this strategy in effective field theory see [31].

zero mode should be excluded since it corresponds to an overall rescaling of the volume.

The next step computes  $\Gamma^{(2)}$  by expanding the local and non-local terms in  $\Gamma$  given by (2) and (3) to second order in  $\hat{\sigma}$ . Restricting to fluctuations which are constant on  $\bar{\Sigma}$ , the result reads

$$\Gamma^{(2)} = \frac{1}{3} [\partial_t^2 - b^2 + \frac{1}{2}\Lambda]. \quad (14)$$

The on-shell condition that  $\bar{\Sigma}$  is a flat torus fixes  $\Lambda = 0$ . The construction of  $\mathfrak{A}_2(t', t)$  then requires the eigen-

values and eigenfunctions of  $\Gamma^{(2)}$ , solving<sup>4</sup>

$$-\phi_n''(t) + b^2\phi_n(t) = \lambda_n\phi_n(t). \quad (15)$$

The solution is readily given in terms of Fourier modes

$$\phi_n(t) = e^{2\pi i n t}, \quad \lambda_n = (2\pi n)^2 + b^2, \quad n \in \mathbb{Z}. \quad (16)$$

Based on the spectrum (16) the propagator can be readily calculated. Carrying out the sum gives

$$\sum_n' \frac{1}{\lambda_n} \phi_n^*(t') \phi_n(t) = \frac{e^{2\pi i(t'-t)}}{2b(4\pi^2 + b^2)} \left[ (b - 2\pi i) {}_2F_1 \left( 1, 1 - \frac{ib}{2\pi}; 2 - \frac{ib}{2\pi} \middle| e^{2\pi i(t'-t)} \right) - (b \rightarrow -b) \right] + c.c. \quad (17)$$

## COMPARISON WITH CDT DATA

We are now in the situation that we can compare to the data obtained from CDT simulations on the torus for bare parameters

$$\kappa_0 = 2.2, \quad \Delta = 0.6, \quad (18)$$

and configurations built from  $N_4 = 160000$  simplices [19]. This point is located well within the de Sitter phase of the CDT phase diagram [21]. Averaging over all times, we can fit the correlator  $\mathfrak{A}_2(t, t + \Delta t)$  to extract the value of Newton's constant  $G_N$  and the mass parameter  $b^2$ . A least squares fit gives

$$G_N = 0.14 a_{\text{CDT}}^2, \quad \ell_{\text{Pl}} = 0.37 a_{\text{CDT}}, \quad b = 6.93/a_{\text{CDT}}. \quad (19)$$

Here  $a_{\text{CDT}}$  is the lattice spacing and  $\ell_{\text{Pl}} \equiv \sqrt{G_N}$  is the Planck length. We display the lattice data (blue dots) and our fit (red line) in Figure 1, and find a very good agreement between the two.

The relation between the lattice spacing and the physical radius  $r$  of the torus can be obtained by fitting the eigenvalues of the covariance matrix to the analytic form (16). Since the higher eigenvalues are less precise, we only fit the lowest three eigenvalues. Demanding that the resulting Newton's constant agrees with the one from the fit, gives a relation between the physical radius and the lattice spacing

$$r = 3.09 a_{\text{CDT}}. \quad (20)$$

The value of  $b$  can also be calculated from the lattice data directly by relating the kinetic and potential terms

of the 3-volume,

$$b_{\text{CDT}} = \sqrt{\frac{\bar{\Gamma}u}{\gamma(1+\gamma)}} \bar{V}_3. \quad (21)$$

Here,  $\bar{\Gamma}$  is related to the kinetic term,  $u$  to the potential term and  $\gamma$  is a critical exponent. For the data point (18), we take  $\bar{\Gamma} = 26.3$ ,  $u = 1.30 \times 10^{-6}$  ("first average then invert"),  $\bar{V}_3 = 2000$  and  $\gamma = 1.16$  given in [19], and find

$$b_{\text{CDT}} = (7.39 \pm 0.84 \pm 0.14\Delta\bar{\Gamma})/a_{\text{CDT}}, \quad (22)$$

which is consistent with the fit value within numerical precision.

Naturally, one would expect that the local part of the quantum effective action also contains higher-derivative terms like  $\int d^4x \sqrt{g} R^2$  containing four (or more) space-time derivatives. On the toroidal background the Hessian  $\Gamma^{(2)}$  then acquires an additional term proportional to  $\partial_t^4$ . Including this contribution in the fitting procedure shows that the related coefficient is negligible though.

Adapting the construction (13) to backgrounds where  $\bar{\Sigma} = S^3$  leads to the eigenproblem studied in [18]. The authors have shown there that the resulting fluctuation spectrum agrees very well with the numerical data. For the configuration (18) a comparison of the lowest eigenvalue of the covariance matrix with the continuum version yields

$$G_N = 0.23 a_{\text{CDT}}^2, \quad \ell_{\text{Pl}} = 0.48 a_{\text{CDT}}. \quad (23)$$

The relation between the lattice spacing and the physical radius  $r = 3.1 a_{\text{CDT}}$  is taken from [18] and agrees with the relation on the torus at the same bare parameters. Thus our continuum approach also reproduces the Monte Carlo results obtained for spherical topology.

Background independence of the quantum effective action suggests that the values for  $G_N$  obtained in different

<sup>4</sup> The mode  $\hat{\sigma}$  comes with a wrong-sign kinetic term. Following the CDT study [18], we consider the negative of the corresponding operator in the sequel.




Figure 1. The correlator  $\mathfrak{V}_2(t, t + \Delta t)$  on the torus in arbitrary units. Blue dots are the lattice data (averaged over all temporal slices), the red line indicates the fitted analytical result.

simulations should agree. The comparison of the Newton's constant found for the toroidal (19) and spherical topology (23) indicates different values though. This puzzle is resolved by the observation that the couplings  $G_N$  obtained from the volume correlations on the torus and spherical background are actually not the same: for non-vanishing background curvature higher-order curvature terms contribute to the  $\partial_t^2$ -terms appearing in  $\Gamma^{(2)}$ , so that the  $G_N$  obtained in (23) is actually a function of the Newton's constant defined in (2) and higher-derivative couplings. Likewise, the fact that the data obtained for  $\bar{\Sigma} = S^3$  suggest  $b = 0$  is not in contradiction with the toroidal results since the correlator of volume fluctuations evaluated on a background with non-zero curvature is not sensitive to this coupling. Instead it probes non-trivial endomorphism terms regulating the inverse Laplacians on a constant curvature background [30], which come with their own couplings.

### INFORMATION FROM COMPLEMENTARY CORRELATORS

The two-point autocorrelation function (7) gives access to some couplings appearing in the quantum effective action. A more complete picture can be developed by either studying correlation functions of different geometrical quantities or higher-order  $n$ -point functions.

*Higher-order  $n$ -point functions* – A natural generalisation of (7) are higher-order correlators involving the fluctuations of spatial volumes at  $n$  time-steps

$$\mathfrak{V}_n(t_1, \dots, t_n) \equiv \langle \delta V_3(t_1) \dots \delta V_3(t_n) \rangle. \quad (24)$$

These correlators can be constructed systematically by taking derivatives of  $\mathfrak{V}_2$  w.r.t. a suitable source. The three-point correlator  $\mathfrak{V}_3$  for instance then involves the

three-point vertex contracted with three propagators. On a flat, toroidal background  $\mathfrak{V}_n(t_1, \dots, t_n)$  carries information on couplings associated with terms built from  $n$  powers of the Riemann tensor (and its contractions).

*2-point functions involving curvatures* – Complementary, one may study the autocorrelation of curvature fluctuations involving the extrinsic or intrinsic curvature. Focusing on one concrete example, we introduce the averaged intrinsic curvature

$$\mathcal{R}_3(t) = \int d^3x \sqrt{\sigma} {}^{(3)}R. \quad (25)$$

The analogue of (9) is then obtained by expanding  $\mathcal{R}_3(t)$  in terms of fluctuations

$$\begin{aligned} \tilde{\sigma}(x, t) &= a(t) \bar{\sigma}^{\mu\nu}(x) \delta \sigma_{\mu\nu}(x, t), \\ \tilde{\sigma}_{\mu\nu}^{\text{TL}}(x, t) &= a(t) \delta \sigma_{\mu\nu}(x, t) - \frac{1}{3} \bar{\sigma}_{\mu\nu}(x) \tilde{\sigma}(x, t). \end{aligned} \quad (26)$$

Dropping integrals over total derivatives, fluctuations of the averaged intrinsic curvature are related to the fluctuation fields by

$$\delta \mathcal{R}_3(t) = \int d^3x \sqrt{\bar{\sigma}} \left( \frac{1}{6} {}^{(3)}\tilde{R} \tilde{\sigma} - {}^{(3)}\tilde{S}^{\mu\nu} \tilde{\sigma}_{\mu\nu}^{\text{TL}} \right), \quad (27)$$

where  ${}^{(3)}\tilde{R} = {}^{(3)}\tilde{R}/a(t)^2$  and  ${}^{(3)}\tilde{S}^{\mu\nu} = {}^{(3)}\tilde{S}^{\mu\nu}/a(t)^4$  indicate the background spatial Ricci scalar and trace-free spatial Ricci tensor, respectively. The autocorrelation function can then again be expressed in terms of the propagators of the fluctuation fields,

$$\begin{aligned} \langle \delta \mathcal{R}_3(t') \delta \mathcal{R}_3(t) \rangle &= \int d^3x \sqrt{\bar{\sigma}} \int d^3y \sqrt{\bar{\sigma}} \left[ \frac{{}^{(3)}\tilde{R}^2}{36} \mathfrak{G}_{\tilde{\sigma}\tilde{\sigma}} \right. \\ &\quad - \frac{1}{3} {}^{(3)}\tilde{R} {}^{(3)}\tilde{S}^{\mu\nu} (\mathfrak{G}_{\tilde{\sigma}\tilde{\sigma}^{\text{TL}}})_{\mu\nu} \\ &\quad \left. + {}^{(3)}\tilde{S}^{\mu\nu} {}^{(3)}\tilde{S}^{\rho\sigma} (\mathfrak{G}_{\tilde{\sigma}^{\text{TL}}\tilde{\sigma}^{\text{TL}}})_{\mu\nu\rho\sigma} \right]. \end{aligned} \quad (28)$$

Notably  $\langle \delta \mathcal{R}_3(t') \delta \mathcal{R}_3(t) \rangle$  vanishes on a toroidal background since it is proportional to the background curvature. This feature is owed to terminating the expansion (27) at leading order in the fluctuation fields. Once terms quadratic in the fluctuations are included, the correlator (28) involves non-zero contributions related to the four-point vertex of the fluctuation fields.

### CONCLUSIONS

We introduced a new research program to reverse-engineer the quantum effective action for gravity from correlation functions. This provides for the first time a direct link between continuum and lattice in quantum gravity beyond abstract quantities like critical exponents [32, 33] and spectral dimensions [34–37]. Where lattice data was available, agreement with an Einstein-Hilbert action, potentially amended by a particular non-local



interaction or higher-order scalar curvature terms, was found.

A particularly intriguing result is that the lattice simulations on the torus suggest the presence of a (non-local) mass term. The authors of [19] argue that the occurrence of this term is a genuine quantum gravity effect. Generically, it is expected that quantum fluctuations of massless particles induce these kind of non-local terms in the quantum effective action. A prototypical example is provided by quantum chromodynamics where such terms correctly describe the non-perturbative gluon propagator in the IR [38–41].

In general, it is conceivable that non-local gravitational interactions provide a dynamical explanation of dark energy, without the need for a fine-tuned cosmological constant [42–44]. In particular, the non-local contribution of (3),

$$\Gamma^{\text{non-local}} = -\frac{b^2}{96\pi G_N} \int d^4x \sqrt{g} R \frac{1}{\square^2} R, \quad (29)$$

forms a key part of the Maggiore-Mancarella cosmological model [43, 45], which has been highly successful in describing the cosmological evolution of the Universe. It is intriguing that the non-classical behaviour seen on the lattice is compatible with such a non-local quantum effect.

In this work, we analysed the arguably simplest non-trivial correlation function describing the autocorrelation of 3-volume fluctuations at two different times (7). The systematic extension to correlation functions of higher order or other structures is evident. Notably, a measurement of the correlators (24) and (28) may be actually feasible within the CDT program, thereby providing further information on the quantum effective action. In particular, correlation functions of the averaged intrinsic curvature may be obtained by summing deficit angles or new sophisticated methods to measure curvature on general quantum spacetimes recently introduced in [46, 47]. We stress that the construction of reverse-engineering the quantum effective action from correlators is not limited to CDT, but applies to any given theory of quantum gravity in which the corresponding quantities can be computed.

We thank J. Ambjørn, A. Görlich and R. Loll for interesting discussions, and A. Görlich for providing us with the numerical data from the CDT simulations. Interesting discussions with J. Donoghue, N. Klitgaard and M. Maggiore are acknowledged. This research is supported by the Netherlands Organisation for Scientific Research (NWO) within the Foundation for Fundamental Research on Matter (FOM) grant 13VP12.

---

\* b.knorr@science.ru.nl

† f.saueressig@science.ru.nl

- [1] K. Becker, M. Becker, and J. H. Schwarz, *String theory and M-theory: A modern introduction* (Cambridge University Press, 2006).
- [2] Volker Schomerus, *A Primer on String Theory* (Cambridge University Press, 2017).
- [3] Thomas Thiemann, “Lectures on loop quantum gravity,” *Proceedings, 271st WE-Heraeus Seminar on Aspects of Quantum Gravity: From Theory to Experiment Search: Bad Honnef, Germany, February 25-March 1, 2002*, *Lect. Notes Phys.* **631**, 41–135 (2003), [41(2002)], arXiv:gr-qc/0210094 [gr-qc].
- [4] Carlo Rovelli, *Quantum gravity* (2004).
- [5] Max Niedermaier and Martin Reuter, “The Asymptotic Safety Scenario in Quantum Gravity,” *Living Rev. Rel.* **9**, 5–173 (2006).
- [6] Martin Reuter and Frank Saueressig, “Quantum Einstein Gravity,” *New J.Phys.* **14**, 055022 (2012), arXiv:1202.2274 [hep-th].
- [7] Robert Percacci, *An Introduction to Covariant Quantum Gravity and Asymptotic Safety*, 100 Years of General Relativity, Vol. 3 (World Scientific, 2017).
- [8] Astrid Eichhorn, “Status of the asymptotic safety paradigm for quantum gravity and matter,” in *Black Holes, Gravitational Waves and Spacetime Singularities Rome, Italy, May 9-12, 2017* (2017) arXiv:1709.03696 [gr-qc].
- [9] J. Ambjørn, A. Goerlich, J. Jurkiewicz, and R. Loll, “Nonperturbative Quantum Gravity,” *Phys. Rept.* **519**, 127–210 (2012), arXiv:1203.3591 [hep-th].
- [10] Rafael D. Sorkin, “Causal sets: Discrete gravity,” in *Lectures on quantum gravity. Proceedings, School of Quantum Gravity, Valdivia, Chile, January 4-14, 2002* (2003) pp. 305–327, arXiv:gr-qc/0309009 [gr-qc].
- [11] Sumati Surya, “Directions in Causal Set Quantum Gravity,” (2011), arXiv:1103.6272 [gr-qc].
- [12] Aristide Baratin and Daniele Oriti, “Group field theory with non-commutative metric variables,” *Phys. Rev. Lett.* **105**, 221302 (2010), arXiv:1002.4723 [hep-th].
- [13] Daniele Oriti, “The microscopic dynamics of quantum space as a group field theory,” in *Proceedings, Foundations of Space and Time: Reflections on Quantum Gravity: Cape Town, South Africa* (2011) pp. 257–320, arXiv:1110.5606 [hep-th].
- [14] Tirthabir Biswas, Anupam Mazumdar, and Warren Siegel, “Bouncing universes in string-inspired gravity,” *JCAP* **0603**, 009 (2006), arXiv:hep-th/0508194 [hep-th].
- [15] Leonardo Modesto, “Super-renormalizable Quantum Gravity,” *Phys. Rev.* **D86**, 044005 (2012), arXiv:1107.2403 [hep-th].
- [16] Leonardo Modesto and Lesław Rachwał, “Nonlocal quantum gravity: A review,” *Int. J. Mod. Phys.* **D26**, 1730020 (2017).
- [17] John F. Donoghue, Mikhail M. Ivanov, and Andrey Shkerin, “EPFL Lectures on General Relativity as a Quantum Field Theory,” (2017), arXiv:1702.00319 [hep-th].
- [18] J. Ambjørn, A. Gorlich, J. Jurkiewicz, and R. Loll, “The Nonperturbative Quantum de Sitter Universe,” *Phys. Rev.* **D78**, 063544 (2008), arXiv:0807.4481 [hep-th].
- [19] J. Ambjørn, Z. Drogosz, Jakub Gizbert-Studnicki, A. Görlich, Jerzy Jurkiewicz, and D. Nemeth, “Impact of topology in causal dynamical triangulations quantum gravity,” *Phys. Rev.* **D94**, 044010 (2016), arXiv:1604.08786 [hep-th].
- [20] Jan Ambjørn, Jakub Gizbert-Studnicki, Andrzej Gör-

- lich, Kevin Grosvenor, and Jerzy Jurkiewicz, “Four-dimensional CDT with toroidal topology,” *Nucl. Phys.* **B922**, 226–246 (2017), [arXiv:1705.07653 \[hep-th\]](#).
- [21] J. Ambjørn, J. Gizbert-Studnicki, A. Görlich, J. Jurkiewicz, and D. Németh, “The phase structure of Causal Dynamical Triangulations with toroidal spatial topology,” (2018), [arXiv:1802.10434 \[hep-th\]](#).
- [22] Jan Ambjorn and R. Loll, “Nonperturbative Lorentzian quantum gravity, causality and topology change,” *Nucl. Phys.* **B536**, 407–434 (1998), [arXiv:hep-th/9805108 \[hep-th\]](#).
- [23] J. Ambjorn, J. Jurkiewicz, and R. Loll, “Emergence of a 4-D world from causal quantum gravity,” *Phys. Rev. Lett.* **93**, 131301 (2004), [arXiv:hep-th/0404156 \[hep-th\]](#).
- [24] Eric Gourgoulhon, “3+1 formalism and bases of numerical relativity,” (2007), [arXiv:gr-qc/0703035 \[GR-QC\]](#).
- [25] Marc H. Goroff and Augusto Sagnotti, “QUANTUM GRAVITY AT TWO LOOPS,” *Phys. Lett.* **B160**, 81 (1985).
- [26] Marc H. Goroff and Augusto Sagnotti, “The Ultraviolet Behavior of Einstein Gravity,” *Nucl. Phys.* **B266**, 709 (1986).
- [27] Holger Gies, Benjamin Knorr, Stefan Lippoldt, and Frank Saueressig, “Gravitational Two-Loop Counterterm Is Asymptotically Safe,” *Phys. Rev. Lett.* **116**, 211302 (2016), [arXiv:1601.01800 \[hep-th\]](#).
- [28] Tirthabir Biswas, Erik Gerwick, Tomi Koivisto, and Anupam Mazumdar, “Towards singularity and ghost free theories of gravity,” *Phys. Rev. Lett.* **108**, 031101 (2012), [arXiv:1110.5249 \[gr-qc\]](#).
- [29] James Edholm, Alexey S. Koshelev, and Anupam Mazumdar, “Behavior of the Newtonian potential for ghost-free gravity and singularity-free gravity,” *Phys. Rev.* **D94**, 104033 (2016), [arXiv:1604.01989 \[gr-qc\]](#).
- [30] A. O. Barvinsky, “Nonlocal gravity, Schwinger-Keldysh technique, dark energy and all that,” in *Proceedings, 17th International Seminar on High Energy Physics (Quarks 2012): Yaroslavl, Russia, June 4-7, 2012*, INR (INR, Moscow, Russia, 2012).
- [31] John F. Donoghue and Basem Kamal El-Menoufi, “Covariant non-local action for massless QED and the curvature expansion,” *JHEP* **10**, 044 (2015), [arXiv:1507.06321 \[hep-th\]](#).
- [32] Kevin Falls, “Critical scaling in quantum gravity from the renormalisation group,” (2015), [arXiv:1503.06233 \[hep-th\]](#).
- [33] Jorn Biemans, Alessia Platania, and Frank Saueressig, “Quantum gravity on foliated spacetimes: Asymptotically safe and sound,” *Phys. Rev.* **D95**, 086013 (2017), [arXiv:1609.04813 \[hep-th\]](#).
- [34] J. Ambjorn, J. Jurkiewicz, and R. Loll, “Spectral dimension of the universe,” *Phys. Rev. Lett.* **95**, 171301 (2005), [arXiv:hep-th/0505113 \[hep-th\]](#).
- [35] O. Lauscher and M. Reuter, “Fractal spacetime structure in asymptotically safe gravity,” *JHEP* **10**, 050 (2005), [arXiv:hep-th/0508202 \[hep-th\]](#).
- [36] Dario Benedetti and Joe Henson, “Spectral geometry as a probe of quantum spacetime,” *Phys. Rev.* **D80**, 124036 (2009), [arXiv:0911.0401 \[hep-th\]](#).
- [37] Giuseppe Clemente and Massimo D’Elia, “Spectrum of the Laplace-Beltrami Operator and the Phase Structure of Causal Dynamical Triangulation,” (2018), [arXiv:1804.02294 \[hep-th\]](#).
- [38] Philippe Boucaud, A. Le Yaouanc, J. P. Leroy, J. Micheli, O. Pene, and J. Rodriguez-Quintero, “Testing Landau gauge OPE on the lattice with a  $\langle A^2 \rangle$  condensate,” *Phys. Rev.* **D63**, 114003 (2001), [arXiv:hep-ph/0101302 \[hep-ph\]](#).
- [39] M. A. L. Capri, D. Dudal, J. A. Gracey, V. E. R. Lemes, R. F. Sobreiro, S. P. Sorella, and H. Verschelde, “A Study of the gauge invariant, nonlocal mass operator  $\text{Tr} \int d^4x F_{\mu\nu} (D^2)^{-1} F_{\mu\nu}$  in Yang-Mills theories,” *Phys. Rev.* **D72**, 105016 (2005), [arXiv:hep-th/0510240 \[hep-th\]](#).
- [40] David Dudal, John A. Gracey, Silvio Paolo Sorella, Nele Vandersickel, and Henri Verschelde, “A Refinement of the Gribov-Zwanziger approach in the Landau gauge: Infrared propagators in harmony with the lattice results,” *Phys. Rev.* **D78**, 065047 (2008), [arXiv:0806.4348 \[hep-th\]](#).
- [41] M. Peláez, M. Tissier, and N. Wschebor, “Two-point correlation functions of QCD in the Landau gauge,” *Phys. Rev.* **D90**, 065031 (2014), [arXiv:1407.2005 \[hep-th\]](#).
- [42] Michele Maggiore, “Phantom dark energy from nonlocal infrared modifications of general relativity,” *Phys. Rev.* **D89**, 043008 (2014), [arXiv:1307.3898 \[hep-th\]](#).
- [43] Michele Maggiore and Michele Mancarella, “Nonlocal gravity and dark energy,” *Phys. Rev.* **D90**, 023005 (2014), [arXiv:1402.0448 \[hep-th\]](#).
- [44] Stefano Foffa, Michele Maggiore, and Ermis Mitsou, “Cosmological dynamics and dark energy from nonlocal infrared modifications of gravity,” *Int. J. Mod. Phys.* **A29**, 1450116 (2014), [arXiv:1311.3435 \[hep-th\]](#).
- [45] Enis Belgacem, Yves Dirian, Stefano Foffa, and Michele Maggiore, “Nonlocal gravity. Conceptual aspects and cosmological predictions,” (2017), [arXiv:1712.07066 \[hep-th\]](#).
- [46] N. Klitgaard and R. Loll, “Introducing Quantum Ricci Curvature,” (2017), [arXiv:1712.08847 \[hep-th\]](#).
- [47] N. Klitgaard and R. Loll, “Quantizing quantum Ricci curvature,” (2018), [arXiv:1802.10524 \[hep-th\]](#).