

# Assessment of in vivo loading history of the patellofemoral joint: a study combining patellar position, tilt, alignment and bone SPECT/CT

Stephan N. Schön · Faik K. Afifi · Helmut Rasch ·  
Felix Amsler · Niklaus F. Friederich ·  
Markus P. Arnold · Michael T. Hirschmann

Received: 30 December 2012 / Accepted: 27 September 2013 / Published online: 11 October 2013  
© Springer-Verlag Berlin Heidelberg 2013

## Abstract

**Purpose** The current study investigates whether patella height and tilt or leg alignment influence the intensity values as well as the distribution pattern of single photon emission computerized tomography/computerized tomography (SPECT/CT) tracer uptake in the patellofemoral joint.

**Methods** 99mTc-HDP-SPECT/CT and radiographs of consecutive 84 knees were prospectively obtained. Lateral radiographs were analyzed in terms of patellar height, Insall-Salvati index and modified Insall-Salvati index. Skyline views were analyzed for Laurin's lateral patellofemoral angle. On long-leg radiographs, the mechanical leg alignment was classified as varus, valgus or neutral. SPECT/CT was analyzed for each anatomical region using a previously validated SPECT/CT localization and grading algorithm. Mean, standard deviation, minimum and maximum of grading for each area of the localization scheme were recorded. Non-parametric Spearman's correlations were used to correlate patellar height, lateral patellar angle and leg alignment with the tracer uptake intensity. Chi-square statistics were used for categorical data ( $p < 0.05$ ).

**Results** A patella baja correlated significantly with higher SPECT/CT tracer uptake in all patellar and lateral femoral regions ( $p < 0.001$ ). A higher lateral patellar tilt correlated significantly with higher tracer uptake in the superior lateral femoral parts and the tibial tubercle. In mechanically varus aligned knees, there was significantly higher SPECT/CT tracer uptake on the medial and in valgus knees on the lateral part of the patellofemoral joint ( $p < 0.05$ ).

**Conclusions** As the intensity and distribution of the SPECT/CT significantly correlated with patella baja and patellar tilt, SPECT/CT might be considered as imaging modality for evaluating patients with patellofemoral disorders and for follow-up of patients after patellofemoral realignment procedures.

**Level of evidence** Diagnostic study, Level II.

**Keywords** Knee · SPECT/CT · Patella · Loading · Tilt · Patella height

## Introduction

Realignment procedures such as tibial tubercle medialization and anteriorization aim to reduce the mechanical loading on the patellofemoral joint [1, 24]. Others such as trochleoplasty strive to improve the stability of the patellofemoral joint without increasing the patellofemoral contact pressure [3, 6]. In vitro studies, a patella infera position has been related with altered contact areas and increased pressure within the patellofemoral joint [26, 31, 34].

To date, there is no optimal imaging modality, which could be equally sensitive and specific to identify changes in patellofemoral joint loading. Recently, the benefits of single photon emission computerized tomography/computerized tomography (SPECT/CT) have been highlighted

---

S. N. Schön · F. K. Afifi · N. F. Friederich ·  
M. P. Arnold · M. T. Hirschmann (✉)  
Department of Orthopaedic Surgery and Traumatology,  
Kantonsspital Baselland-Bruderholz, 4101 Bruderholz,  
Switzerland  
e-mail: michael.hirschmann@ksbh.ch;  
michael.hirschmann@unibas.ch

H. Rasch  
Institute for Radiology and Nuclear Medicine, Kantonsspital  
Baselland-Bruderholz, 4101 Bruderholz, Switzerland

F. Amsler  
Amsler Consulting, Basel, Switzerland

in orthopaedic patients with knee problems [10–17, 19, 32]. SPECT/CT is a hybrid imaging, which combines 3D scintigraphy (SPECT) and CT into one imaging modality. The most commonly used tracers are diphosphonates, which target active osteoblasts. SPECT/CT offers the benefits of combined anatomical, mechanical (CT) and functional (SPECT) imaging [10–15, 19]. In a previous study, Hirschmann et al. [18] found that the intensity and distribution of SPECT/CT tracer uptake within the tibiofemoral joint reflect the loading pattern of the knee joint with regard to the mechanical and anatomical alignment.

SPECT and SPECT/CT are able to give valuable information about in vivo joint loading of the tibiofemoral joint [18, 25]. Biomechanically the patellofemoral joint is not only exposed to the joint reaction and loading forces but also shear forces. Hence, it was unclear whether SPECT/CT is also a useful imaging modality for the evaluation of in vivo joint loading of the patellofemoral joint. In this case, it could be established as novel imaging modality evaluating patients with patellofemoral disorders and for follow-up of patients after patellofemoral realignment procedures.

The purpose of this study was to investigate whether the position of the patella as well as the mechanical alignment


influences the intensity and the distribution pattern of bone SPECT/CT tracer uptake in the patellofemoral joint. The hypotheses were that an increased  $^{99m}\text{Tc}$ -HDP tracer uptake in SPECT/CT images would be present within the patellofemoral compartment in situations of altered patella position such as patella infera, patella alta, overly tilted patella or in varus or valgus malalignment of the femoro-tibial joint (Fig. 1).

## Materials and methods

$^{99m}\text{Tc}$ -HDP-SPECT/CT and conventional radiographs of consecutive 84 knees ( $n = 71$  patients, male:female = 33:38, mean age  $48 \pm 16$ ) were prospectively obtained. The patients underwent these imaging modalities due to knee pain. Exclusion criteria were a known history of avascular necrosis of the knee, a tumour, Paget's disease, joint infection, periarticular fracture, neuropathic arthropathy, reactive arthritis, gout, arthroscopic surgery within the last 3 months.

Skyline views as well as lateral radiographs were obtained. Lateral radiographs were then analyzed in terms of patellar height, the Insall-Salvati index (normal range

**Fig. 1** The localization scheme, which allows grading and identification of increased SPECT/CT tracer uptake in each patellofemoral area of interest




0.8–1.2) and the modified Insall-Salvati index (normal range <2) [9, 20]. The Caton-Deschamps index was also measured on lateral radiographs [5]. Skyline views were analyzed measuring the lateral patellofemoral angle according to Laurin [28] (Fig. 2).

Long-leg radiographs (femur to ankle) were obtained in standing position with the tibial tubercle facing forward. The X-ray beam was centred at the knee at a distance of 2.4 m. A setting of 100–300 mA/s and 80–90 kV was used. These radiographs were used to assess the mechanical leg alignment, which was then classified as varus, valgus or neutral. Mechanical alignment was measured as the angle at the intersection of a line connecting femoral head and intercondylar notch centre with a line connecting talar surface centre and tibial eminence sulcus base. Knee alignment of  $180^\circ \pm 1$  from the full-limb radiograph was taken as neutral axis. Angles of less than  $179^\circ$  were labelled varus, and angles greater than  $181^\circ$  were labelled valgus alignment. In addition, three different patellar tilt angles were measured on CT slices (Fig. 3).

The patellar tilt angle according to Grelsamer was measured as the angle between the mid-patellar line and the horizontal line [8]. The patellar tilt angle according to Sasaki [33] was measured as the angle between the mid-patellar line and the anterior trochlear line. The patellar tilt angle modified by Fulkerson [7] was measured as the angle between the mid-patellar line and the posterior condyle line. The angle was indicated as positive when it was open medially.

The tibial tuberosity trochlear groove (TT-TG) distance was measured. It was indicated as positive when the tibial tuberosity was lateral in relation to the trochlear groove.

All SPECT/CTs were performed using a Symbia T16 (Siemens, Erlangen, Germany). The CT collimation was  $16 \times 0.75$  mm. Planar scintigraphic images were taken in three phases, the perfusion phase (immediately after injection), the soft tissue phase (1–5 min after injection) and the delayed metabolic phase (2–3 h after injection). A commercial 700 MBq Tc-99m-HDP tracer (Mallinckrodt,




**Fig. 3** Schematically transaxial CT slices of a distal right femur (1–3) and a transaxial slice in overlay technique of a distal right femur and a proximal tibia. This figure shows three methods of measuring the patella tilt angle described by Grelsamer (1), Sasaki (2) and Fulkerson (3). Nr. 4 illustrates the measurement of the TT-TG distance in overlay technique. [7, 8, 33]

Wollerau, Switzerland) was used. Delayed SPECT images were obtained with a matrix size of  $128 \times 128$ , an angle step of 32, and a time per frame of 25 s 2–3 h after injection.

Data were processed by interactive reconstruction and images were displayed in transaxial, coronal and sagittal planes (Syngo, Siemens, Erlangen, Germany).

The SPECT/CT images were analyzed for the patellofemoral regions using a previously validated SPECT/CT localization and grading algorithm (Fig. 1) [18]. The localization and grading scheme, which showed high inter- and intra-observer reliability, defines 8 tibial, 9 femoral and 4 patellar regions. The anatomical area (femur, tibia, patella) is indicated with capital letters (F, T, P). The femur (F) is divided into nine zones which include one shaft and eight distal femoral zones. Each

**Fig. 2** The upper part of the figure shows the measurement of the lateral patellofemoral angle according to Laurin. The lateral patellofemoral angle is formed by the lines A–A and B–B. Normally, the angle is open laterally. In knees with an abnormal lateral angle, the lines are parallel or the angle is open medially. [28]



distal femoral zone is represented with a number (1 for medial, 2 for lateral) and two small letters (a-anterior, p-posterior and s-superior, i-inferior). The tibia (T) is divided into eight zones which include one shaft region, one region of the tibial tubercle and six tibial regions. Each tibial zone is represented with a number (1 for medial, 2 for lateral, 3 for mid zone) and a small letter (a-anterior, p-posterior). The patella (P) is divided into four zones (superomedial, superolateral, inferomedial and inferolateral).

Mean, standard deviation, minimum and maximum of grading for each area of the localization scheme were recorded using a semiquantitative colour-coded grading scale (0–10). The inter-observer reliability and intra-observer reliability quantified by intraclass correlation coefficients (ICC) were all  $>0.85$ . The inter-observer agreement and intra-observer agreement for measurements on radiographs were described previously by Specogna et al. [35]. The study was approved by the local ethical committee of Basel (EKBB EK 91/10) (Fig. 4).

#### Statistical analysis

Data were analyzed using SPSS 17.0 (SPSS, Chicago, USA). Nonparametric Spearman's correlation coefficients were used to correlate the patella height, the lateral patellar angle and leg alignment measurements with the intensity of tracer uptake in each area of interest. The values of the patella height and the lateral patellar angle measurements were also correlated categorically using chi-square statistics. The normal range for each index was set as normal, the values below and above the normal range as other

categories. The level of statistical significance was defined as  $p < 0.05$ .

#### Results

Patella height measurements were available of 84 knees and categorized in Table 1. Lateral patellar angle measurements according to Laurin were mean  $5.54 \pm 3.18^\circ$ . On long-leg radiographs, 16 (19 %) knees showed valgus, 34 (40.5 %) varus and 34 (40.5 %) neutral mechanical alignment. The mean patellar tilt angle according to Grelsamer was  $7 \pm 9^\circ$ , the mean patellar tilt angle according to Sasaki was  $15 \pm 7^\circ$  and the mean patellar tilt angle modified by Fulkerson was  $8 \pm 7^\circ$ . The mean tibial tuberosity trochlear groove distance was  $9 \pm 11$  mm.

A lower patella position correlated significantly with higher  $^{99m}\text{Tc}$ -HDP-SPECT/CT tracer uptake in all patellar and lateral femoral regions ( $p < 0.001$ ). A higher lateral patellar angle in radiographs correlated significantly with higher  $^{99m}\text{Tc}$ -HDP tracer uptake in the superior femoral parts such as F1s, F1sa, F1sp, F2sp and the tibial tubercle. In CT, a higher patellofemoral tilt angle according to Sasaki and Fulkerson correlated significantly with increased tracer uptake in the lateral superior zones (F2s  $p < 0.01$ , F2sa  $p < 0.01$ ) and the lateral superior and inferior patella (P2s, P2i). An increased TT-TG was significantly correlated with decreased tracer uptake in the tibial tuberosity and the lateral superior patellar zone (P2s  $p < 0.05$ ).


Univariate Spearman's correlations between  $^{99m}\text{Tc}$ -SPECT/CT tracer uptake in each patellofemoral region of interest and analysis of patella position are presented in Table 2.

The intensity of  $^{99m}\text{Tc}$ -HDP-SPECT/CT tracer uptake on the medial part of the patellofemoral joint significantly correlated with mechanical varus alignment of the knee ( $p < 0.05$ ). The intensity of  $^{99m}\text{Tc}$ -HDP tracer uptake on the lateral part of the patellofemoral joint significantly correlated with mechanical valgus alignment of the knee ( $p < 0.05$ ) (Table 3).

#### Discussion

This present study investigated whether patellar tilt, patellar height or mechanical alignment influence the intensity and the distribution pattern of SPECT/CT tracer uptake in the patellofemoral joint. The most important findings of this study were threefold:

Firstly, a lower patella position significantly correlated with higher  $^{99m}\text{Tc}$ -HDP-SPECT/CT tracer uptake within the patellofemoral compartment of the knee. Interestingly, a patella alta condition did not lead to an increase in



**Fig. 4** Lateral radiograph (left) and  $^{99m}\text{Tc}$ -SPECT/CT (right) of a 25-year-old patient's left knee. In this case, the Insall-Salvati index (1.15) was within the normal range (0.8–1.2). The SPECT/CT showed increased tracer uptake in the lateral patellar facet

**Table 1** The patellar height documented using the Insall-Salvati index and the modified Insall-Salvati index categorized into three classes (−1 = below the neutral range, 0 = within the neutral range, 1 = above neutral range)

	Insall-Salvati (neutral 0.8–1.2)		Modified Insall-Salvati (neutral <2)		Caton-Deschamps (neutral 0.6–1.2)	
	Frequency	Percentage	Frequency	Percentage	Frequency	Percentage
−1	9	11.3	–	–	1	1.3
0	61	76.3	77	96.3	70	87.5
1	10	12.5	3	3.8	9	11.3
Total	80	100	80	100	80	100

**Table 2** Univariate Spearman's correlations of 99mTc-SPECT/CT tracer uptake in each patellofemoral region of interest and patella measurements (\* =  $p < 0.05$ , \*\* =  $p < 0.01$ )

	Insall-Salvati >1.0	Insall-Salvati <1.0	Modified Insall-Salvati	Caton-Deschamps >0.9	Caton-Deschamps <0.9	Caton-Deschamps <0.9	Lateral patellar angle °
F1s	−0.05	0.19	−0.10	−0.07	−0.07	0.05	0.41**
F1i	−0.09	0.09	0.03	−0.09	−0.12	−0.01	−0.13
F2s	0.01	0.29**	−0.28*	−0.15	−0.12	0.17	0.22
F2i	−0.03	0.36**	−0.21	−0.12	−0.12	0.12	0.09
F1sa	−0.04	0.21	−0.09	0	−0.01	−0.02	0.37*
F1ia	−0.08	0.18	−0.05	−0.02	−0.06	−0.05	−0.01
F1sp	−0.07	0.15	−0.04	−0.14	−0.15	0.07	0.36*
F1ip	−0.15	0.05	0.05	−0.19	−0.23*	0	−0.11
F2sa	−0.02	0.21	−0.21	−0.03	−0.08	−0.04	0.21
F2ia	−0.01	0.24*	−0.17	−0.09	−0.13	0.03	−0.03
F2sp	−0.04	0.28*	−0.09	−0.14	−0.09	0.2	0.33*
F2ip	−0.02	0.29**	−0.03	−0.18	−0.15	0.21	0.24
F	−0.08	0.3**	−0.15	−0.12	−0.14	0.07	0.27
T Tub	−0.02	0.27*	0.02	−0.1	−0.09	0.08	0.35*
P1s	−0.10	0.26*	−0.18	−0.05	−0.13	−0.06	0.05
P1i	−0.08	0.31**	−0.23*	−0.07	−0.11	0	0.18
P2s	−0.11	0.23*	−0.19	−0.06	−0.13	−0.05	0.01
P2i	−0.1	0.25*	−0.27*	−0.04	−0.09	−0.04	0.17
P	−0.11	0.27*	−0.24*	−0.06	−0.13	−0.04	0.11

SPECT/CT tracer uptake within the patellofemoral joint. To date, there has not been any study showing a relationship between patellofemoral alignment and SPECT or SPECT/CT tracer uptake. Only other imaging modalities such as MRI and conventional radiographs were used to assess this relationship [21–23, 34].

Our findings are at least partially in contrast with others showing correlations between osteoarthritic changes and a patella alta [36]. Singerman et al. [34] investigated in an in vitro study the effects of patella alta and infera position on patellofemoral contact forces. They found that in a patella alta position, the magnitude of the patellofemoral contact continued to increase with increasing flexion angle [34]. A patella infera position resulted in a decrease in the patellofemoral contact force [34]. The medially directed

component of the contact force acting on the patella increased with superior displacement of the patella [34]. The resultant contact force migrated superiorly with inferior displacement of the patella [34].

It has been shown by Kalichman [21–23] that for MRI patellar alignment measurements such as Insall-Salvati ratio, sulcus angle and lateral patellar angle are related to cartilage loss and bone marrow oedema. A patella alta position was significantly associated with increasing cartilage loss in both medial and lateral compartments of the patellofemoral joint and with bone marrow oedema in the lateral compartment [21–23].

Secondly, a higher lateral patellar angle in radiographs and CT correlated significantly with higher 99mTc-HDP tracer uptake in the superior lateral femoral zones, the



**Table 3** Univariate Spearman's correlations between  $^{99m}\text{Tc}$ -SPECT/CT tracer uptake in each patellofemoral region of interest and the upper and lower half of leg alignment measurements

	Mechanical alignment $>4^\circ$	Mechanical alignment $<4^\circ$
F1 s	0.02	-0.12
F1i	-0.18	0.16
F2 s	0.14	-0.27*
F2i	0.25*	-0.29**
F1sa	0.08	-0.17
F1ia	-0.11	0.08
F1sp	0.06	-0.12
F1ip	-0.11	0.15
F2sa	0.10	-0.22*
F2ia	0.21	-0.28**
F2sp	0.16	-0.28**
F2ip	0.25*	-0.26*
F	0.08	-0.17
T Tub	0.07	-0.17
P1s	0.17	-0.20
P1i	0.25*	-0.27*
P2s	0.14	-0.17
P2i	0.21	-0.24*
P	0.21	-0.22*

Increased valgus when alignment  $>4^\circ$ , increased varus when alignment  $<4^\circ$ . (\* =  $p < 0.05$ , \*\* =  $p < 0.01$ )

lateral superior and inferior patella zones and the tibial tubercle. An increased TT-TG was significantly correlated with decreased tracer uptake in the tibial tuberosity and the lateral superior patellar zone.

Physiologically it is understood that in normally aligned knees with normally shaped patellofemoral joints, the loading is bigger on the lateral than on the medial facet. This is due to the physiological valgization of the femorotibial joint [2].

Kalichman reported that increased sulcus angle and lateral patellar tilt angle were related to increased cartilage loss and bone marrow oedema [22, 23]. Numerous studies investigated the relationship of bone marrow oedema in MRI and patellofemoral osteoarthritis or pain level [22, 23]. The question whether areas of present bone marrow oedema in MRI represent areas of increased SPECT/CT tracer uptake has not sufficiently been answered. However, Buck et al. [4] reported findings that all patients with bone marrow oedema in MRI showed abnormal bone scintigraphy and vice versa. Approximately one out of ten patients showed no bone marrow oedema but tracer uptake in bone scintigraphy. It seems that bone marrow oedema does not represent the identical pathophysiology. Interestingly, Lo et al. [29] found that the bone mineral density correlated with the occurrence of bone marrow lesions in MRI.

Thirdly, mechanical alignment significantly influenced  $^{99m}\text{Tc}$ -HDP tracer uptake of the patellofemoral compartment of the knee. Patients with more mechanical valgus alignment had significantly increased values of the SPECT/CT tracer uptake for the medial patellofemoral region. Patients with more varus mechanical alignment had significantly decreased values of the SPECT/CT tracer uptake of the medial and lateral patellar regions.

In case of patellofemoral pain, chronic instability or localized patellofemoral osteoarthritis realignment procedures such as tibial tubercle anteriorization or medialization are part of the surgeon's armamentarium [1, 24, 27]. These aim to unload areas of cartilage lesions and overloading [1, 24, 27]. One problem before and after these procedures is that one does not want to increase the load in these regions [37].

Importantly to date, there is no optimal imaging modality, which is able to visualize areas of increased loading and stress. Lorberboym et al. [30] pointed out that SPECT was 100 % sensitive and 64 % specific in detecting patellofemoral lesions when compared with arthroscopy. SPECT/CT has been increasingly recognized as promising imaging modality not only for patients with patellofemoral problems [10–15].

The knowledge established in the present study about the influence of mechanical as well as patellofemoral alignment on SPECT/CT tracer uptake distribution is of paramount importance. SPECT/CT could be used as an adjunct imaging modality in patients before and after patellofemoral surgery. In the daily clinical work, SPECT/CT could then help to determine which knee compartment is overloaded and subsequent treatment can be minimized to the affected compartment.

A number of limitations have to be considered. In our study, rotational alignment of the femur or tibia was not assessed, but could significantly influence the distribution of SPECT/CT tracer uptake. In addition, there might be a difference in SPECT/CT tracer uptake in symptomatic and asymptomatic patients. The influence of the degree of osteoarthritis present has been investigated in a previous study. Here it was shown that it is correlated, but not the decisive factor.

## Conclusion

SPECT/CT reflects the in vivo loading within the patellofemoral joint. The intensity and distribution of the SPECT/CT significantly correlated with patella infera and patellar tilt, measured in conventional radiographs. Based on these findings, SPECT/CT is promising for evaluating patients with patellofemoral disorders and for follow-up of patients after patellofemoral realignment procedures.

**Acknowledgments** We greatly thank the Deutsche Arthrose-Hilfe e.V., Frankfurt am Main, Germany, the Gottfried und Julia Bangerter-Rhyner-Stiftung, Berne, Switzerland for their support to our research.

## References

- AL-Sayyad M, Cameron JC (2002) Functional outcome after tibial tubercle transfer for the painful patella alta. *Clin Orthop Relat Res* 396:152–162
- Amis AA, Farahmand F (1996) Biomechanics masterclass: extensor mechanism of the knee. *Curr Orthop* 10(2):102–109
- Amis AA, Oguz C, Bull AM, Senavongse W, Dejour D (2008) The effect of trochleoplasty on patellar stability and kinematics: a biomechanical study in vitro. *J Bone Joint Surg Br* 90(7):864–869
- Buck FM, Hoffmann A, Hofer B, Pfirrmann CW, Allgayer B (2009) Chronic medial knee pain without history of prior trauma: correlation of pain at rest and during exercise using bone scintigraphy and MR imaging. *Skelet Radiol* 38(4):339–347
- Caton J (1989) Method of measuring the height of the patella. *Acta Orthop Belg* 55(3):385–386
- Dejour D, Saggin P (2010) The sulcus deepening trochleoplasty—the Lyon’s procedure. *Int Orthop* 34(2):311–316
- Fulkerson JP, Schutzer SF, Ramsby GR, Bernstein RA (1987) Computerized tomography of the patellofemoral joint before and after lateral release or realignment. *Arthroscopy* 3(1):19–24
- Grelsamer RP, Bazos AN, Proctor CS (1993) Radiographic analysis of patellar tilt. *J Bone Joint Surg Br* 75(5):822–824
- Grelsamer RP, Meadows S (1992) The modified Insall-Salvati ratio for assessment of patellar height. *Clin Orthop Relat Res* 282:170–176
- Hirschmann MT, Adler T, Rasch H, Hugli RW, Friederich NF, Arnold MP (2010) Painful knee joint after ACL reconstruction using biodegradable interference screws—SPECT/CT a valuable diagnostic tool? A case report. *Sports Med Arthrosc Rehabil Ther Technol* 2:24
- Hirschmann MT, Davda K, Iranpour F, Rasch H, Friederich NF (2011) Combined single photon emission computerised tomography and conventional computerised tomography (SPECT/CT) in patellofemoral disorders: a clinical review. *Int Orthop* 35(5):675–680
- Hirschmann MT, Davda K, Rasch H, Arnold MP, Friederich NF (2011) Clinical value of combined single photon emission computerized tomography and conventional computer tomography (SPECT/CT) in sports medicine. *Sports Med Arthrosc* 19(2):174–181
- Hirschmann MT, Iranpour F, Davda K, Rasch H, Hugli R, Friederich NF (2010) Combined single-photon emission computerized tomography and conventional computerized tomography (SPECT/CT): clinical value for the knee surgeons? *Knee Surg Sports Traumatol Arthrosc* 18(3):341–345
- Hirschmann MT, Iranpour F, Konala P, Kerner A, Rasch H, Cobb JP, Friederich NF (2010) A novel standardized algorithm for evaluating patients with painful total knee arthroplasty using combined single photon emission tomography and conventional computerized tomography. *Knee Surg Sports Traumatol Arthrosc* 18(7):939–944
- Hirschmann MT, Konala P, Iranpour F, Kerner A, Rasch H, Friederich NF (2011) Clinical value of SPECT/CT for evaluation of patients with painful knees after total knee arthroplasty—a new dimension of diagnostics? *BMC Musculoskelet Disord* 12(1):36
- Hirschmann MT, Mathis D, Afifi FK, Rasch H, Henckel J, Amsler F, Wagner CR, Friederich NF, Arnold MP (2013) Single photon emission computerized tomography and conventional computerized tomography (SPECT/CT) for evaluation of patients after anterior cruciate ligament reconstruction: a novel standardized algorithm combining mechanical and metabolic information. *Knee Surg Sports Traumatol Arthrosc* 21(4):965–974
- Hirschmann MT, Mathis D, Rasch H, Amsler F, Friederich NF, Arnold MP (2013) SPECT/CT tracer uptake is influenced by tunnel orientation and position of the femoral and tibial ACL graft insertion site. *Int Orthop* 37(2):301–309
- Hirschmann MT, Schon S, Afifi FK, Amsler F, Rasch H, Friederich NF, Arnold MP (2013) Assessment of loading history of compartments in the knee using bone SPECT/CT: a study combining alignment and 99mTc-HDP tracer uptake/distribution patterns. *J Orthop Res* 31(2):268–274
- Hirschmann MT, Wagner CR, Rasch H, Henckel J (2012) Standardized volumetric 3D-analysis of SPECT/CT imaging in orthopaedics: overcoming the limitations of qualitative 2D analysis. *BMC Med Imaging* 12(1):5
- Insall J, Salvati E (1971) Patella position in the normal knee joint. *Radiology* 101(1):101–104
- Kalichman L, Zhang Y, Niu J, Goggins J, Gale D, Felson DT, Hunter D (2007) The association between patellar alignment and patellofemoral joint osteoarthritis features—an MRI study. *Rheumatology (Oxford)* 46(8):1303–1308
- Kalichman L, Zhang Y, Niu J, Goggins J, Gale D, Zhu Y, Felson DT, Hunter DJ (2007) The association between patellar alignment on magnetic resonance imaging and radiographic manifestations of knee osteoarthritis. *Arthritis Res Ther* 9(2):R26
- Kalichman L, Zhu Y, Zhang Y, Niu J, Gale D, Felson DT, Hunter D (2007) The association between patella alignment and knee pain and function: an MRI study in persons with symptomatic knee osteoarthritis. *Osteoarthritis Cartilage* 15(11):1235–1240
- Karlsson J, Bunketorp O, Lansinger O, Romanus B, Sward L (1986) Lowering of the patella secondary to anterior advancement of the tibial tubercle for the patellofemoral pain syndrome. *Arch Orthop Trauma Surg* 105(1):40–45
- Kraus VB, McDaniel G, Worrell TW, Feng S, Vail TP, Varju G, Coleman RE (2009) Association of bone scintigraphic abnormalities with knee malalignment and pain. *Ann Rheum Dis* 68(11):1673–1679
- Lancourt JE, Cristini JA (1975) Patella alta and patella infera. Their etiological role in patellar dislocation, chondromalacia, and apophysitis of the tibial tubercle. *J Bone Joint Surg Am* 57(8):1112–1115
- Laprade J, Culham E (2003) Radiographic measures in subjects who are asymptomatic and subjects with patellofemoral pain syndrome. *Clin Orthop Relat Res* 414:172–182
- Laurin CA, Levesque HP, Dussault R, Labelle H, Peides JP (1978) The abnormal lateral patellofemoral angle: a diagnostic roentgenographic sign of recurrent patellar subluxation. *J Bone Joint Surg Am* 60(1):55–60
- Lo GH, Hunter DJ, Zhang Y, McLennan CE, Lavalley MP, Kiel DP, McLean RR, Genant HK, Guermazi A, Felson DT (2005) Bone marrow lesions in the knee are associated with increased local bone density. *Arthritis Rheum* 52(9):2814–2821
- Lorberboym M, Ami DB, Zin D, Nikolov G, Adar E (2003) Incremental diagnostic value of 99mTc methylene diphosphonate bone SPECT in patients with patellofemoral pain disorders. *Nucl Med Commun* 24(4):403–410
- Meyer SA, Brown TD, Pedersen DR, Albright JP (1997) Retropatellar contact stress in simulated patella infera. *Am J Knee Surg* 10(3):129–138
- Rasch H, Falkowski AL, Forrer F, Henckel J, Hirschmann MT (2013) 4D-SPECT/CT in orthopaedics: a new method of combined quantitative volumetric 3D analysis of SPECT/CT tracer uptake and component position measurements in patients after total knee arthroplasty. *Skeletal Radiol* 42(9):1215–1223
- Sasaki T, Yagi T (1986) Subluxation of the patella Investigation by computerized tomography. *Int Orthop* 10(2):115–120

34. Singerman R, Davy DT, Goldberg VM (1994) Effects of patella alta and patella infera on patellofemoral contact forces. *J Biomech* 27(8):1059–1065
35. Specogna AV, Birmingham TB, Hunt MA, Jones IC, Jenkyn TR, Fowler PJ, Giffin JR (2007) Radiographic measures of knee alignment in patients with varus gonarthrosis: effect of weight-bearing status and associations with dynamic joint load. *Am J Sports Med* 35(1):65–70
36. Stefanik JJ, Zhu Y, Zumwalt AC, Gross KD, Clancy M, Lynch JA, Frey Law LA, Lewis CE, Roemer FW, Powers CM, Guermazi A, Felson DT (2010) Association between patella alta and the prevalence and worsening of structural features of patellofemoral joint osteoarthritis: the multicenter osteoarthritis study. *Arthritis Care Res (Hoboken)* 62(9):1258–1265
37. Wirth CJ, Zichner L, Kohn D (2005) Knie. In: Kohn D (ed) *Orthopädie und Orthopädische Chirurgie*. Thieme, Stuttgart, pp 332–343