

The algebra of essential relations on a finite set

By *Serge Bouc* at Amiens and *Jacques Thévenaz* at Lausanne

Abstract. Let X be a finite set and let k be a commutative ring. We consider the k -algebra of the monoid of all relations on X , modulo the ideal generated by the relations factorizing through a set of cardinality strictly smaller than $\text{Card}(X)$, called inessential relations. This quotient is called the essential algebra associated to X . We then define a suitable nilpotent ideal of the essential algebra and describe completely the structure of the corresponding quotient, a product of matrix algebras over suitable group algebras. In particular, we obtain a description of all the simple modules for the essential algebra.

1. Introduction

Let X and Y be finite sets. A *correspondence* between X and Y is a subset R of $X \times Y$. In case $X = Y$, we say that R is a *relation* on X . Correspondences can be composed as follows. If $R \subseteq X \times Y$ and $S \subseteq Y \times Z$, then RS is the correspondence between X and Z defined by

$$RS = \{(x, z) \in X \times Z \mid \text{there exists } y \in Y \text{ such that } (x, y) \in R \text{ and } (y, z) \in S\}.$$

In particular the set of all relations on X is a monoid. Given a commutative ring k and a finite set X , let \mathcal{R} be the k -algebra of the monoid of all relations on X (having this monoid as a k -basis).

Throughout this paper, X will denote a finite set of cardinality n . We say that a relation R on X is *inessential* if there exists a set Y with $\text{Card}(Y) < \text{Card}(X)$ and two relations $S \subseteq X \times Y$ and $T \subseteq Y \times X$ such that $R = ST$. Otherwise, R is called *essential*. The set of all inessential relations on X spans a two-sided ideal I of \mathcal{R} . We define $\mathcal{E} = \mathcal{R}/I$. It is clear that \mathcal{E} is a k -algebra having as a k -basis the set of all essential relations on X . The purpose of this paper is to explore the concept of essential relation and to study the structure of \mathcal{E} .

We shall define a nilpotent ideal N of \mathcal{E} and describe completely the quotient $\mathcal{P} = \mathcal{E}/N$. More precisely, \mathcal{P} is isomorphic to a product of matrix algebras over suitable group algebras, the product being indexed by the set of all (partial) order relations on X , up to permutation. Consequently, we find all the simple \mathcal{E} -modules.

The motivation for this work comes from various sources. First, it is clear that the notion of relation on a set is of fundamental importance in mathematics. Moreover, the fact that relations can be composed leads naturally to the monoid of all relations on a finite set. This monoid has been studied by several authors: see for instance [13–15], and more recently [8]

(which contains many references), [5, 7, 12]. But the algebra \mathcal{R} still needs to be better understood. The dimension of \mathcal{R} grows extremely fast (as a function of the cardinality of the set X), so \mathcal{R} seems quite difficult to handle.

The idea of passing to the quotient by all elements obtained from something smaller is quite elementary and is a first way for decreasing the size of the algebra and simplifying its structure (although the quotient \mathcal{E} still has a very large dimension). Analogous ideas are widely used, in particular in the representation theory of finite groups. In the theory of G -algebras, the notion of Brauer quotient is of this kind (see [17]). In the more recent development of the theory of biset functors for finite groups (see [3]), the same idea plays a key role in [4]. The use of a similar idea for sets instead of groups yields the notion of essential relation, which does not seem to have been studied. It is one of the purposes of this paper to fill this gap.

The essential algebra \mathcal{E} is a very natural quotient of \mathcal{R} . It is therefore remarkable that a lot can be said about the structure of \mathcal{E} , despite its very large dimension, and that all simple \mathcal{E} -modules can be described. We view this as an interesting phenomenon which motivates further investigation.

Finally, another motivation for studying \mathcal{R} and \mathcal{E} comes from the representation theory of the category \mathcal{C} of all finite sets, where morphisms in \mathcal{C} are all correspondences between finite sets. It is known that the simple functors $\mathcal{C} \rightarrow k\text{-Mod}$ are parametrized by all pairs consisting of a finite set and a simple module for the corresponding algebra \mathcal{E} . Thus the knowledge of all simple \mathcal{E} -modules gives fundamental information on such simple functors. A deeper investigation of this representation theory will appear in a future paper.

2. Essential relations

Given a correspondence $R \subseteq X \times Y$ between a set X and a set Y , then for every $a \in X$ and $b \in Y$ we write

$${}_aR = \{y \in Y \mid (a, y) \in R\} \quad \text{and} \quad R_b = \{x \in X \mid (x, b) \in R\}.$$

We call ${}_aR$ a *column* of R and R_b a *row* of R .

We first characterize inessential relations. Any subset of $X \times Y$ of the form $U \times V$ will be called a *block* (where $U \subseteq X$ and $V \subseteq Y$).

Lemma 2.1. *Let X, Y, Z be finite sets.*

- (a) *Let $R \subseteq X \times Z$ be a correspondence between X and Z . Then R factorizes through Y if and only if R can be decomposed as a union of blocks indexed by the set Y .*
- (b) *Let R be a relation on X , where X has cardinality n . Then R is inessential if and only if R can be decomposed as a union of at most $n - 1$ blocks.*

Proof. (a) If R factorizes through Y , then $R = ST$, where $S \subseteq X \times Y$ and $T \subseteq Y \times Z$. Then we can write

$$R = \bigcup_{y \in Y} S_y \times_y T,$$

as required.

Suppose conversely that $R = \bigcup_{y \in Y} U_y \times V_y$, where each U_y is a subset of X and each V_y is a subset of Z . Then we define

$$S = \bigcup_{y \in Y} U_y \times \{y\} \subseteq X \times Y \quad \text{and} \quad T = \bigcup_{y \in Y} \{y\} \times V_y \subseteq Y \times Z.$$

Then S is a correspondence between X and Y , and $U_y = S_y$, the y -th row of S . Similarly, T is a correspondence between Y and Z , and $V_y = {}_yT$, the y -th column of T .

We now claim that $R = ST$. If $(x, z) \in R$, then there exists an element $y \in Y$ such that $(x, z) \in U_y \times V_y$. It follows that $(x, y) \in S$ and $(y, z) \in T$, hence $(x, z) \in ST$, proving that $R \subseteq ST$. If now $(x, z) \in ST$, then there exists an element $y \in Y$ such that $(x, y) \in S$ and $(y, z) \in T$. It follows that $x \in S_y = U_y$ and $z \in {}_yT = V_y$, hence $(x, z) \in U_y \times V_y \subseteq R$, proving that $ST \subseteq R$. We have shown that $R = ST$, proving the claim.

(b) This follows immediately from (a). □

Corollary 2.2. *Let R be a relation on X . If two rows of R are equal, then R is inessential. If two columns of R are equal, then R is inessential.*

Proof. Suppose that ${}_aR = {}_bR = V$. Then

$$R = (\{a, b\} \times V) \cup \left(\bigcup_{\substack{c \in X \\ c \neq a, c \neq b}} \{c\} \times {}_cR \right),$$

a union of $n - 1$ blocks, where $n = \text{Card}(X)$. The proof for rows is similar. □

Corollary 2.3. *Let R be a relation on X . If a row of R is empty, then R is inessential. If a column of R is empty, then R is inessential.*

Proof. Assume that ${}_aR = \emptyset$. Then

$$R = \bigcup_{\substack{c \in X \\ c \neq a}} \{c\} \times {}_cR,$$

a union of $n - 1$ blocks, where $n = \text{Card}(X)$. The proof for rows is similar. □

Corollary 2.4. *Let R be a relation on X . If R is an equivalence relation different from the equality relation (i.e. $R \neq \Delta$ where Δ is the diagonal of $X \times X$), then R is inessential.*

Proof. Suppose that a and b are equivalent and $a \neq b$. Then the rows R_a and R_b are equal and Corollary 2.2 applies. □

We need a few basic facts about reflexive relations. Recall that a relation S on X is reflexive if S contains $\Delta = \{(x, x) \mid x \in X\}$. Moreover, a *preorder* is a relation which is reflexive and transitive, while an *order* is a preorder which is moreover antisymmetric. (Note that, throughout this paper, the word “order” stands for “partial order”.) Associated to a preorder R , there is an equivalence relation \sim_R defined by $x \sim_R y$ if and only if $(x, y) \in R$ and $(y, x) \in R$. Then \sim_R is the equality relation if and only if R is an order.

We will often use the containment of relations on X (as subsets of $X \times X$). Note that if $R \subseteq S$, then $RT \subseteq ST$ and $TR \subseteq TS$ for any relation T on X . If S is a reflexive relation, then $S \subseteq S^2 \subseteq S^3 \subseteq \dots$ and there exists an $m \in \mathbb{N}$ such that $S^m = S^{m+1}$, hence $S^m = S^N$ for all $N \geq m$. The relation $\bar{S} := S^m$ is called the *transitive closure* of S . It is reflexive and transitive, that is, a preorder. Note that any preorder is an idempotent relation.

Proposition 2.5. *Let R be a preorder on a finite set X of cardinality n .*

- (a) *If R is not an order, then R is inessential.*
- (b) *If R is an order and if Q is a reflexive relation contained in R , then Q is essential. In particular, if R is an order, then R is essential.*
- (c) *If R is a total order, then R is maximal among essential relations.*

Proof. (a) If R is not an order, then the associated equivalence relation \sim_R is not the equality relation. Let a and b be equivalent under \sim_R with $a \neq b$. Then, by transitivity of R , the rows R_a and R_b are equal. By Corollary 2.2, R is inessential.

(b) Suppose now that R is an order and that Q is reflexive with $Q \subseteq R$. We claim that, if $a \neq b$, then (a, a) and (b, b) cannot belong to a block contained in Q . This is because, if $(a, a), (b, b) \in U \times V \subseteq Q$, then we have $(a, b) \in U \times V$ (because $a \in U$ and $b \in V$) and $(b, a) \in U \times V$ (because $b \in U$ and $a \in V$), and therefore $(a, b) \in Q$ and $(b, a) \in Q$, hence $(a, b) \in R$ and $(b, a) \in R$, contrary to antisymmetry. It follows that, in any expression of Q as a union of blocks, the diagonal elements (a, a) all lie in different blocks, so that the number of blocks is at least n . This shows that Q is essential.

(c) Without loss of generality, we can assume that the total order R is the usual total order on the set $X = \{1, 2, \dots, n\}$, i.e. $(x, y) \in R \iff x \leq y$. Let S be a relation strictly containing R . Then $S - R \neq \emptyset$ and we choose $(j, i) \in S - R$ with i maximal, and then j maximal among all x with $(x, i) \in S - R$. In other words, $(j, i) \in S$, but $j > i$ because $(j, i) \notin R$, and moreover

$$(x, y) \in S - R \implies y \leq i \quad \text{and} \quad (x, i) \in S - R \implies x \leq j.$$

If $i = j - 1$, then the rows S_i and S_j are equal, so S is inessential by Corollary 2.2.

Assume now that $j - 1 > i$. Then we claim that

$$S = (S_i \times \{i, j\}) \cup (S_{j-1} \times \{j-1, j\}) \cup \left(\bigcup_{k \neq i, j-1, j} S_k \times \{k\} \right).$$

To show that the first block is contained in S , let $x \in S_i$. Then $x \leq j$ if $(x, i) \in S - R$, and also $x \leq j$ if $(x, i) \in R$, i.e. $x \leq i$. Hence $x \leq j$ in both cases, and thus $(x, j) \in R \subset S$. This shows that $S_i \times \{j\} \subseteq S$.

To show that the second block is contained in S , let $x \in S_{j-1}$. Then $(x, j-1)$ cannot belong to $S - R$, by maximality of i (because $j-1 > i$). Thus $(x, j-1) \in R$, that is, $x \leq j-1$. Then $x < j$, hence $(x, j) \in S$. This shows that $S_{j-1} \times \{j\} \subseteq S$.

Next we show that S is contained in the union of the blocks above. This is clear for any $(x, y) \in S$ such that $y \neq j$. Now take $(x, j) \in S$. By maximality of i and since $j > i$, we have $(x, j) \in R$, that is, $x \leq j$. If $x = j$, then $(j, j) \in S_i \times \{i, j\}$ because $(j, i) \in S$, that is, $j \in S_i$. If $x < j$, then $x \leq j-1$, hence $(x, j-1) \in S$, that is, $x \in S_{j-1}$, and therefore we have $(x, j) \in S_{j-1} \times \{j-1, j\}$.

This proves the claim about the block decomposition. It follows that S is a union of $n - 1$ blocks, so S is inessential. \square

Example 2.6. Let $n = \text{Card}(X)$. Let Δ be the diagonal of the product $X \times X$ and let $R = (X \times X) - \Delta$. It is not difficult to see that R is essential if $2 \leq n \leq 4$. However, for $n \geq 5$, we prove that R is inessential. Without loss of generality, we can choose $X = \{1, \dots, n\}$. If $U \subseteq X$, we write U^c for the complement of U in X . Then it is easy to prove that

$$R = \left(\bigcup_{i=1}^{n-3} \{i, i+3\}^c \times \{i, i+3\} \right) \cup (\{n-2, n-1, n\}^c \times \{n-2, n-1, n\}) \cup (\{1, 2, 3\}^c \times \{1, 2, 3\}).$$

This is a union of $n - 1$ blocks, so R is inessential.

3. Permutations

As before, X denotes a finite set. We let Σ be the symmetric group on X , that is, the group of all permutations of X . For any $\sigma \in \Sigma$, we define

$$\Delta_\sigma = \{(\sigma(x), x) \in X \times X \mid x \in X\}.$$

This is actually the graph of the map σ^{-1} , but the choice is made so that $\Delta_\sigma \Delta_\tau = \Delta_{\sigma\tau}$ for all $\sigma, \tau \in \Sigma$. With a slight abuse, we shall often call Δ_σ a permutation. We also write $\Delta = \Delta_{\text{id}}$.

The group Σ has a left action on the set of all relations, σ acting via left multiplication by Δ_σ . Similarly, Σ also acts on the right on the set of relations. It is useful to note how multiplication by Δ_σ behaves. Given any relation R on X ,

$$(x, y) \in R \iff (\sigma(x), y) \in \Delta_\sigma R \iff (x, \sigma^{-1}(y)) \in R \Delta_\sigma.$$

Lemma 3.1. *Let R be a relation on X and let Δ_σ be a permutation.*

- (a) R is essential if and only if $\Delta_\sigma R$ is essential.
- (b) Δ_σ is essential.
- (c) The left action of Σ on the set of all essential relations is free.

Proof. (a) If R factorizes through a set of cardinality smaller than $\text{Card}(X)$, then so does $\Delta_\sigma R$. The converse follows similarly using multiplication by $\Delta_{\sigma^{-1}}$.

(b) This follows from (a) by taking $R = \Delta$ (which is essential by Proposition 2.5 because it is an order).

(c) Suppose that $\Delta_\sigma R = R$ for some $\sigma \neq \text{id}$. Then

$$(x, y) \in R \iff (\sigma(x), y) \in R,$$

hence we have ${}_x R = {}_{\sigma(x)} R$. Since $\sigma \neq \text{id}$, two columns of R are equal and so R is inessential by Corollary 2.2. Thus if R is essential, $\Delta_\sigma R \neq R$ for all $\sigma \neq \text{id}$. \square

Our next result will be essential in our analysis of essential relations.

Theorem 3.2. *Any essential relation contains a permutation.*

We shall provide two different proofs. The first is direct, while the second uses a theorem of Philip Hall. In fact, a relation containing a permutation is called a Hall relation in a paper of Schwarz [15], because of Hall's theorem, so Theorem 3.2 asserts that any essential relation is a Hall relation.

First proof. Let R be a relation on X and let $n = \text{Card}(X)$. We have to prove that, if R is essential, then there exists a $\sigma \in \Sigma$ such that R contains $\Delta_{\sigma^{-1}}$, that is, $R\Delta_\sigma$ contains Δ (or in other words $R\Delta_\sigma$ is reflexive). Let $D_\sigma = R\Delta_\sigma \cap \Delta$ and suppose that $\text{Card}(D_\sigma) < n$ for all $\sigma \in \Sigma$. Then we have to prove that R is inessential.

We choose $\tau \in \Sigma$ such that $\text{Card}(D_\sigma) \leq \text{Card}(D_\tau)$ for all $\sigma \in \Sigma$. We let $S = R\Delta_\tau$ and we aim to prove that S is inessential (hence R too by Lemma 3.1). Note that $D_\tau \subseteq S$ by construction. Define

$$A = \{a \in X \mid (a, a) \in D_\tau\}, \quad \text{in other words} \quad D_\tau = \{(a, a) \mid a \in A\}.$$

In particular $\text{Card}(A) = \text{Card}(D_\tau) < n$. By maximality of D_τ , we have the following property:

$$(*) \quad \text{Card}(S\Delta_\sigma \cap \Delta) \leq \text{Card}(A) \quad \text{for all } \sigma \in \Sigma.$$

Given $x, y \in X$, define a *path* from x to y to be a sequence x_0, x_1, \dots, x_r of elements of X such that $x_0 = x$, $x_r = y$, and $(x_i, x_{i+1}) \in S$ for all $i = 0, \dots, r-1$. We write $x \rightsquigarrow y$ to indicate that there is a path from x to y , and also $x \rightarrow y$ whenever $(x, y) \in S$ (path of length 1). Define A^c to be the complement of A in X (so A^c is nonempty by assumption). Define also

$$\begin{aligned} A_1 &= \{a \in A \mid \text{there exists } z \in A^c \text{ and a path } z \rightsquigarrow a\}, \\ A_2 &= \{a \in A \mid \text{there exists } z \in A^c \text{ and a path } a \rightsquigarrow z\}. \end{aligned}$$

We claim that there is no path from an element of A_1 to an element of A_2 . Suppose by contradiction that there is a path $a_1 \rightsquigarrow a_2$ with $a_i \in A_i$. Then there exist elements $z_i \in A^c$ and paths $z_1 \rightsquigarrow a_1 \rightsquigarrow a_2 \rightsquigarrow z_2$, in particular $z_1 \rightsquigarrow a \rightsquigarrow z_2$ with $a \in A$. In the path $z_1 \rightsquigarrow a$, let w_1 be the element of A^c closest to a , so that the path $w_1 \rightsquigarrow a$ does not contain any element of A^c except w_1 . Similarly, let w_2 be the element of A^c closest to a in the path $a \rightsquigarrow z_2$, so that the path $a \rightsquigarrow w_2$ does not contain any element of A^c except w_2 . We obtain a path $w_1 \rightsquigarrow a \rightsquigarrow w_2$ having all its elements in A except the two extremities w_1 and w_2 . By suppressing cycles within A , we can assume that all elements of A in this path are distinct. We end up with a path

$$w_1 \rightarrow x_1 \rightarrow \dots \rightarrow x_r \rightarrow w_2$$

where $x_1, \dots, x_r \in A$ are all distinct.

Let $\sigma \in \Sigma$ be the cycle defined as follows: $\sigma(w_1) = x_1$, $\sigma(x_i) = x_{i+1}$ for $1 \leq i \leq r-1$, $\sigma(x_r) = w_2$, and finally $\sigma(w_2) = w_1$ in case $w_2 \neq w_1$. In case $w_2 = w_1$, then $\sigma(w_2)$ is already defined to be $\sigma(w_2) = \sigma(w_1) = x_1$. We emphasize that $\sigma(y) = y$ for all the other elements $y \in X$. Then we obtain

$$\begin{aligned} (w_1, x_1) \in S & \quad \text{hence} \quad (w_1, w_1) \in S\Delta_\sigma, \\ (x_i, x_{i+1}) \in S & \quad \text{hence} \quad (x_i, x_i) \in S\Delta_\sigma, \\ (x_r, w_2) \in S & \quad \text{hence} \quad (x_r, x_r) \in S\Delta_\sigma, \\ (y, y) \in S & \quad \text{hence} \quad (y, y) \in S\Delta_\sigma \quad \text{for all } y \in A - \{x_1, \dots, x_r\}. \end{aligned}$$

Thus we obtain $(a, a) \in S\Delta_\sigma$ for all $a \in A$, but also $(w_1, w_1) \in S\Delta_\sigma$. Therefore

$$\text{Card}(S\Delta_\sigma \cap \Delta) > \text{Card}(A),$$

contrary to property (*). This proves the claim that there is no path from A_1 to A_2 .

In particular, $A_1 \cap A_2 = \emptyset$ because if $a \in A_1 \cap A_2$, we would have a path of length 0 from A_1 to A_2 (since $(a, a) \in S$). Let A_3 be the complement of $A_1 \cup A_2$ in A . Thus X is the disjoint union of the four subsets A^c, A_1, A_2 and A_3 .

We now claim the following:

- (a) There is no relation between A^c and A^c , that is, $S \cap (A^c \times A^c) = \emptyset$.
- (b) There is no relation between A^c and $A_2 \cup A_3$, that is, we have $S \cap (A^c \times A_2) = \emptyset$ and $S \cap (A^c \times A_3) = \emptyset$.
- (c) There is no relation between A_1 and A^c , that is, $S \cap (A_1 \times A^c) = \emptyset$.
- (d) There is no relation between A_1 and A_2 , that is, $S \cap (A_1 \times A_2) = \emptyset$.
- (e) There is no relation between A_1 and A_3 , that is, $S \cap (A_1 \times A_3) = \emptyset$.

To prove (a), suppose that $(w, z) \in S$ where $w, z \in A^c$. Choose $\rho \in \Sigma$ such that $\rho(a) = a$ for all $a \in A$ and $\rho(w) = z$. Then $(a, a) \in S\Delta_\rho$ for all $a \in A$ but also $(w, w) \in S\Delta_\rho$, contrary to property (*).

To prove (b), we note that the definition of A_1 implies that, if $(z, a) \in S$ with $z \in A^c$ and $a \in A$, then $a \in A_1$. Thus $a \notin A_2 \cup A_3$.

To prove (c), suppose that $(a, z) \in S$ with $a \in A$ and $z \in A^c$. Then $a \in A_2$ by the definition of A_2 and in particular $a \notin A_1$.

Property (d) follows immediately from the previous claim that there is no path from A_1 to A_2 .

To prove (e), suppose that $(a_1, a_3) \in S$ with $a_1 \in A_1$ and $a_3 \in A_3$. Then by the definition of A_1 , there is a path $z \rightsquigarrow a_1 \rightarrow a_3$ where $z \in A^c$, but this means that $a_3 \in A_1$, a contradiction.

It follows that the relation S has the property that there is no relation between $A^c \cup A_1$ and $A^c \cup A_2 \cup A_3$. Therefore S is the union of the columns indexed by $(A^c \cup A_1)^c = A_2 \cup A_3$ and the lines indexed by $(A^c \cup A_2 \cup A_3)^c = A_1$, that is,

$$S = \left(\bigcup_{b \in A_2 \cup A_3} bS \right) \cup \left(\bigcup_{a \in A_1} S_a \right).$$

Since $\text{Card}(A_2 \cup A_3) + \text{Card}(A_1) = \text{Card}(A)$, we obtain a union of $\text{Card}(A)$ blocks. But we have $\text{Card}(A) < n$ by assumption, so S is inessential, as was to be shown. \square

Second proof. Let R be an essential relation on X . For any subset A of X , define

$$R_A = \{x \in X \mid \text{there exists } a \in A \text{ such that } (x, a) \in R\} = \bigcup_{a \in A} R_a.$$

Then R decomposes as a union of blocks

$$R = \left(\bigcup_{y \notin A} (R_y \times \{y\}) \right) \cup \left(\bigcup_{x \in R_A} (\{x\} \times {}_xR) \right).$$

Since R is essential, $\text{Card}(X - A) + \text{Card}(R_A)$ cannot be strictly smaller than $\text{Card}(X)$. Therefore $\text{Card}(R_A) \geq \text{Card}(A)$ for all subsets A of X , that is,

$$\text{Card}\left(\bigcup_{a \in A} R_a\right) \geq \text{Card}(A).$$

This is precisely the assumption in a theorem of Philip Hall (see [9, Theorem 5.1.1], or [10] for the original version which is slightly different). The conclusion is that there exist elements $x_y \in R_y$, where y runs over X , which are all distinct. In other words $\sigma : y \mapsto x_y$ is a permutation and

$$(\sigma(y), y) = (x_y, y) \in R \quad \text{for all } y \in X.$$

This means that R contains Δ_σ , as required. \square

Corollary 3.3. *Let R be an essential relation on X . Then there exists an $m \in \mathbb{N}$ such that R^m is a preorder.*

Proof. By Theorem 3.2, R contains Δ_σ for some $\sigma \in \Sigma$. If σ has order k in the group Σ , then R^k contains $\Delta_{\sigma^k} = \Delta$, so R^k is reflexive. Then the transitive closure of R^k is some power R^{kt} . This is reflexive and transitive, that is, a preorder. \square

We know that any order is an essential relation (Proposition 2.5), hence contains a permutation (Theorem 3.2). But in fact, we have a more precise result.

Lemma 3.4. *If R is an order on X , then R contains a unique permutation, namely Δ .*

Proof. Suppose that R is reflexive and transitive and contains a nontrivial permutation Δ_σ . Then σ contains a nontrivial k -cycle, say on x_1, \dots, x_k , for some $k \geq 2$. It follows that $(x_{i+1}, x_i) \in R$ for $1 \leq i \leq k-1$, hence $(x_k, x_1) \in R$ by transitivity of R . Now we also have $(x_1, x_k) \in R$ because $\sigma(x_k) = x_1$. Thus the relation R is not antisymmetric, hence cannot be an order. \square

In the same vein, we have the following more general result.

Lemma 3.5. *Let R be an order and let S, S' be two relations on X . The following two conditions are equivalent:*

- (a) $\Delta \subseteq S'S \subseteq R$.
- (b) *There exists a permutation Δ_σ such that $\Delta \subseteq \Delta_{\sigma^{-1}}S \subseteq R$ and $\Delta \subseteq S'\Delta_\sigma \subseteq R$.*

Moreover, in condition (b), the permutation σ is unique.

Proof. If (b) holds, then

$$\Delta = \Delta^2 \subseteq (S'\Delta_\sigma)(\Delta_{\sigma^{-1}}S) = S'S \subseteq R^2 = R,$$

so (a) holds.

If (a) holds, then $S'S$ is essential, by Proposition 2.5. It follows that S is essential, and therefore S contains a permutation Δ_σ , by Theorem 3.2. Then we obtain

$$\Delta \subseteq \Delta_{\sigma^{-1}}S \quad \text{and} \quad S'\Delta_\sigma \subseteq S'S \subseteq R.$$

Similarly, S' is essential, hence contains a permutation Δ_τ , and we obtain

$$\Delta \subseteq S' \Delta_{\tau^{-1}} \quad \text{and} \quad \Delta_\tau S \subseteq S' S \subseteq R.$$

Now R contains $\Delta_\tau \Delta_\sigma = \Delta_{\tau\sigma}$ and Lemma 3.4 implies that $\tau\sigma = \text{id}$, that is, we have $\tau = \sigma^{-1}$. Then (b) follows.

If moreover S contains a permutation Δ_ρ (that is, $\Delta \subseteq \Delta_{\rho^{-1}} S$), then

$$\Delta_{\sigma^{-1}\rho} = \Delta_{\sigma^{-1}} \Delta_\rho \subseteq S' S \subseteq R,$$

and so $\sigma^{-1}\rho = \text{id}$ by Lemma 3.4, proving the uniqueness of σ . □

4. The essential algebra

Let X be a finite set and let k be a commutative ring. We shall be mainly interested in the cases where k is either the ring \mathbb{Z} of integers or a field, but it is convenient to work with an arbitrary commutative ring.

Let \mathcal{R} be the k -algebra of the monoid of all relations on X . This monoid is a k -basis of \mathcal{R} and the product in the monoid defines the algebra structure. The set of all inessential relations on X spans a two-sided ideal I of \mathcal{R} . We define $\mathcal{E} = \mathcal{R}/I$ and call it the *essential algebra*. It is clear that \mathcal{E} is a k -algebra having as a k -basis the set of all essential relations on X . Moreover, if R and S are essential relations but RS is inessential, then $RS = 0$ in \mathcal{E} .

Both \mathcal{R} and \mathcal{E} have an anti-automorphism, defined on the basis elements by $R \mapsto R^{\text{op}}$, where $(x, y) \in R^{\text{op}}$ if and only if $(y, x) \in R$. It is easy to see that $(RS)^{\text{op}} = S^{\text{op}} R^{\text{op}}$.

As before, we let Σ be the symmetric group of all permutations of X . We first describe an obvious quotient of \mathcal{E} .

Lemma 4.1. *Let H be the k -submodule of the essential algebra \mathcal{E} spanned by the set of all essential relations which strictly contain a permutation. Then H is a two-sided ideal of \mathcal{E} and $\mathcal{E}/H \cong k\Sigma$, the group algebra of the symmetric group Σ .*

Proof. Let us write \subset for the strict containment relation. Let R be an essential relation such that $\Delta_\sigma \subset R$ and let S be any essential relation. Then S contains a permutation Δ_τ , by Theorem 3.2. We obtain $\Delta_{\sigma\tau} = \Delta_\sigma \Delta_\tau \subset R \Delta_\tau \subseteq RS$, showing that $RS \in H$. Similarly we find that $SR \in H$ and therefore H is a two-sided ideal of \mathcal{E} .

The quotient \mathcal{E}/H has a k -basis consisting of all the permutations Δ_σ , for $\sigma \in \Sigma$. Moreover, they multiply in the same way as permutations, so \mathcal{E}/H is isomorphic to the group algebra of the symmetric group Σ . □

If k is a field, it follows, not surprisingly, that every irreducible representation of the symmetric group Σ gives rise to a simple \mathcal{E} -module. In short, the representation theory of the symmetric group Σ is part of the representation theory of \mathcal{E} .

We now want to describe another \mathcal{E} -module, which is simple when k is a field. We fix a total order T on X (e.g. the usual total order on $X = \{1, \dots, n\}$). Then any other total order on X is obtained by permuting the elements of X . Since permuting via σ corresponds to conjugation by Δ_σ , we see that $\{T_\sigma := \Delta_\sigma T \Delta_{\sigma^{-1}} \mid \sigma \in \Sigma\}$ is the set of all total orders on X . All of them are maximal essential relations on X , by Proposition 2.5.

Lemma 4.2. *Let T be a total order on X .*

- (a) *If $\rho \in \Sigma$, then $T\Delta_\rho T = 0$ in \mathcal{E} if $\rho \neq \text{id}$ and otherwise $T\Delta T = T^2 = T$.*
 (b) *The set $\{T_\sigma := \Delta_\sigma T \Delta_{\sigma^{-1}} \mid \sigma \in \Sigma\}$ is a set of pairwise orthogonal idempotents of \mathcal{E} .*

Proof. (a) Observe that TT_ρ contains both T and T_ρ (because both T and T_ρ contain Δ). Since $T \neq T_\rho$ if $\rho \neq \text{id}$, the product TT_ρ contains strictly T and is therefore inessential by Proposition 2.5. Thus $T\Delta_\rho T$ is also inessential, that is, $T\Delta_\rho T = 0$. On the other hand $T^2 = T$ because any preorder is idempotent.

(b) It follows from (a) that

$$T_\sigma T_\tau = \Delta_\sigma T \Delta_{\sigma^{-1}\tau} T \Delta_{\tau^{-1}} = \begin{cases} 0 & \text{if } \sigma \neq \tau, \\ \Delta_\sigma T \Delta_{\sigma^{-1}} = T_\sigma & \text{if } \sigma = \tau, \end{cases}$$

as was to be shown. □

Proposition 4.3. *Fix a total order T on X . Let L be the k -submodule of the essential algebra \mathcal{E} spanned by the set $\{\Delta_\sigma T \mid \sigma \in \Sigma\}$. Then L is a left ideal of \mathcal{E} and is free of rank $n!$ as a k -module, where $n = \text{Card}(X)$. If k is a field, then L is a simple \mathcal{E} -module of dimension $n!$.*

Proof. Write $S_\sigma = \Delta_\sigma T$ for all $\sigma \in \Sigma$. Let R be an essential relation on X . Then we have $\Delta_\tau \subseteq R$ for some $\tau \in \Sigma$ by Theorem 3.2. Therefore $\Delta \subseteq \Delta_{\tau^{-1}} R$ and this implies that

$$S_\sigma = \Delta S_\sigma \subseteq \Delta_{\tau^{-1}} R S_\sigma$$

and

$$T = \Delta_{\sigma^{-1}} S_\sigma \subseteq \Delta_{\sigma^{-1}} \Delta_{\tau^{-1}} R S_\sigma.$$

If this containment is strict, then $\Delta_{\sigma^{-1}} \Delta_{\tau^{-1}} R S_\sigma$ is inessential (by Proposition 2.5) and so $R S_\sigma$ is inessential too (by Lemma 3.1). Otherwise

$$S_\sigma = \Delta_{\tau^{-1}} R S_\sigma,$$

hence

$$R S_\sigma = \Delta_\tau S_\sigma = \Delta_{\tau\sigma} T = S_{\tau\sigma}.$$

Therefore, in the algebra \mathcal{E} , either $R S_\sigma = 0$ or $R S_\sigma = S_{\tau\sigma}$. This proves that L is a left ideal of \mathcal{E} . Clearly L has rank $n!$ with basis $\{S_\sigma \mid \sigma \in \Sigma\}$. The action of \mathcal{E} on L induces a k -algebra map

$$\phi : \mathcal{E} \rightarrow M_{n!}(k)$$

and L can be viewed as an $M_{n!}(k)$ -module (consisting of column vectors with entries in k). By Lemma 4.2, the action of $\Delta_\tau T \Delta_{\rho^{-1}}$ on basis elements is given by

$$(\Delta_\tau T \Delta_{\rho^{-1}}) \cdot S_\sigma = (\Delta_\tau T \Delta_{\rho^{-1}}) \cdot \Delta_\sigma T = \begin{cases} 0 & \text{if } \rho \neq \sigma, \\ S_\tau & \text{if } \rho = \sigma. \end{cases}$$

This means that $\phi(\Delta_\tau T \Delta_{\rho^{-1}})$ is the elementary matrix with a single nonzero entry 1 in position (τ, ρ) . Therefore the map ϕ is surjective. This implies that, if k is a field, the module L is simple as an \mathcal{E} -module, because the space of column vectors is a simple $M_{n!}(k)$ -module. □

5. A nilpotent ideal

The purpose of this section is to construct a suitable nilpotent ideal N of the essential algebra \mathcal{E} . We shall later pass to the quotient by N and describe the quotient \mathcal{E}/N . In order to find nilpotent ideals, the following well-known result is often useful.

Lemma 5.1. *Let k be a commutative ring and let \mathcal{B} be a k -algebra which is finitely generated as a k -module. Let I be a two-sided ideal of \mathcal{B} which is k -linearly spanned by a set of nilpotent elements of \mathcal{B} .*

- (a) *If k is a field, then I is a nilpotent ideal of \mathcal{B} .*
- (b) *If $k = \mathbb{Z}$ and if \mathcal{B} is a finitely generated free \mathbb{Z} -module, then I is a nilpotent ideal of \mathcal{B} .*
- (c) *Suppose that \mathcal{B} is defined over \mathbb{Z} , that is, $\mathcal{B} \cong k \otimes_{\mathbb{Z}} \mathcal{B}_{\mathbb{Z}}$ for some \mathbb{Z} -algebra $\mathcal{B}_{\mathbb{Z}}$ which is finitely generated free as a \mathbb{Z} -module. Suppose that I is defined over \mathbb{Z} , that is, $I \cong k \otimes_{\mathbb{Z}} I_{\mathbb{Z}}$, where $I_{\mathbb{Z}}$ is a two-sided ideal of $\mathcal{B}_{\mathbb{Z}}$ which is \mathbb{Z} -linearly spanned by a set of nilpotent elements of $\mathcal{B}_{\mathbb{Z}}$. Then I is a nilpotent ideal of \mathcal{B} .*

Proof. (a) The assumption still holds after extending scalars to an algebraic closure of k . Therefore we can assume that k is algebraically closed. Let $J(\mathcal{B})$ be the Jacobson radical of \mathcal{B} . Then $J(\mathcal{B}) = \bigcap_{i=1}^r M_i$, where M_i is a maximal two-sided ideal of \mathcal{B} . Moreover, by Wedderburn's theorem, \mathcal{B}/M_i is isomorphic to a matrix algebra $M_{n_i}(k)$, because k is algebraically closed. We will show that $I \subseteq M_i$ for all $i = 1, \dots, r$. It then follows that $I \subseteq J(\mathcal{B})$, so I is nilpotent (because it is well known that the Jacobson radical of a finite-dimensional k -algebra is nilpotent).

Let \bar{I} be the image of I in \mathcal{B}/M_i . Then \bar{I} is spanned by nilpotent elements of $M_{n_i}(k)$. But any nilpotent matrix has trace zero (because its characteristic polynomial is X^{n_i} and the coefficient of X^{n_i-1} is the trace, up to sign). It follows that \bar{I} is contained in $\text{Ker}(\text{tr})$, which is a proper subspace of $M_{n_i}(k)$. Now \bar{I} is a two-sided proper ideal of the simple algebra $M_{n_i}(k)$, hence $\bar{I} = \{0\}$, proving that $I \subseteq M_i$.

(b) Let F be a basis of \mathcal{B} as a \mathbb{Z} -module. Extending scalars to \mathbb{Q} , we see that F is a \mathbb{Q} -basis of the \mathbb{Q} -algebra $\mathbb{Q} \otimes_{\mathbb{Z}} \mathcal{B}$ and \mathcal{B} embeds in $\mathbb{Q} \otimes_{\mathbb{Z}} \mathcal{B}$. By part (a), the ideal $\mathbb{Q} \otimes_{\mathbb{Z}} I$ is nilpotent in $\mathbb{Q} \otimes_{\mathbb{Z}} \mathcal{B}$. Since I embeds in $\mathbb{Q} \otimes_{\mathbb{Z}} I$, it follows that I is nilpotent.

(c) By part (b), $I_{\mathbb{Z}}$ is a nilpotent ideal of $\mathcal{B}_{\mathbb{Z}}$. Extending scalars to k , we see that I is a nilpotent ideal of \mathcal{B} . □

Recall that Σ denotes the symmetric group on X and that, if R is a reflexive relation, then \bar{R} denotes the transitive closure of R .

Lemma 5.2. *If $S = \Delta_{\tau^{-1}} R \Delta_{\tau}$ where $\tau \in \Sigma$ and R is a reflexive relation, then*

$$\bar{S} = \Delta_{\tau^{-1}} \bar{R} \Delta_{\tau}.$$

Proof. We have $\bar{R} = R^m$ for some m and we obtain

$$S^m = (\Delta_{\tau^{-1}} R \Delta_{\tau})^m = \Delta_{\tau^{-1}} R^m \Delta_{\tau} = \Delta_{\tau^{-1}} \bar{R} \Delta_{\tau}.$$

Therefore S^m is a preorder, because it is conjugate to a preorder, and so $\bar{S} = S^m = \Delta_{\tau^{-1}} \bar{R} \Delta_{\tau}$, as claimed. □

Theorem 5.3. *Let \mathcal{F} be the set of all reflexive essential relations on X . Let N be the k -submodule of the essential algebra \mathcal{E} generated by all elements of the form $(S - \overline{S})\Delta_\sigma$ with $S \in \mathcal{F}$ and $\sigma \in \Sigma$ (where \overline{S} denotes the transitive closure of S).*

- (a) *N is a nilpotent two-sided ideal of \mathcal{E} . In particular, N is contained in the Jacobson radical $J(\mathcal{E})$.*
- (b) *The quotient algebra $\mathcal{P} = \mathcal{E}/N$ has a k -basis consisting of all elements of the form $S\Delta_\sigma$, where S runs over the set of all orders on X and σ runs over the symmetric group Σ .*

Proof. (a) Let \mathcal{E}_1 be the subalgebra of \mathcal{E} which is k -linearly generated by the set \mathcal{F} of all reflexive essential relations. It is clearly a subalgebra since the product of two reflexive relations is reflexive. Let N_1 be the k -submodule of \mathcal{E}_1 generated by all elements of the form $S - \overline{S}$ with $S \in \mathcal{F}$. We claim that N_1 is a two-sided ideal of \mathcal{E}_1 .

If $T \in \mathcal{F}$, then $T\overline{S} = \overline{TS}$ (because \overline{TS} contains both T and \overline{S} , hence \overline{TS} , and \overline{TS} contains both T and S , hence \overline{TS}). Therefore

$$T(S - \overline{S}) = (TS - \overline{TS}) - (T\overline{S} - \overline{TS}) = (TS - \overline{TS}) - (T\overline{S} - \overline{TS}).$$

Note that if TS is inessential (hence zero in \mathcal{E}_1), then its transitive closure \overline{TS} cannot be an order by Proposition 2.5 and is therefore also zero in \mathcal{E}_1 (again by Proposition 2.5). Thus, in the expression above, we obtain either generators of N_1 or zero. The same argument works for right multiplication by T (or use the anti-automorphism of \mathcal{E}) and this proves the claim.

The ideal N_1 is invariant under conjugation by Σ because, for every $\sigma \in \Sigma$,

$$\Delta_{\sigma^{-1}}(S - \overline{S})\Delta_\sigma = \Delta_{\sigma^{-1}}S\Delta_\sigma - \Delta_{\sigma^{-1}}\overline{S}\Delta_\sigma = \Delta_{\sigma^{-1}}S\Delta_\sigma - \overline{\Delta_{\sigma^{-1}}S\Delta_\sigma}$$

by Lemma 5.2. Therefore the generators of N can also be written $\Delta_\sigma(S' - \overline{S'})$ with $S' \in \mathcal{F}$ and $\sigma \in \Sigma$ (namely $S' = \Delta_{\sigma^{-1}}S\Delta_\sigma$). It follows that $N = N_1\Delta_\Sigma = \Delta_\Sigma N_1$, where we write for simplicity $\Delta_\Sigma = \{\Delta_\sigma \mid \sigma \in \Sigma\}$.

If R is an essential relation on X , then R contains a permutation Δ_σ (for some $\sigma \in \Sigma$) by Theorem 3.2, so $R = Q\Delta_\sigma$ with $Q \in \mathcal{F}$, and also $R = \Delta_\sigma Q'$ where $Q' = \Delta_{\sigma^{-1}}Q\Delta_\sigma$. Since N_1 is an ideal of \mathcal{E}_1 , it follows that N is invariant by right and left multiplication by R . Thus N is a two-sided ideal of \mathcal{E} .

The generators of N_1 are nilpotent, because if $\overline{S} = S^m$, then

$$\begin{aligned} (S - \overline{S})^m &= (S - S^m)^m = \sum_{j=0}^m \binom{m}{j} (-1)^j S^{m-j} S^{mj} \\ &= \left(\sum_{j=0}^m \binom{m}{j} (-1)^j \right) S^m = (1 - 1)^m S^m = 0. \end{aligned}$$

Thus N_1 is a nilpotent ideal of \mathcal{E}_1 , by Lemma 5.1 (because clearly \mathcal{E}_1 and N_1 are defined over \mathbb{Z}). Since N_1 is invariant under conjugation by Σ , we obtain $N^n = (N_1\Delta_\Sigma)^n = N_1^n\Delta_\Sigma$ for every $n \in \mathbb{N}$. Since $N_1^m = 0$ for some m , the ideal N is nilpotent.

(b) In the quotient algebra \mathcal{E}/N , any reflexive relation Q is identified with its transitive closure \overline{Q} . Moreover, by Theorem 3.2, any essential relation R on X can be written $R = Q\Delta_\sigma$, with Q reflexive, and $Q\Delta_\sigma$ is identified with $\overline{Q}\Delta_\sigma$ in the quotient algebra \mathcal{E}/N . Note that \overline{Q} is a preorder and that \overline{Q} is zero in \mathcal{E} if it is not an order, by Proposition 2.5.

On each basis element R of \mathcal{E} , the effect of passing to the quotient by N consists of just two possibilities.

- If R can be written $R = Q\Delta_\sigma$, with Q reflexive and \overline{Q} is not an order, then R is identified with $\overline{Q}\Delta_\sigma$, so R is zero in \mathcal{E}/N because \overline{Q} is zero.
- If R can be written $R = Q\Delta_\sigma$, with Q reflexive and \overline{Q} is an order, then R is identified with an element of the form $S\Delta_\sigma$ where S is an order (namely $S = \overline{Q}$).

In the second case, Δ_σ is the unique permutation contained in R (or in other words the expression $R = Q\Delta_\sigma$ is the unique decomposition of R as a product of a reflexive relation and a permutation). This is because if $\Delta_{\sigma'} \subseteq R$, we obtain

$$\Delta_{\sigma'\sigma^{-1}} = \Delta_{\sigma'}\Delta_{\sigma^{-1}} \subseteq R\Delta_{\sigma^{-1}} = Q \subseteq \overline{Q},$$

so that $\sigma'\sigma^{-1} = \text{id}$ since \overline{Q} is an order (Lemma 3.4). Thus $\sigma' = \sigma$. This uniqueness property shows, on the one hand, that both possibilities cannot occur simultaneously and, on the other hand, that in the second case the order $S = \overline{Q}$ is uniquely determined by R .

It follows that the nonzero images in \mathcal{E}/N of the basis elements of \mathcal{E} form a k -basis of \mathcal{E}/N consisting of (the images of) the elements $S\Delta_\sigma$, where S is an order and $\sigma \in \Sigma$. \square

The quotient algebra $\mathcal{P} = \mathcal{E}/N$ will be called the *algebra of permuted orders* on X , because every basis element $S\Delta_\sigma$ is obtained from the order S by applying a permutation σ to the rows of S . Moreover, Δ_σ is the unique permutation contained in $S\Delta_\sigma$, because Δ is the unique permutation contained in S by Lemma 3.4. This defines a Σ -grading on \mathcal{P} :

$$\mathcal{P} = \bigoplus_{\sigma \in \Sigma} \mathcal{P}_\sigma,$$

where \mathcal{P}_σ is spanned by the set of all permuted orders containing Δ_σ . Clearly $\mathcal{P}_\sigma \cdot \mathcal{P}_\tau = \mathcal{P}_{\sigma\tau}$, so we have indeed a Σ -grading. We also write $\mathcal{P}_1 := \mathcal{P}_{\text{id}}$ and call it the *algebra of orders* on X . Moreover,

$$\mathcal{P}_\sigma = \Delta_\sigma \mathcal{P}_1 = \mathcal{P}_1 \Delta_\sigma,$$

so that the product in \mathcal{P} is completely determined by the product in the subalgebra \mathcal{P}_1 , the product in the symmetric group Σ , and the action of Σ on \mathcal{P}_1 . Hence we first need to understand the subalgebra \mathcal{P}_1 .

6. The algebra of orders

Let \mathcal{P}_1 be the algebra of orders on X defined above. It has a k -basis \mathcal{O} consisting of all orders on X . The product of basis elements $R, S \in \mathcal{O}$ will be written $R \cdot S$ and is described as follows.

Lemma 6.1. *Let \cdot be the product in the k -algebra \mathcal{P}_1 .*

- Let $R, S \in \mathcal{O}$. Then the product $R \cdot S$ is equal to the transitive closure of $R \cup S$ if this closure is an order, and zero otherwise.*
- The product \cdot is commutative.*

Proof. (a) By definition of the ideal N , the product RS in the algebra $\mathcal{P} = \mathcal{E}/N$ is identified with the transitive closure \overline{RS} , which is also the transitive closure of $R \cup S$, because the inclusions

$$R \cup S \subseteq RS \subseteq (R \cup S)^2 \subseteq \overline{R \cup S} \subseteq \overline{RS}$$

force the equality $\overline{R \cup S} = \overline{RS}$. Now \overline{RS} is a preorder. If this is an order, then $R \cdot S = \overline{RS}$. If this preorder is not an order, then it is zero in \mathcal{E} (by Proposition 2.5), hence also zero in \mathcal{P}_1 .

(b) This follows from (a) and the fact that $R \cup S = S \cup R$. \square

Theorem 6.2. *The following statements hold.*

- (a) *There exists a k -basis $\{f_R \mid R \in \mathcal{O}\}$ of \mathcal{P}_1 consisting of mutually orthogonal idempotents whose sum is 1 and such that, for every $R \in \mathcal{O}$, the ideal generated by f_R is free of rank one as a k -module.*
- (b) *\mathcal{P}_1 is isomorphic to a product of copies of k , indexed by \mathcal{O} ,*

$$\mathcal{P}_1 \cong \prod_{R \in \mathcal{O}} k \cdot f_R.$$

Proof. We know that \mathcal{P}_1 is commutative, with a basis \mathcal{O} consisting of all orders on X . Any such basis element is idempotent. Moreover \mathcal{O} is a partially ordered set with respect to the containment relation and we make it a lattice by adding an element ∞ and defining $R \vee S = \infty$ whenever the transitive closure of $R \cup S$ is not an order, while $R \vee S$ is the transitive closure of $R \cup S$ otherwise. The greatest lower bound of R and S is just the intersection $R \cap S$.

Now define $g_R = R$ if $R \in \mathcal{O}$ and $g_\infty = 0$. By Lemma 6.1, these elements satisfy the condition $g_R \cdot g_S = g_{R \vee S}$. Therefore Theorem A.1 of the appendix applies. We let

$$f_R = \sum_{\substack{S \in \mathcal{O} \\ R \subseteq S}} \mu(R, S) S,$$

where μ denotes the Möbius function of the poset \mathcal{O} , so by Möbius inversion, we have

$$R = \sum_{\substack{S \in \mathcal{O} \\ R \subseteq S}} f_S.$$

The transition matrix from $\{R \in \mathcal{O}\}$ to $\{f_R \mid R \in \mathcal{O}\}$ is upper-triangular, with 1 along the main diagonal, hence invertible over \mathbb{Z} . It follows that $\{f_R \mid R \in \mathcal{O}\}$ is a k -basis of \mathcal{P}_1 . By Theorem A.1 of the appendix, $\{f_R \mid R \in \mathcal{O}\}$ is a set of mutually orthogonal idempotents in \mathcal{P}_1 whose sum is 1. Moreover, by the same theorem,

$$f_R \cdot T = \begin{cases} f_R & \text{if } T \subseteq R, \\ 0 & \text{if } T \not\subseteq R. \end{cases}$$

Since T runs over a k -basis of \mathcal{P}_1 , this proves that the ideal $\mathcal{P}_1 f_R$ generated by f_R is equal to the rank one submodule $k \cdot f_R$ spanned by f_R . Thus we obtain

$$\mathcal{P}_1 \cong \prod_{R \in \mathcal{O}} \mathcal{P}_1 f_R = \prod_{R \in \mathcal{O}} k \cdot f_R,$$

as was to be shown. \square

Note that if k is a field, then each idempotent f_R is primitive.

7. The algebra of permuted orders

We know from the end of Section 5 that the algebra \mathcal{P} of permuted orders is Σ -graded,

$$\mathcal{P} = \bigoplus_{\sigma \in \Sigma} \mathcal{P}_\sigma.$$

If $R, S \in \mathcal{O}$ and $\sigma, \tau \in \Sigma$, then the product in \mathcal{P} satisfies

$$(R\Delta_\sigma)(S\Delta_\tau) = (R \cdot (\Delta_\sigma S \Delta_{\sigma^{-1}}))\Delta_\sigma \Delta_\tau = (R \cdot (\Delta_\sigma S \Delta_{\sigma^{-1}}))\Delta_{\sigma\tau},$$

where \cdot denotes the product in \mathcal{P}_1 described in Lemma 6.1. Note that this definition makes sense because $\Delta_\sigma S \Delta_{\sigma^{-1}}$ is an order, since S is. Note also that we can write the basis elements as $\Delta_\sigma S$ with $S \in \mathcal{O}$, because $R\Delta_\sigma = \Delta_\sigma(\Delta_{\sigma^{-1}} R \Delta_\sigma)$ and $\Delta_{\sigma^{-1}} R \Delta_\sigma \in \mathcal{O}$.

Instead of \mathcal{O} , we can use the basis $\{f_R \mid R \in \mathcal{O}\}$ of \mathcal{P}_1 , consisting of the idempotents of \mathcal{P}_1 defined in Theorem 6.2. The group Σ acts by conjugation on the set \mathcal{O} of all orders, hence also on the set $\{f_R \mid R \in \mathcal{O}\}$. We first record the following easy observation.

Lemma 7.1. *Let R be an order and let f_R be the corresponding idempotent of \mathcal{P}_1 . For every $\sigma \in \Sigma$,*

$$\Delta_\sigma f_R \Delta_{\sigma^{-1}} = f^{\sigma R},$$

where ${}^\sigma R := \Delta_\sigma R \Delta_{\sigma^{-1}}$.

Proof. This follows immediately from the definition of f_R in Section 6. □

Since $\mathcal{P} = \bigoplus_{\sigma \in \Sigma} \mathcal{P}_\sigma$, this has a k -basis $\{\Delta_\sigma f_R \mid \sigma \in \Sigma, R \in \mathcal{O}\}$. We now describe the product in \mathcal{P} with respect to this basis.

Lemma 7.2. *The product of basis elements of \mathcal{P} is given by:*

$$(\Delta_\tau f_S)(\Delta_\sigma f_R) = \begin{cases} 0 & \text{if } S \neq {}^\sigma R, \\ \Delta_{\tau\sigma} f_R & \text{if } S = {}^\sigma R, \end{cases}$$

for all $S, R \in \mathcal{O}$ and all $\tau, \sigma \in \Sigma$.

Proof. We have

$$(\Delta_\tau f_S)(\Delta_\sigma f_R) = \Delta_\tau f_S f^{\sigma R} \Delta_\sigma.$$

This is zero if $S \neq {}^\sigma R$. Otherwise we obtain $\Delta_\tau f^{\sigma R} \Delta_\sigma = \Delta_\tau \Delta_\sigma f_R = \Delta_{\tau\sigma} f_R$. □

Corollary 7.3. *Let R be an order and let f_R be the corresponding idempotent of \mathcal{P}_1 . The left ideal $\mathcal{P}f_R$ has a k -basis $\{\Delta_\sigma f_R \mid \sigma \in \Sigma\}$.*

Proof. We have

$$(\Delta_\tau f_S)f_R = \begin{cases} 0 & \text{if } S \neq {}^\sigma R, \\ \Delta_\tau f_R & \text{if } S = {}^\sigma R, \end{cases}$$

and we know that the set $\{\Delta_\sigma f_R \mid \sigma \in \Sigma\}$ is part of the basis of \mathcal{P} . □

Now we introduce central idempotents in \mathcal{P} . Let R be an order, let Σ_R be the stabilizer of R in Σ , and denote by $[\Sigma/\Sigma_R]$ a set of coset representatives. By Lemma 7.1, Σ_R is also the stabilizer of f_R and we define

$$e_R = \sum_{\sigma \in [\Sigma/\Sigma_R]} \Delta_\sigma f_R \Delta_{\sigma^{-1}} = \sum_{\sigma \in [\Sigma/\Sigma_R]} f_{\sigma R},$$

the sum of the Σ -orbit of f_R .

Lemma 7.4. *Let $[\Sigma \setminus \mathcal{O}]$ be a set of representatives of the Σ -orbits in the set \mathcal{O} . Then the set $\{e_R \mid R \in [\Sigma \setminus \mathcal{O}]\}$ is a set of orthogonal central idempotents of \mathcal{P} , whose sum is $1_{\mathcal{P}} = \Delta$.*

Proof. We compute

$$\Delta_\tau f_S e_R = \sum_{\sigma \in [\Sigma/\Sigma_R]} \Delta_\tau f_S f_{\sigma R} = \begin{cases} 0 & \text{if } S \text{ does not belong to the orbit of } R, \\ \Delta_\tau f_S & \text{if } S = \sigma R. \end{cases}$$

On the other hand

$$e_R \Delta_\tau f_S = \sum_{\sigma \in [\Sigma/\Sigma_R]} f_{\sigma R} \Delta_\tau f_S = \sum_{\sigma \in [\Sigma/\Sigma_R]} f_{\sigma R} f_{\tau S} \Delta_\tau.$$

This is zero if τS does not belong to the Σ -orbit of σR , that is, if S does not belong to the Σ -orbit of R , while if $\tau S = \sigma R$, then we get $f_{\tau S} \Delta_\tau = \Delta_\tau f_S$. This shows that e_R is central.

We know that $\{f_R \mid R \in \mathcal{O}\}$ is a set of orthogonal idempotents with sum 1. Since we have just grouped together the Σ -orbits, it is clear that the set $\{e_R \mid R \in [\Sigma \setminus \mathcal{O}]\}$ is also a set of orthogonal idempotents of \mathcal{P} , whose sum is $1_{\mathcal{P}} = \Delta$. □

It follows from Lemma 7.4 that

$$\mathcal{P} \cong \prod_{R \in [\Sigma \setminus \mathcal{O}]} \mathcal{P} e_R$$

and we have to understand the structure of each term.

Theorem 7.5. *Let R be an order on X and let Σ_R be its stabilizer in Σ . Then*

$$\mathcal{P} e_R \cong M_{|\Sigma:\Sigma_R|}(k\Sigma_R),$$

a matrix algebra of size $|\Sigma : \Sigma_R|$ on the group algebra $k\Sigma_R$. In other words

$$\mathcal{P} \cong \prod_{R \in [\Sigma \setminus \mathcal{O}]} M_{|\Sigma:\Sigma_R|}(k\Sigma_R).$$

Proof. By Corollary 7.3, the left ideal $\mathcal{P} f_R$ is a free k -submodule of \mathcal{P} spanned by the set $\{\Delta_\sigma f_R \mid \sigma \in \Sigma\}$. The group Σ_R acts on the right on this set, because $f_R \Delta_h = \Delta_h f_R$ for every $h \in \Sigma_R$. Therefore $\mathcal{P} f_R$ is a free right $k\Sigma_R$ -module with basis $\{\Delta_\sigma f_R \mid \sigma \in [\Sigma/\Sigma_R]\}$.

Clearly, the left action of \mathcal{P} commutes with the right action of $k\Sigma_R$. The left action of \mathcal{P} on this free right $k\Sigma_R$ -module induces a k -algebra map

$$\phi_R : \mathcal{P} \rightarrow M_{|\Sigma:\Sigma_R|}(k\Sigma_R).$$

By Lemma 7.2, e_R acts as the identity on $\mathcal{P}f_R$, while e_S acts by zero if S does not belong to the Σ -orbit of R . Therefore we get a k -algebra map

$$\phi_R : \mathcal{P}e_R \rightarrow M_{|\Sigma:\Sigma_R|}(k\Sigma_R),$$

because $\phi_R(e_S) = 0$ whenever S does not belong to the Σ -orbit of R . Putting all these maps together, we obtain a k -algebra map

$$\phi = \prod_{R \in [\Sigma \setminus \mathcal{O}]} \phi_R : \mathcal{P} \rightarrow \prod_{R \in [\Sigma \setminus \mathcal{O}]} M_{|\Sigma:\Sigma_R|}(k\Sigma_R).$$

Lemma 7.2 shows that, if $\rho, \sigma, \tau \in [\Sigma/\Sigma_R]$ and $g \in \Sigma_R$, we have

$$(\Delta_\tau f_R \Delta_g \Delta_{\rho^{-1}}) \cdot \Delta_\sigma f_R = \begin{cases} 0 & \text{if } \rho \neq \sigma, \\ \Delta_\tau f_R \Delta_g & \text{if } \rho = \sigma, \end{cases}$$

using the fact that $f_R \Delta_g = \Delta_g f_R$. This means that $\phi_R(\Delta_\tau f_R \Delta_g \Delta_{\rho^{-1}})$ is the elementary matrix with a single nonzero entry equal to Δ_g in position (τ, ρ) . Moreover, we also have

$$\phi_S(\Delta_\tau f_R \Delta_g \Delta_{\rho^{-1}}) = 0$$

whenever S does not belong to the Σ -orbit of R . Therefore the map ϕ is surjective.

Finally, we prove that ϕ is an isomorphism. It suffices to do this in the case where $k = \mathbb{Z}$, because all the algebras are defined over \mathbb{Z} (that is, they are obtained by extending scalars from \mathbb{Z} to k) and the algebra map ϕ is also defined over \mathbb{Z} . Now if $k = \mathbb{Z}$, then all algebras under consideration are finitely generated free \mathbb{Z} -modules and we know that the map ϕ is surjective. So it suffices to show that the source and the target of ϕ have the same rank as \mathbb{Z} -modules. The rank of \mathcal{P} is $|\Sigma| \text{Card}(\mathcal{O})$. On the other hand,

$$\text{rank}(M_{|\Sigma:\Sigma_R|}(k\Sigma_R)) = |\Sigma : \Sigma_R|^2 |\Sigma_R| = |\Sigma : \Sigma_R| |\Sigma|.$$

Summing over $R \in [\Sigma \setminus \mathcal{O}]$, we obtain

$$\sum_{R \in [\Sigma \setminus \mathcal{O}]} |\Sigma : \Sigma_R| |\Sigma| = |\Sigma| \sum_{R \in [\Sigma \setminus \mathcal{O}]} \text{Card}(\text{orbit of } R) = |\Sigma| \text{Card}(\mathcal{O}),$$

as was to be shown. □

Remark 7.6. Since a matrix algebra $M_r(A)$ is Morita equivalent to A (for any k -algebra A), it follows from Theorem 7.5 that the algebra \mathcal{P} is Morita equivalent to a product of group algebras, namely $B = \prod_{R \in [\Sigma \setminus \mathcal{O}]} k\Sigma_R$. The bimodule which provides the Morita equivalence is

$$M = \bigoplus_{R \in [\Sigma \setminus \mathcal{O}]} \mathcal{P}f_R,$$

which is clearly a left \mathcal{P} -module by left multiplication, and a right module for each group algebra $k\Sigma_R$, acting by right multiplication on the summand $\mathcal{P}f_R$, and acting by zero on the other summands $\mathcal{P}f_S$, where $S \neq R$ in $[\Sigma \setminus \mathcal{O}]$. Notice that $\mathcal{P}f_R$ is the bimodule appearing in the proof of Theorem 7.5.

Then the bimodule inducing the inverse Morita equivalence is $M^\vee = \bigoplus_{R \in [\Sigma \setminus \mathcal{O}]} f_R \mathcal{P}$. Indeed, we obtain

$$M \otimes_B M^\vee \cong \bigoplus_{S \in [\Sigma \setminus \mathcal{O}]} \bigoplus_{R \in [\Sigma \setminus \mathcal{O}]} \mathcal{P}f_S f_R \mathcal{P} = \bigoplus_{R \in [\Sigma \setminus \mathcal{O}]} \mathcal{P}f_R \mathcal{P} = \mathcal{P}$$

and on the other hand

$$\begin{aligned} M^\vee \otimes_{\mathcal{P}} M &\cong \bigoplus_{S \in [\Sigma \setminus \mathcal{O}]} \bigoplus_{R \in [\Sigma \setminus \mathcal{O}]} f_S \mathcal{P} f_R = \bigoplus_{R \in [\Sigma \setminus \mathcal{O}]} f_R \mathcal{P} f_R \\ &= \bigoplus_{R \in [\Sigma \setminus \mathcal{O}]} k \Sigma_R \cdot f_R \cong \bigoplus_{R \in [\Sigma \setminus \mathcal{O}]} k \Sigma_R = B. \end{aligned}$$

Each module $\mathcal{P}f_R$ has rank $|\Sigma|$, because it has a k -basis $\{\Delta_\sigma f_R \mid \sigma \in \Sigma\}$, but we view it as a free right $k \Sigma_R$ -module of rank $|\Sigma : \Sigma_R|$.

8. Simple modules for the essential algebra

By standard commutative algebra, we know any simple \mathcal{E} -module is actually a module over $k/m \otimes_k \mathcal{E}$, where m is a maximal ideal of k . Replacing k by the quotient k/m , we assume from now on that k is a field. Let \mathcal{E} be the essential algebra of Section 5 and let $\mathcal{P} = \mathcal{E}/N$ be the algebra of permuted orders. Since N is a nilpotent ideal and since nilpotent ideals act by zero on simple modules, any simple \mathcal{E} -module can be viewed as a simple \mathcal{P} -module. So we work with \mathcal{P} and we wish to describe all simple left \mathcal{P} -modules.

There is a general procedure for constructing all simple modules for the algebra of a semigroup S , using equivalence classes of maximal subgroups of S , see [6, Theorem 5.33], or [11, Section 3] for a short presentation. But our previous results allow for a very direct and easy approach, so we do not need to follow the method of [6].

First notice that the simple \mathcal{P}_1 -modules are easy to describe, because \mathcal{P}_1 is a product of copies of k (by Theorem 6.2). More precisely, $\mathcal{P}_1 \cong \prod_{R \in \mathcal{O}} k \cdot f_R$ and each one-dimensional space $k \cdot f_R$ is a simple \mathcal{P}_1 -module (where R runs through the set \mathcal{O} of all orders).

Theorem 8.1. *Assume that k is a field. Let \mathcal{W} be the set of all pairs (R, V) , where R is an order on X and V is a simple $k \Sigma_R$ -module up to isomorphism. The group Σ acts on \mathcal{W} via ${}^\sigma(R, V) := (\sigma R, {}^\sigma V)$, where ${}^\sigma R = \Delta_\sigma R \Delta_{\sigma^{-1}}$ and ${}^\sigma V$ is the conjugate module, a module for the group algebra $k \Sigma_{\sigma R} = k[\sigma \Sigma_R \sigma^{-1}]$.*

- (a) *The set of isomorphism classes of simple \mathcal{P} -modules is parametrized by the set $\Sigma \setminus \mathcal{W}$ of Σ -conjugacy classes of pairs $(R, V) \in \mathcal{W}$.*
- (b) *The simple module corresponding to (R, V) under the parametrization of part (a) is*

$$S_{R,V} = W_R \otimes_k V,$$

where W_R is the unique (up to isomorphism) simple module for the matrix algebra $M_{|\Sigma : \Sigma_R|}(k)$ and $W_R \otimes V$ is viewed as a module for the algebra

$$M_{|\Sigma : \Sigma_R|}(k) \otimes_k k \Sigma_R \cong M_{|\Sigma : \Sigma_R|}(k \Sigma_R),$$

which is one of the factors of the decomposition of \mathcal{P} in Theorem 7.5.

- (c) *The simple \mathcal{P} -module $S_{R,V}$ is also isomorphic to $\mathcal{P}f_R \otimes_{k \Sigma_R} V$, with its natural structure of \mathcal{P} -module under left multiplication.*
- (d) *The simple \mathcal{P} -module $S_{R,V}$ has dimension $|\Sigma : \Sigma_R| \cdot \dim(V)$.*

Proof. By Theorem 7.5, any simple \mathcal{P} -module is a simple module for one of the factors $M_{|\Sigma:\Sigma_R|}(k\Sigma_R)$, where R belongs to a set $[\Sigma \setminus \mathcal{O}]$ of representatives of Σ -orbits in \mathcal{O} . In view of the isomorphism

$$M_{|\Sigma:\Sigma_R|}(k\Sigma_R) \cong M_{|\Sigma:\Sigma_R|}(k) \otimes_k k\Sigma_R,$$

any such simple module is isomorphic to a tensor product $W_R \otimes_k V$ as in the statement. This proves (a) and (b).

By Theorem 7.5, $\mathcal{P}e_R \cong M_{|\Sigma:\Sigma_R|}(k\Sigma_R)$ and its identity element e_R decomposes as a sum of orthogonal idempotents

$$e_R = \sum_{\sigma \in [\Sigma/\Sigma_R]} f_{\sigma R}.$$

Cutting by the idempotent f_R , we obtain the left ideal $\mathcal{P}f_R$, which is a free right $k\Sigma_R$ -module, isomorphic to the space of column vectors with coefficients in $k\Sigma_R$. Now $\mathcal{P}f_R$ is the bimodule providing the Morita equivalence between $M_{|\Sigma:\Sigma_R|}(k\Sigma_R)$ and $k\Sigma_R$ (see Remark 7.6). Therefore, it follows that, for any simple left $k\Sigma_R$ -module V , the corresponding simple module for $\mathcal{P}e_R \cong M_{|\Sigma:\Sigma_R|}(k\Sigma_R)$ is the left \mathcal{P} -module $\mathcal{P}f_R \otimes_{k\Sigma_R} V$. Since $\mathcal{P}f_R$ is the space of column vectors with coefficients in $k\Sigma_R$, while W_R is the space of column vectors with coefficients in k , we get $\mathcal{P}f_R \cong W_R \otimes_k k\Sigma_R$. Therefore our simple \mathcal{P} -module is

$$\mathcal{P}f_R \otimes_{k\Sigma_R} V \cong W_R \otimes_k k\Sigma_R \otimes_{k\Sigma_R} V \cong W_R \otimes_k V \cong S_{R,V},$$

proving (c).

Finally, the dimension is

$$\dim(S_{R,V}) = \dim(W_R \otimes_k V) = \dim(W_R) \cdot \dim(V) = |\Sigma : \Sigma_R| \cdot \dim(V),$$

proving (d). □

Example 8.2. Consider the trivial order Δ . Then $\Sigma_\Delta = \Sigma$ and the matrix algebra reduces to

$$M_{|\Sigma:\Sigma_\Delta|}(k\Sigma_\Delta) \cong M_1(k) \otimes_k k\Sigma \cong k\Sigma.$$

The simple module W_Δ for the algebra $M_1(k)$ is just k and the simple module

$$S_{\Delta,V} = W_\Delta \otimes_k V \cong V$$

is just a simple $k\Sigma$ -module. In that case, the central idempotent e_S of \mathcal{P} acts by zero on V for any order $S \neq \Delta$, hence f_S too (because $f_S e_S = f_S$). Then $R = \sum_{R \subseteq S} f_S$ also acts by zero for any order $R \neq \Delta$. For any essential reflexive relation Q with $Q \neq \Delta$, the action of Q is equal to the action of \overline{Q} (because $Q - \overline{Q}$ belongs to the nilpotent ideal N), and therefore Q also acts by zero on V . Then so does the action of the essential relation $\Delta_\sigma Q$ containing the permutation Δ_σ . This shows that the simple modules $S_{\Delta,V} \cong V$ are just the modules for $k\Sigma$ viewed as a quotient algebra as in Lemma 4.1.

Example 8.3. Consider a total order T . Then $\Sigma_T = \{\text{id}\}$ and the matrix algebra reduces to

$$M_{|\Sigma:\Sigma_T|}(k\Sigma_T) \cong M_n!(k) \otimes_k k \cong M_n!(k).$$

In that case, there is a unique simple $k\Sigma_T$ -module, namely $V = k$, the trivial module for the trivial group. We obtain the single simple module

$$S_{T,k} = W_T \otimes_k k \cong W_T$$

for the algebra $M_{n!}(k)$. Equivalently, with the approach of part (c) of Theorem 8.1, we have $f_T = T$ (by maximality of T in \mathcal{O}) and so

$$S_{T,k} = \mathcal{P}f_T \otimes_{k\Sigma_T} V = \mathcal{P}T \otimes_k k \cong \mathcal{P}T.$$

So we obtain just the left ideal $\mathcal{P}f_T = \mathcal{P}T$, which turns out to be simple in that case. But it is also the left ideal L appearing in Proposition 4.3. So we have recovered the simple module of Proposition 4.3.

We also mention another byproduct of Theorem 7.5.

Theorem 8.4. *If the characteristic of the field k is zero or greater than n , then \mathcal{P} is a semi-simple k -algebra.*

Proof. It suffices to see that each factor in the decomposition of Theorem 7.5 is semi-simple. Now we have the isomorphism

$$M_{|\Sigma:\Sigma_R|}(k\Sigma_R) \cong M_{|\Sigma:\Sigma_R|}(k) \otimes_k k\Sigma_R,$$

and $M_{|\Sigma:\Sigma_R|}(k)$ is a simple algebra. Moreover the group algebra $k\Sigma_R$ is semi-simple by Maschke’s theorem, because the characteristic of k does not divide the order of the group Σ_R , by assumption. The result follows. \square

Every simple \mathcal{P} -module $S_{R,V}$ is a simple \mathcal{R} -module, because of the successive quotients $\mathcal{R} \rightarrow \mathcal{E} \rightarrow \mathcal{P}$. We now give a direct description of the action on $S_{R,V}$ of an arbitrary relation in \mathcal{R} . Since $S_{R,V} \cong \mathcal{P}f_R \otimes_{k\Sigma_R} V$, it suffices to describe the action on $\mathcal{P}f_R$, and for this we can work again with an arbitrary commutative base ring k . Recall that $\mathcal{P}f_R$ has a basis $\{\Delta_\sigma f_R \mid \sigma \in \Sigma\}$.

Proposition 8.5. *Let k be a commutative ring. Let R be an order on X and let Q be an arbitrary relation (in the k -algebra \mathcal{R}). The action of Q on $\mathcal{P}f_R$ is described on the basis elements as follows:*

$$Q \cdot \Delta_\sigma f_R = \begin{cases} \Delta_{\tau\sigma} f_R & \text{if there exists } \tau \in \Sigma \text{ such that } \Delta \subseteq \Delta_{\tau^{-1}} Q \subseteq \sigma R, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Suppose first that S is an order. By Lemma 7.2, the action of f_S is given by

$$f_S \cdot \Delta_\sigma f_R = \begin{cases} \Delta_\sigma f_R & \text{if } S = \sigma R, \\ 0 & \text{otherwise.} \end{cases}$$

Now

$$S = \sum_{\substack{T \in \mathcal{O} \\ S \subseteq T}} f_T$$

and the action of f_T is nonzero only if $T = {}^\sigma R$. So we obtain the action of S as follows:

$$S \cdot \Delta_\sigma f_R = \begin{cases} \Delta_\sigma f_R & \text{if } S \subseteq {}^\sigma R, \\ 0 & \text{otherwise.} \end{cases}$$

Next we suppose that S is reflexive and that its transitive closure \bar{S} is an order. Since $S - \bar{S}$ belongs to the nilpotent ideal N of Section 5, which acts by zero because $\mathcal{P} = \mathcal{E}/N$, the action of S coincides with the action of \bar{S} . Moreover, the condition $\bar{S} \subseteq {}^\sigma R$ is equivalent to $S \subseteq {}^\sigma R$, because ${}^\sigma R$ is transitive. Therefore the action of S is the following:

$$S \cdot \Delta_\sigma f_R = \begin{cases} \Delta_\sigma f_R & \text{if } S \subseteq {}^\sigma R, \\ 0 & \text{otherwise.} \end{cases}$$

Now suppose that S is reflexive and that \bar{S} is not an order. Then \bar{S} is inessential by Proposition 2.5, hence zero in \mathcal{E} . So \bar{S} acts by zero, and since $S - \bar{S}$ acts by zero, the action of S is also zero. On the other hand, S cannot be contained in ${}^\sigma R$, otherwise $\bar{S} \subseteq {}^\sigma R$, which would force \bar{S} to be an order since ${}^\sigma R$ is an order. Therefore the condition $S \subseteq {}^\sigma R$ is never satisfied in that case. So the previous formula still holds, because we have zero on both sides:

$$S \cdot \Delta_\sigma f_R = \begin{cases} \Delta_\sigma f_R & \text{if } S \subseteq {}^\sigma R, \\ 0 & \text{otherwise.} \end{cases}$$

Now suppose that Q contains a permutation Δ_τ . Then $S = \Delta_{\tau^{-1}} Q$ is reflexive and $Q = \Delta_\tau S$. Thus the action of Q is

$$\begin{aligned} Q \cdot \Delta_\sigma f_R &= \begin{cases} \Delta_\tau \Delta_\sigma f_R & \text{if } S \subseteq {}^\sigma R, \\ 0 & \text{otherwise,} \end{cases} \\ &= \begin{cases} \Delta_{\tau\sigma} f_R & \text{if } \Delta_{\tau^{-1}} Q \subseteq {}^\sigma R, \\ 0 & \text{otherwise,} \end{cases} \\ &= \begin{cases} \Delta_{\tau\sigma} f_R & \text{if } \Delta \subseteq \Delta_{\tau^{-1}} Q \subseteq {}^\sigma R, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

The last equality holds because the condition $\Delta_{\tau^{-1}} Q \subseteq {}^\sigma R$ is equivalent to $\Delta \subseteq \Delta_{\tau^{-1}} Q \subseteq {}^\sigma R$, since $S = \Delta_{\tau^{-1}} Q$ is reflexive. This proves the result for such a relation Q .

Finally if Q does not contain a permutation, then Q is inessential by Theorem 3.2, hence acts by zero. On the other hand the condition that there exists $\tau \in \Sigma$ such that $\Delta \subseteq \Delta_{\tau^{-1}} Q$ cannot be satisfied since Q does not contain a permutation. Therefore the previous formula still holds, because we have zero on both sides:

$$Q \cdot \Delta_\sigma f_R = \begin{cases} \Delta_{\tau\sigma} f_R & \text{if } \Delta \subseteq \Delta_{\tau^{-1}} Q \subseteq {}^\sigma R, \\ 0 & \text{otherwise.} \end{cases}$$

This proves the result in all cases. □

Remark 8.6. In the description of the algebra \mathcal{P} (Theorem 7.5) and in the description of its simple modules (Theorem 8.1), we may wonder which groups appear as Σ_R for some order R . The answer is that the group Σ_R is arbitrary. More precisely, by a theorem of Birkhoff [2]

(and further improvements by Thornton [18] and Barmak–Minian [1]), any finite group is isomorphic to Σ_R for some order R on a suitable finite set X . However, for a given finite set X , it is not clear which isomorphism classes of groups Σ_R appear.

Another question is to determine whether or not the simple modules of Theorem 8.1 are absolutely simple. But again this depends on the group Σ_R , because the field k may or may not be a splitting field for the group algebra $k\Sigma_R$.

9. A branching rule

In this section, we let $X = \{1, \dots, n\}$ for simplicity. In order to let n vary, we use a superscript (n) for all objects depending on n , such as $X^{(n)}$ for the set X , $\Sigma^{(n)}$ for the symmetric group on $X^{(n)}$, $\mathcal{O}^{(n)}$ for the set of all orders on $X^{(n)}$, $\mathcal{P}_1^{(n)}$ for the algebra of orders on $X^{(n)}$, $\mathcal{P}^{(n)}$ for the algebra of permuted orders on $X^{(n)}$, etc.

In the representation theory of the symmetric group $\Sigma^{(n)}$, there are well-known branching rules, describing the restriction of simple modules to the subgroup $\Sigma^{(n-1)}$ of all permutations of $X^{(n-1)}$, and the induction of simple modules from $\Sigma^{(n-1)}$ to $\Sigma^{(n)}$. In a similar fashion, working again over an arbitrary commutative base ring k , we will describe how modules for $\mathcal{P}^{(n-1)}$ behave under induction to $\mathcal{P}^{(n)}$. For this we need to view the former as a subalgebra of the latter. We first define

$$\phi : \mathcal{P}_1^{(n-1)} \rightarrow \mathcal{P}_1^{(n)}, \quad \phi(R) = R \cup \{(n, n)\},$$

for any order R on $X^{(n-1)}$. It is clear that $\phi(R)$ is an order on $X^{(n)}$. Since $\Sigma^{(n-1)}$ is a subgroup of $\Sigma^{(n)}$ (by fixing the last letter n), the map ϕ clearly extends to an injective algebra homomorphism $\phi : \mathcal{P}^{(n-1)} \rightarrow \mathcal{P}^{(n)}$.

Now we want to compute the image under ϕ of the idempotents $f_R^{(n-1)}$. For a given order R on $X^{(n-1)}$, we define

$$\mathfrak{S}_R = \{S \in \mathcal{O}^{(n)} \mid S \cap (X^{(n-1)} \times X^{(n-1)}) = R\}.$$

Lemma 9.1. *Let R be an order on $X^{(n-1)}$ and let $f_R^{(n-1)}$ be the corresponding idempotent of $\mathcal{P}_1^{(n-1)}$.*

(a) *If S is an order on $X^{(n)}$ and if $f_S^{(n)}$ is the corresponding idempotent of $\mathcal{P}_1^{(n)}$, then*

$$\phi(f_R^{(n-1)}) \cdot f_S^{(n)} = \begin{cases} f_S^{(n)} & \text{if } S \in \mathfrak{S}_R, \\ 0 & \text{otherwise.} \end{cases}$$

(b) *We have*

$$\phi(f_R^{(n-1)}) = \sum_{S \in \mathfrak{S}_R} f_S^{(n)}.$$

Proof. We have

$$f_R^{(n-1)} = \sum_{\substack{Y \in \mathcal{O}^{(n-1)} \\ R \subseteq Y}} \mu(R, Y)Y$$

and

$$\phi(Y) = Y \cup \{(n, n)\}.$$

Therefore, using part (a) of Theorem A.1, we obtain

$$\begin{aligned} \phi(f_R^{(n-1)}) \cdot f_S^{(n)} &= \sum_{\substack{Y \in \mathcal{O}^{(n-1)} \\ R \subseteq Y}} \mu(R, Y)(Y \cup \{(n, n)\}) \cdot f_S^{(n)} \\ &= \left(\sum_{\substack{Y \in \mathcal{O}^{(n-1)} \\ R \subseteq Y, Y \cup \{(n, n)\} \subseteq S}} \mu(R, Y) \right) f_S^{(n)} \\ &= \left(\sum_{\substack{Y \in \mathcal{O}^{(n-1)} \\ R \subseteq Y \subseteq S \cap (X^{(n-1)} \times X^{(n-1)})}} \mu(R, Y) \right) f_S^{(n)}. \end{aligned}$$

We get zero if $R \not\subseteq S \cap (X^{(n-1)} \times X^{(n-1)})$ and also if $R \subset S \cap (X^{(n-1)} \times X^{(n-1)})$ (by the definition of the Möbius function). If now $R = S \cap (X^{(n-1)} \times X^{(n-1)})$, that is, if $S \in \mathfrak{S}_R$, then the sum reduces to $\mu(R, R) = 1$ and we obtain $f_S^{(n)}$, proving (a).

Now we have

$$\phi(f_R^{(n-1)}) = \sum_{S \in \mathcal{O}^{(n)}} \phi(f_R^{(n-1)}) \cdot f_S^{(n)} = \sum_{S \in \mathfrak{S}_R} f_S^{(n)},$$

proving (b). □

Theorem 9.2. *Let R be an order on $X^{(n-1)}$, let $f_R^{(n-1)}$ be the corresponding idempotent of $\mathcal{P}_1^{(n-1)}$, and let V be a $k\Sigma_R^{(n-1)}$ -module. Then, inducing to $\mathcal{P}^{(n)}$ the $\mathcal{P}^{(n-1)}$ -module*

$$S_{R,V} = \mathcal{P}^{(n-1)} f_R^{(n-1)} \otimes_{k\Sigma_R^{(n-1)}} V,$$

we obtain

$$\begin{aligned} &\mathcal{P}^{(n)} \otimes_{\mathcal{P}^{(n-1)}} (\mathcal{P}^{(n-1)} f_R^{(n-1)} \otimes_{k\Sigma_R^{(n-1)}} V) \\ &\cong \bigoplus_{S \in \mathfrak{S}_R} \mathcal{P}^{(n)} f_S^{(n)} \otimes_{k\Sigma_S^{(n)}} \text{Ind}_{\Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}}^{\Sigma_S^{(n)}} \text{Res}_{\Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}}^{\Sigma_R^{(n-1)}} V. \end{aligned}$$

Proof. Using Lemma 9.1, we obtain

$$\begin{aligned} &\mathcal{P}^{(n)} \otimes_{\mathcal{P}^{(n-1)}} (\mathcal{P}^{(n-1)} f_R^{(n-1)} \otimes_{k\Sigma_R^{(n-1)}} V) \\ &= \mathcal{P}^{(n)} \phi(f_R^{(n-1)}) \otimes_{k\Sigma_R^{(n-1)}} V \\ &\cong \bigoplus_{S \in \mathfrak{S}_R} \mathcal{P}^{(n)} f_S^{(n)} \otimes_{k\Sigma_R^{(n-1)}} V \\ &\cong \bigoplus_{S \in \mathfrak{S}_R} \mathcal{P}^{(n)} f_S^{(n)} \otimes_{k\Sigma_S^{(n)}} k\Sigma_S^{(n)} \otimes_{k[\Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}]} k[\Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}] \otimes_{k\Sigma_R^{(n-1)}} V \\ &\cong \bigoplus_{S \in \mathfrak{S}_R} \mathcal{P}^{(n)} f_S^{(n)} \otimes_{k\Sigma_S^{(n)}} \text{Ind}_{\Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}}^{\Sigma_S^{(n)}} \text{Res}_{\Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}}^{\Sigma_R^{(n-1)}} V, \end{aligned}$$

proving the result. □

Assume for simplicity that the base ring k is a field of characteristic zero and let V be a simple $k\Sigma_R^{(n-1)}$ -module. The $k\Sigma_S^{(n)}$ -module

$$\text{Ind}_{\Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}}^{\Sigma_S^{(n)}} \text{Res}_{\Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}}^{\Sigma_R^{(n-1)}} V$$

is a direct sum of simple modules W , and each W gives rise to a simple $\mathcal{P}^{(n)}$ -module

$$\mathcal{P}^{(n)} f_S^{(n)} \otimes_{k\Sigma_S^{(n)}} W.$$

Moreover, every such simple $\mathcal{P}^{(n)}$ -module occurs with multiplicities, appearing for instance whenever we have σ running in a set of representatives of cosets

$$[\Sigma_R^{(n-1)} / \Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}].$$

For any such σ , we have ${}^\sigma S \in \mathcal{S}_R$ and ${}^\sigma V \cong V$, since V is a $k\Sigma_R^{(n-1)}$ -module and $\sigma \in \Sigma_R^{(n-1)}$. Therefore the corresponding term in the direct sum is

$$\mathcal{P}^{(n)} f_{{}^\sigma S}^{(n)} \otimes_{k\Sigma_R^{(n-1)}} V \cong \mathcal{P}^{(n)} f_{{}^\sigma S}^{(n)} \otimes_{k\Sigma_R^{(n-1)}} {}^\sigma V,$$

but this gives rise to the same simple $\mathcal{P}^{(n)}$ -modules as the ones coming from S , by Theorem 8.1. Thus the multiplicity of these simple $\mathcal{P}^{(n)}$ -modules is at least $|\Sigma_R^{(n-1)} / \Sigma_R^{(n-1)} \cap \Sigma_S^{(n)}|$.

A. Appendix: Möbius inversion

In this appendix, we prove a general result on Möbius inversion involving idempotents in a ring. This was already used by the first author in other contexts (see [3, Section 6.2]) and can be of independent interest.

Let (P, \leq) be a finite lattice. Write 0 for the minimal element of P and write $x \vee y$ for the least upper bound of x and y in P .

Theorem A.1. *Let P be a finite lattice. Let $\{g_x \mid x \in P\}$ be a family of elements in a ring A such that $g_0 = 1$ and $g_x g_y = g_{x \vee y}$ for all $x, y \in P$. For every $x \in P$, define*

$$f_x = \sum_{\substack{y \in P \\ x \leq y}} \mu(x, y) g_y,$$

where μ denotes the Möbius function of the poset P .

(a) *For all $x, y \in P$, we have*

$$g_z f_x = f_x g_z = \begin{cases} f_x & \text{if } z \leq x, \\ 0 & \text{if } z \not\leq x. \end{cases}$$

(b) *The set $\{f_x \mid x \in P\}$ is a set of mutually orthogonal idempotents in P whose sum is 1.*

Note that our assumption implies that every g_x is idempotent, because $x \vee x = x$.

Proof. By Möbius inversion, we have

$$g_x = \sum_{\substack{y \in P \\ x \leq y}} f_y,$$

and in particular $1 = g_0 = \sum_{y \in P} f_y$.

Next we compute products. If $x, z \in P$, then

$$\begin{aligned} f_x g_z &= \left(\sum_{\substack{y \in P \\ x \leq y}} \mu(x, y) g_y \right) g_z = \sum_{\substack{y \in P \\ x \leq y}} \mu(x, y) g_y g_z \\ &= \sum_{\substack{y \in P \\ x \leq y}} \mu(x, y) g_{y \vee z} \\ &= \sum_{\substack{w \in P \\ x \vee z \leq w}} \left(\sum_{\substack{y \in P \\ x \leq y, y \vee z = w}} \mu(x, y) \right) g_w. \end{aligned}$$

Note that $g_z f_x = f_x g_z$ since $g_z g_y = g_{z \vee y} = g_{y \vee z} = g_y g_z$. If x is strictly smaller than $x \vee z$, the inner sum runs over the set of all elements y in the interval $[x, w] := \{v \in P \mid x \leq v \leq w\}$ such that $y \vee (x \vee z) = w$. But we have

$$\sum_{\substack{x \leq y \\ y \vee z = w}} \mu(x, y) = \sum_{\substack{x \leq y \\ y \vee (x \vee z) = w}} \mu(x, y) = 0,$$

by a well-known property of the Möbius function ([16, Corollary 3.9.3]). Thus $f_x g_z = 0$ if x is strictly smaller than $x \vee z$, that is, if $z \not\leq x$.

If now $x = x \vee z$, that is, $z \leq x$, we get $y = y \vee z$ (because $z \leq x \leq y$), hence $y = w$, so that the inner sum has a single term for $y = w$. In that case, we get

$$f_x g_z = \sum_{\substack{w \in P \\ x \leq w}} \mu(x, w) g_w = f_x.$$

Therefore

$$f_x g_z = \begin{cases} f_x & \text{if } z \leq x, \\ 0 & \text{if } z \not\leq x, \end{cases}$$

proving (a).

If now $x, u \in P$, then

$$f_x f_u = \sum_{\substack{y \in P \\ u \leq y}} \mu(u, y) f_x g_y = \sum_{\substack{y \in P \\ u \leq y \leq x}} \mu(u, y) f_x.$$

If $u \not\leq x$, the sum is empty and we get zero. If $u < x$, then

$$\sum_{u \leq y \leq x} \mu(u, y) = 0$$

by the very definition of the Möbius function. This shows that $f_x f_u = 0$ if $u \neq x$. Finally, if $u = x$, then we get $f_x f_x = f_x$, thus f_x is idempotent, and the proof is complete. \square

Corollary A.2. *Let P be a finite lattice. Write t for the maximal element of P and write $x \wedge y$ for the greatest lower bound of x and y in P . Let $\{g_x \mid x \in P\}$ be a family of elements in a ring A such that $g_t = 1$ and $g_x g_y = g_{x \wedge y}$ for all $x, y \in P$. For every $x \in P$, define*

$$f_x = \sum_{\substack{y \in P \\ y \leq x}} \mu(y, x) g_y,$$

where μ denotes the Möbius function of the poset P . Then the set $\{f_x \mid x \in P\}$ is a set of mutually orthogonal idempotents in P whose sum is 1.

Proof. This follows from Theorem A.1 by using the opposite ordering on P . □

References

- [1] J. A. Barmak and E. G. Minian, Automorphism groups of finite posets, *Discrete Math.* **309** (2009), 3424–3426.
- [2] G. Birkhoff, On groups of automorphisms, *Rev. Un. Mat. Argentina* **11** (1946), 155–157.
- [3] S. Bouc, Biset functors for finite groups, *Lecture Notes in Math.* **1990**, Springer-Verlag, Berlin 2010.
- [4] S. Bouc, R. Stancu and J. Thévenaz, Simple biset functors and double Burnside ring, *J. Pure Appl. Algebra* **217** (2013), 546–566.
- [5] M. R. Bremner and M. El Bachraoui, On the semigroup algebra of binary relations, *Comm. Algebra* **38** (2010), 3499–3505.
- [6] A. H. Clifford and G. B. Preston, Algebraic theory of semigroups. Vol. I, *Math. Surveys* **7**, American Mathematical Society, Providence 1961.
- [7] B. Coecke and E. O. Paquette, Categories for the practising physicist, in: *New structures for physics*, *Lecture Notes in Phys.* **813**, Springer-Verlag, Heidelberg (2011), 173–286.
- [8] Y. I. Diasamidze, Complete semigroups of binary relations, *J. Math. Sci. (N. Y.)* **117** (2003), no. 4, 4271–4319.
- [9] M. Hall, *Combinatorial theory*, John Wiley & Sons, New York 1986.
- [10] P. Hall, On representatives of subsets, *J. Lond. Math. Soc.* **10** (1935), 26–30.
- [11] J. C. Harris and N. J. Kuhn, Stable decompositions of classifying spaces of finite abelian p -groups, *Math. Proc. Cambridge Philos. Soc.* **103** (1988), 427–449.
- [12] Y. Huang and B. Liu, On a conjecture for fully indecomposable exponent and Hall exponent, *Linear Multilinear Algebra* **58** (2010), no. 5–6, 699–710.
- [13] R. J. Plemmons and M. T. West, On the semigroup of binary relations, *Pacific J. Math.* **35** (1970), 743–753.
- [14] S. Schwarz, On the semigroup of binary relations on a finite set, *Czechoslovak Math. J.* **20(95)** (1970), 632–679.
- [15] S. Schwarz, The semigroup of fully indecomposable relations and Hall relations, *Czechoslovak Math. J.* **23(98)** (1973), 151–163.
- [16] R. P. Stanley, *Enumerative combinatorics. Vol. I*, Wadsworth & Brooks/Cole, Monterey 1986.
- [17] J. Thévenaz, *G-algebras and modular representation theory*, Oxford Science Publications, Oxford 1995.
- [18] M. C. Thornton, Spaces with given homeomorphism groups, *Proc. Amer. Math. Soc.* **33** (1972), 127–131.

Serge Bouc, CNRS-LAMFA, Université de Picardie - Jules Verne,
33, rue St Leu, F-80039 Amiens Cedex 1, France
e-mail: serge.bouc@u-picardie.fr

Jacques Thévenaz, Section de mathématiques, EPFL,
Station 8, CH-1015 Lausanne, Switzerland
e-mail: Jacques.Thevenaz@epfl.ch

Eingegangen 29. Oktober 2013, in revidierter Fassung 26. Februar 2014