

METHANE EMISSION INVENTORY AND FORECASTING IN MALAYSIA

RAFIU OLASUNKANMI YUSUF

A thesis submitted in fulfilment of the
requirement for the award of the degree of
Doctor of Philosophy (Environmental Engineering)

Faculty of Chemical Engineering
Universiti Teknologi Malaysia

SEPTEMBER 2013

ABSTRACT

The increase in global surface temperature by 0.74 ± 0.18 °C between 1901 and 2000 as a result of global warming has become a serious threat. It is caused by the emission of greenhouse gases into the atmosphere due to human activities. The major greenhouse gases are carbon dioxide, methane and nitrous oxide. Records show that only carbon dioxide received detailed investigation but not methane, hence the motive behind this study. This study examined the emission of methane from six main sources in Malaysia. Data for the inventories of the production of these six sources were taken from 1980 – 2011 and were used to forecast emissions from 2012 – 2020. The data were sourced from Ministries, Departments and International Agencies. Six categories of animals were studied under livestock with their corresponding methane emissions from 1980 – 2011 computed as follows: cattle: 1993Gg (6.13%), buffaloes: 341Gg (10.8%), sheep: 24Gg (0.8%), goats: 55Gg (1.8%), horses: 3Gg (0.1%), poultry: 161Gg (5.1%), and pigs: 579Gg (18.3%). Methane emissions from the other sources from 1980 to 2011 are rice production: 1617Gg (0.02%), crude oil production: 8016636Gg (99.8%), Wastewater (POME): 11362Gg (0.14%), municipal solid waste landfills: 3294Gg (0.04%), coal mining: 14Gg (0.0002%). Forecasting of methane emissions from 2012 to 2020 were carried out using the Box-Jenkins ARIMA method. There were close similarities between the observed and forecast values. In the year 2020 predicted methane emissions will be cattle: 113Gg (72.2%), buffaloes: 8.0Gg (5.1%), sheep: 1.2Gg (0.8%), goats: 4.2 Gg (2.7%), horses: 0.2Gg (0.1%), pigs: 13.2Gg (8.4%), and poultry: 16.8Gg (10.7%) for the livestock sector. For other sectors the forecast will be wastewater: 836Gg for wastewater, 4.7 Gg for coal production, 503,208 Gg for crude oil production, 50.6 Gg for rice production, and 167 Gg from municipal solid waste landfills. Population and GDP will rise to 33.26 million and 329US \$ billion by 2020, respectively. Optimisation was carried out after running a linear regression to determine the significant parameters. The equation developed was a nonlinear programming problem and was solved using sequential quadratic programming (SQL) and implemented on MATLAB environment. Sensitivity analysis carried out on the constraints showed the need to maintain the present livestock and rice production levels. The amount of meat protein currently available far exceeds the dietary protein requirement by more than five times. Several mitigation measures aimed towards reducing future methane emissions in Malaysia were also suggested for the various sources. These are in line with the country's commitment to reduce greenhouse gas emissions by 40% over the 2005 level by 2020. The use of renewable energy in the energy mix was suggested in line with the government's five fuel policy and increase in the number of vehicles using gas was also proposed.

ABSTRAK

Peningkatan suhu pada permukaan global dengan $0.74 \pm 0.18^{\circ}\text{C}$ di antara tahun 1901 dan 2000 merupakan akibat pemanasan global telah menjadi satu ancaman yang serius. Ia adalah disebabkan oleh pelepasan gas rumah hijau ke atmosfera akibat daripada aktiviti manusia. Gas rumah hijau yang utama adalah karbon dioksida, metana dan nitrus oksida. Rekod menunjukkan bahawa hanya karbon dioksida sahaja yang menerima siasatan terperinci tetapi tiada siasatan dilakukan atas gas metana. Maka, motif di sebalik kajian ini ialah untuk meneliti pelepasan metana dari enam sumber utama di Malaysia. Data bagi inventori pengeluaran dari enam sumber diambil diantara tahun 1980 - 2011 dan telah digunakan untuk meramal pengeluaran dari tahun 2012-2020. Data diperolehi daripada Kementerian, Jabatan dan Agensi Antarabangsa. Pelepasan gas metana dari enam kategori haiwan ternakan telah dikaji dari tahun 1980 - 2011 telah dikira seperti berikut: lembu: 1993Gg (6.13%), kerbau: 341Gg (10.8%), kambing biri-biri: 24Gg (0.8%), kambing: 55Gg (1.8 %), kuda: 3Gg (0.1%), ayam: 161Gg (5.1%), dan khinzir: 579Gg (18.3%). Pelepasan metana dari sumber-sumber lain dari tahun 1980-2011 adalah pengeluaran beras: 1617Gg (0.02%), pengeluaran minyak mentah: 8016636Gg (99.8%), Air sisa (POME): 11362Gg (0.14%), tapak pelupusan sisa pepejal perbandaran: 3294Gg (0.04%), perlombongan arang batu: 14Gg (0.0002%). Ramalan pelepasan metana 2012-2020 telah dijalankan dengan menggunakan Kaedah Box-Jenkins ARIMA. Terdapat persamaan yang rapat antara nilai-nilai yang telah diperhatikan dan diramalkan. Pada tahun 2020, pelepasan metana yang diramalkan bagi seksor ternakan adalah sepasi berikut: 113Gg (72.2%), kerbau: 8.0Gg (5.1%), biri-biri: 1.2Gg (0.8%), kambing: 4.2 Gg (2.7%), kuda: 0.2Gg (0.1 %), khinzir: 13.2Gg (8.4%), dan ayam: 16.8Gg (10.7%). Bagi sektor-sektor lain, ramalan adalah air: 836Gg untuk air sisa, 4.7 Gg untuk pengeluaran arang batu, 503208 Gg bagi pengeluaran minyak mentah, 50.6 Gg untuk pengeluaran beras, dan 167 Gg dari tapak pelupusan sisa pepejal perbandaran. Jumlah penduduk dan KDNK masing-masing akan meningkat kepada 33.26 juta orang dan US \$329 bilion pada tahun 2020. Pengoptimuman dilakukan selepas menjalankan regresi linear parameter yang penting. Persamaan dibangunkan adalah pengaturcaraan masalah bukan linear dan telah diselesaikan dengan menggunakan pengaturcaraan kuadratik berjujukan (SQL) dan dilaksanakan pada persekitaran MATLAB. Analisis kepekaan dijalankan ke atas kekangan menunjukkan keperluan untuk mengekalkan tahap terkni penternakan dan pengeluaran beras. Jumlah protein daging sekarang didapati melebihi keperluan protein pemakanan sebanyak lima kali. Beberapa langkah-langkah pengawalan dicadangkan yang bertujuan untuk mengurangkan pelepasan metana dari pelbagai sumber pada masa depan di Malaysia. Ini adalah selaras dengan komitmen negara untuk mengurangkan pelepasan gas rumah hijau sebanyak 40% berbanding tahun 2005 pada tahun 2020. Penggunaan tenaga yang boleh diperbaharui dalam pencampuran tenaga telah dicadangkan selaras dengan polisi kerajaan dalam lima bahan api dan peningkatan bilangan kenderaan yang menggunakan gas juga telah dicadangkan.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xii
	LIST OF FIGURES	xiv
	LIST OF ABBREVIATIONS	xvii
	LIST OF APPENDICES	xxi
1	INTRODUCTION	1
	1.1 Introduction	1
	1.2 Problem Statement	4
	1.3 Aims of the Research	6
	1.4 Objectives of the Research	7
	1.5 Scope of the Research	7
	1.6 Output/Benefits of the Research	8
	1.7 Structure of the Thesis	10
2	LITERATURE REVIEW	12
	2.1 Global Warming and Climate Change	12
	2.2 Effects of Climate Change	14

2.3	Greenhouse Gases and Greenhouse Gas Effect	16
2.4	Methane	18
2.4.1	Sources of Methane	21
2.5	Global Warming Potential (GWP)	24
2.6	Sources of Methane Emissions in Malaysia	25
2.6.1	Municipal Solid Waste Landfill	27
2.6.1.1	Municipal solid waste characterisation in Malaysia	29
2.6.2	Livestock Production and its Methane Emission	31
2.6.2.1	Livestock production in Malaysia	33
2.6.3	Rice Paddies and Methane Emission	34
2.6.3.1	Rice cultivation in Malaysia	36
2.6.4	Coal Mining and Methane Emission	39
2.6.4.1	Coal Mining in Malaysia	40
2.6.5	Anaerobic Wastewater Treatment and Methane Emission	41
2.6.5.1	Malaysia palm oil mill effluent (POME)	46
2.6.6	Oil and Gas Production and Methane Emission	47
2.6.6.1	Oil and Gas Production in Malaysia	48
2.7	Greenhouse Gases Emission Inventory in Selected Countries	51
2.8	Methane Emission Estimation Methods and Techniques	53
2.8.1	Default Methodology	53
2.8.2	First Order Decay (FOD) Method	54
2.8.3	The Static or Closed Chamber Technique	55
2.8.4	The Open or Dynamic Chamber Technique	57
2.8.5	The Optical Remote Sensing Method	57
2.8.6	The Voronoi Method	58

2.8.7	The Triangular Method	59
2.8.8	The LandGEM Method	60
2.8.9	The Mass-Balance Method (MBM)	60
2.8.10	Mobile Plume Method	61
2.8.11	The German EPER Model	62
2.8.12	The GasSim Method	62
2.8.13	The SEMEN Method	63
2.8.14	The MICROGEN-MGM Model	63
2.8.15	The Scholl Canyon Models	63
2.8.16	The GASFILL Model	64
2.8.17	Advantages and disadvantages of the models	65
2.9	Tools for Forecasting Time Series	66
2.9.1	The Auto-Regressive Integrated Moving Average (ARIMA) model	68
2.9.1.1	Estimation and Validation Periods	71
2.10.2	Optimisation	72
3	RESEARCH METHODOLOGY	76
3.1	Introduction	76
3.2	Data Collection	77
3.3	Emission Calculations from Inventory	78
3.3.1	Municipal Solid Waste Emission Calculation	79
3.3.1.1	The First order decay (FOD) method	79
3.3.2	Livestock Production	82
3.3.3	Emission Calculation from Rice Cultivation	84
3.3.4	Emission Calculation from Coal Mining	87
3.3.4.1	Surface mining	87
3.3.4.2	Post-mining	88
3.3.5	Emission Calculation from Wastewater	

	Treatment	
3.3.6	Emission Calculation from Crude Oil	91
	Production	
3.4	Emission Forecasting	93
3.4.1	The ARIMA Model	93
3.4.2	Optimisation	94
3.4.2.1	Definition of variables used	95
3.4.2.2	Model equations for methane emissions	95
3.4.2.3	Total methane emissions	99
3.4.2.4	Optimisation model	99
3.4.2.5	Constraints	99
3.5	Limitations to the study	101
4	RESULTS AND DISCUSSION	102
4.1	Inventories	102
4.1.1	Livestock and Poultry Production and Methane Emissions	102
4.1.2	Inventory of Municipal Solid Waste Management and Methane Emission	106
4.1.3	Crude Oil and Coal Production and Associated Methane Emissions	109
4.1.4	Rice Planted Area and Methane Emissions	111
4.1.5	Wastewater Production (POME)	113
4.2	Emission Forecasting using ARIMA Model	115
4.2.1	ARIMA Model for Livestock Methane Emissions Forecasting	115
4.2.2	ARIMA Model for all Sources	126
4.2.3	Total emissions	137
4.3	Optimisation	137
4.3.1	Post-Optimisation Operation	140
4.3.2	Optimisation of MSW emissions and Mitigation Methods	141

4.3.3	Crude Oil Production and Mitigation Methods	142
4.3.4	Livestock Production and Mitigation Measures	143
4.3.5	Rice Production and Mitigation Measures	145
4.3.6	Wastewater (POME) Treatment and Mitigation Measures	146
4.3.7	Coal Mining and Mitigation Measures	147
4.3.8	Overall Methane Emission Reductions	148
4.3.9	Renewable Energy (RE) Policy	149
5	CONCLUSION AND RECOMMENDATIONS	152
5.1	Conclusion	152
5.2	Achievement and Reflection	152
5.2.1	Revisiting Research Objective 1	153
5.2.2	Revisiting Research Objective 2	153
5.2.3	Revisiting Research Objective 3	154
5.2.4	Revisiting Research Objective 4	154
5.2.5	Researcher's Reflection	155
5.3	Research Contribution	156
5.3.1	Theoretical Contribution	156
5.3.2	Practical Contribution	156
5.4	Recommendations	157
5.4.1	MSW Landfill Gas (LFG) Collection and Use	157
5.4.2	Renewable Energy Pursuit	158
5.4.3	Biogas from Wastewater Treatment	158
5.4.4	Livestock Production	158
5.4.5	Field Measurement	159
5.4.6	Site Visitations	159
	REFERENCES	160
	Appendices A – M	194 – 208

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	UNFCCC Annex I countries - (Developed nations and nations with economies in transition)	13
2.2	Natural and man-made sources of methane	21
2.3	Current studies on methane inventories and emissions	22
2.4	Composition of 100-year GWP of the greenhouse gases	24
2.5	Annual waste generation in Malaysia	28
2.6	Typical characterisation of Malaysian MSW	29
2.7	Domestic rice self-sufficiency level for Malaysia	38
2.8	Methane emission potentials during treatment of wastewater and sludge	45
2.9	Malaysia's responses to climate change and energy policies	50
2.10	Greenhouse gas emissions in Gg CO ₂ eq (without LULUCF) in Annex I countries	52
2.11	Advantages and disadvantages of the emission models	65
2.12	Time series models	67
2.13	Advantages and disadvantages of the ARIMA model	71
2.14	Application of optimisation techniques	74
3.1	Sources and types of data obtained	77
3.2	Emissions factors for enteric fermentation using Tier 1 method	83
3.3	Default emission scaling factors for water regimes	85
3.4	Default conversion factors for different organic amendments	86
3.5	Default MCF values for industrial wastewater	90
3.6	Classification of set of variables	96
4.1	Livestock production in Malaysia (1980 – 2010)	103

4.2	Annual livestock methane emission (1980 – 2011)	104
4.3	Solid waste generation and methane emission in Malaysia (1980 – 2010)	107
4.4	Inventory and emission from crude oil and coal production (1980 – 2011)	109
4.5	Rice planted area methane emissions (1980 – 2011)	111
4.6	Palm oil production and methane emission from POME (wastewater) (1980 – 2011)	113
4.7	Livestock model description	115
4.8	Livestock model fit statistics	116
4.9	Ljung-Box Q statistics for livestock model validation	118
4.10	Forecast of methane emissions from livestock (2012 – 2020)	119
4.11	Comparison of observed and fitted methane emissions from livestock	124
4.12	ARIMA model description for all sources	126
4.13	Model fit statistics for all sources	127
4.14	Ljung-Box Q statistics for model validation	129
4.15	Methane emissions forecast (2012 – 2020)	129
4.16	Comparison of observed and fitted values of methane emissions	134
4.17	ARIMA model parameters	136
4.18	Optimised methane emissions	138
4.19	Electricity production and demand (2005 – 2010)	149
4.20	Gas supply scenario (2010 – 2025)	150

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
1.1	Global surface temperature from 1880 to 2010 and trend from 1880 to 2000	2
1.2	The major greenhouse gases	3
1.3	Past and projected CO ₂ emissions for four sectors in Malaysia	4
2.1	Southern Oscillation Index (SOI)	15
2.2	The greenhouse gas phenomenon	17
2.3	Methane molecule	19
2.4	Sources of methane emissions in Malaysia	25
2.5	Methane emission from each sector in Malaysia	26
2.6	Percentage contribution of each source	26
2.7	Municipal solid waste landfill	27
2.8	Daily waste generation by states in Malaysia	39
2.9	Average composition of MSW generated in Malaysia	40
2.10	Livestock farm	31
2.11	Paddy rice farm	35
2.12	Methane emission process in a paddy field	35
2.13	Harvested areas from different ecologies	37
2.14	Paddy planted area in Malaysia, 1961–2007	37
2.15	Wastewater treatment systems and discharge pathways	42
2.16	Anaerobic wastewater treatment plant	43
2.17	A palm oil plantation in Malaysia	46
2.18	Oil and gas production facility	48
2.19	Greenhouse gas emissions changes in Annex I countries	51
2.20	The static or closed chamber	56

2.21	Schematic of the triangular model	59
2.25	The ARIMA algorithm	69
3.1	Projected plan of the present research	76
4.1	Methane emission from livestock and poultry	105
4.2	Annual methane emission from MSW landfills	108
4.3	Annual methane emission from oil and coal production	110
4.4	Methane emissions from rice production	112
4.5	Methane emissions from wastewater treatment	114
4.6	Residual ACF and PACF for livestock	117
4.7	Observed, fitted and forecast methane emission for cattle	120
4.8	Observed, fitted and forecast methane emission for buffaloes	120
4.9	Observed, fitted and forecast methane emission for sheep	121
4.10	Observed, fitted and forecast methane emission for goats	121
4.11	Observed, fitted and forecast methane emission for horses	122
4.12	Observed, fitted and forecast methane emission for pigs	123
4.13	Observed, fitted and forecast methane emission for poultry	123
4.14	Residual ACF and PACF for all sources	128
4.15	Observed, fitted and forecast methane emission for rice	130
4.16	Observed, fitted and forecast methane emission for livestock	131
4.17	Observed, fitted and forecast methane emission for coal	131
4.18	Observed, fitted and forecast methane emission for crude oil	132
4.19	Observed, fitted and forecast methane emission for wastewater	133
4.20	Observed, fitted and forecast methane emission for MSW	133
4.21	Observed and fitted emissions (1980 – 2011)	137
4.22	Sensitivity analysis at 10%. 20% and 30%	141
4.23	Methane emission comparison for MSW landfills	142
4.24	Comparison of methane emissions from crude oil production	143
4.25	Comparison of livestock methane emissions	144
4.26	Methane emission comparison for rice production	145

4.27	Methane emission comparison for wastewater	146
4.28	Methane emission comparison for coal mining	147
4.29	Comparison of optimised and uncontrolled (BAU) emissions	148

LIST OF ABBREVIATIONS

<i>ANN</i>	-	Artificial Neural Network
<i>ACT</i>	-	Australian Capital Territory
<i>API</i>	-	American Petroleum Institute
<i>AR4</i>	-	Fourth Assessment Report
<i>ARIMA</i>	-	Auto-Regressive Integrated Moving Average
<i>ATSDR</i>	-	Agency for Toxic Substances and Disease Registry
<i>BAU</i>	-	Business as usual
<i>BIC</i>	-	Bayesian information criterion
<i>BOD</i>	-	Biochemical oxygen demand
<i>C₂F₆</i>	-	Carbon hexafluoride
<i>CDM</i>	-	Clean development mechanism
<i>CE</i>	-	Conservation efficiency
<i>CF₄</i>	-	Carbon tetrafluoride
<i>CH₄</i>	-	Methane
<i>CO</i>	-	Carbon monoxide
<i>CO₂</i>	-	Carbon dioxide
<i>COD</i>	-	Chemical oxygen demand
<i>COP</i>	-	Conference of Parties (of the UNFCCC)
<i>CFC</i>	-	Chlorofluorocarbon
<i>CPO</i>	-	Crude palm oil
<i>DDOC</i>	-	Decomposable degradable organic carbon
<i>DOC</i>	-	Degradable organic content
<i>DOS</i>	-	Department of Statistics
<i>ECER</i>	-	East Coast Economic Region
<i>EPU</i>	-	Economic Planning Unit
<i>EF</i>	-	Enteric fermentation, or Emission factor

<i>EIA</i>	-	Energy Information Administration
<i>EIT</i>	-	Economies in transition
<i>ENSO</i>	-	El Nino Southern Oscillation
<i>EPER</i>	-	European Pollutants Emission Register
<i>EPI</i>	-	Environmental performance index
<i>ETP</i>	-	Economic transformation programme
<i>FAO</i>	-	Food and Agriculture Organisation
<i>FELDA</i>	-	Federal Land Development Agency
<i>FFB</i>	-	Fresh fruit bunch
<i>FOD</i>	-	First-order decay
<i>GDP</i>	-	Gross Domestic Product
<i>GHG</i>	-	Greenhouse gas
<i>GMI</i>	-	Global Methane Initiative
<i>GOR</i>	-	Gas-to-oil ratio
<i>GWP</i>	-	Global warming potential
<i>H₂O</i>	-	Water
<i>H₂S</i>	-	Hydrogen sulphide
<i>HFC</i>	-	Hydroflouorocarbon
<i>IAEA</i>	-	International Atomic Energy Agency
<i>ICU</i>	-	Implementation and Coordination Unit
<i>INC</i>	-	Initial National Communication
<i>IPCC</i>	-	Intergovernmental Panel on Climate Change
<i>IPP</i>	-	Independent Power Plant
<i>IRRI</i>	-	International Rice Research Institute
<i>JPSPN</i>	-	Jabatan Pengurusan Sisa Pepejal Negera
<i>LandGEM</i>	-	Landfill gas emission model
<i>LFG</i>	-	Landfill gas
<i>LULUCF</i>	-	Land use and land use change and forestry
<i>MAPE</i>	-	Mean absolute percentage error
<i>MaxAPE</i>	-	Maximum absolute percentage error
<i>MBM</i>	-	Mass balance method
<i>MCF</i>	-	Methane correction factor
<i>MDA</i>	-	Ministries, Departments and Agencies

<i>MM</i>	-	Manure management
<i>MPOB</i>	-	Malaysian Palm Oil Board
<i>MSW</i>	-	Municipal solid waste
<i>N₂O</i>	-	Nitrous oxide
<i>NaCl</i>	-	Sodium chloride
<i>NaOCl</i>	-	Sodium hypochlorite (bleach)
<i>NaOH</i>	-	Sodium hydroxide
<i>NC2</i>	-	Second National Communication
<i>NCl₃</i>	-	Nitrogen trichloride
<i>NMOC</i>	-	Non-methane organic compounds
<i>NO₂</i>	-	Nitrogen dioxide
<i>NSCCC</i>	-	National Steering Committee on Climate Change
<i>OP-FTIR</i>	-	Open-path Fourier Transform Infrared Radiation
<i>PASW</i>	-	Predictive Analysis Software
<i>POME</i>	-	Palm oil mill effluent
<i>ppb</i>	-	Parts per billion
<i>ppm</i>	-	Parts per million
<i>PSO</i>	-	Particle swarm optimisation
<i>RBW</i>	-	Rapidly biodegradable waste
<i>RE</i>	-	Renewable energy
<i>SAR</i>	-	Second Assessment Report
<i>SBW</i>	-	Slowly biodegradable waste
<i>SEMEN</i>	-	Semi-automated empirical methane emission model
<i>SF₆</i>	-	Sulphur hexafluoride
<i>SOI</i>	-	Southern Oscillation Index
<i>SPSS</i>	-	Statistical Processes for the Social Sciences
<i>SREP</i>	-	Small Renewable Energy Programme
<i>ST</i>	-	Surahanjaya Tenaga (Energy Commission)
<i>TAR</i>	-	Third Assessment Report
<i>TNB</i>	-	Tenaga Nasional Berhad
<i>UNCED</i>	-	United Nations Conference on Environment and Development
<i>UNFCCC</i>	-	United Nations Framework Convention on Climate Change
<i>USEPA</i>	-	United States Environmental Protection Agency

<i>VFA</i>	-	Volatile fatty acids
<i>VOC</i>	-	Volatile organic compounds
<i>WMO</i>	-	World Meteorological Organisation

LIST OF APPENDICES


APPENDIX	TITLE	PAGE
A	Livestock inventory compilation	194
B	Inventory compilation for all emission sources	195
C	Rice planted area from Department of Statistics	196
D	Palm oil production data from Department of Statistics (1974-1999)	197
E	Palm oil production data from Department of Statistics (2000-2010)	198
F	Palm oil planted area (ha) from Department of Statistics	199
G	Crude oil and natural gas production data from Department of Statistics (1963-2010)	200
H	Population data from Department of Statistics (1974-1999)	201
I	Population data from Department of Statistics (2000-2012)	202
J	Malaysia GDP (US\$) (1960-2011)	203
K	Predicted livestock quantities	204
L	Predicted emission values	206
M	Municipal solid waste data from JPSPN	208

CHAPTER 1


INTRODUCTION

1.1 Introduction

The rising trend in the temperature of the earth has become a global threat. This is as a result of global warming. Global warming is caused by the emission of greenhouse gases into the atmosphere and it has had a significant impact on the world's climate (Boakye-Agyei, 2011; Bulkeley and Newell, 2010; Calabrò, 2009; Doria *et al.*, 2009; Halady and Rao, 2010; Wong *et al.*, 2010). There is increase in global surface temperature by 0.74 ± 0.18 °C between the start and the end of the 20th century and is expected to increase by 1.1 to 6.4 °C in the 21st century (Karthik, 2011). Another evidence of global warming is the increasing heat content of the oceans and sea level rise (Trenberth, 2010). The planet is said to be heating at a faster rate than at any time in the last 10,000 years. Moreover, eleven of the hottest years on record have occurred since 1983 with the decade of the 1990s being the hottest in the 20th century. The global mean surface temperature in 1998 is the highest on record since 1860 and is followed by 2005 (Hansen *et al.*, 2006; Kaufmann *et al.*, 2006). Figure 1.1(a) shows the global surface temperature change from 1880 to 2010 and Figure 1.1(b) shows the temperature trend from 1880 to 2000 (VijayaVenkataRaman *et al.*, 2012).



(a)



(b)

Figure 1.1: Global surface temperature from 1880 to 2010 and trend from 1880 to 2000 (VijayaVenkataRaman *et al.*, 2012)

Naturally greenhouse gases are 1– 2 % of the earth’s atmosphere and form a shield that absorbs some of the solar radiation which would otherwise have been radiated into space (Houghton *et al.*, 2001). This helps to keep the planet warm to a comfortable and conducive temperature range of around 14°C (57°F). Without this natural greenhouse effect, the average temperature on earth would be approximately –18°C (–2°F).

Climate change is largely a result of human activities, especially the combustion of fossil fuels, which lead to increase in the atmospheric concentrations of greenhouse gases – carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), and other gases (Figure 1.2) (Boakye-Agyei, 2011; Radojevic *et al.*, 2010;

Ramanathan and Feng, 2009). It is a global concern and its continuation is significantly impacting on people, environment, and economic conditions globally (Al-Amin *et al.*, 2010; Kaijage, 2010; Liu and Sweeney, 2012; Nursey-Bray, 2010).




Figure 1.2: The major greenhouse gases (Ramanathan and Feng, 2009)

The combustion of fossil fuels and other human activities have increased the atmospheric concentrations of greenhouse gases (GHGs) since the beginning of the industrial revolution, and have increased the heat-trapping capability of the earth's atmosphere. The major greenhouse gases (GHGs) are water vapour (36–70%), carbon dioxide (9–26%), methane (4–9%), and nitrous oxide (3–7%) (Scheutz *et al.*, 2009). Since the pre-industrial era, atmospheric concentrations of CO₂ and CH₄ have gone up by nearly 30% and more than 100% respectively because of human activities through the burning of fossil fuels (Boakye-Agyei, 2011).

1.2 Problem Statement

In managing the world's climate, the primary attention has focused on reducing the emission of carbon dioxide or CO₂ (Nusbaum, 2010). Carbon dioxide is a powerful greenhouse gas. It is most often blamed for causing global warming and climate change and has been the main target of emission control. It is a product of combustion of any carbon-based fuel (mainly the fossil fuels), and is produced in large quantities. All measures to reduce global warming due to the greenhouse effect tend to focus on CO₂ emissions reduction from combustion of fossil fuels (Nusbaum, 2010). However, it has been reported that the largest cuts in CO₂ emission would not be felt in decades, if it is felt at all (Clinkard, 2010). This is because CO₂ has a very long atmospheric life of 50 – 200 years and more importantly, it is not the only gas that contributes to global climate change.

Many studies have been carried out on the negative effect of CO₂ in Malaysia (Abushammala *et al.*, 2011; Afroz *et al.*, 2003; Awang *et al.*, 2000; Hashim *et al.*, 2005; Mahlia *et al.*, 2001; Safaai *et al.*, 2011). Figure 1.3 shows the emissions of CO₂ from the year 2000 and its projected emission till 2025 (Safaai *et al.*, 2011).




Figure 1.3: Past and projected CO₂ emissions for four sectors in Malaysia (Safaai *et al.*, 2011)

There is no record to show that such a comprehensive study has been carried out on CH₄ inventory, estimation and projection. Compared with studies of CO₂ emissions, there is scarcity of literature on CH₄ emissions in Malaysia and there is neither a comprehensive nor systematic tool to predict these emissions. The recorded study on methane emissions in Malaysia are from municipal solid waste landfills (Abushammala *et al.*, 2010; Chua *et al.*, 2011; Kathiravale *et al.*, 2003; Kathirvale *et al.*, 2004) and wastewater treatment (El-Fadel and Massoud, 2001; Hassan *et al.*, 2011; Sumathi *et al.*, 2008; Yacob *et al.*, 2005, 2006a). The emissions of methane from the other sources have not been adequately investigated.

Methane (CH₄) is another important greenhouse gas and is also a significant contributor to global warming (Xiaoli *et al.*, 2010; Zhang and Chen, 2010). Although annual CH₄ emissions around the world are significantly smaller than CO₂ emissions, and CH₄ concentrations in the atmosphere are about 200 times lower than those of CO₂ (Mackie and Cooper, 2009), but CH₄ still accounts for about 20% of global warming (Adushkin and Kudryavtsev, 2010; Lelieveld *et al.*, 2009; Szemesova and Gera, 2010). On an equivalent mass basis, CH₄ is 21–25 times more powerful greenhouse gas than CO₂ (Abichou *et al.*, 2011; Adushkin and Kudryavtsev, 2010). It is even postulated that the Global Warming Potential (GWP) of CH₄ could be greater than previously stated (Shindell *et al.*, 2009). Because of its shorter atmospheric life span of 12 – 17 years, reduction in methane emissions will have a much more immediate impact on climate, and its implementation will be cheaper (Clinkard, 2010).

Methane is emitted from various man-made (anthropogenic) and natural sources including municipal solid wastes (MSW) landfills, cattle ranching, rice paddies, coal mining, oil and gas drilling and processing, wetlands, termites, wildfires (Mackie and Cooper, 2009). It is a greenhouse gas ‘second only to carbon dioxide in enhanced climate forcing from the pre–industrial era (1750) to the present’ (Hofmann *et al.*, 2006).

Hence the focus of this research will be to take stock of CH₄ emission inventory from 1980 and to project future emissions in Malaysia up to 2020. There is a need to forecast and predict future emission of this gas in order to plan adequately on how to maximally utilise its vast potential as a renewable source of energy. The forecast will provide relevant and reliable information for policy makers for sound planning and to make important decisions. It will also keep all stakeholders alert so as to be able to face the challenges that will arise and to protect the environment. This is more so in view of the drive and plan of the government to make renewable energy the fifth part of the energy mix as enshrined in the Tenth Malaysian Plan. Emission inventories are prepared to determine the contribution from different sources.

The determination of an emission inventory is a useful tool in air quality management. Combined with forecasting, an emission inventory is used to assess the impact of specific human activities and the main sources responsible for such emissions and also to develop and assess the results of specific mitigation strategies (Karl *et al.*, 2009; Winiwarter *et al.*, 2009).

1.3 Aims of the research

The focus of this research will be to take stock of methane emission inventory in Malaysia from 1980 – 2011 and to project the emission from 2012 – 2020.

1.4 Objectives of the Research

The main objectives of this study are outlined below.

1. To take stock of methane inventory from all the likely sources in Malaysia from 1980 to 2011.
2. To forecast methane emissions from these sources from 2012 to 2020.
3. To develop an optimisation model that will lead to reductions in methane emissions in Malaysia.
4. To propose mitigation measures in line with Malaysian government policies.

1.5 Scope of the Research

To achieve the objective of the research, the following will be the scope of study.

The study will be limited to methane emissions from six identified sectors: coal mining, oil and gas production, livestock and poultry activities, rice paddies, wastewater treatment (palm oil mill effluent) and municipal solid waste management in Malaysia.

Palm oil mill effluent (POME) will be used to represent wastewater because it is the highest source of wastewater generation. Other sources like domestic wastewater and other industrial wastewater will be excluded.

Data will be sourced from relevant government Ministries, Departments and similar international organisations. Majority of the data will be from the Economic Planning Unit (EPU) and the Department of Statistics (DOS)

The projection of emissions will be from 2012 - 2020. The time interval was chosen to cover two periods of the national plans (2011-2015 and 2016-2020). Moreover, a longer time horizon reduces accuracy. Box-Jenkins ARIMA model will be used for the forecasting.

Mitigation options will be provided based on sensitivity analysis for some sectors that can be controlled.

The emission factors to be used will be the ones approved by the Intergovernmental Panel on Climate Change, IPCC (IPCC, 2006). These emission factors will be suitable for Tier 1 emissions calculations.

1.6 Output/Benefits of the Research

The study will be beneficial in many respects. These include:

The advantage of using methane as a good source of renewable energy from which small-capacity power generating plants of about 5–10 MW could be built will be highlighted.

It will expose the inherent risks (fire hazard) associated with some of the sources of methane emissions and will enhance environmental awareness with

respect to the dangers associated with increased anthropogenic activities leading to increased greenhouse gas concentrations.

The economic potentials of some of the emissions sources will be shown. Municipal solid waste landfills are known to be potential sources of renewable energy that are cheaper and cleaner than energy from conventional fossil fuels.

The energy diversification programme of the Malaysian government will be boosted as the quantified potentials will reveal the actual amount of available energy. The heating value of methane which is 55.5 MJ/kg is equivalent to 1.2 kg of diesel or 3.7 kg of wood (Fountoulakis and Manios, 2009).

There will be attainment of sustainable development by relying less on non-renewable fossil fuels that will bring about a reduction in environmental pollution. This will further boost the Clean Development Mechanisms (CDM) being canvassed by the Malaysian government and will be a boost for the five fuel energy mix policy of the government.

Job opportunities will be created through the proliferation of small-capacity generating plants.

It will provide the basis for future policy framework that will address greenhouse gas emissions.

1.7 Structure of the Thesis

The thesis is divided into five (5) chapters. The thrust of each chapter is summarised below.

The **first chapter** gives a general introduction to the subject matter which is climate change and its damaging consequences. The chapter also mentions the greenhouse gases (GHG) and their effects including their global warming potentials. The problem statement is mentioned while the aims, objective and scope of the research are highlighted. The expected outputs and benefits of the research are mentioned while the chapter ends with an overview of the thesis arrangement

Chapter two presents a detailed review of global warming; climate change, greenhouse gases and their effects on the environment are discussed extensively. The major GHGs are introduced and methane gas/emission is given extensive review. The emission of methane in Malaysia is also discussed in this chapter. Detailed discussions of all sources of methane (natural and anthropogenic) are described as well as their estimation methods. The chapter also introduces the ARIMA method and the concept of optimisation.

The research design and the methodology to be employed in carrying out the research are highlighted in **chapter three**. The sources of data used for the inventories and for all computations of methane emissions are mentioned. Methane emission calculation methods, as given by the Intergovernmental Panel on Climate Change (IPCC), are also shown. Forecasting for the years 2012 – 2020 were carried out using the Box-Jenkins Auto-Regressive Integrated Moving Average (ARIMA) from the SPSS-PASW 18 software. The chapter concludes by highlighting the method used in carrying out the optimisation and the development of the model for methane emission reduction.

The results of all the computations are given in **Chapter four**. All inventories taken from 1980 – 2012 and the corresponding amounts of methane that would be emitted from the inventories are also given. Methane emissions forecasts were made for 2012 – 2020 for all the sources. The observed and predicted methane emissions values were compared with each other to determine the accuracy of the model. The comparisons are made in graphical forms. The optimisation aspect of the research was also carried out in this chapter. Optimised and uncontrolled (business-as-usual, BAU) emissions were compared and savings to be made from reduced emissions are shown. The chapter ends with different mitigation methods on how to reduce methane emissions from all the sources.

Chapter five presents the conclusions of the study. The research objectives are revisited and the theoretical and practical contributions of the study are mentioned. Recommendations are made in line with the policy direction of the government to make renewable energy the 5th component of the energy mix.

REFERENCES

- Abbaspour, M., Rahmani, A. M. and Teshnehlab, M. (2005). Carbon monoxide prediction using novel intelligent network. *International Journal of Environmental Science and Technology*, 1(4), 257-264.
- Abichou, T., Clark, J. and Chanton, J. (2011). Reporting central tendencies of chamber measured surface emission and oxidation. *Waste Management*, 31(5), 1002-1008.
- Abushammala, M. F. M., Ahmad Basri, N. E., Basri, H., El-Shafie, A. H. and Kadhum, A. A. H. (2011). Regional landfills methane emission inventory in Malaysia. *Waste Management & Research*, 29(8), 863-873.
- Abushammala, M. F. M., Basri, N. E. A., Basri, H., Kadhum, A. A. H. and El-Shafie, A. H. (2010). Estimation of methane emission from landfills in Malaysia using the IPCC 2006 FOD model. *Journal of Applied Sciences*, 10(15), 1603-1609.
- Adushkin, V. V. and Kudryavtsev, V. P. (2010). Global methane flux into the atmosphere and its seasonal variations. *Izvestiya, Physics of the Solid Earth*, 46(4), 350–357.
- Afroz, R., Hassan, M. N. and Ibrahim, N. A. (2003). Review of air pollution and health impacts in Malaysia. *Environmental Research*, 92(2), 71-77.
- Agamuthu, P. and Fauziah, S. H. (2009). Solid waste: Environmental factors and health, from <http://www.ea-swmc.org/download/postconf/Agamuthu.pdf>
- Agamuthu, P., Tan, E. L. and Shaifal, A. A. (1986). Effect of aeration and soil inoculum on the composition of palm oil effluent (POME). *Agricultural Wastes*, 15, 121–132.
- Agarwal, R., Gupta, R. and Garg, J. K. (2009). A hierarchical model for estimating methane emission from wetlands using MODIS data and ARIMA modeling. [Research article]. *Journal of the Indian Society of Remote Sensing* 37, 473-481.

- Agarwal, V. and Garg, J. K. (2009). Methane emission modelling from wetlands and waterlogged areas using MODIS data. *Current Science*, 96(1), 36-40.
- Agustin, M. B., Sengpracha, W. P. and Phutdhawong, W. (2008). Electrocoagulation of palm oil mill effluent. *International Journal of Environmental Research and Public Health*, 5(3), 177-180.
- Ahmad, S., Kadir, M. Z. A. and Shafie, S. (2011). Current perspective of the renewable energy development in Malaysia. *Renewable and Sustainable Energy Reviews*, 15(2), 897-904.
- Akkoyunlu, A., Yetilmezsoy, K., Erturk, F. and Oztemel, E. (2010). A neural network-based approach for the prediction of urban SO₂ concentrations in the Istanbul metropolitan area *International Journal of Environment and Pollution*, 40, 301–321.
- Al-Amin, A. Q., Jaafar, A. H. and Siwar, C. (2010). Climate change mitigation and policy concern for prioritization. *International Journal of Climate Change Strategies and Management*, 2(4), 418 - 425.
- Al-Hamad, K. K., Nassehi, V. and Khan, A. R. (2008). Impact of greenhouse gases (GHG) emissions from oil production facilities at Northern Kuwait oilfields: Simulated results. *American Journal of Environmental Sciences*, 4(5), 491-501.
- Al-Shiab, M. (2006). The predictability of the Amman Stock Exchange using the univariate autoregressive integrated moving average (ARIMA) model. *Journal of Economic & Administrative Sciences*, 17-35.
- Alanis, P., Ashkan, S., Krauter, C., Campbell, S. and Hasson, A. S. (2010). Emissions of volatile fatty acids from feed at dairy facilities. *Atmospheric Environment*, 44(39), 5084-5092.
- Alemu, A. W., Dijkstra, J., Bannink, A., France, J. and Kebreab, E. (2011). Rumen stoichiometric models and their contribution and challenges in predicting enteric methane production. *Animal Feed Science and Technology* 166-167(0), 761-778.
- Allahdadi, M. and Nehi, H. M. (2012). The optimal solution set of the interval linear programming problems. *Optimization Letters*, 1-19.

- Altunkaynak, A., Ozger, M. and Cakmakci, M. (2005). Fuzzy logic modeling of the dissolved oxygen fluctuations in Golden Horn. *Ecological Modelling*, 189, 436–446.
- Aluwong, T., Wuyep, P. A. and Allam, L. (2011). Livestock-environment interactions: Methane emissions from ruminants. *African Journal of Biotechnology*, 10(8), 1265-1269.
- Apine, L. (2011). Residents' attitude towards possible adaptation measures to the sea coast erosion in Latvia. *International Journal of Climate Change Strategies and Management*, 3(3), 238 - 249.
- Ariffin, A., Shatat, R. S. A., Nik Norulaini, A. R. and Mohd Omar, A. K. (2005). Synthetic polyelectrolytes of varying charge densities but similar molar mass based on acrylamide and their applications on palm oil mill effluent treatment. *Desalination*, 173(3), 201-208.
- Aronica, S., Bonanno, A., Piazza, V., Pignato, L. and Trapani, S. (2009). Estimation of biogas produced by the landfill of Palermo, applying a Gaussian model. *Waste Management*, 29, 233-239.
- Asadi, S., Tavakoli, A. and Hejazi, S. R. (2012). A new hybrid for improvement of auto-regressive integrated moving average models applying particle swarm optimization. *Expert Systems with Applications*, 39(5), 5332-5337.
- Asem, S. O. and Roy, W. Y. (2010). Biodiversity and climate change in Kuwait. *International Journal of Climate Change Strategies and Management*, 2(1), 68 - 83.
- Assis, L., Maravilha, A., Vivas, A., Campelo, F. and Ramírez, J. (2012). Multiobjective vehicle routing problem with fixed delivery and optional collections. *Optimization Letters*, 1-13.
- ATSDR. (2001). Landfill gas primer: An overview for environmental health professionals (A. f. T. S. a. D. Registry, Trans.): Department of Health and Human Services, Division of Health Assessment and Consultation, Agency for Toxic Substances and Disease Registry.
- Aulakh, M. S., Khera, T. S., Doran, J. W., Singh, K. and Singh, B. (2000). Yields and nitrogen dynamics in a rice-wheat system using green manure and inorganic fertilizer. *Soil Science Society of America Journal*., 64, 1867-1876.

- Awang, M. B., Jaafar, A. B., Abdullah, A. M., Ismail, M. B., Hassan, M. N., Abdullah, R., *et al.* (2000). Air quality in Malaysia: Impacts, management issues and future challenges. *Respirology*, 5(2), 183-196.
- Aworanti, O. A., Agarry, S. E., Arinkoola, A. O. and Adeniyi, V. (2011). Mathematical modelling for the conversion of animal waste to methane in batch bioreactor. *International Journal of Engineering Science and Technology*, 3 (1), 573-581.
- Babu, Y. J., Nayak, D. R. and Adhya, T. K. (2006). Potassium application reduces methane emission from a flooded field planted to rice. *Biology and Fertility of Soils* 42, 532-541.
- Banerji, G. and Basu, S. (2009). Adapting to climate change in Himalayan cold deserts. *International Journal of Climate Change Strategies and Management*, 2(4), 426 - 448.
- Bannink, A., Smits, M. C. J., Kebreab, E., Mills, J. A. N., Ellis, J. L., Klop, A., *et al.* (2010). Simulating the effects of grassland management and grass ensiling on methane emission from lactating cows. [Modelling animal systems]. *Journal of Agricultural Science* 148, 55–72.
- Barker-Read, G. R. and Radchenko, S. A. (1989). Methane emission from coal and associated strata samples. *International Journal of Mining and Geological Engineering*, 7, 101-126.
- Basturk, A. and Akay, R. (2012). Parallel implementation of synchronous type artificial bee colony algorithm for global optimization. *Journal of Optimization Theory and Applications*, 155(3), 1095-1104.
- Batool, S. A. and Chuadhry, M. N. (2009). The impact of municipal solid waste treatment methods on greenhouse gas emissions in Lahore, Pakistan. *Waste Management*, 29(1), 63-69.
- Bauer, A., Bösch, P., Friedl, A. and Amon, T. (2009). Analysis of methane potentials of steam-exploded wheat straw and estimation of energy yields of combined ethanol and methane production. *Journal of Biotechnology*, 142, 50-55.
- Beauchemin, K. A. and McGinn, S. M. (2009). Reducing methane in dairy and beef cattle operations: What is feasible? Retrieved 12 Jan, 2012, 2012, from http://www.prairiesoilsandcrops.ca/articles/Issue-1_Article_3.pdf

- Beckett, M. (2006). Ministerial address. In H. J. Schellnhuber, Cramer, W., Nakicenovic, N., Wigley, T., and Yohe, G. (Ed.), *Avoiding climate change*: Cambridge University Press, Cambridge, UK.
- Beukes, P. C., Gregorini, P., Romera, A. J., Levy, G. and Waghorn, G. C. (2010). Improving production efficiency as a strategy to mitigate greenhouse gas emissions on pastoral dairy farms in New Zealand. *Agriculture, Ecosystems and Environment*, 136(3–4), 358-365.
- Björkman, M. and Holmström, K. (1999). Global optimization using the DIRECT algorithm in MATLAB. *Advanced Modeling and Optimization*, 1(2), 17-37.
- BMI. (2008). Business Monitor International, Malaysia Power Report Q2 2008: Business Monitor International, London, UK.
- Boakye-Agyei, K. (2011). Approaching climate adjusted environmental due diligence for multilateral financial institutions. *International Journal of Climate Change Strategies and Management*, 3(3), 264 - 274.
- Boo, S. L. (2010). FELDA unit to spend RM688m for cattle project, *The Malaysian Insider*.
- Borges, A. V., Abril, G., Delille, B., Descy, J.-P. and Darchambeau, F. (2011). Diffusive methane emissions to the atmosphere from Lake Kivu (Eastern Africa). *Journal of Geophysical Research*, 116, G03032.
- Boston, J. and Lempp, F. (2011). Climate change: Explaining and solving the mismatch between scientific urgency and political inertia. *Accounting, Auditing & Accountability Journal*, 24(8), 1000 - 1021.
- Bouchard, B. and Nutz, M. (2012). Weak dynamic programming for generalized state constraints. *SIAM Journal of Control and Optimization*, 50(6), 3344–3373.
- Box, G. E. P., Jenkins, G. M. and Reinsel, G. E. (2008). *Time Series Analysis, Forecasting and Control* (4th ed.): John Wiley & Sons, San Francisco.
- BP. (2012). BP Statistical review of world energy - 2012 Retrieved Dec 2, 2012, from bp.com/statisticalreview
- Brockwell, P. J. and Davis, R. A. (2009). *Time series: Theory and methods, 2nd ed.* : Springer.
- Brown, E. G., Anderson, R. C., Carstens, G. E., Gutierrez-Bañuelos, H., McReynolds, J. L., Slay, L. J., *et al.* (2011). Effects of oral nitroethane

- administration on enteric methane emissions and ruminal fermentation in cattle. *Animal Feed Science and Technology*, 166-167(0), 275-281.
- Brown, W. H., Foote, C. S., Iverson, B. L. and Anslyn, E. V. (2009). *Organic Chemistry* (5th ed.): Brooks/Cole Cengage Learning, Belmont, CA, USA.
- Buddle, B. M., Denis, M., Attwood, G. T., Altermann, E., Janssen, P. H., Ronimus, R. S., *et al.* (2011). Strategies to reduce methane emissions from farmed ruminants grazing on pasture. *The Veterinary Journal*, 188(1), 11-17.
- Bulkeley, H. and Newell, P. (2010). *Governing climate change*: Routledge, London.
- Cadenas, E. and Rivera, W. (2010). Wind speed forecasting in three different regions of Mexico, using a hybrid ARIMA-ANN model. *Renewable Energy*, 35(12), 2732-2738.
- Cai, Y., Liu, D., Yao, Y., Li, J. and Qiu, Y. (2011). Geological controls on prediction of coalbed methane of No. 3 coal seam in Southern Qinshui Basin, North China. *International Journal of Coal Geology*, 88(2-3), 101-112.
- Cakmakci, M. (2007). Adaptive neuro-fuzzy modelling of anaerobic digestion of primary sedimentation sludge. *Bioprocess and Biosystems Engineering*, 30(5), 349-357.
- Cakmakci, M., Kinaci, C., Bayramoglu, M. and Yildirim, Y. (2010). A modeling approach for iron concentration in sand filtration effluent using adaptive neuro-fuzzy mode. *Expert Systems with Applications*, 37, 1369-1373.
- Calabrò, P. S. (2009). Greenhouse gases emission from municipal waste management: The role of separate collection. *Waste Management*, 29, 2178-2187.
- Calabrò, P. S. (2010). The effect of separate collection of municipal solid waste on the lower calorific value of the residual waste. *Waste Management & Research*, 28(8), 754-758.
- Cao, Y., Takahashi, T., Horiguchi, K.-i., Yoshida, N. and Cai, Y. (2010). Methane emissions from sheep fed fermented or non-fermented total mixed ration containing whole-crop rice and rice bran. *Animal Feed Science and Technology*, 157(1-2), 72-78.
- Chakraborty, M., Sharma, C., Pandey, J., Singh, N. and Gupta, P. K. (2011). Methane emission estimation from landfills in Delhi: A comparative

- assessment of different methodologies. *Atmospheric Environment*, 45(39), 7135-7142.
- Chandra, R., Takeuchi, H. and Hasegawa, T. (2012). Methane production from lignocellulosic agricultural crop wastes: A review in context to second generation of biofuel production. *Renewable and Sustainable Energy Reviews*, 16(3), 1462-1476.
- Chang, N.-B., Qi, C., Islam, K. and Hossain, F. (2012). Comparisons between global warming potential and cost-benefit criteria for optimal planning of a municipal solid waste management system. *Journal of Cleaner Production*, 20(1), 1-13.
- Chen, H., Zhu, Q. a., Peng, C., Wu, N., Wang, Y., Fang, X., *et al.* (2013). Methane emissions from rice paddies, natural wetlands, and lakes in China: synthesis and new estimate. *Global Change Biology*, 19(1), 19-32.
- Chen, S.-M. and Chen, C.-D. (2011). Handling forecasting problems based on high-order fuzzy logical relationships. *Expert Systems with Applications*, 38(4), 3857-3864.
- Chen, S.-M. and Tanuwijaya, K. (2011). Fuzzy forecasting based on high-order fuzzy logical relationships and automatic clustering techniques. *Expert Systems with Applications*, 38(12), 15425-15437.
- Chen, Y., Yang, G., Sweeney, S. and Feng, Y. (2010). Household biogas use in rural China: A study of opportunities and constraints. *Renewable and Sustainable Energy Reviews*, 14(1), 545-549.
- Cheng, W. and Li, D. (2012). An active set modified Polak-Ribière-Polyak method for large-scale nonlinear bound constrained optimization. *Journal of Optimization Theory and Applications*, 155(3), 1084-1094.
- Chigusa, K., Hasegawa, T., Yamamoto, N. and Watanabe, Y. (1996). Treatment of waste water from oil manufacture plant by yeasts. *Water Science and Technology*, 34, 51-58.
- Choi, H.-L., How, J. and Barton, P. (2012). An outer-approximation approach for information-maximizing sensor selection. *Optimization Letters*, 1-20.
- Chua, K. H., Sahid, E. J. M. and Leong, Y. P. (2011). Sustainable municipal solid waste management and GHG abatement in Malaysia. *Paper presented at the Green & Energy Management, Malaysia.*

<http://www.uniten.edu.my/newhome/uploaded/admin/research/centres/iepre/2012/papers>

- Clinkard, J. (2010). Why is methane the almost-forgotten greenhouse gas? Retrieved Nov 18, 2012, from <http://www.reedconstructiondata.com/market-intelligence/articles/why-is-methane-the-almost-forgotten-greenhouse-gas/>
- Cooper, C. D. and Alley, F. C. (2002). *Air pollution control: A design approach* (Third ed.): Waveland Press Inc., Long Grove, IL.
- Datta, A., Nayak, D. R., Sinhababu, D. P. and Adhya, T. K. (2009). Methane and nitrous oxide emissions from an integrated rainfed rice–fish farming system of Eastern India. *Agriculture, Ecosystems and Environment*, 129, 228-237.
- DCCEE. (2012). Australian National Greenhouse Gas Accounts - Quarterly Update of Australia's National Greenhouse Gas Inventory, Department of Climate Change and Energy Efficiency.
- Dellana, S. A. and West, D. (2009). Predictive modeling for wastewater applications: Linear and nonlinear approaches. *Environmental Modelling & Software*, 24(1), 96-106.
- Demirbas, M. F., Balat, M. and Balat, H. (2009). Potential contribution of biomass to the sustainable energy development. *Energy Conversion and Management*, 50(7), 1746-1760.
- DESR. (2007). Department of Electricity Supply Regulation. Electricity supply iIndustry in Malaysia - Performance and statistical information 2006: Suruhanjaya Tenaga (Energy Commission).
- Díaz-Robles, L. A., Ortega, J. C., Fu, J. S., Reed, G. D., Chow, J. C., Watson, J. G., *et al.* (2008). A hybrid ARIMA and artificial neural networks model to forecast particulate matter in urban areas: The case of Temuco, Chile. *Atmospheric Environment*, 42(35), 8331-8340.
- Dijkstra, J., Oenema, O. and Bannink, A. (2011). Dietary strategies to reducing N excretion from cattle: implications for methane emissions. *Current Opinion in Environmental Sustainability*, 3, 414–422.
- Dlugokencky, E. J., Houweling, S., Bruhwiler, L., Masarie, K. A., Lang, P. M., Miller, J. B., *et al.* (2003). Atmospheric methane levels off: Temporary pause or a new steady-state? *Geophysical Research Letters*, 30(19), n/a-n/a.

- Doagooei, A. R. and Mohebi, H. (2012). Optimization of the difference of topical functions. *Journal of Global Optimization*, 1-10.
- Dobre, I. and Alexandru, A. A. (2008). Modelling unemployment rate using Box-Jenkins procedure. *Journal of Applied Quantitative Methods*, 3(2), 156-166.
- Dong, H., Zhu, Z., Zhou, Z., Xin, H. and Chen, Y. (2011). Greenhouse gas emissions from swine manure stored at different stack heights. *Animal Feed Science and Technology*, 166–167(0), 557-561.
- Doria, M. F., Boyd, E., Tompkins, E. L. and Adger, W. N. (2009). Using expert elicitation to define successful adaptation to climate change. *Environmental Science and Policy*, 12(7), 810-819.
- Dubey, S. K. (2005). Microbial ecology of methane emission in rice agroecosystem: A review. *Applied Ecology and Environmental Research*, 3(2), 1-27.
- Eckard, R. J., Grainger, C. and de Klein, C. A. M. (2010). Options for the abatement of methane and nitrous oxide from ruminant production: A review. *Livestock Science*, 130(1–3), 47-56.
- EIA. (2011). Malaysia energy data, statistics and analysis - oil, gas, electricity, coal: US Energy Information Administration, Washington, DC.
- El-Fadel, M. and Massoud, M. (2001). Methane emissions from wastewater management. *Environmental Pollution*, 114, 177-185.
- Ellis, J. L., Bannink, A., France, J., Kebreab, E. and Dijkstra, J. (2010). Evaluation of enteric methane prediction equations for dairy cows used in whole farm models. *Global Change Biology*, 16, 3246–3256.
- EPU. (2010). Economic Planning Unit, Tenth Malaysia Plan: Economic Planning Unit, Prime Ministers Department, Putrajaya, Malaysia.
- Epule, E. T., Peng, C. and Mafany, N. M. (2011). Methane emissions from paddy rice fields: Strategies towards achieving a win-win sustainability scenario between rice production and methane emission reduction. *Journal of Sustainable Development*, 4(6), 188-196.
- Erickson, P., Heaps, C. and Lazarus, M. (2009). Greenhouse gas mitigation in developing countries (pp. 116): Stockholm Environment Institute.
- Eusuf, M. A., Che Omar, C. M., Mohd. Din, S. A. and Ibrahim, M. (2007). An overview on waste generation characteristics in some selected local authorities in Malaysia. Proceedings of the 2007 *International Conference on*

- Sustainable Solid Waste Management*. 5 - 7 September 2007. Chennai, India 118-125.
- Falk, M. (2011). *A First Course on Time Series Analysis - Examples with SAS*: Chair of Statistics, University of Würzburg.
- FAO. (2010). FAOSTATS, Agricultural database of the Food and Agriculture Organization Retrieved November 14, 2012, from <http://www.fao.org/DOCREP/005/Y4347E/y4347e14.htm>
- Fernandez, R. and Robinson, D. R. (2005). Projects that achieve large methane emissions reductions in oil and gas operations. *Paper presented at the Oil and Gas Methane Emissions Reduction Workshop*, Tomsk, Russia, 14-16 September 2005
- Ferreira, P. A. V. and Machado, M. E. S. (1996). Solving multiple-objective problems in the objective space. *Journal of Optimization Theory and Applications*, 89(3), 659-680.
- Figuroa, V. K., Cooper, C. D. and Mackie, K. R. (2009a). Estimating landfill greenhouse gas emissions from measured ambient methane concentrations and dispersion modeling. Paper no: 327.
- Figuroa, V. K., Mackie, K. R., Cooper, C. D. and Guarriello, N. (2009b). A robust method for estimating landfill methane emissions. *Journal of the Air and Waste Management Association.*, 9(8), 925-935.
- Findikakis, A. N., Papelis, C., Halvadakis, C. P. and Leckie, J. O. (1988). Modeling gas production in managed sanitary landfills. *Waste Management and Research*, 6, 115–123.
- Foley, J., Yuan, Z. and Lant, P. (2009). Dissolved methane in rising main sewer systems: field measurements and simple model development for estimating greenhouse gas emissions. *Journal Water Science and Technology*, 60(11), 2963-2971.
- Fountoulakis, M. S. and Manios, T. (2009). Enhanced methane and hydrogen production from municipal solid waste and agro-industrial by-products co-digested with crude glycerol. *Bioresource Technology*, 100, 3043-3047.
- Franklin, P. (2009). Methane from coal mines presents opportunity to recover energy and generate revenues *Coalbed Methane Outreach Program* US Environmental Protection Agency, Climate Change Division

- Fredenslund, A. M., Scheutz, C. and Kjeldsen, P. (2010). Tracer method to measure landfill gas emissions from leachate collection systems. *Waste Management*, 30(11), 2146-2152.
- Garnsworthy, P. C., Craigon, J., Hernandez-Medrano, J. H. and Saunders, N. (2012). On-farm methane measurements during milking correlate with total methane production by individual dairy cows. *Journal of Dairy Science*, 95(6), 3166-3180.
- Georgakarakos, S., Koutsoubas, D. and Valavanis, V. (2006). Time series analysis and forecasting techniques applied on loliginid and ommastrephid landings in Greek waters. *Fisheries Research*, 78(1), 55-71.
- Ghiassi, M. and Nangoy, S. (2009). A dynamic artificial neural network model for forecasting nonlinear processes. *Computers & Industrial Engineering*, 57(1), 287-297.
- GMI. (2008). Global Methane Initiative. Underground coal mine methane recovery and use opportunities Retrieved Feb. 8, 2012, from http://www.globalmethane.org/documents/coal_fs_eng.pdf
- GMI. (2011). Global Methane Initiative. Coal mine methane: Reducing emissions, advancing recovery and use opportunities Retrieved Aug 18, 2012, from http://www.globalmethane.org/documents/coal_fs_eng.pdf
- Goyal, P., Chan, A. T. and Jaiswal, N. (2006). Statistical models for the prediction of respirable suspended particulate matter in urban cities. *Atmospheric Environment*, 40(11), 2068-2077.
- Grainger, C. and Beauchemin, K. A. (2011). Can enteric methane emissions from ruminants be lowered without lowering their production? *Animal Feed Science and Technology*, 166 -167, 308– 320.
- Gregory, R. G., Attenborough, M. G., Hall, C. D. and Deed, C. (2003). The validation and development of an integrated landfill gas risk assessment model GasSim. Proceedings of the 2003 *Sardinia Proceedings* Cagliari, Italy,
- Grinham, A., Dunbabin, M., Gale, D. and Udy, J. (2011). Quantification of ebullitive and diffusive methane release to atmosphere from a water storage. *Atmospheric Environment*, 45(39), 7166-7173.

- Guisasola, A., Sharma, K. R., Keller, J. and Yuan, Z. (2009). Development of a model for assessing methane formation in rising main sewers. *Water Research*, 43, 2874–2884.
- Gutiérrez-Estrada, J. C., Vasconcelos, R. and Costa, M. J. (2008). Estimating fish community diversity from environmental features in the Tagus estuary (Portugal): multiple linear regression and artificial neural network approaches. *Journal of Applied Ichthyology*, 24, 50–162.
- Halady, I. R. and Rao, P. H. (2010). Does awareness to climate change lead to behavioral change? *International Journal of Climate Change Strategies and Management*, 2(1), 6 - 22.
- Hann, W. C. (2012). Malaysia oil & gas: The golden age of gas? *Sector Update: Maybank*, PP16832/01/2013 (031128), 21 February 2012.
- Hansen, J., Sato, M., Ruedy, R., Lo, K., Lea, D. W. and Medina-Elizade, M. (2006). Global temperature change. *PNAS*, 103(39), 14288–14293.
- Hashim, H., Douglas, P., Elkamel, A. and Croiset, E. (2005). Optimization model for energy planning with CO₂ emission considerations. *Industrial and Engineering Chemistry Research*, 44, 879–890.
- Hasni, A., Chikr-el-Mezouar, Z., Draoui, B. and Boulard, T. (2011). Applying time series analysis model to temperature data in greenhouses. *Sensors & Transducers Journal*, 126(3), 119-124.
- Hassan, M. N. A., Jaramillo, P. and Griffin, W. M. (2011). Life cycle GHG emissions from Malaysian oil palm bioenergy development: The impact on transportation sector's energy security. *Energy Policy*, 39, 2615–2625.
- Hensen, A. and Scharff, H. (2001). Methane emission estimates from landfills obtained with dynamic plume measurements. *Water, Air, and Soil Pollution: Focus*, 1(5-6), 455-464.
- Hepp, S., Augustenborg, C., Dieterich, B., Hochstrasser, T. and Mueller, C. (2010). Impacts of soil moisture on trace gas emissions from grassland: a case study on grassland in Northern Ireland. *Paper presented at the 19th World Congress of Soil Science, Soil Solutions for a Changing World*, Brisbane, Australia, 1 – 6 August 2010.
- Hiriart-Urruty, J.-B. (2012). When only global optimization matters. *Journal of Global Optimization*, 1-3.

- Hofmann, D. J., Butler, J. H., Dlugokencky, E. J., Elkins, J. W., Masarie, K., Montzka, S. A., *et al.* (2006). The role of carbon dioxide in climate forcing from 1979 to 2004: Introduction of the Annual Greenhouse Gas Index. *Tellus*, 58B, 614-619.
- Hoh, R. (2012). Malaysia - Grain and feed annual: Global Agricultural Information Network (GAIN), USDA, Foreign Agricultural Service, MY2001, Washington, DC.
- Hong, S.-T., Lee, A.-K., Lee, H.-H., Park, N.-S. and Lee, S.-H. (2012). Application of neuro-fuzzy PID controller for effective post-chlorination in water treatment plant. *Desalination and Water Treatment*, 47(1-3), 211-220.
- Hoseini, M., Hosseinpour, H. and Bastae, B. (2012). A new multi objective optimization approach in distribution systems. *Optimization Letters*, 1-19.
- Hosseinelou, M. H. and Sohrabi, M. (2009). Predicting and identifying traffic hot spots applying neuro-fuzzy systems in intercity roads. *International Journal of Environmental Science and Technology* 6(2), 309-314.
- Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, N., van der Linden, P. J., Xiaosu, D., *et al.* (2001). *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*: Cambridge University Press, Cambridge.
- Howarth, R., Santoro, R. and Ingraffea, A. (2011). Methane and the greenhouse-gas footprint of natural gas from shale formations. *Climatic Change*, 106(4), 679-690.
- Hsu, Y.-K., VanCuren, T., Park, S., Jakober, C., Herner, J., FitzGibbon, M., *et al.* (2010). Methane emissions inventory verification in southern California. *Atmospheric Environment* 44, 1-7.
- Huang, Y. and He, Q. (2008). Study on the status of output and utilization of landfill gas in China *Journal of Sichuan University of Science and Engineering (natural science edition)* 1, 117-120. .
- Huarte, A., Cifuentes, V., Gratton, R. and Clausse, A. (2010). Correlation of methane emissions with cattle population in Argentine Pampas. *Atmospheric Environment*, 44(23), 2780-2786.

- Hussain, A., Ani, F. N., Silaiman, N. and Adnan, M. F. (2006). Combustion modelling of an industrial municipal waste combustor in Malaysia. *International Journal of Environmental Studies*, 63(3), 313–329.
- INC. (2000). Malaysia: Initial National Communication. Kuala Lumpur: Ministry of Science, Technology and Environment, Malaysia.
- IPCC. (2001). Climate Change 2001: The Scientific Basis. In Y. D. J.T. Houghton, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, C.A. Johnson, and K. Maskell (Ed.). Cambridge, United Kingdom: Intergovernmental Panel on Climate Change.
- IPCC. (2006a). IPCC Guidelines for National Greenhouse Gas Inventories (Vol. 1–5): Intergovernmental Panel on Climate Change.
- IPCC. (2006b). IPCC guidelines for national greenhouse gas inventories (Vol. 1–5): Intergovernmental Panel on Climate Change.
- IPCC. (2007). Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. In D. Q. S. Solomon , M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (Ed.). Cambridge, United Kingdom: Intergovernmental Panel on Climate Change.
- Islami, N., Taib, S., Yusoff, I. and Abdul Ghani, A. (2011). Time lapse chemical fertilizer monitoring in agriculture sandy soil. *International Journal of Environmental Science and Technology* 8(4), 765-780.
- Ismail, N. N., Bono, A., Valintinus, A. C. R., Nilus, S. and Chng, L. M. (2010). Optimization of reaction conditions for preparing carboxymethylcellulose. *Journal of Applied Sciences*, 10(21), 2530-2536.
- Jamal, P., Tompong, M. F. and Alam, M. Z. (2009). Optimization of media composition for the production of bioprotein from pineapple skins by liquid-state bioconversion. *Journal of Applied Sciences*, 9(17), 3104-3109.
- Johansson, D. J. A., Persson, U. M. and Azar, C. (2005). The cost of using global warming potentials: analyzing the trade off between CO₂, CH₄ and N₂O. *Climate Change*, 77, 291-309.
- Johari, A., Ahmed, S. I., Hashim, H., Alkali, H. and Ramli, M. (2012). Economic and environmental benefits of landfill gas from municipal solid waste in Malaysia. *Renewable and Sustainable Energy Reviews*, 16(5), 2907-2912.

- Jones, F. M., Phillips, F. A., Naylor, T. and Mercer, N. B. (2011). Methane emissions from grazing Angus beef cows selected for divergent residual feed intake. *Animal Feed Science and Technology*, 166-167(0), 302-307.
- Jouquet, P., Traoré, S., Choosai, C., Hartmann, C. and Bignell, D. (2011). Influence of termites on ecosystem functioning. Ecosystem services provided by termites. *European Journal of Soil Biology*, 47(4), 215-222.
- JPSPN. (2010). Landfill sites distribution in Malaysia: Jabatan Pengurusan Sisa Pepejal Negara, Kuala Lumpur, Malaysia.
- Kaijage, H. R. (2010). A basis for climate change adaptation in Africa: burdens ahead and policy options. *International Journal of Climate Change Strategies and Management*, 4(2), 52 - 160.
- Kamal, S. A., Md Jahim, J., Anuar, N., Hassan, O., Wan Daud, W. R., Mansor, M. F., *et al.* (2012). Pre-treatment effect of palm oil mill effluent (POME) during hydrogen production by a local isolate *Clostridium butyricum*. *International Journal on Advanced Science Engineering Information Technology*, 2(4), 54-60.
- Kamarudin, W. N. B. (2008). The CDM/sustainable energy market in Malaysia: Malaysian Energy Center (PTM), Kuala Lumpur, Malaysia.
- Kanadasan, G., Mashitah, M. D. and Vadivelu, V. M. (2010). Fixed bed adsorption of methylene blue by using palm oil mill effluent waste activated sludge. *Paper presented at the 3rd IWA Asia Pacific Young Water Professional Conference*, Singapore. 21-24 November, 2010.
- Karaca, F. and Ozkaya, B. (2006). NN-LEAP: a neural network-based model for controlling leachate flow-rate in a municipal solid waste landfill site. *Environmental Modeling Software*, 21, 1190–1197.
- Karacan, C. Ö. and Goodman, G. V. R. (2012). A CART technique to adjust production from longwall coal operations under ventilation constraints. *Safety Science*, 50(3), 510-522.
- Karacan, C. Ö., Ruiz, F. A., Cotè, M. and Phipps, S. (2011). Coal mine methane: A review of capture and utilization practices with benefits to mining safety and to greenhouse gas reduction. *International Journal of Coal Geology*, 86(2–3), 121-156.

- Karakurt, I., Aydin, G. and Aydiner, K. (2012). Sources and mitigation of methane emissions by sectors: A critical review. *Renewable Energy*, 39(1), 40-48.
- Karl, M., Guenther, A., Koble, R., Leip, A. and Seufert, G. (2009). A new European plant-specific emission inventory of biogenic volatile organic compounds for use in atmospheric transport models. *Biogeosciences*, 6(6), 1059–1087.
- Karthik, M. (2011). Impact of methane emissions from wastewater sector in India through a case study of an effluent treatment plant. Proceedings of the 2011 *National Research Conference on Climate Change*, IIT Delhi November 5-6, 2011
- Kasuya, H. and Takahashi, J. (2010). Methane emissions from dry cows fed grass or legume silage. *Asian-Australasian Journal of Animal Science*, 23(5), 563 - 566.
- Kathiravale, S., Muhd Yunusa, M. N., Sopian, K., Samsuddin, A. H. and Rahman, R. A. (2003). Modeling the heating value of municipal solid waste. *Fuel*, 82, 1119–1125.
- Kathirvale, S., Muhd Yunus, M. N., Sopian, K. and Samsuddin, A. H. (2004). Energy potential from municipal solid waste in Malaysia. *Renewable Energy*, 29(4), 559-567.
- Katimon, A. and Demun, A. S. (2004). Water use trend at Universiti Teknologi Malaysia: Application of ARIMA Model. *Technology Journal*, 41, 47-56.
- Kaufmann, R. K., Kauppi, H. and Stock, J. H. (2006). Emissions, concentrations, and temperature: A time series analysis *Climate Change*, 77(3-4), 249-278.
- Ke, J., Singh, D. and Chen, S. (2011). Aromatic compound degradation by the wood-feeding termite *Coptotermes formosanus* (Shiraki). *International Biodeterioration & Biodegradation*, 65(6), 744-756.
- Khashei, M. and Bijari, M. (2010). An artificial neural network (p,d,q) model for timeseries forecasting. *Expert Systems with Applications*, 37(1), 479-489.
- Khashei, M. and Bijari, M. (2011). A novel hybridization of artificial neural networks and ARIMA models for time series forecasting. *Applied Soft Computing*, 11(2), 2664-2675.
- Khashei, M. and Bijari, M. (2012). A new class of hybrid models for time series forecasting. *Expert Systems with Applications*, 39(4), 4344-4357.

- Khashei, M., Bijari, M. and Raissi Ardali, G. A. (2009). Improvement of autoregressive integrated moving average models using fuzzy logic and artificial neural networks (ANNs). *Neurocomputing*, 72(4–6), 956-967.
- Koch, J., Dayan, U. and Amir, S. (2003). Stabilization of atmospheric concentrations of greenhouse gases. *Climatic Change*, 57, 227–241.
- Kornboonraksa, T., Chiemchaisri, C., Chiemchaisri, W., Towprayoon, S. and Visvanathan, C. (2004). Determination of methane gas emissions from waste disposal sites in Thailand with geographical information system application. Proceedings of the 2004 *The Joint International Conference on “Sustainable Energy and Environment (SEE)*. 1-3 December 2004. Hua Hin, Thailand, 293-297.
- Kumar, S., Gaikwad, S. A., Shekdar, A. V., Kshirsagar, P. S. and Singh, R. N. (2004). Estimation method for national methane emission from solid waste landfills. *Atmospheric Environment*, 38, 3481-3487.
- Lam, M. K. and Lee, K. T. (2011). Renewable and sustainable bioenergies production from palm oil mill effluent (POME): Win-win strategies toward better environmental protection. *Biotechnology Advances*, 29(1), 124-141.
- Larbani, M. and Yu, P. (2012). Decision Making and Optimization in Changeable Spaces, a New Paradigm. *Journal of Optimization Theory and Applications*, 155(3), 727-761.
- Lascano, C. E. and Cárdenas, E. (2010). Alternatives for methane emission mitigation in livestock systems. *Revista Brasileira de Zootecnia*, 39, 175-182.
- Latif, M. A., Ahmad, A., Ghufuran, R. and Wahid, Z. A. (2011). Effect of temperature and organic loading rate on upflow anaerobic sludge blanket reactor and CH₄ production by treating liquidized food waste. *Environmental Progress & Sustainable Energy*, n/a-n/a.
- Legesse, G., Small, J. A., Scott, S. L., Crow, G. H., Block, H. C., Alemu, A. W., *et al.* (2011). Predictions of enteric methane emissions for various summer pasture and winter feeding strategies for cow calf production. *Animal Feed Science and Technology* 166 - 167, 678– 687.

- Lelieveld, J., Hoor, P., Jöckel, P., Pozzer, A., Hadjinicolaou, P., Cammas, J.-P., *et al.* (2009). Severe ozone air pollution in the Persian Gulf region. *Atmospheric Chemistry and Physics*, 9(4), 1393–1406.
- Li, T., Huang, Y., Zhang, W. and Song, C. (2010). CH₄MOD_{wetland}: A biogeophysical model for simulating methane emissions from natural wetlands. *Ecological Modelling*, 221, 666–680.
- Li, X. (2009). Applying GLM model and ARIMA model to the analysis of monthly temperature of Stockholm *Essay in Statistics*: Department of Economics and Society, Dalarna University, Sweden.
- Li, Z.-S., Yang, L., Qu, X.-Y. and Sui, Y.-M. (2009). Municipal solid waste management in Beijing City. *Waste Management*, 29, 2596–2599.
- Lin, Q. G., Huang, G. H., Bass, B. and Qin, X. S. (2009). IFTEM: An interval-fuzzy two-stage stochastic optimization model for regional energy systems planning under uncertainty. *Energy Policy*, 37(3), 868-878.
- Lin, Y., Bolca, S., Vandevijvere, S., Van Oyen, H., Van Camp, J., De Backer, G., *et al.* (2011). Dietary sources of animal and plant protein intake among Flemish preschool children and the association with socio-economic and lifestyle-related factors. *Nutrition Journal*, 10(1), 1-12.
- Liu, X. and Sweeney, J. (2012). The impacts of climate change on domestic natural gas consumption in the Greater Dublin Region. *International Journal of Climate Change Strategies and Management*, 4(2), 161 - 178.
- LoRESTANI, A. A. Z. (2006). *Biological treatment of palm oil mill effluent (POME) using an up-flow anaerobic sludge fixed film (UASFF) bioreactor*. PhD Thesis, Universiti Sains Malaysia (USM).
- LoRESTANI, A. A. Z., Mohamed, A. R., Mashitah, M. D., Abdullah, A. Z. and Hasnain Isa, M. (2006). Effects of organic loading rate on palm oil mill effluent treatment in an up-flow anaerobic sludge fixed film bioreactor. *Environmental Engineering and Management Journal*, 5, 337-350.
- Lu, J., Vecchi, G. A. and Reichler, T. (2007). Expansion of the Hadley cell under global warming. *Geophysical Research Letters*, 34, L06805.
- Luo, D. and Dai, Y. (2009). Economic evaluation of coalbed methane production in China. *Energy Policy*, 37, 3883–3889.

- Ma, J., Xu, H. and Cai, Z. (2010). Effect of fertilization on methane emissions from rice fields. *Soils*, 42(2), 153-163.
- Machado, S. L., Carvalho, M. F., Gourc, J.-P., Vilar, O. M. and do Nascimento, J. C. F. (2009). Methane generation in tropical landfills: Simplified methods and field results. *Waste Management*, 29, 153–161.
- Mackie, K. R. and Cooper, C. D. (2009). Landfill gas emission prediction using Voronoi diagrams and importance sampling. *Environmental Modelling and Software* 24, 1223-1232.
- MAHA. (2008). The growth and sustainability of the livestock industry in Malaysia: The Malaysian Experience. *Paper presented at the Austrade Symposium: MAHA 2008*, Serdang, Selangor, Malaysia.
- Mahlia, T. M., Abdulmuin, M. Z., Alamsyah, T. M. and Mukhlisien, D. (2001). An alternative energy source from palm wastes industry for Malaysia and Indonesia. *Energy conversion and Management*, 42, 2109-2118.
- Maia, A. L. S. and de Carvalho, F. d. A. T. (2011). Holt's exponential smoothing and neural network models for forecasting interval-valued time series. *International Journal of Forecasting*, 27(3), 740-759.
- Manaf, L. A., Samah, M. A. A. and Zukki, N. I. M. (2009). Municipal solid waste management in Malaysia: Practices and challenges. *Waste Management*, 29(11), 2902-2906.
- McGeough, E. J., O'Kiely, P., Foley, P. A., Hart, K. J., Boland, T. M. and Kenny, D. A. (2010). Methane emissions, feed intake, and performance of finishing beef cattle offered maize silage harvested at 4 different stages of maturity. *J. Anim. Sci.*, 88, 1479–1491.
- McGinn, S. M., Turner, D., Tomkins, N., Charmley, E., Bishop-Hurley, G. and Chen, D. (2011). Methane emissions from grazing cattle using point-source dispersion. *Journal of Environmental Quality*, 40(1), 22-27.
- Mendes, P., Santos, A. C., Perna, F. and Ribau Teixeira, M. (2012). The balanced scorecard as an integrated model applied to the Portuguese public service: a case study in the waste sector. *Journal of Cleaner Production*, 24(0), 20-29.
- MfE. (2012). New Zealand's greenhouse gas inventory: 1990–2010: Ministry fo the Environment.

- Minamikawa, K. and Sakai, N. (2005). The effect of water management based on soil redox potential on methane emission from two kinds of paddy soils in Japan. *Agriculture, Ecosystems and Environment* 107, 397–407.
- Mirza, U. K., Ahmad, N. and Majeed, T. (2008). An overview of biomass energy utilization in Pakistan. *Renewable and Sustainable Energy Reviews*, 12(7), 1988-1996.
- Mitra, S., Majumdar, D. and Wassmann, R. (2012). Methane production and emission in surface and subsurface rice soils and their blends. *Agriculture, Ecosystems & Environment*, 158(0), 94-102.
- MMD. (2012). El Niño/La Niña: Its impact on weather in Malaysia: Malaysian Meteorological Department, Ministry of Science, Technology and Innovation (MOSTI), Petaling Jaya, Selangor, Malaysia.
- Modarres, R. and Dehkordi, A. K. (2005). Daily air pollution time series analysis of Isfahan City. *International Journal of Environmental Science and Technology* 2(3), 259-267.
- Molinuevo-Salces, B., García-González, M. C., González-Fernández, C., Cuetos, M. J., Morán, A. and Gómez, X. (2010). Anaerobic co-digestion of livestock wastes with vegetable processing wastes: A statistical analysis. *Bioresource Technology*, 101(24), 9479-9485.
- Montazar, A. (2011). A decision tool for optimal irrigated crop planning and water resources sustainability. *Journal of Global Optimization*, 1-14.
- Moore, S., Freund, P., Riemer, P. and Smith, A. (1998). Abatement of methane emissions. *Greenhouse Gas R&D Programme*: International Energy Agency.
- Mordukhovich, B. S. and Nghia, T. T. A. (2012). DC optimization approach to metric regularity of convex multifunctions with applications to infinite systems. *Journal of Optimization Theory and Applications*, 155(3), 762-784.
- Mori, A. and Hojito, M. (2011). Nitrous oxide and methane emissions from grassland treated with bark- or sawdust-containing manure at different rates. *Soil Science and Plant Nutrition*, 57(1), 138-149.
- Morris, J. W. F., Crest, M., Barlaz, M. A., Spokas, K. A., Åkerman, A. and Yuan, L. (2012). Improved methodology to assess modification and completion of landfill gas management in the aftercare period. *Waste Management*, 32(12), 2364-2373.

- MPOB. (2008). Malaysian Palm Oil Board. Economics and Industrial Development Division Retrieved 7 July, 2012, from www.mpob.gov.my
- Mukherjee, R. and Sarkar, U. (2008). Development of a micrometeorological model for the estimation of methane flux from paddy fields: Validation with standard direct measurements. *Environmental Modelling & Software*, 23, 1229-1239.
- Nahlik, A. M. and Mitsch, W. J. (2011). Methane emissions from tropical freshwater wetlands located in different climatic zones of Costa Rica. *Global Change Biology*, 17, 321–1334.
- Najim, M. M. M., Lee, T. S., Haque, M. A. and Esham, M. (2007). Sustainability of rice production: A Malaysian perspective. *The Journal of Agricultural Sciences*, 3(1), 1-12.
- Nanthakumar, L. and Ibrahim, Y. (2010). Forecasting international tourism demand in Malaysia using Box-Jenkins SARIMA application. *South Asian Journal of Tourism and Heritage*, 3(2), 50-60.
- Naqvi, S. M. K. and Sejian, V. (2011). Global climate change: Role of livestock. *Asian Journal of Agricultural Sciences* 3(1), 19-25.
- Nasir, A. A. (2007). Institutionalizing solid waste management in Malaysia. Kuala Lumpur: Department of National Solid Waste Management (JPSPN), Ministry of Housing and Local Government, Malaysia.
- NC2. (2010). Malaysia's second national communication (NC2) submitted to the United Nations Framework Convention on Climate Change (UNFCCC). Putrajaya: Ministry of Natural Resources and Environment, Malaysia.
- NCCFN. (2010). Malaysian dietary guidelines, National Coordinating Committee on Food and Nutrition, Ministry of Health, Putrajaya, Malaysia Retrieved Jan 7, 2013, from <http://www.moh.gov.my/images/gallery/Garispanduan/diet/introduction.pdf>
- Noor, Z. Z., Yusuf, R. O., Abba, A. H., Abu Hassan, M. A. and Mohd Din, M. F. (2012). An overview for energy recovery from municipal solid wastes (MSW) in Malaysia scenario. *Renewable and Sustainable Energy Reviews*.
- Noor, Z. Z., Yusuf, R. O., Abba, A. H., Abu Hassan, M. A. and Mohd Din, M. F. (2013). An overview for energy recovery from municipal solid wastes

- (MSW) in Malaysia scenario. *Renewable and Sustainable Energy Reviews*, 20(0), 378-384.
- Nowakowski, A. and Popa, A. (2012). A dynamic programming approach for approximate optimal control for cancer therapy. *Journal of Optimization Theory and Applications*, 1-15.
- Nurdiyana, H. and Siti Mazlina, M. K. (2009). Optimization of protein extraction from fish waste using response surface methodology. *Journal of Applied Sciences*, 9(17), 3121-3125.
- Nurse-Bray, M. (2010). Climate change adaptation in Australia: Education, training and achieving social and political outcomes. *International Journal of Climate Change Strategies and Management*, 2(4), 393 - 402.
- Nusbaum, N. J. (2010). Dairy livestock methane remediation and global warming. *J Community Health*, 35, 500–502.
- Oh, T. H., Pang, S. Y. and Chua, S. C. (2010). Energy policy and alternative energy in Malaysia: Issues and challenges for sustainable growth. *Renewable and Sustainable Energy Reviews*, 14(4), 1241-1252.
- Omran, A., Mahmood, A., Abdul Aziz, H. and Robinson, G. M. (2009). Investigating households attitude toward recycling of solid waste in Malaysia: A case study. *International Journal of Environmental Research*, 3(2), 275-288.
- Ong, H. C., Mahlia, T. M. I. and Masjuki, H. H. (2011). A review on energy scenario and sustainable energy in Malaysia. *Renewable and Sustainable Energy Reviews*, 15(1), 639-647.
- Oonk, H. and Boom, T. (2000). Landfill gas emission measurements using a mass-balance method. Proceedings of the 2000 *Intercontinental Landfill Research Symposium*. 11–13 Dec. 2000. Lulea University of Technology, Lulea, Sweden,
- Osada, T., Takada, R. and Shinzato, I. (2011). Potential reduction of greenhouse gas emission from swine manure by using a low-protein diet supplemented with synthetic amino acids. *Animal Feed Science and Technology*, 166–167(0), 562-574.
- Othman, J. and Jafari, Y. (2012). Accounting for depletion of oil and gas resources in Malaysia. *Natural Resources Research*, 21(4), 483-494.

- Ozdemir, E. (2009). Modeling of coal bed methane (CBM) production and CO₂ sequestration in coal seams. *International Journal of Coal Geology* 77, 145–152.
- Park, K. H., Jeon, J. H., Jeon, K. H., Kwag, J. H. and Choi, D. Y. (2011). Low greenhouse gas emissions during composting of solid swine manure. *Animal Feed Science and Technology*, 166–167(0), 550-556.
- Park, S., Brown, K. W., Thomas, J. C., Lee, I.-C. and Sung, K. (2010). Comparison study of methane emissions from landfills with different landfill covers. *Environmental Earth Sciences*, 60, 933–941.
- Parsopoulos, K. E. and Vrahatis, M. N. (2010). *Particle Swarm Optimization and Intelligence: Advances and Applications*. New York: Information Science Reference.
- Peñalba, L. M., Elazegui, D. D., Pulhin, J. M. and Cruz, R. V. O. (2012). Social and institutional dimensions of climate change adaptation. *International Journal of Climate Change Strategies and Management*, 4(3), 308 - 322.
- Pennock, D., Yates, T., Bedard-Haughn, A., Phipps, K., Farrell, R. and McDougal, R. (2010). Landscape controls on N₂O and CH₄ emissions from freshwater mineral soil wetlands of the Canadian Prairie Pothole region. *Geoderma*, 155(3–4), 308-319.
- Percival, D. B. and Walden, A. T. (1993). *Spectral analysis for physical applications*: Cambridge University Press.
- Perera, M. D. N., Hettiaratchi, J. P. A. and Achari, G. (2002). A mathematical modeling approach to improve the point estimation of landfill gas surface emissions using the flux chamber technique. *Journal of Environmental Engineering Science*, 1, 451-463.
- Peterson, C., Barrera, C. and Azizova, Z. (2010). Waste and the World Bank. *Waste Management World*, 14-21.
- Poh, P. E. and Chong, M. F. (2009). Development of anaerobic digestion methods for palm oil mill effluent (POME) treatment. [Review]. *Bioresource Technology*, 100(1), 1-9.
- Pratt, C., Walcroft, A. S., Tate, K. R., Ross, D. J., Roy, R., Reid, M. H., *et al.* (2012). Biofiltration of methane emissions from a dairy farm effluent pond. *Agriculture, Ecosystems & Environment*, 152(0), 33-39.

- Préndez, M. and Lara-González, S. (2008). Application of strategies for sanitation management in wastewater treatment plants in order to control/reduce greenhouse gas emissions. *Journal of Environmental Management*, 88(4), 658-664.
- Prista, N., Diawara, N., Costa, M. J. and Jones, C. (2011). Use of SARIMA models to assess data-poor fisheries: A case study with a sciaenid fishery off Portugal. *Fishing Bulletin*, 109, 170–185.
- Raco, B., Battaglini, R. and Lelli, M. (2010). Gas emission into the atmosphere from controlled landfills: an example from Legoli landfill (Tuscany, Italy). *Environmental Science and Pollution Research*, 17(6), 1197-1206.
- Radojevic, B. D., Breil, P. and Chocat, B. (2010). Assessing impact of global change on flood regimes. *International Journal of Climate Change Strategies and Management*, 2(2), 167 - 179.
- Rahim, K. A. and Liwan, A. (2012). Oil and gas trends and implications in Malaysia. *Energy Policy*, 50, 262-271.
- Ramanathan, V. and Feng, Y. (2009). Air pollution, greenhouse gases and climate change: Global and regional perspectives. *Atmospheric Environment* 43, 37-50.
- Ramli, N. N., Shamsudin, M. N., Mohamed, Z. and Radam, A. (2012). The impact of fertilizer subsidy on Malaysia paddy/rice industry using a system dynamics approach. *International Journal of Social Science and Humanity*, 2(3), 213-219.
- Reay, D., Smith, P. and van Amstel, A. (2010). *Methane and Climate Change*: Earthscan, London.
- Robinson, D. R., Fernandez, R. and Kantamaneni, R. (2003). Methane emissions mitigation options in the global oil and natural gas industries. Proceedings of the 2003 3rd International Methane & Nitrous Oxide Mitigation Conference Nov. 17-21, 2003.
- Rodríguez, R. and Lombardía, C. (2010). Analysis of methane emissions in a tunnel excavated through Carboniferous strata based on underground coal mining experience. *Tunnelling and Underground Space Technology*, 25(4), 456-468.
- Rojey, A. (2009). *Energy and Climate: how to achieve a successful energy transition*: Society of Chemical Industry and John Wiley & Sons, Ltd.

- Royset, J. O. and Pee, E. Y. (2012). Rate of convergence analysis of discretization and smoothing algorithms for semiinfinite minimax problems. *Journal of Optimization Theory and Applications*, 155(3), 855-882.
- Rupani, P. F., Singh, R. P., Ibrahim, M. H. and Esa, N. (2010). Review of current palm oil mill effluent (POME) treatment methods: Vermicomposting as a sustainable practice. *World Applied Sciences Journal*, 11(1), 70-81.
- Rutishauser, I. H. E. (2005). Dietary intake measurements. *Public Health Nutrition*, 8(7A), 1100-1107.
- Sadr, M. H. and Bargh, H. G. (2012). Optimization of laminated composite plates for maximum fundamental frequency using Elitist-Genetic algorithm and finite strip method. *Journal of Global Optimization*, 54(4), 707-728.
- Saeed, M. O., Hassan, M. N. and Abdul Mujeebu, M. (2009). Assessment of municipal solid waste generation and recyclable materials potential in Kuala Lumpur, Malaysia. *Waste Management* 29, 2209-2213.
- Safaai, N. S. M., Noor, Z. Z., Hashim, H., Ujang, Z. and Talib, J. (2011). Projection of CO₂ emissions in Malaysia. *Environmental Progress & Sustainable Energy*, 30(4), 658-665.
- Safitri, A., Gao, X. and Mannan, M. S. (2011). Dispersion modeling approach for quantification of methane emission rates from natural gas fugitive leaks detected by infrared imaging technique. *Journal of Loss Prevention in the Process Industries*, 24(2), 138-145.
- Saleh, A. F., Kamarudin, E., Yaacob, A. B., Yussof, A. W. and Abdullah, M. A. (2011). Optimization of biomethane production by anaerobic digestion of palm oil mill effluent using response surface methodology. *Asia-Pacific Journal of Chemical Engineering*, from http://eprints.utp.edu.my/6379/1/APJ_550.pdf
- Sass, R. L. (2005). Methane emissions from rice paddies - Factors affecting. In R. Lal (Ed.), *Encyclopedia of Soil Science*: CRC Press.
- Scharff, H. (2005, October 2005). Landfill gas production and emission on former landfills Retrieved October 18, 2011, from <http://www.docstoc.com/docs/43112955/LANDFILL-GAS-PRODUCTION-AND-EMISSION-ON-FORMER-LANDFILLS>

- Scharff, H. and Jacobs, J. (2006). Applying guidance for methane emission estimation for landfills. *Waste Management*, 26, 417–429.
- Scheutz, C., Kjeldsen, P. and Gentil, E. (2009). Greenhouse gases, radiative forcing, global warming potential and waste management — an introduction. *Waste Management & Research*, 27(8), 716-723.
- Schubert, C. J., Diem, T. and Eugster, W. (2012). Methane emissions from a small wind shielded lake determined by eddy covariance, flux chambers, anchored funnels, and boundary model calculations: A comparison. *Environmental Science & Technology*, 46(8), 4515-4522.
- Searle, K. and Gow, K. (2010). Do concerns about climate change lead to distress? *International Journal of Climate Change Strategies and Management*, 2(4), 362 - 379.
- Sejian, V., Lal, R., Lakritz, J. and Ezeji, T. (2011). Measurement and prediction of enteric methane emission. *International Journal of Biometeorology*, 55(1), 1-16.
- Senevirathna, D. G. M., Achari, G. and Hettiaratchi, J. P. A. (2006). A laboratory evaluation of errors associate with the determination of landfill gas emissions. *Canada Journal of Civil Engineering*, 33, 240-244.
- Shafie, S. M., Mahlia, T. M. I., Masjuki, H. H. and Andriyana, A. (2011). Current energy usage and sustainable energy in Malaysia: A review. *Renewable and Sustainable Energy Reviews*, 15(9), 4370-4377.
- Shahabadi, M. B., Yerushalmi, L. and Haghghat, F. (2010). Estimation of greenhouse gas generation in wastewater treatment plants – Model development and application. *Chemosphere* 78, 1085–1092.
- Shibata, M. and Terada, F. (2010). Factors affecting methane production and mitigation in ruminants. *Animal Science Journal* 81, 2–10.
- Shindell, D. T., Faluvegi, G., Koch, D. M., Schmidt, G. A., Unger, N. and Bauer, S. E. (2009). Improved attribution of climate forcing to emissions. *Science*, 326, 716–718.
- Singh, S. N., Tyagi, L. and Tiwari, S. (2009). Attenuating methane emission from paddy fields S. N. Singh (Ed.) *Climate Change and Crops*
- Siraj, M. (2006). Waste reduction: no longer an option bur a necessity *In Bernam*. Kuala Lumpur, Malaysia.

- Smith, S. J. (2003). The evaluation of greenhouse gas indices. *Climate Change*, 58, 261-265.
- Solaimani, K., Mohammadi, H., Ahmadi, M. Z. and Habibnejad, M. (2005). Flood occurrence hazard forecasting based on geographical information system. *International Journal of Environmental Science and Technology* 2(3), 253-258.
- Solomon, S., Plattner, G. K., Knutti, R. and Friedlingstein, P. (2009). Irreversible climate change due to carbon dioxide emissions. *Proceedings of the National Academy of Sciences*, 106(6), 1704-1709.
- Srivastava, A. K. and Nema, A. K. (2008). Forecasting of solid waste composition using fuzzy regression approach: a case of Delhi. *International Journal of Environment and Waste Management*, 2(1-2), 65-74.
- ST. (2010). Suruhanjaya Tenaga. Suruhanjaya Tenaga Annual Report 2009: Suruhanjaya Tenaga (Energy Commission).
- Stepanenko, V., Machul'skaya, E., Glagolev, M. and Lykossov, V. (2011). Numerical modeling of methane emissions from lakes in the permafrost zone. *Izvestiya, Atmospheric and Oceanic Physics*, 47(2), 252-264.
- Su, S., Adhikary, D., Worrall, R. and Gabeva, D. (2009). Study on coal mine methane resources and potential project development *CSIRO Exploration and Mining Report P2009/423* (Vol. P2009/423).
- Su, S. and Agnew, J. (2006). Catalytic combustion of coal mine ventilation air methane. *Fuel*, 85, Issue: 9, Publisher: Elsevier, Pages: 1201-1210(9), 1201-1210.
- Su, S., Han, J., Wu, J., Li, H., Worrall, R., Guo, H., *et al.* (2011). Fugitive coal mine methane emissions at five mining areas in China. *Atmospheric Environment*, 45(13), 2220-2232.
- Sumathi, S., Chai, S. P. and Mohamed, A. R. (2008). Utilization of oil palm as a source of renewable energy in Malaysia. *Renewable and Sustainable Energy Reviews*, 12(9), 2404-2421.
- Szemesova, J. and Gera, M. (2010). Uncertainty analysis for estimation of landfill emissions and data sensitivity for the input variation. *Climatic Change*, 103, 37-54.

- Tanaka, K., O'Neill, B. C., Rokityanskiy, D., Obersteiner, M. and Tol, R. S. J. (2009). Evaluating global warming potentials with historical temperature. *Climatic Change*, 96, 443-466.
- Tarmudi, Z., Abdullah, M. L. and Tap, A. O. M. (2009). An overview of municipal solid wastes generation in Malaysia. *Jurnal Teknologi*, 51(F), 1-15.
- Tchobanoglous, G., Theisen, H. and Vigil, S. A. (1993). *Integrated solid waste management: Engineering principles and management issues*. New York.: McGraw Hill.
- Tecele, D., Lee, J. and Hasan, S. (2009). Quantitative analysis of physical and geotechnical factors affecting methane emission in municipal solid waste landfill. *Environmental Geology* 56, 1135-1143.
- Tey, J. Y.-S., Darham, S., Mohd Noh, A. F. and Idris, N. (2010). Acreage response of paddy in Malaysia. *Agricultural Economics – Czech*, 3, 135-140.
- Tey, Y.-S., Shamsudin, M. N., Mohamed, Z., Abdullah, A. M. and Radam, A. (2008). Demand for meat products in Malaysia: Munich Personal RePEc Archive (MPRA).
- Thoma, E. D., Shores, R. C., Thompson, E. L., Harris, D. B., Thorneloe, S. A., Varma, R. M., *et al.* (2005). Open-path tunable diode laser absorption spectroscopy for acquisition of fugitive emission flux data. *Journal of the Air and Waste Management Association*, 55, 658-668.
- Thompson, S., Sawyer, J., Bonam, R. and Valdivia, J. E. (2009). Building a better methane generation model: Validating models with methane recovery rates from 35 Canadian landfills. *Waste Management*, 29, 2085-2091.
- Tian, G. M., He, Y. F. and Li, Y. X. (2002). Effect of water and fertilization management on emission of CH₄ and N₂O in paddy soil. *Soil and Environmental Sciences*, 11, 294-298.
- Todd, R. W., Cole, N. A., Casey, K. D., Hagevoort, R. and Auvermann, B. W. (2011). Methane emissions from southern High Plains dairy wastewater lagoons in the summer. *Animal Feed Science and Technology*, 166-167(0), 575-580.
- Tol, R. S. J. (1999). The marginal costs of greenhouse gas emissions. *Energy Journal*, 20, 61-81.

- Toraño, J., Torno, S., Menendez, M., Gent, M. and Velasco, J. (2009). Models of methane behaviour in auxiliary ventilation of underground coal mining. *International Journal of Coal Geology* 80, 35–43.
- Towprayoon, S., Smakgahn, K. and Poonkaew, S. (2005). Mitigation of methane and nitrous oxide emissions from drained irrigated rice fields. *Chemosphere*, 59, 1547–1556.
- Observations of climate change: The 2007 IPCC Assessment*, United States House of Representatives (2007).
- Trenberth, K. E. (2010). Global change: The ocean is warming, isn't it? *Nature*, 465(7296), 304-304.
- Tsaur, R.-C. and Kuo, T.-C. (2011). The adaptive fuzzy time series model with an application to Taiwan's tourism demand. *Expert Systems with Applications*, 38(8), 9164-9171.
- Tseng, F.-M. and Tzeng, G.-H. (2002). A fuzzy seasonal ARIMA model for forecasting. *Fuzzy Sets and Systems*, 126(3), 367-376.
- Tsuchiya, T. (2012). Global optimization of polynomial-expressed nonlinear optimal control problems with semidefinite programming relaxation. *Journal of Global Optimization*, 54(4), 831-854.
- Turkdogan-Aydinol, F. I. and Yetilmezsoy, K. (2010). A fuzzy-logic-based model to predict biogas and methane production rates in a pilot-scale mesophilic UASB reactor treating molasses wastewater. *Journal of Hazardous Materials*, 182(1–3), 460-471.
- Tyatyushkin, A. (2011). A multimethod technique for solving optimal control problem. *Optimization Letters*, 1-13.
- Umezawa, T., Aoki, S., Kim, Y., Morimoto, S. and Nakazawa, T. (2011). Carbon and hydrogen stable isotopic ratios of methane emitted from wetlands and wildfires in Alaska: Aircraft observations and bonfire experiments. *Journal of Geophysical Research*, 116(D15), D15305.
- UNFCCC. (1992). United Nations Framework Convention on Climate Change. *Paper presented at the United Nations Conference on Environment and Development (UNCED)*, Rio de Janeiro, Brazil.

- UNFCCC. (2010). Greenhouse gas emission changes in Annex I countries Retrieved June, 25, 2012, from http://unfccc.int/files/inc/graphics/image/jpeg/trends_2010.jpg
- UNFCCC. (2012). Greenhouse gas emissions of Annex I countries Retrieved 25 June, 2012, from http://unfccc.int/ghg_data/ghg_data_unfccc/ghg_profiles/items/4625.php
- USDA. (2012). Malaysia milled rice domestic consumption by year: United States Department of Agriculture.
- USEPA. (1999). U.S. methane emissions 1990–2020: Inventories, projections, and opportunities for reductions: U.S. Environmental Protection Agency (USEPA). EPA 430-R-99-013, Washington, DC. USEPA Office of Air and Radiation.
- USEPA. (2006). Global anthropogenic non-CO₂ greenhouse gas emissions: 1990–2020 Office of Atmospheric Programs, Climate Change Division. US Environmental Protection Agency, Washington D.C.
- USEPA. (2010). Inventory of U.S. greenhouse gas emissions and sinks: 1990 – 2008: U.S. Environmental Protection Agency.
- USEPA. (2011). Inventory of U.S. greenhouse gas emissions and sinks: 1990 – 2009. Washington, DC 20460, USA: U.S. Environmental Protection Agency, EPA 430-R-11-005.
- van Zijderveld, S. M., Dijkstra, J., Perdok, H. B., Newbold, J. R. and Gerrits, W. J. J. (2011a). Dietary inclusion of diallyl disulfide, yucca powder, calcium fumarate, an extruded linseed product, or medium-chain fatty acids does not affect methane production in lactating dairy cows. *Journal of Dairy Science*, 94(6), 3094-3104.
- van Zijderveld, S. M., Fonken, B., Dijkstra, J., Gerrits, W. J. J., Perdok, H. B., Fokkink, W., *et al.* (2011b). Effects of a combination of feed additives on methane production, diet digestibility, and animal performance in lactating dairy cows. *Journal of Dairy Science*, 94(3), 1445-1454.
- Vellinga, T. V., de Haan, M. H. A., Schils, R. L. M., Evers, A. and van den Pol–van Dasselaar, A. (2011). Implementation of GHG mitigation on intensive dairy farms: Farmers' preferences and variation in cost effectiveness. *Livestock Science*, 137(1–3), 185-195.

- Vellinga, T. V. and Hovin, I. E. (2011). Maize silage for dairy cows: mitigation of methane emissions can be offset by land use change. *Nutrient Cycling in Agroecosystems* 89(3), 413–426.
- Vibol, S. and Towprayoon, S. (2010). Estimation of methane and nitrous oxide emissions from rice field with rice straw management in Cambodia. *Environmental Monitoring and Assessment*, 161, 301–313.
- Vijayaraghavan, K. and Ahmad, D. (2006). Biohydrogen generation from palm oil mill effluent using anaerobic contact filter. *International Journal of Hydrogen Energy*, 31, 284–1291.
- VijayaVenkataRaman, S., Iniyana, S. and Goic, R. (2012). A review of climate change, mitigation and adaptation. *Renewable and Sustainable Energy Reviews*, 16(1), 878-897.
- Wang, J., Duan, C., Ji, Y. and Sun, Y. (2010a). Methane emissions during storage of different treatments from cattle manure in Tianjin. *Journal of Environmental Sciences*, 22(10), 1564-1569.
- Wang, J., Sui, J., Guo, L., Karney, B. W. and Jüpner, R. (2010b). Forecast of water level and ice jam thickness using the back propagation neural network and support vector machine methods. *International Journal of Environmental Science and Technology* 7(2), 215-224,.
- Wang, S., Yang, X., Lin, X., Hu, Y., Luo, C., Xu, G., *et al.* (2009). Methane emission by plant communities in an alpine meadow on the Qinghai-Tibetan Plateau: a new experimental study of alpine meadows and oat pasture. *Biology Letters* 5, 535-538.
- Wang, T., Geng, A., Li, X., Wang, H., Wang, Z. and Qiufen, L. (2011). A prediction model of oil cracked gas resources and Its application in the gas pools of Feixianguan Formation in NE Sichuan Basin, SW China. [Research]. *Journal of Geological Research*, 2011, 11 pages, Article ID 592567.
- Wang, Y. H. (2009). Coal demand and coal industry development in China. *Paper presented at the 5th Meeting of the Australia-China Bilateral Dialogue on Resources Cooperation*, Sydney, Australia. 3 December 2009.
- Wangyao, K., Towprayoon, S., Chiemchaisri, C., Gheewala, S. H. and Nopharatana, A. (2010a). Application of the IPCC Waste Model to solid waste disposal

- sites in tropical countries: case study of Thailand. *Environmental Monitoring and Assessment* 164, 249–261.
- Wangyao, K., Yamada, M., Endo, K., Ishigaki, T., Naruoka, T., Towprayoon, S., *et al.* (2010b). Methane generation rate constant in tropical landfill. *Journal of Sustainable Energy & Environment*, 1, 181-184.
- Warmuzinski, K. (2008). Harnessing methane emissions from coal mining. *Process Safety and Environmental Protection*, 86(5), 315-320.
- Weart, S. (2008). *The discovery of global warming*: Harvard University Press.
- Wielenga, K. (2003). Personal communication.
- Wilkinson, P. (2009). Sea level rise could cost port cities \$28 billion, *CNN: Going green*.
- Winiwarter, W., Kuhlbusch, T. A. J., Viana, M. and Hitzenberger, R. (2009). Quality considerations of European PM emission inventories. *Atmospheric Environment*, 43(25), 3819–3828.
- Witting, K., Ober-Blöbaum, S. and Dellnitz, M. (2012). A variational approach to define robustness for parametric multiobjective optimization problems. *Journal of Global Optimization*, 1-15.
- Wong, H.-L., Tu, Y.-H. and Wang, C.-C. (2010). Application of fuzzy time series models for forecasting the amount of Taiwan export. *Expert Systems with Applications*, 37(2), 1465-1470.
- Wu, T. Y., Mohammad, A. W., Jahim, J. M. and Anuar, N. (2010). Pollution control technologies for the treatment of palm oil mill effluent (POME) through end-of-pipe processes. *Journal of Environmental Management*, 91, 1467–1490.
- Xiaoli, C., Ziyang, L., Shimaoka, T., Nakayama, H., Ying, Z., Xiaoyan, C., *et al.* (2010). Characteristics of environmental factors and their effects on CH₄ and CO₂ emissions from a closed landfill: An ecological case study of Shanghai. *Waste Management*, 30(3), 446–451.
- Yacob, S., Hassan, M. A., Shirai, Y., Wakisaka, M. and Subash, S. (2005). Baseline study of methane emission from open digesting tanks of palm oil mill effluent treatment. *Chemosphere*, 59(11), 1575-1581.
- Yacob, S., Hassan, M. A., Shirai, Y., Wakisaka, M. and Subash, S. (2006a). Baseline study of methane emission from anaerobic ponds of palm oil mill effluent treatment. *Science of The Total Environment*, 366(1), 187-196.

- Yacob, S., Shirai, Y., Hassan, M. A., Wakisaka, M. and Subash, S. (2006b). Start-up operation of semi-commercial closed anaerobic digester for palm oil mill effluent treatment. *Process Biochemistry*, 41(4), 962-964.
- Yan, X., Yagi, K., Akiyama, H. and Akimoto, H. (2005). Statistical analysis of the major variables controlling methane emission from rice fields. *Global Change Biology*, 11(7), 1131-1141.
- Yang, X., Shang, Q., Wu, P., Liu, J., Shen, Q., Guo, S., *et al.* (2010). Methane emissions from double rice agriculture under long-term fertilizing systems in Hunan, China. *Agriculture, Ecosystems & Environment*, 137(3-4), 308-316.
- Yechiel, A. and Shevah, Y. (2012). Optimization of energy costs for SWRO desalination plants. *Desalination and Water Treatment*, 46 (1-3), 304-311.
- Yetilmezsoy, K. and Demirel, S. (2008). Artificial neural network (ANN) approach for modeling of Pb(II) adsorption from aqueous solution by *Antep pistachio* (*Pistacia Vera L.*) shells. *Journal of Hazardous Materials*, 153(3), 1288-1300.
- Yu, T. H.-K. and Huarng, K.-H. (2010). A neural network-based fuzzy time series model to improve forecasting. *Expert Systems with Applications*, 37(4), 3366-3372.
- Yuan, W.-L., Cao, C.-G., C-F., L., Zhan, M., Cai, M.-L. and Wang, J.-P. (2009). Methane and nitrous oxide emissions from rice-duck and rice-fish complex ecosystems and the evaluation of their economic significance. *Agricultural Sciences in China*, 42(6), 2052-2060.
- Yusof, N. M., Abdu-Rashid, R. S. and Mohamed, Z. (2010). Malaysia crude oil production estimation: An application of ARIMA model. Proceedings of the 2010 *International Conference on Science and Social Research (CSSR 2010)*. Dec 5 - 7, 2010. Kuala Lumpur, Malaysia, 1255-1259.
- Yusuf, R. O., Noor, Z. Z., Abba, A. H., Hassan, M. A. A. and Din, M., F.M. (2012a). Greenhouse gas emissions: Quantifying methane emissions from livestock. *American Journal of Engineering and Applied Sciences*, 5(1), 1-8.
- Yusuf, R. O., Noor, Z. Z., Abba, A. H., Hassan, M. A. A. and Din, M. F. M. (2012b). Methane emission by sectors: A comprehensive review of emission sources and mitigation methods. *Renewable and Sustainable Energy Reviews*, 16(7), 5059-5070.

- Zalilah, M. S., Khor, G. L., Mirnalini, K., Norimah, A. K. and Ang, M. (2006). Dietary intake, physical activity and energy expenditure of Malaysian adolescents. *Singapore Medical Journal*, 47(6), 491-498.
- Zavala, V. M., Constantinescu, E. M., Krause, T. and Anitescu, M. (2009). On-line economic optimization of energy systems using weather forecast information. *Journal of Process Control*, 19(10), 1725-1736.
- Zeng, Y., Trauth, K. M., Peyton, R. L. and Banerji, S. K. (2005). Characterization of solid waste disposed at Columbia Sanitary Landfill in Missouri. *Waste Management & Research*, 23, 62–71.
- Zhang, B. and Chen, G. Q. (2010). Methane emissions by Chinese economy: Inventory and embodiment analysis. *Energy Policy*, 38(8), 4304-4316.
- Zhang, G. P. (2003). Time series forecasting using a hybrid ARIMA and neural network model. *Neurocomputing*, 50(0), 159-175.
- Zhang, L., Yu, D., Shi, X., Weindorf, D. C., Zhao, L., Ding, W., *et al.* (2009). Simulation of global warming potential (GWP) from rice fields in the Tai-Lake region, China by coupling 1:50,000 soil database with DNDC model. *Atmospheric Environment* 43, 2737–2746.
- Zhao, X., He, J. and Cao, J. (2011). Study on mitigation strategies of methane emission from rice paddies in the implementation of ecological agriculture. *Energy Procedia*, 5(0), 2474-2480.
- Zhou, M., Hernandez-Sanabria, E. and Guan, L. L. (2009). Assessment of the microbial ecology of ruminal methanogens in cattle with different feed efficiencies. *Applied Environmental Microbiology*, 75(20), 6524-6533.
- Zinatizadeh, A. A. L., Mohamed, A. R., Abdullah, A. Z., Mashitah, M. D., Isa, M. H. and Najafpour, G. D. (2006). Process modeling and analysis of palm oil mill effluent treatment in an up-flow anaerobic sludge fixed film bioreactor using response surface methodology (RSM). *Water Research*, 40(17), 3193-3208.