

Rogala Dorota, Kornowska Joanna, Ziolkowska Mirosława. Clitoridectomy, excision, infibulation- female circumcision ritual and its consequences for women's health. *Journal of Education, Health and Sport*. 2018;8(11):583-593. eISSN 2391-8306. DOI

<http://dx.doi.org/10.5281/zenodo.2533136>

<http://ojs.ukw.edu.pl/index.php/johs/article/view/6451>

<https://pbn.nauka.gov.pl/sedno-webapp/works/896357>

The journal has had 7 points in Ministry of Science and Higher Education parametric evaluation. Part B item 1223 (26/01/2017).
1223 Journal of Education, Health and Sport eISSN 2391-8306 7

© The Authors 2018:

This article is published with open access at Licensee Open Journal Systems of Kazimierz Wielki University in Bydgoszcz, Poland
Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license Share alike.
(<http://creativecommons.org/licenses/by-nc-sa/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 26.11.2018. Revised: 30.11.2018. Accepted: 30.11.2018.

Clitoridectomy, excision, infibulation- female circumcision ritual and its consequences for women's health

Dorota Rogala ¹, Joanna Kornowska ², Mirosława Ziolkowska ³

¹ Department of Oncology, Radiotherapy and Gynecologic Oncology, Faculty of Health Sciences, Collegium Medicum, Nicolaus Copernicus University, Toruń, Poland.

² Department of Health Care R. Czerwiakowski in Brodnica, Poland.

³ Department of Psychiatric Nursing, Department of Conservative Nursing. Faculty of Health Sciences, Collegium Medicum, Nicolaus Copernicus University, Toruń, Poland.

Abstract

The issue of female mutilation is important for modern gynecology and obstetrics. The article discusses the concepts of ritual, tradition and initiation as the moment of introduction into the world of adulthood. The approach to circumcision was explained for the sake of religion. The essence of the sense of one's sexuality and femininity is also shown. Describes the definition of female genital mutilation as a procedure contributing to the partial or complete elimination of the external genital organs in women. The paper presents recommendations for perinatal care for a circumcised patient developed by the Royal College of Obstetricians and Gynecologist, the South Australian Perinatal Practice Guidelines and the London Safeguarding Children Board. The perspectives of protecting and stopping female mutilation practices that violate a number of human rights are explained.

Key words: FMG, female circumcision, clitoridectomy, infibulation, excision.

Introduction

Female genital mutilation (Female Genital Mutilation/Circumcision), also known as circumcision, includes procedures that contribute to the partial or complete removal of external genital organs in women, or other damage to female genitalia for non-medical reasons [1]. Definition (abbreviated FGM/C or FGM) was first adopted by the African Committee for Traditional Practices Affecting Women's and Children's Health in 1990 to highlight the gender-specific nature of female genital mutilation [2].

In 1995 the WHO has developed a terminology (the last update of 2007) according to which four types of FGM are distinguished:

I - clitoridectomy/sunnic circumcision - partial or complete removal of the entire clitoris, clitoris glans and / or foreskin,

II - excision - resection of clitoris and labia minora with or without labia majora,

III - infibulation/pharaonic circumcision - narrowing of the vaginal opening by incision and compression of the labia majora and / or labia minora, with excision or leaving the clitoris. In most cases, the edges remaining after excision of the labia majora are stitched, leaving a small gap. The skin fold completely covers the urethral meatus, completely or partially obscures the entrance of the vagina. To enable sexual intercourse or childbirth, the scar must be open.

IV - so-called symbolic circumcision - all harmful procedures regarding mutilation of female genitalia for purposes other than medical, for example piercing, scraping, cutting, burning, stretching, rubbing caustic substances, herbs, etc. (Fig.1) [1].




Figure 1. Illustration of unaltered external female genitalia and female genital mutilation/cutting (FGM/C) types I-III. From top left: unaltered external female genitalia, type I (clitoridectomy), type II (excision) and type III (infibulation).

The course of circumcision

Girls are most often circumcised before the age of 10 (6-12 years). The role of tools is played by pieces of glass, razors, scissors, knives, razor blades, can covers, etc. The procedure is performed by a woman specializing in circumcision, a woman/grandmother/a charlatan from a village who has a high social position. Other women help immobilize the girl, which in turn leads to fractures of the collarbone and thigh bone. In the hole left after the procedure, a plastic straw or match is introduced to prevent the complete fusion during healing and allow urine output after the procedure, and in the future, outflow of menstrual blood. To allow the labia to adhere to each other and can even grow together, the girl is bound with a rope or rags for a period of about 1 month. A paste of myrrh is put on the wound that stops bleeding and compress of herbs, mud, cow excrements [2-4].

Epidemiology

It is estimated that the number of women circumcised around the world is about 125 – 140 million, of which there are about 500.000 in the EU. Each year, another 180 thousand girls are at risk FGM [5].

According to UNICEF data from 2013. the highest percentage of circumcised women comes from Africa. Prevails Somalia- 98% (type III circumcision 63%), Guinea- 96%, Djibouti 93% and Egypt 91%; Eritrea, Malaysia, Sierra Leone and Sudan-88%. Rarely FGM is conducted in Iraq, Ghana and Togo - below 10%, in Nigeria, Cameroon and Uganda - about 1-2% [2]

Consequences of FGM

On the basis of the available studies, there is clear evidence that female genital mutilation has negative consequences for the physical, mental and social health of a woman who has performed the procedure as an infant or child. However, accurate determination of the frequency of complications is not possible [6].

The most frequently mentioned consequences of FGM include:

1. Short-term complications after the procedure:

- bleeding, haemorrhage
- damage to the urethra, bladder, anal sphincter, vagina,
- infection (tetanus, gas gangrene, hepatitis type C, HIV),
- septic shock,
- fracture of the femur and collarbone etc. [7-11].

2. Long-term consequences after the procedure:

- keloids,
- fistulas (recto-vaginal and bladder-vaginal),
- recurrent urinary tract infections,
- bacterial vaginosis,
- chronic pain [7,9,11-25].

3. Obstetric complications:

- prolonged delivery,
- higher percentage of instrumental deliveries,
- higher percentage of cesarean sections,
- extensive crotch injuries,
- bleeding,
- increased risk of shoulder dystocia [14, 15, 23, 25, 27-49].

4. Consequences for sexual functioning

- dyspareunia,
- vaginismus,
- contact bleeding,
- reduced vaginal lubrication,
- inability to penetrate due to scarring or as a consequence of untreated inflammations in the pelvic area [13, 16-18, 23, 50].

In addition to physical consequences, circumcised women have problems with mental and social functioning. Such behaviors as apathy, constant tension, fear, guilt, shame, feelings of helplessness, nightmares, isolation from the environment. There is often a frustration and anger towards men [51].

Among the socio-cultural consequences, there is a risk of marginalization of the position of an uncircumcised woman and even exclusion from society [52].

Obstetrics and gynecology care

In recent years, there has been an increasing migration of people from countries where FGM is practiced. There is a need to develop procedures for dealing with women after circumcision in the aspect of gynecological care, and especially perinatal care. Teachers, pedagogues, basic health care doctors who have contact with a migrant girl can also play an

important role. The answer to two questions: which girl is burdened with the risk of FGM and what may suggest that circumcision has already taken place will allow indicate a group of children requiring special attention and care.

To the endangered procedure of FGM may belong female child, who:

- comes from countries with a high FGM level,
- is born in a family in which women have been subjected to this procedure,
- have mother, sister / sisters after the surgery,
- it is isolated from the local/pre-school/school community,
- defends against understanding her body and her basic rights,
- drew the attention of the educator / doctor to overheard conversations of adults about the "ceremony", "special procedure", which awaits them in the near future,
- she says about a trip to his home country, where the FGM ratio is high [53].

You can suspect the circumcision:

- problems with sitting or long walks,
- problems with physical exercises,
- spending more time in the toilet,
- frequent absences from school caused by painful menstruation, diseases of the urinary tract,
- symptoms suggestive of depression or other psychiatric disorders [53].

Most often, the first contact of a circumcised woman with a gynecologist or midwife takes place during pregnancy. An interview may take place in the presence of an interpreter, provided that he is not a family member, nor is he not related to the family, community or country of the patient [54].

According to the recommendations of the Royal College of Obstetricians and Gynaecologists [55], the London Safeguarding Children Board [56] and South Australian Perinatal Practice Guidelines [57] in the gynecological examination should identify changes in the woman's genitals, determine the type of FGM and describe in the documentation and inform the woman about the need for deinfibulation.

Deinfibulation/defibulation is the surgical opening of the genitalia covered by the skin fold. They are performed only in case of infibulated patients - type III. In the case of pregnancy under gynecological care, it is recommended to perform deinfibulation in anesthesia, in hospital conditions about 20-30 weeks of pregnancy. In the situation when the

diagnosis of FGM occurs in a hospital at the time of delivery, deinfibulation should be performed in the first stage, or in the second stage of delivery.

Childbirth should be carried out by a qualified midwife; the presence of apprentices/ students is not recommended. In the puerperal period, gynecological consultation is recommended 2 weeks after delivery (assessment of crotch condition), avoiding sexual intercourse until full healing, return to physical activity depending on the condition of the crotch. Every woman should get an educational package regarding Kegel muscle exercises recommendations. The midwife should be informed about the circumcised woman. The patient and her family should also be informed about the harmfulness of the use of circumcision in the case of the birth of a female child and of the law consequences associated with it.

Perspectives to protect and stop female mutilation practices

A circumcised patient is not only a challenge for healthcare professionals. Female genital mutilation is an act of violence against women, which is opposed by leading international organizations.

The first law against FGM was adopted in Sweden in 1982 [58]. Following it went the United Kingdom [59], Belgium [60], France [61], Germany [62], Canada [63].

The European Parliament in the resolution of 2001: *The resolution on female genital mutilation* radically opposes the practice of FGM, claiming that it is a violation of fundamental human rights. Stresses the essence of the harmonious action of European states, opposes the medication of FGM, urges "prosecution, condemnation and punishment of conducting these practices", recognizing them as a separate offense and "punishing anyone who helps, encourages, advises and provides support to anyone to conduct any from these acts on the body of a woman or a girl " with the possibility of punishing parents who, for the procedure, went with their child outside the border of the inhabited state [64].

General Assembly of the United Nations December 20, 2012 has adopted a resolution to intensify global forces to eliminate the practice of female mutilation. This resolution recommends a coordinated approach to the subject, and above all developing uniform methods for collecting data on all forms of violence and discrimination against girls, especially undocumented forms such as FGM [2].

November 26, 2006 the most influential Islamic clerics - the Great Sheikh Al-Azhar and the Great Mufti of Egypt - announced that FGM is a practice that is not a reflection of the

religion of Islam. This is a big step towards change, because religion is one of the most common arguments in justifying the continued practice of FGM [64].

February 6 was declared "International Day of Non-Tolerance for Female Genital Mutilation" and November 29 "An International Day Against Female Genital Mutilation" [64].

References

1. WHO. Eliminating female genital mutilation: an interagency statement. Geneva: World Health Organization 2008,1-47.
2. Report of UNICEF: United Nations Children's Fund, Female Genital Mutilation/Cutting: A statistical overview and exploration of the dynamics of change. UNICEF 2013,1-194.
3. Henrion R. Mutilations génitales féminines, mariages forcés et grossesses précoces. Academie Nationale Médecine 2003,1051-66.
4. Chmielowska D, Grabowska B, Machut-Mendecka E. Być kobietą w Oriencie, Wydawnictwo Akademickie Dialog, Warszawa 2008.
5. <http://www.uefgm.org> – [data dostępu 01.02.2018]
6. Berg RC, Underland V, Odgaard- Jensen J, et al. Effects of female genital cutting on physical health outcomes: a systematic review and metaanalysis. *BMJ Open* 2014;4: e006316. doi:10.1136/bmjopen-2014-006316.
7. El Dareer A. Complications of female circumcision in the Sudan. *Trop Doct* 1983;13:131–3.
8. Ministère de l'Économie, du Plan et de la Coopération Internationale, Macro International Inc. Enquête démographique et de santé, République Centrafricaine 1994–95. Calverton, Maryland: Direction des Statistiques Démographiques et Sociales, Macro International Inc, 1995.
9. Lovel H, Gettigan C, Mohammed Z, et al. A systematic review of the health complications of female genital mutilation including sequelae in childbirth. *WHO* 2000,1-181.
10. Institut National de la Stastique, de Études Économiques et Démographiques (INSEED). Ministère de l'Économie, du Plan et dela Coopération, ORC Macro. Enquête démographique et de santé Tchad 2004. Calverton, Maryland: INSEED, ORC Macro, 2005.
11. Diouf K, Nour N. Female Genital Cutting and HIV transmission: is there an association? *AM J Reprod Immunol* 2013;69,1:45-50.

12. Shandall AA. Circumcision and infibulation of females: a general consideration of the problem and a clinical study of the complications in Sudanese women. *Sudan Med J* 1967;5:178–212.
13. Rushwan H, Slot CJM, El Dareer A, et al. Female circumcision in the Sudan. Prevalence, complications, attitudes and change. A report of a study conducted by the faculty of medicine, University of Khartoum, Sudan (1977–1982). Khartoum: University of Khartoum, 1983.
14. De Silva S. Obstetric sequelae of female circumcision. *Eur J Obstet Gynecol Reprod Biol* 1989;32:233–40.
15. Jones H, Diop N, Askew I, et al. Female genital cutting practices in Burkina Faso and Mali and their negative health outcomes. *Stud Fam Plann* 1999;30:219–30.
16. el Defrawi MH, Lotfy G, Dandash KF, et al. Female genital mutilation and its psychosexual impact. *J Sex Marital Ther* 2001;27:465–73.
17. Morison L, Scherf C, Ekpo G, et al. The long-term reproductive health consequences of female genital cutting in rural Gambia: a community-based survey. *Trop Med Int Health* 2001;6:643–53.
18. Okonofu FE, Larsen U, Oronsaye F, et al. The association between female genital cutting and correlates of sexual and gynaecological morbidity in Edo State, Nigeria. *BJOG* 2002;109:1089–96.
19. Msuya SE, Mbizvo E, Hussain A, et al. Female genital cutting in Kilimanjaro, Tanzania: changing attitudes? *Trop Med Int Health* 2002;7:159–65.
20. Almroth L, Bedri H, El Musharaf S, et al. Urogenital complications among girls with genital mutilation: a hospital based study in Khartoum. *Afr J Reprod Health* 2005;9:118–24.
21. Elmusharaf S, Elkhidir I, Hoffmann S, et al. A case-control study on the association between female genital mutilation and sexually transmitted infections in Sudan. *BJOG* 2006;113:469–74.
22. Nwajei SD, Otiono AI. Female genital mutilation: implications for female sexuality. *Women Stud Int Forum* 2003;26:575–80.
23. Elnashar A, Abdelhady R. The impact of female genital cutting on health of newly married women. *Int J Gynaecol Obstet* 2007;97:238–44.
24. Fillo GF, Leone T. Female genital cutting, reproductive tract infections and perinatal outcome in Burkina Faso. Paper presented 2007-3-29 at the Annual Meeting of the Population Association of America; New York.

25. Browning A, Allsworth JE, Wall LL. The relationship between female genital cutting and obstetric fistulae. *Obstet Gynecol* 2010;115:578–83.
26. Kaplan A, Hechavarría S, Martin M, et al. Health consequences of female genital mutilation/cutting in the Gambia, evidence into action. *Reprod Health* 2011;8:26.
27. Berardi JC, Teillet JF, Godard J, et al. Consequences obstétricales de l'excision féminine. Etude chez 71 femmes africaines excisées. *J Gynecol Obstet Biol Reprod* 1985;14:743–6.
28. Bohoussou KM, Anongba S, Djanhan Y, et al. Complications gynécologiques médicales et obstétricales de l'excision rituelle. *Afr Med* 1986;25:160–2.
29. Adinma JI. Current status of female circumcision among Nigerian Igbos. *West Afr J Med* 1997;16:227–31.
30. National Statistics Office (NSO). Macro International. Eritrea demographic and health survey 1995. Calverton, Maryland: NSO, Macro International, 1997.
31. Lupo VR, Marcotte KL. Obstetric complications of Somali female circumcision. *Obstet Gynecol* 1999;93:19S.
32. Hakim LY. Impact of female genital mutilation on maternal and neonatal outcomes during parturition. *East Afr Med J* 2001;78:255–8.
33. Larsen U, Okonofua FE. Female circumcision and obstetric complications. *Int J Gynaecol Obstet* 2002;77:255–65.
34. Slinger TE, Snow RC, Okonofua FE. The impact of female genital cutting on first delivery in southwest Nigeria. *Stud Fam Plann* 2002;33:173–84.
35. Vangen S, Stoltenberg C, Johansen RE, et al. Perinatal complications among ethnic Somalis in Norway. *Acta Obstet Gynecol Scand* 2002;81:317–22.
36. National Statistics and Evaluation Office (NSEO). ORC Macro. Eritrea demographic and health survey 2002. Calverton, Maryland: NSEO, ORC Macro, 2003.
37. Essen B, Sjöberg NO, Gudmundsson S, et al. No association between female circumcision and prolonged labour: a case control study of immigrant women giving birth in Sweden. *Eur J Obstet Gynecol Reprod Biol* 2005;121:182–5.
38. Johnson EB, Reed SD, Hitti J, et al. Increased risk of adverse pregnancy outcome among Somali immigrants in Washington state. *Am J Obstet Gynecol* 2005;193:475–82.
39. Oduro A, Ansah P, Hodgson A, et al. Trends in the prevalence of female genital mutilation and its effect on delivery outcomes in the Kassena-Aankana district of northern Ghana. *Ghana Med J* 2006;40:87–92.
40. Orji EO, Babalola A. Correlates of female genital mutilation and its impact on safe motherhood. *JTGGA* 2006;7:319–24.

41. Banks E, Meirik O, Farley T et al. Female genital mutilation and obstetric outcome: WHO collaborative prospective study in six African countries. *Lancet* 2006;367:1835–41.
42. Yount KM, Carrera JS. Female genital cutting and reproductive experience in Minya, Egypt. *Med Anthropol Q* 2006;20:182–211.
43. Brewer DD, Potterat JJ, Roberts JM Jr, et al. Male and female circumcision associated with prevalent HIV infection in virgins and adolescents in Kenya, Lesotho, and Tanzania. *Ann Epidemiol* 2007;17:217–26.
44. Millogo-Traore F, Kaba ST, Thieba B, et al. Pronostic maternel et foetal au cours de l'accouchement chez la femme excisee. *J Gynecol Obstet Biol Reprod* 2007;36:393–8.
45. Yount KM, Abraham BK. Female genital cutting and HIV/AIDS among Kenyan women. *Stud Fam Plann* 2007;38:73–88.
46. Small R, Gagnon A, Gissler M, et al. Somali women and their pregnancy outcomes postmigration: data from six receiving countries. *BJOG* 2008;115:1630–40.
47. Ndiaye P, Diongue M, Faye A, et al. Mutilation genitale feminine et complications de l'accouchement dans la province de Gourma (Burkina Faso). *Sante Publique* 2010;22:563–70.
48. Chibber R, El-Saleh E, El-Harmi J. Female circumcision: obstetrical and psychological sequelae continues unabated in 21st century. *The Journal of Maternal-Fetal and Neonatal Medicine* 2011;24,6:833-6.
49. Rigmor CB. *The Obstetric Consequences of FGM: A Systematic Review and Meta-Analysis*. Hindawi Publishing Corporation, Obstetrics and Gynecology International 2013,1-15.
50. Odoi A, Brody SP, Elkins TE. Female genital mutilation in rural Ghana, West Africa. *Int J Gynaecol Obstet* 1997;56:179–80.
51. Vloeberghs E, Kwaak A, Knipscheer J, et al. Copic and chronic psychosocial consequences of female genital mutilation in the Netherlands. *Ethnicity & Health* 2012;17,6, 677-95.
52. Shabila N, Saleh A, Jawad R. Women's perspectives of FGM: Q-methodology. *Biomed Central* 2014, 1-21.
53. <http://www.gov.uk> - HM Government, Multi-Agency Practice Guidelines: Female Genital Mutilation, 2011 – [data dostępu 10.03.2018]
54. Department of Gender, Women and Health (GWH) Department of Reproductive Health and Research (RHR) Family and Community Health (FCH) and World Health

- Organization (WHO), Management of pregnancy, childbirth and the postpartum period in the presence of female genital mutilation, publication of WHO Switzerland, 2001,1-48.
55. Royal College of Obstetricians and Gynaecologists, Female Genital Mutilation and its Management, RCOG Green-top Guideline, 2009;53:1-17.
 56. London Safeguarding Children Board, London Female Genital Mutilation – Resource Pack, Globe Academy Southwark, 2009,1-58.
 57. South Australian Perinatal Practice Guidelines, Female Genital Mutilation, Department of Health, Department of South Australia, 2015,1-9.
 58. Johndotter S, The FGM Legislation Implemented: Experiences from Sweden, Malmo University, 2009,1-12.
 59. <http://www.legislation.gov.uk/asp/2005/8> - [data dostępu 12.03.2018]
 60. <http://www.hsph.harvard.edu/population/fgm/Belgium.fgm.00.htm> - [data dostępu 12.03.2018]
 61. <http://www.hsph.harvard.edu/population/fgm/france.fgm.htm> - [data dostępu 12.03.2018]
 62. <http://www.bundesgesundheitsministerium.de/fileadmin/dateien/Downloads/G/Genitalverstueummelung/Recommendations-Patients-History-Genital-Mutilation.pdf> - [data dostępu 22.03.2018]
 63. Vissandjee B, Denetto S, Migliardi P, et al. FGC and the ethics of care: community engagement and cultural sensivity at the interface of migration experiences. Biomed Central 2014,1-19.
 64. Kane J. Szkodliwe praktyki tradycyjne, Program Daphne Komisji Europejskiej DG ds. Sprawiedliwości, Wolności i Bezpieczeństwa, Bruksela, 2008,1-24.