

Szafraniec-Porada Aneta, Porada Dominik, Konopelko Michał, Przybylska Daria, Puculek Małgorzata. The influence of selenium on thyroid gland. *Journal of Education, Health and Sport*. 2018;8(9):1501-1508. eISSN 2391-8306. DOI <http://dx.doi.org/10.5281/zenodo.1435937>  
<http://ojs.ukw.edu.pl/index.php/johs/article/view/6111>

The journal has had 7 points in Ministry of Science and Higher Education parametric evaluation. Part B item 1223 (26/01/2017).  
1223 Journal of Education, Health and Sport eISSN 2391-8306 7

© The Authors 2018;

This article is published with open access at Licensee Open Journal Systems of Kazimierz Wielki University in Bydgoszcz, Poland  
Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license Share alike. (<http://creativecommons.org/licenses/by-nc-sa/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 22.08.2018. Revised: 28.08.2018. Accepted: 15.09.2018.

## The influence of selenium on thyroid gland

Aneta Szafraniec-Porada<sup>1</sup>, Dominik Porada<sup>1</sup>, Michał Konopelko<sup>2</sup>, Daria Przybylska<sup>2</sup>  
Małgorzata Puculek<sup>3</sup>

<sup>1</sup> Chair and Department of Endocrinology, Medical University of Lublin

<sup>2</sup> Chair and Department of Hygiene, Medical University of Lublin

<sup>3</sup> Chair and Department of Human Anatomy, Medical University of Lublin

### Abstract

The proper functioning of the thyroid gland depends on many factors, which include adequate supply of various trace elements, including iodine, but also selenium or iron. The purpose of this article was to attempt to summarize the current knowledge on the effects of selenium on the thyroid gland and the clinical implications associated with it.

Selenium affects the homeostasis of the entire human body, and its main function in the thyroid gland is to counteract damage of oxidative origin. This microelement is part of selenoproteins, which include glutathione peroxidase and thioredoxin reductase. Both selenium deficiency and excess can negatively affect the thyroid gland. The research results show that selenium supplementation may bring benefits in some morbidities, eg Graves-Basedow disease. Currently, I advise recommending semi-annual selenium supplementation in a mild form of thyroid orbitopathy. However, no clinical benefit has been demonstrated in increasing the supply of selenium in the diet of other patients with thyroid disease, including Hashimoto's disease. In conclusion, despite the numerous studies already conducted, further work is necessary to fully explain the mechanisms of action of this element on the thyroid

gland, but also to deepen knowledge about the benefits and risks of selenium supplementation in various types of thyroid structural disorders and, most importantly, to establish safe guidelines for possible treatment.

**Key words:** selenium, thyroid gland, selenoblasts, thyroid hormones

## **Introduction**

Selenium (Se) is a trace element necessary to maintain homeostasis of the body. It is a component of many proteins, the so-called selenoproteins, including: glutathione peroxidase, iodothyronine 5 thiodiodate, thioredoxin reductase, selenoprotein P or selenioproprotein W [1-3]. Demand for selenium differs depending on age and sex, and increases during pregnancy and lactation. The currently recommended daily dose for healthy adults should not exceed 70 µg. Increased Se consumption is allowed, however higher doses may be toxic. The range of normal selenium concentration in the blood serum ranges between 75-140 µg / l. [2-6]. In many areas of the world there is hyposelenic soil, including in Poland, which results in insufficient amount of this element in food products, both animal and vegetable [7].

**Purpose of work:** This article is an attempt to summarize current knowledge about the influence of selenium on the thyroid gland. The analysis of scientific literature concerned both the deficiency and excess selenium in the body and possible supplementation opportunities.

## **The role of selenoblasts in the thyroid gland**

The highest concentration of selenium is observed in the thyroid gland, even when it is deficient in the body, which confirms its important role in the proper functioning of this gland [8-10]. Among the most important selenoid proteins found in the thyroid gland are: glutathione peroxidase, thioredoxin reductase and iodothyronine deiodinase. Thyroid peroxidase uses hydrogen peroxide to oxidize iodide ions during the production of thyroid hormones, decomposing it into water and oxygen, thus protecting thyrocytes against free radicals. Reactive oxygen species in the thyroid are also neutralized by thioredoxin reductase, which is an electron donor for other oxidoreductive enzymes. On the other hand,

iodothyronine deiodinase plays a role in the thyroid hormone metabolism by transforming thyroxine (T4) to triiodothyronine (T3) and reverse triiodothyronine (rT3) [ 2.3, 11-16].

### **Selenium deficiency**

Selenium deficiency is widespread and affects different populations in many countries of the world [17]. The effects of deficiency of this element affect the entire human body, and result with disorders associated with proteins containing selenium [18]. Selenium deficiency may result in an increase in incidences of cancer (including colon, thyroid, prostate, lung and breast) or cardiovascular morbidities (heart failure, arrhythmia, dilated cardiomyopathy) [19, 20]. It can also lead to the progression of many inflammatory diseases (including Kashin-Beck's disease, systemic lupus erythematosus) and metabolic (diabetes), as well as worsening existing degenerative diseases of the central nervous system [21-25]. Additionally, in the case of selenium deficiency, the risk of strokes, male infertility, nephropathy increases and the symptoms of autoimmune diseases may increase [26-28]. It was found that low selenium concentration correlates with an increased risk of thyroid disease [29]. Inadequate supply of selenium may cause disorders of thyroid hormone synthesis, both as a consequence of abnormalities associated with the functioning of selenoproteins, as well as by disturbances of the metabolism of iodine in the body [27].

### **Excess of selenium**

Increased selenium intake may also have negative health effects. Toxic action of high doses of selenium consists of inactivation of proteins responsible for the repair of damaged DNA and may lead to increased production of free radicals [30]. Mild symptoms of selenium intoxication are often non-specific, and the most common include malaise, gastrointestinal complaints (such as nausea, vomiting or diarrhea), as well as neurological disorders [31, 32]. In the case of chronic consumption of high doses of selenium, both through diet and dietary supplements / drugs may lead to selenosis, which results in liver damage and damage to the hematopoietic system, as well as skin lesions and serious dysfunctions of the nervous system [31, 33]. Excess selenium may also result in disruption of thyroid hormone synthesis and increase the risk of developing type 2 diabetes [31, 34].

### **Effect of selenium on thyroid disease**

Disorders of the structure and / or function of the thyroid gland are one of the most common endocrine diseases. Adequate supply of selenium affects the proper functioning of the thyroid by optimizing the synthesis and metabolism of thyroid hormones, but also reducing the production of pro-inflammatory factors, including proinflammatory cytokines, leukotrienes or prostaglandins. As previously stated, it protects the thyroid against oxidative stress. In addition, due to its regulatory effect on the immune system, both T and B lymphocytes cause a decrease in thyroid antibodies [35]. It was found that selenium deficiency promotes thyroid volume increase (in women) [36], as well as the development of focal lesions in the thyroid gland [37]. However, the importance of the supply of selenium in the context of thyroid cancer is mixed [38, 39]. In the literature on the subject, attention is drawn to the significant effect of selenium on thyroid diseases of autoimmune etiology. It has been demonstrated that the supply of selenium suppresses the immune cellular response, which is important for the occurrence and severity of chronic lymphocytic thyroiditis [40]. In patients with Hashimoto's disease, there is a significant reduction in selenium concentration compared to a healthy population [41]. In numerous studies on the effect of selenium supplementation in people with autoimmune thyroiditis, it has been demonstrated that the use of selenium preparations results in the lowering of anti-TPO antibody concentration and often anti-TG antibodies and improvement of thyroid gland echogenicity. The best effects were observed with a dose of 200 µg / day [42-46]. However, a meta-analysis carried out by van Zuuren et al. showed that despite current research, there is currently no clear scientific evidence for the effectiveness of selenium supplementation in Hashimoto's disease, therefore there is no indication for the general use of selenium preparations in this group of patients, and any possible treatment with selenium preparations should be approached with caution [47].

In patients with Graves' disease, lower blood selenium levels were also observed in comparison with the control group. In addition, in a state of remission the level of this microelement is significantly higher than in patients with overt hyperactivity [48, 49]. The use of selenium supplements in this group of patients tended to normalize the concentration of free thyroid hormones and TSH and lower the concentration of antibodies against the receptors for TSH, and results in a higher remission rate [50]. The best documented and currently recommended medical indication for selenium supplementation is mild thyroid orbitopathy. It was found that the use of increased doses of selenium causes regression of ailments and prevents the progression of eye changes in Graves' disease, which was reflected

in the recommendations of the European Thyroid Society, which recommends 6-month selenium therapy (200 µg in a divided dose) [51, 52]. Selenium therapy cannot replace thyrostatic treatment, but selenium supplementation during treatment may shorten the time needed for clinical and biochemical euthyroidism [53-55].

### Summary

Selenium is a trace element, for which both excess and deficiency result in disorders of the body's homeostasis, in particular thyroid function. Selenium supplementation in areas with deficiency of this element brings benefits to patients with mild orbitopathy in the course of Graves-Basedow disease. Currently, there is no reliable scientific data on the legitimacy of selenium supplementation in other thyroid diseases. The use of this element cannot replace the previously used pharmacological treatment, but the possibility of adding selenium preparations is still an interesting therapeutic option in thyroid diseases. One should remember about the negative impact of large quantities of selenium on the human body. Further research is necessary to obtain reliable data that will allow a full explanation of the mechanisms of action of this element on the thyroid gland in pathological conditions, but also to establish safe and effective guidelines for possible treatment.

### References

1. Kryukov G.V. et al. Characterization of mammalian selenoproteomes. *Science*, 2003; 300: 1439-1443.
2. Lobanov A.V., Hatfield D.L., Gladyshev V.N. Eukaryotic selenoproteins and selenoproteomes. *Biochim. Biophys. Acta*, 2009; 1790: 1424-1428
3. Imai H., Nakagawa Y. Biological significance of phospholipid hydroperoxide glutathione peroxidase (PHGPx, GPx4) in mammalian cells. *Free Radic. Biol. Med.*, 2003; 34: 145-169.
4. Jarosz M., Bułhak-Jachymczyk B. Normy żywienia człowieka. Podstawy prewencji otyłości i chorób niezakaźnych. [Wydawnictwo Lekarskie PZWL](#) 2008.
5. Kieliszek M, Błażej S. Selenium: Significance, and outlook for supplementation. *Nutrition*. 2013; 29(5): 713–718, 5.
6. Slencu BG, Ciobanu C, Cuciureanu R. Selenium content in foodstuffs and its nutritional requirement for humans. *Clujul Med.* ; 2012: 139–145
7. Combs G.F.Jr. Selenium in global food systems. *Br. J. Nutr.*, 2001; 85: 517-547.
8. Cardoso B.R. et al. Nutritional status of selenium in Alzheimer's disease patients. *Br. J. Nutr.*, 2010; 103: 803-806.
9. Tiran B., Karpf E., Tiran A. Age dependency of selenium and cadmium content in human liver, kidney, and thyroid. *Arch. Environ. Health*, 1995; 50: 242-246.

10. Katoh Y., Sato T., Yamamoto Y. Determination of multielement concentrations in normal human organs from the Japanese. *Biol.Trace Elem. Res.*, 2002; 90: 57-70.
11. Talas Z.S. et al. Antioxidative role of selenium against the toxic effect of heavy metals (Cd<sup>2+</sup>, Cr<sup>3+</sup>) on liver of rainbow trout (*Oncorhynchus mykiss* Walbaum 1792). *Fish Physiol. Biochem.*, 2008; 34: 217-222.
12. Mondal S., Raja K., Schweizer U., Mugesh G. Chemistry and Biology in the Biosynthesis and Action of Thyroid Hormones. *Angew Chem Int Ed Engl.* 2016 Jun 27;55(27):7606-30.
13. Lacka K., Szeliga A. Significance of selenium in thyroid physiology and pathology. *Pol Merkur Lekarski.* 2015 Jun;38(228):348-53.
14. Drutel A., Archambeaud F., Caron P. Selenium and the thyroid gland: More good news for clinicians. *Clin. Endocrinol.*, 2013; 78: 155-164.
15. Arnér E.S., Holmgren A. Physiological functions of thioredoxin and thioredoxin reductase. *Eur. J. Biochem.*, 2000; 267: 6102-6109.
16. Behne D, Kyriakopoulos A, Meinhold H, et al. Identification of type I iodothyronine 5'-deiodinase as a selenoenzyme. *Biochem Biophys Res Commun.* 1990; 173(3): 1143-1149.
17. Combs G.F.Jr. Selenium in global food systems. *Br. J. Nutr.*, 2001; 85: 517-547.
18. Pedrero Z, Madrid Y. Novel approaches for selenium speciation in foodstuffs and biological specimens: a review. *Anal Chim Acta.* 2009; 634(2): 135-152.
19. Micke O., Schomburg L., Buentzel J., Kisters K., Muecke R. Selenium in oncology - an update. *Trace Elem. Electrol.*, 2010; 27: 250-257.
20. Yoo M.H., Xu X.M., Carlson B.A., Gladyshev V.N., Hatfield D.L. Thioredoxin reductase 1 deficiency reverses tumor phenotype and tumorigenicity of lung carcinoma cells. *J. Biol. Chem.*, 2006; 281: 13005-13008.
21. Chen J. An original discovery: selenium deficiency and Keshan disease (an endemic heart disease). *Asia Pac J Clin Nutr.* 2012; 21(3): 320-326.
22. Vunta H. et al. The anti-inflammatory effects of selenium are mediated through 15-deoxy-delta12,14-prostaglandin J2 in macrophages. *J. Biol. Chem.*, 2007; 282: 17964-17973.
23. Zhang S., Rocourt C., Cheng W.H. Selenoproteins and the aging brain. *Mech. Ageing. Dev.*, 2010; 131: 253-260.
24. Hill K.E., Zhou J., McMahan W.J., Motley A.K., Burk R.F. Neurological dysfunction occurs in mice with targeted deletion of the selenoprotein P gene. *J. Nutr.*, 2004; 134: 157-161.
25. Cardoso B.R. et al. Nutritional status of selenium in Alzheimer's disease patients. *Br. J. Nutr.*, 2010; 103: 803-806.
26. Ruseva B, Himcheva I, Nankova D. Importance of selenoproteins for the function of the thyroid gland. *Medicine.* 2013; 3: 60-64.
27. Roman M, Jitaru P, Barbante C. Selenium biochemistry and its role for human health. *Metallomics.* 2014; 6(1): 25-54,
28. Mistry HD., Broughton Pipkin F., et al. Selenium in reproductive health. *Am J Obstet Gynecol.* 2012; 206(1): 21-30.
29. Wilmar M. Wiersinga.: Clinical Relevance of Environmental Factors in the Pathogenesis of Autoimmune Thyroid Disease. *Endocrinol Metab* 2016 Forthcoming.
30. Letavayová L, Vlasáková D, Spallholz J, et al. Toxicity and mutagenicity of selenium compounds in *Saccharomyces cerevisiae*. *Mutation Research/ Fundamental and Molecular Mechanisms of Mutagenesis.* 2008; 638(1-2): 1-10,

31. Navarro-Alarcon M, Cabrera-Vique C. Selenium in food and the human body: a review. *Sci Total Environ.* 2008; 400(1-3): 115–141.
32. Fordyce F. Selenium Deficiency and Toxicity in the Environment. *Essentials of Medical Geology.* 2012: 375–416.
33. Li S, Xiao T, Zheng B. Medical geology of arsenic, selenium and thallium in China. *Sci Total Environ.* 2012; 421-422: 31–40.
34. Drutel A, Archambeaud F, Caron P. Selenium and the thyroid gland: more good news for clinicians. *Clin Endocrinol (Oxf).* 2013; 78(2): 155–164.
35. Schomburg L. Selenium, selenoproteins and the thyroid gland: interactions in health and disease. *Nat Rev Endocrinol.* 2011; 8(3): 160–171,
36. Derumeaux H, Valeix P, Castetbon K, et al. Association of selenium with thyroid volume and echostructure in 35- to 60-year-old French adults. *Eur J Endocrinol.* 2003; 148(3): 309–315.
37. Rasmussen LB, Schomburg L, Köhrle J, et al. Selenium status, thyroid volume, and multiple nodule formation in an area with mild iodine deficiency. *Eur J Endocrinol.* 2011; 164(4): 585–590
38. Shen F, Cai WS, Li JL, et al. The Association Between Serum Levels of Selenium, Copper, and Magnesium with Thyroid Cancer: a Meta-analysis. *Biol Trace Elem Res.* 2015; 167(2): 225–235.
39. Baltaci AK, Dundar TK, Aksoy F, et al. Changes in the Serum Levels of Trace Elements Before and After the Operation in Thyroid Cancer Patients. *Biol Trace Elem Res.* 2017; 175(1): 57–64
40. Tan L, Sang ZNa, Shen J, et al. Selenium supplementation alleviates autoimmune thyroiditis by regulating expression of TH1/TH2 cytokines. *Biomed Environ Sci.* 2013; 26(11): 920–925
41. Socha K., et al. Nawyki żywieniowe a stężenie selenu w surowicy u pacjentów z chorobą Hashimoto, *Probl Hig Epidemiol* 2012, 93(4): 824-827.
42. Vasiliu I., et al. SELENIUM STATUS IN AUTOIMMUNE THYROIDITIS. *Rev Med Chir Soc Med Nat Iasi.* 2015 Oct-Dec;119(4):1037-44.
43. Krysiak R., Okopien B.: The effect of levothyroxine and selenomethionine on lymphocyte and monocyte cytokine release in women with Hashimoto's thyroiditis. *J. Clin. Endocrinol. Metab.*, 2011; 96: 2206-2215.
44. Nacamulli D., et al. Influence of physiological dietary selenium supplementation on the natural course of autoimmune thyroiditis. *Clin. Endocrinol.*, 2010; 73: 535-539.
45. Turker O., Kumanlioglu K., Karapolat I. Selenium treatment in autoimmune thyroiditis: 9-month follow-up with variable doses. *J. Endocrinol.*, 2006; 190: 151-156.
46. Gärtner R., Gasnier B.C., Dietrich J.W., Krebs B., Angstwurm M.W. Selenium supplementation in patients with autoimmune thyroiditis decreases thyroid peroxidase antibodies concentrations. *J. Clin. Endocrinol. Metab.*, 2002; 87: 1687-1691.
47. van Zuuren EJ, Albusta AY, Fedorowicz Z, Carter B, Pijl H. Selenium supplementation for Hashimoto's thyroiditis, *Cochrane Library* 2013, Issue 6.
48. Reglinski J., Smith W., Wilson R., Halls D., McKillop J., Thomson J. Selenium in Graves' disease. *Clin. Chim. Acta*, 1992; 211: 189-190.
49. Wertenbruch T, et al. Serum selenium levels in patients with remission and relapse of graves' disease. *Med. Chem.*, 2007; 3: 281-284.
50. Wang L., et al. Effect of Selenium Supplementation on Recurrent Hyperthyroidism Caused by Graves' Disease: A Prospective Pilot Study. *Horm Metab Res.* 2016 Jul 8.

51. Dehina N., Hofmann PJ., Behrends T., Eckstein A., Schomburg L. Lack of Association between Selenium Status and Disease Severity and Activity in Patients with Graves' Ophthalmopathy. *Eur Thyroid J.* 2016 Mar;5(1):57-64.
52. Bartalena L. et al. European Group on Graves' Orbitopathy (EUGOGO).: The 2016 European Thyroid Association/European Group on Graves' Orbitopathy Guidelines for the Management of Graves' Orbitopathy. *Eur Thyroid J.* 2016 Mar;5(1):9-26.
53. Guerra L., et al. Antioxidants in the treatment of Graves' disease. *IUBMB Life*, 2001; 51: 105-109.
54. Roy G., Muges G. Selenium analogues of antithyroid drugs – recent developments. *Chem. Biodivers.*, 2008; 5: 414-439.
55. Vrca V.B., Skreb F., Cepelak I., Romic Z., Mayer L. Supplementation with antioxidants in the treatment of Graves' disease; the effect on glutathione peroxidase activity and concentration of selenium. *Clin. Chim. Acta*, 2004; 341: 55-63.