

UPLanD

Journal of Urban Planning, Landscape & environmental Design



Research & experimentation
Ricerca e sperimentazione

THE ALGAL DIVERSITY IN THE PHLEGREAN FIELDS (CAMPANIA, ITALY) ARCHEOLOGICAL DISTRICTS: AN OVERVIEW

Paola Cennamo^a, Claudia Ciniglia^b

^a *Facoltà di Lettere, University of Naples, Suor Orsola Benincasa, IT*

^b *Department of Environmental, Biological and Pharmaceutical Science and Technology, University of Campania Luigi Vanvitelli, Caserta, IT*

HIGHLIGHTS

- Microbial communities in Phlegrean Fields.
- Different microhabitats in Phlegrean Fields.
- Algal diversity in the archaeological and geothermal areas.

ABSTRACT

The coexistence of several ecological conditions (hot springs, streams, mud, rock walls), characterized by different pH, temperature values, water potential and mineralogical makes the Phlegrean Fields as an ideal area for the analysis of the microalgal assemblages and their relationships with the different microhabitats occurring in the sites. Here we report an overview of the algal diversity within the archaeological and geothermal areas of Phlegrean Fields and the adaptability for most of them to acidic and thermal conditions in the volcanic soils.

ARTICLE HISTORY

Received: February 12, 2017
Reviewed: March 19, 2017
Accepted: June 05, 2017
On line: July 31, 2017

KEYWORDS

Phlegrean Fields
Hot springs
Archaeological district
Extreme environments
Microbial communities

1. INTRODUCTION

Study on the biogeography of species provide clues on those mechanisms which are at the base of the genesis and the maintenance of biodiversity, like speciation, extinction, dispersion and interactions among species. The geographic distribution of plants and animals has been study object for the biologists since 1900 while research on biogeography of microorganisms is more recent. The application of more advanced biomolecular methodologies has allowed to analyze the real biodiversity of a microbial community. The small dimensions, the high abundance and the metabolic plasticity of the unicellular organisms would confer them a very wide dispersion capacity, not affected by ecological or geographic parameters. The endolithic and cryptoendolithic environments are ubiquitous habitat for microorganisms. These microorganisms form an important interface between biological and geological ecosystems. The biogeochemical characteristics, such as the physical and chemical properties of the rocks, climate, direction of exposure, influence the specific microbial composition of the endolithic biofilms (Walker and Pace, 2007). Algae are the most frequent organisms found on rock surfaces, after cyanobacteria and bacteria. However, microorganisms are able to obtain different elements for their metabolism by biosolubilization of the rock material. Such microbial biosolubilization involves the production of organic and inorganic acids by the metabolic activity of algae, lichens, fungi and bacteria. Phlegrean Fields represent an important archeological and volcanic area situated northwest of Naples, Italy, of undeniable fascination; one of the most famous archeological site of Cuma, and one of the forty volcanoes located in Phlegrean Fields, the worldwide known Solfatara and Pisciarelli; it is an ancient volcanic crater still active but quiescent that for nearly two millennia retains activity of sulphur dioxide fumaroles, boiling mud jets and high soil temperature (Chidini et al. 1996). What makes these two Phlegrean Fields even more fascinating, along with their history, is that both of them preserve the most ancient life forms which are at the base of the photosynthetic life. At the same time, considering that the algae living in these habitats cannot be considered solely as a threat, but also as an important source of biodiversity. In this review we reported the microalgal flora occurring in some Phlegrean sites, namely, Pisciarelli (Agnano, Italy), Sybil Cave (Cuma, Italy), Piscina Mirabilis (Bacoli, Italy), along with the type of mineral substratum, and exploited to examine the physico-chemical interactions between microorganisms and substrate and to provide an evaluation of the biodeteriogenic and/or bioprotective action carried out by the microorganisms based on experimental measurements, occurrence locations and references. The data gathered together here will be useful for the study of biological colonization in archaeological district and extreme environmental, and should provide a basis for ecological studies.

2. ENVIRONMENTS

2.1 Volcanic district

Solfatara and Pisciarelli (Figure 1) are a hydrothermal system located in the central part of the Campi Flegrei caldera (Ciniglia et al 2009). At Solfatara, volcanites are basically represented by incoherent products often with trachytic occurrences deposited on the trachytic lavas of Monte Olibano dated before 4000 years (Di Girolamo et al 1984). The geothermal system is vapour-dominant, with the hottest fumaroles reaching temperatures of 140-160°C, according to Cioni et al. (1984), Bolognesi et al. (1986) and Chiodini et al. (1996).



Figure 1: Pisciarelli, Phlegrean Fields. *Source: photo by Claudia Ciniglia*

2.2 Archeological district

A wide research programme on the archaeological area of Southern Italy has revealed a valuable cultural heritage of monuments built in tuff rock. Tuff is a type of rock consisting of consolidated volcanic ash ejected from vents during a volcanic eruption (Sgobbo & Moccia, 2016). This type of rock has a high porosity and a high water capacity. These characteristics make it a perfect substrate for the colonization and the formation of biofilms. Two archeological sites, the Sybil Cave and the Piscina Mirabilis (Figure 2), are artificial caves dug in the yellow tuff and used during antiquity for various purposes (Mangoni & Sgobbo, 2013). Phlegrean Fields has been inhabited since prehistoric times and became an important centre of civilization during antiquity. Different underground habitats have been dug in the district, and two of them have remarkable dimensions. The most ancient is the so-called Sybil Cave (fourth century BCE), near Cuma, which is a long gallery cut into the soft tuff, lighted by multiple openings to the surface. The other large cave (dating back to the first century CE) is the Piscina Mirabilis, located in Bacoli, a cistern excavated in tuff, used as storage for drinking water. In addition, both caves were dug in the same geological material originated from the eruption of the Neapolitan Yellow Tuff (dated at 15 ka BP; Insinga et al., 2004), which produced the caldera collapse of the bay of Pozzuoli (Scarpati et al., 1993).

2.3 Mineralogical analysis

The products of the Phlegraean activity range in composition from K-basalt to phonolite: K-basalt (3%), trachybasalt (8%), latite (10%), trachyte (27%), alkali-trachyte (45%) and phonolite (7%) (Di Girolamo et al., 1984). The geothermal system is liquid-dominant with spring water temperatures up to 90°C. The geothermal waters are fed by steam which has a composition similar to that of the fumaroles emerging in the Solfatara crater (Chiodini et al. 1996). In both localities, gaseous sulphur in the fluids is totally represented by H₂S. Other gases in Solfatara fumaroles include major CO₂ and minor N₂, H₂, CH₄, Ar and He (Dall' Aglio et al., 1972; Tedesco et al., 1990; Martini et al., 1991; Allard et al., 1991; Chiodini et al., 1996).




Figure 2: (A) Sybil Cave ; (B) Piscina Mirabilis; (C) exemple of algal communities. Paola Cennamo & Claudia Ciniglia

3. ALGAL COMMUNITIES

Table 1 and 2 lists the microorganisms present and isolated in the Phlegrean area, together with the substratum. A total of 28 species were identified by molecular methods (18S rDNA); the largest group was represented by Cyanobacteria, followed by Chlorophyta, Rhodophyta and Bacillariophyta (Ciniglia et al. 2005; Ciniglia et al. 2007, Pollio et al 2005; Cennamo et al 2012).

Tabel 1: List of prokaryota microorganism in the sampling sites of the Phlegrean Fields

Species	Sampling Sites	Reference
PROKARYOTA		
CYANOPHYTA		
<i>Aphanotecae</i> sp.	Sybil cave	Ciniglia et al. 2005; Cennamo et al. 2012
<i>Chroococcus lithophilus</i>	Sybil cave	Ciniglia et al. 2005; Cennamo et al. 2012
<i>Chroococcus varius</i>	Sybil cave; Piscina Mirabilis	Ciniglia et al. 2005; Cennamo et al. 2012
<i>Gloeocapsa</i> sp.	Sybil cave	Ciniglia et al. 2005; Cennamo et al. 2012
<i>Leptolyngbya foveolarum</i>	Piscina Mirabilis	Ciniglia et al. 2005; Cennamo et al. 2012
<i>Leptolyngbya fragilis</i>	Piscina Mirabilis	Ciniglia et al. 2005; Cennamo et al. 2012
<i>Lyngbya</i> sp.	Piscina Mirabilis	Ciniglia et al. 2005; Cennamo et al. 2012
<i>Nostoc commune</i>	Piscina Mirabilis	Ciniglia et al. 2005; Cennamo et al. 2012
<i>Oscillatoria</i> sp.	Sybil cave; Piscina Mirabilis	Ciniglia et al. 2005; Cennamo et al. 2012

3.1 Cyanophyta

Aphanotecae sp. is a coccoid cyanobacterium growing in the litholitic substratum along with *Chroococcus lithophilus*, *Chroococcus varius* and *Gloeocapsa* sp.; the genus *Leptolungbya* was the most widespread filamentous cyanobacterium, the species isolated were *Leptolungbya foveolarum*, *Leptolungbya fragilis*, *Lyngbya* sp. Other cyanobacteria were *Nostoc commune*, *Oscillatoria* sp., *Synechococcus bacillaris*, *Synechococcus* sp. All these cyanobacterial species were able to colonize different stone substrata and are typical of many archeological sites (Cennamo et al. 2009; Cennamo et al. 2012; Cennamo et al. 2013; Cennamo et al. 2015). Since these genera are ubiquitous and therefore their presence is not strictly related to a specific lithic substratum or climate, all reference reported in the literature did not establish a clear relationship between organisms and the substratum. The tuff stone is considered as the best substratum for the growth of the cyanobacteria because of the porosity, roughness, highly hygroscopic, and capillary water absorption, which strongly influence water availability for micro-organisms. All these cyanobacteria are unable to grow on scarcely porous substrata and a low pH typical of acidic fumarole sites.

3.2 Bacillariophyceae

There are many studies on diatoms that live in acidic environments, Negro reported the presence of *Nitzschia* and *Pinnularia* near fumaroles where the temperature reached 50 ° C. *Pinnularia obscura* Krasske is frequently present in acidic environments (De Nicola D., 2000). In Italy the presence of *Pinnularia* has been repeatedly reported in acidic areas of a different nature (Pinto et al., 1992; Ciniglia et al. 2007). In recent studies Cennamo et al. 2015 examined the biofilm of the volcanic substrata (tuff and Piperno) and species belonging to *Pinnularia obscura* and *Navicula* sp. were abundant.

3.3 Chlorophyta

The Chlorophyta are the most abundant algal group present in the Italian acidic environment (Pinto et al. 1992, Huss et al. 2001; Pollio et al. 2005); *Chlorella saccharophila* var. *saccharophila* (Chlorococcales), *Auxenochlorella protothecoides* var. *acidicola*, (Huss et al. 2001); *Viridiella fridericiana* (Albertano et al., 1991, Huss et al. 2001), *Pseudococcomyxa simplex* (Albertano et al., 1990), *Stichococcus bacillaris* (Pinto et al. 1992; Cennamo et al. 2012), all these species are also present in the Phlegrean Fields. *Stichococcus* sp., *Gloeocystis* sp. *Scenedesmus arcuatus* and *Scotiellopsis terrestris* (Cennamo et al. 2012) are present only in the archeological area. *Chlorella* and *Pseudococcomyxa* are ubiquitous algae able to live both on the soil near fumaroles and in the archeological areas. Among the Chlorophyta, *Chlamydomonas acidophila* Negro and *Dunaliella acidophila*, were frequently isolated from thermoacidic sites (Pollio et al. 2005).

3.4 Chrysophyta

Ochromonas, along with *Chlamydomonas*, is one of the algal species commonly colonizing acidic lakes, where in many cases represent more than 50% of the entire algal population. The genus *Ochromonas* (Chrysophyta) includes at least 70 species widely distributed in heterotrophic waters (Ettl et al., 1978; Ettl et al., 1999). *Ochromonas* spp. cells can contain bacterial cells thus acting as mixotrophic and phagotrophic organisms; many genera sampled in acidic lakes in Lusatia (Germany) were described as phagotrophic (Sanders & Porter, 1988). *Ochromonas* species was detected on volcanic soils from Kurili Island, at pH 1.0 and 42°C, successively identified as *O. vulcania*. The presence of *Ochromonas* sp. in Italian acidic soils was reported for the first time in 1994 by Albertano et al. Morphological and

ecophysiological analyses successively confirmed the identification of this strain as *Ochromonas vulcania* (Gromov et al. 1991).

3.5 Rhodophyta

One of the most intriguing groups of photosynthetic microorganisms is represented by Cyanidiophyceae (Rhodophyta), a group of red unicellular algae perfectly suited to the environmental extremes offered by the volcanic and post volcanic areas, where temperatures rise above 50°C, and high sulphuric acid concentrations, generated by the oxidation of sulphur gaseous emissions, greatly reduce the pH to values prohibitive for the majority of eukaryotic life forms (pH 0.5-3.0). Phylogenetic tools, and genome analyses of different algal taxa have shown that Cyanidiophyceae represent one of the ancient and largest algal groups which have diversified; their long evolutionary history began around 1.5 MYA, just before the formation of the supercontinent Rodinia (1.2 MYA), which resulted in an increase in volcanic activity that would have favored the diversification and dispersal of these thermoacidophilic algae.

Table 2: List of eukaryota microorganism in the sampling sites of the Phlegrean Fields

Species	Sampling Sites	Reference
EUKARYOTA		
BACILLARIOPHYTA		
<i>Navicula</i> sp.	Sybil cave; Piscina Mirabilis	Cennamo et al. 2012; Ciniglia et al. 2005
<i>Pinnularia obscura</i>	Sybil cave ; Pisciarelli	Cennamo et al. 2012; Cennamo et al. 2015; Ciniglia et al. 2005; Ciniglia et al. 2007; Pinto et al., 1992
CHLOROPHYTA		
<i>Chlorella</i> sp.	Sybil cave; Piscina Mirabilis	Cennamo 2005; Cennamo et al. 2012; Ciniglia et al. 2005
<i>Chlorella saccharophila</i> var. <i>saccharophila</i>	Sybil cave; Piscina Mirabilis; Pisciarelli	Cennamo et al. 2012; Cennamo et al. 2015; Ciniglia et al. 2005
<i>Auxenochlorella protothecoides</i> var. <i>acidicola</i>	Pisciarelli	Ciniglia et al. 2005; Huss et al. 2001
<i>Chlamydomonas acidophila</i>	Pisciarelli	Pollio et al. 2005
<i>Dunaliella acidophila</i>	Pisciarelli	Ciniglia et al. 2005
<i>Gloeocystis</i> sp.	Sybil cave; Piscina Mirabilis	Cennamo et al. 2012; Ciniglia et al. 2005
<i>Pseudococcomyxa simplex</i>	Sybil cave; Piscina Mirabilis; Pisciarelli	Albertano et al. 1990; Cennamo et al. 2012; Ciniglia et al. 2005
<i>Stichococcus</i> sp.	Sybil cave; Piscina Mirabilis	Cennamo et al. 2012; Ciniglia et al. 2005
<i>Stichococcus bacillaris</i>	Pisciarelli	Pinto et al. 1992
<i>Scenedesmus arcuatus</i>	Sybil cave; Piscina Mirabilis	Cennamo et al. 2012; Ciniglia et al. 2005
<i>Scotiellopsis terrestris</i>	Sybil cave; Piscina Mirabilis	Cennamo et al. 2012; Ciniglia et al. 2005
<i>Viridiella fridericiana</i>	Pisciarelli	Albertano et al. 1991; Huss et al. 2001
CHRYSOPHYCEAE		
<i>Ochromonas</i> sp.	Pisciarelli	Albertano et al. 1994
RHODOPHYTA		
<i>Cyanidium chilense</i>	Sybil cave; Piscina Mirabilis	Cennamo et al. 2012
<i>Cyanidioschzoon merole</i>	Pisciarelli	Ciniglia et al. 2005; Pinto et al. 2003; Pinto et al. 2006
<i>Galdieria sulphuraria</i>	Pisciarelli	Ciniglia et al. 2005; Pinto et al. 2003; Pinto et al. 2006
<i>Cianidium caldarium</i>	Pisciarelli	Ciniglia et al. 2005; Pinto et al. 2003; Pinto et al. 2006
<i>Phragmonema sordidum</i>	Pisciarelli	Cennamo et al. 2012

Surprisingly, around 600 million years ago, a mesophilic lineage, named as *Cyanidium chilense*, evolved from the extremophilic ones, colonizing caves, which can be considered as extreme environments, as well, where nutrient input, light intensity, temperature and humidity are limiting factors for microorganisms. Members of the genus *Cyanidium* have been frequently found in caves worldwide occurring, and enthusiastically it has been also detected inside the archeological park of Cuma (Naples, Italy), specifically in the Sybil cave (Cennamo et al. 2012). *Cyanidium chilense* lives intermingled with a microbial community composed by bacteria, cyanobacteria, green and brown unicellular algae and fungi colonizing both the tuff cave walls of Sybil Cave and the tracheite cave walls and the natural fumarole caves in Solfatara (Ciniglia et al. 2005; Cennamo et al. 2012).

The hydrothermal system of Pisciarelli is located on the eastern edge of the Solfatara crater in the central part of the Phlegrean Fields Caldera (Napoli, Italy), characterized by thermal pools fed by meteoric waters intersected by fumarolic gases rising from a deep boiling aquifer. The gaseous sulphur in the fumarolic fluids is totally represented by H₂S. The microbial community is composed of different groups of prokaryotes, such as Archaea (Kvist et al., 2005), chemotrophic sulphur bacteria, anoxygenic phototrophic bacteria and eukaryotic algae (Brock, 1978); these latter are mainly represented by Cyanidiophytina assemblages, growing diffusely over the entire site. The sulphur fumes favored the development of a large, thick algal mat characterized by algal communities in which only members of the Cyanidiophytina were observed; green patches, formed exclusively by Cyanidiophytina, were distributed along the station at temperatures ranging from 33° C to 45°C and at very low pH values (0.5–1.5). Among them, the relative abundance of *Galdieria sulphuraria* varied from about 20% to 50%, whereas *Cyanidium caldarium* was the dominant species, occurring at a relative percentage of 48–67%. Finally *Cyanidioschizon merolae* was the least represented species in the Cyanidiophytina assemblages (Pinto et al. 2003; Pinto et al. 2006).

4. CONCLUSION

Microbial communities are selected by their habitat and by the interactions within the community. They are typically characterized by few dominating organisms and are often formed in extreme environments, acidic hydrothermal habitats being a classical example. A rich nutrient or energy source is a prerequisite for communities formation, and the Phlegrean area these are present. Lithological formation of Pisciarelli/Solfatara area are the result of the water-rock interaction and the microorganisms-rock interaction. The water-rock interaction is enhanced by high temperatures and low pHs of the fluids and strongly influenced by the direct or indirect mediation of microorganisms. The process causes a colonization only of thermophilic and acidophilic organisms can be considered the most important agents of rocks alteration. Cryptoendolithic algae, exclusively made of *Galdieria*, collected in the altered side of local volcanic rocks. As previously reported (Hoffmann, 1989; Bell, 1993; Gross et al., 1998), the cryptoendolithic growth is probably a strategy adopted by the algae to avoid desiccation, thanks to the alunite-opale layers of the altered trachytic volcanic rock in which only few fissures are present. *Dunaliella* has been found in this site since 1970 and in no other sites; probably, the absence of pools and water stream with variable pH values have given such stable environmental conditions that allowed *Dunaliella* to persist in this site, while *Ochromonas* has never been found.

The interaction between microorganisms (algae and cyanobacteria) and vulcanites or volcanic rocks (tuff) is also detectable in the archeological sites. The major microbial biodiversity on the tuff rock depends on the climatic parameter, are susceptible of variations as a consequence of weather seasonal fluctuations. Here, "opportunistic" green unicellular algae grow. The development of extremely different biological communities has been observed in different microenvironments: from the

extremophilic algae "sensu stricto", -Cyanidiophytina, Diatoms and flagellate green unicellular algae (*Chlamydomonas*, *Dunaliella*)- to more tolerant ones (Chlorophyceae).

Certain organisms found only in geothermally heated environments are quite possibly relicts of the earliest biosphere that have remained essentially unchanged by the vagaries of time and evolution for more than three billion years. Several extreme thermophiles have optimal growth rates near the boiling point of water. Some are found in geothermal aquifers hundreds of meters below the surface, or in deep oceanic volcanic vents. algal communities. Subsequently, sampling point sites will be identified where the algal communities will be collected.

REFERENCES

- Albertano, P., Pollio, A., & Taddei, R. (1991). Viridiella fridericiana (Chlorococcales, Chlorophyta), a new species isolated from extremely acid environments. *Phycologia* 30(4), 346-354.
- Albertano, P., Pinto, G., Pollio, A., & Taddei, R. (1990). Morphology, Ultrastructure and ecology of an acidophilic alga *Pseudococcomyxa Simplex* (Mainx) Fott (Chlorococcales). *Archiv Fur Hydrobiologie, Algological Studies*, 37, 401- 408.
- Albertano, P., Pollio, A., & Pinto, G. (1994). Ecophysiology and Ultrastructure of an Acidophilic Species of *Ochromonas* (Chrysophyceae, Ochromonadales). *Archiv für Protistenkunde* , 144, 75–82.
- Allard, P., Maiorani, A., Tedesco, D., Cortecchi, G., & Turi, B. (1991). Isotopic study of the origin of sulfur and carbon in Solfatara fumaroles, Campi Flegrei caldera. *Journal of Volcanology and Geothermal Research*, 48, 139-159.
- Bell, R. A. (1993). Cryptoendolithic algae of hot semiarid land and deserts. *Journal of Phycology*, 29, 133–139.
- Bolognesi, L., Noto, P., & Nuti, S. (1986). Studio chimico della Solfatara di Pozzuoli: ipotesi sulle origini e sulle temperature profonde dei fluidi. *Rendiconti della Societa Italiana di Mineralogia e Petrologia*, 42, 281-295.
- Brock, T.D. (1978). The genus *Cyanidium*. In: P.M. Starr (Ed), *Thermophilic microorganisms and life at high temperatures*. (pp. 255- 301) New York, USA: Springer-Verlag
- Cennamo, P., & De Luca, P. (2004). *Phragmonema sordidum* Zopf var. *desanctisianum* Cennamo et De Luca (Phragmonemataceae, Porphydiales), a new red algal variety from tuff cisterns. *Delpinoa*, 46,17-22.
- Cennamo, P. (2005). Caratterizzazione morfologica e molecolare degli organismi algali presenti all'interno del Parco Archeologico di Paestum. *Informatore Botanico Italiano*, 37.
- Cennamo, P., Marzano, C., Giorgio, A., Caputo, P., & Moretti, A. (2009). Bioincrustations in the church of saint Maria of Piedigrotta (Pizzo Calabro, Italy). *Science and technology for the Safeguard Cultural Heritage in the Mediterranean Basin:Session C1 -Genetics, Etno Anthropological Heritage*, I, 304-308.
- Cennamo, P., Marzano, C., Ciniglia, C., Pinto, G., Cappelletti, P., Caputo, P., & Pollio, A. (2012). A survey of the algal flora of anthropogenic caves of Campi Flegrei (Naples, Italy) archeological district. *Journal Of Caves And Karst Studies*,74(3),243-250. doi: 10.4311/ 2011JCKS0194
- Cennamo, P., Caputo, P., Giorgio, A., Moretti, A., & Pasquino, N. (2013). Biofilms on Tuff Stones at Historical Sites: Identification and Removal by Nonthermal Effects of Radiofrequencies. *Microbial Ecology*, 66 (3), 659-668.doi:10.1007/s00248-013-0247-7
- Chiodini, G., Cioni, R., Magro, G., Marini, L., Panichi, C. Raco, B., Russo, M., & Taddeucci, G. (1996). Chemical and isotopic variation of Bocca Grande Fumarole (Solfatara Volcano, Phlegraena Fields). *Acta Vulcanologica*, 8, 129-138.
- Ciniglia, C., Yoon, H.S., Pollio, A., Pinto, G., & Bhattacharya, D. (2004). Hidden biodiversity of the extremophilic Cyanidiales red algae. *Molecular Ecology*, 13, 18-27.doi:10.1111/j.1365-294X.2004.02180.x

- Ciniglia, C., Valentino, G. M., Cennamo, P., De Stefano, M., Stanzione, D., Pinto, G., & Pollio, A. (2005). Influences of geochemical and mineralogical constraints on algal distribution in acidic hydrothermal environments: Pisciarelli (Naples, Italy) as a model site. *Archiv für Hydrobiologie*, *162*, 121-142. doi: 10.1127/0003-9136/2005/0162-0121
- Ciniglia, C., Cennamo, P., De Stefano, M., Pinto, G., Caputo, P., & Pollio, A. (2007). Pinnularia species (Bacillariophyceae, Bacillariophyta) from acidic environments: morphological, ultrastructural and phylogenetic characterization. *Fundamental And Applied Limnology*, *170*, 29-47. doi:10.1127/1863-9135/2007/0170-0029
- Ciniglia, C., Cennamo, P., Pollio, P., & Pinto, G. (2009). Phylogenetic relationships and taxonomic position of Dunaliella acidophila isolates from acidic and thermal environments. *Phycologia*, *48*, 21-22.
- Cioni, R., Corazza, E., & Marini, L. (1984). Gas steam ratio as indicator of heat transfer at Solfatarà fumaroles, Phlegraean Fields (Italy). *Bulletin of Volcanology*, *47* (2), 295-302.
- Dall'Aglio, M., Martini, M., & Tonani, F. (1972). Rilevamento geochimico delle emanazioni vulcaniche dei Campi Flegrei. *Quaderni De "La Ricerca Scientifica" CNR*, *83*, 152-181.
- De Nicola, D. M. (2000). A review of diatoms found in highly acidic environments. *Hydrobiologia*, *433*, 111-122.
- Di Girolamo, P., Ghiara M.R., Lirer, L., Munno, R., Rolandi G., & Stanzione D. (1984). Vulcanologia e petrografia dei Campi Flegrei. *Bulletin, Geological Society of Italy*, *103*, 349-413.
- Ettl, H. (1978). Die Gattung Chlamydomonas Ehrenberg (Chlamydomonas und die nächsterwandten Gattungen II). *Beih Nova Hedwigia* *60*, 1-1122.
- Ettl, H. (1988). Unterschiedliche Teilungsverläufe bei den Phytomonaden (Chlorophyta). *Archiv Protistenkd* *135*, 81-101.
- Gromov, B. V., Vepritskiy, A.A., Titova, N.N., Mamkayeva, K.A., & Alexandrova, O.V. (1991). Production of the antibiotic cyanobacterin LU-1 by Nostoc linckia Calv 892 (Cyanobacterium). *Journal of Applied Phycology*, *3*, 55-59.
- Gross, W., Kuver, J., Tischendorf, G., Bouchaala, N., & Busch, W. (1998). Cryptoendolithic growth of the red alga Galdieria sulphuraria in volcanic areas. *European Journal of Phycology*, *33*, 25-31.
- Gross, W. (1999). Revision of comparative traits for the acido- and thermophilic red algae Cyanidium and Galdieria. In J. Seckbach (Ed.), *Enigmatic Microorganisms and life in Extreme Environments* (pp. 439-445). London, UK: Kluwer Academic Publishers.
- Hoffmann, L. (2002). Caves and other low-light environments: aerophitic photoautotrophic microorganisms. In: G. Bitton (Ed), *Encyclopedia of environmental microbiology* (pp. 835-843). New York: John Wiley & Sons.
- Huss, V.A., Ciniglia, C., Cennamo P., Cozzolino, S., Pinto, G., & Pollio, A. (2002). Phylogenetic relationships and taxonomic position of Chlorella-like isolates from low pH environments (pH < 3.0). *BMC Evolutionary Biology*, *26*, 1-13.
- Insinga, D., Calvert, A., D'Argenio, B., Fedele, L., Lanphere, M., Morra, V., Perrotta, A., Sacchi, M., & Scarpati, C. (2004). ⁴⁰Ar/³⁹Ar dating of the Neapolitan Yellow Tuff eruption (Campi Flegrei, southern Italy): volcanological and chronostratigraphic implications. *European Geophysical Union 1st General Assembly*, Nice.
- Kvist, L., Broggi, J., Illera, J.C., & Koivula, K. (2005). Colonisation and diversification of the blue tits (Parus caeruleus teneriffae-group) in the Canary Islands. *Molecular and Phylogenetic Evolution*, *34*, 501-511. doi: 10.1016/j.ympev.2004.11.017
- Mangoni, F., & Sgobbo, A. (2013). *Pianificare per lo sviluppo. Un nuovo insediamento ai margini della metropoli*. Napoli, IT: Edizioni Scientifiche Italiane.
- Martini, M., Giannini, L., Buccianti, A., Prati, F., Cellini Legittimo, P., Iozzelli, P., & Capaccioni, B. (1991). Ten years of geochemical investigation at Phlegraean Fields (Italy). *Journal of Volcanology and Geothermal Research*, *48*, 161-171.
- Pinto, G., & Taddei, R. (1977). Le alghe delle acque e dei suoli acidi italiani. *Delpinoa* *18-19*, 77-106.

- Pinto, G., Pollio, A., & Taddei, R. (1992) - List of algae from low pH environments cultivated at the University "Federico II" at Naples (Italy). *Bollettino della Società Adriatica di Scienze*, LXXII (1), 5-24.
- Pinto, G., Albertano, P., Ciniglia, C., Cozzolino, S., Pollio, A., Yoon, H.S., & Battacharya, D. (2003). Comparative approaches to the taxonomy of the genus *Galdieria merolae*. *Cryptogamie Algologie*, 24(1), 13-32.
- Pinto, G., Ciniglia, C., Cascone, C., & Pollio, A. (2006). Species composition of Cyanidiales assemblages in Pisciarelli (Campi Flegrei, Italy), and description of *Galdieria phlegrea* sp. In J. Seckbach (Ed), *Algae and Cyanobacteria In Extreme Environments* (pp 487-502). New York, USA: Springer-Verlag
- Pollio, A., Cennamo, P., Ciniglia, C., De Stefano, M., Pinto, G., & Huss, V. A. R (2005). *Chlamydomonas pitschmannii* Ettl, a Little Known Species from Thermoacidic Environments. *Protist*, 156, 210-237.
- Sanders & Porter, (1988). Phagotrophic phytoflagellates. *Advances in Microbial Ecology*, 10, 167-192.
- Scarpato, C., Cole, P., & Perrotta, A. (1993). The Neapolitan Yellow Tuff—A large volume multiphase eruption from Campi Flegrei, Southern Italy. *Bulletin of Volcanology*, 55, 343-356.
- Sgobbo, A., & Moccia, F. D. (2016). Synergetic Temporary Use for the Enhancement of Historic Centers: The Pilot Project for the Naples Waterfront. *TECHNE Journal of Technology for Architecture and Environment*, 12, 253-260. doi:10.13128/Techne-19360
- Sgobbo, A. (2016). Recycling, waste management and urban vegetable gardens. *WIT Transactions on Ecology and The Environment*, 202, 61-72. doi:10.2495/WM160071
- Tedesco, D., Allard, P., Sano, Y., Wakita, H., & Pece, R. (1990). Helium-3 in subaerial and submarine fumaroles of Campi Flegrei caldera, Italy. *Geochimica and Cosmochimica Acta* 54, 1105-1116.
- Walker, J.J. & Pace, N.R. (2007). Endolithic microbial ecosystems, *Annual review of Microbiology*, 61, 331-347.