
SPECIAL ARTICLE

Particulate Matter 2.5 and Obstetric Complications

Vorapong Phupong, M.D.*.

* Department of Obstetrics and Gynecology, Faculty of Medicine, Chulalongkorn University, Rama IV Road, Pathumwan, Bangkok 10330, Thailand

ABSTRACT

Particulate matter (PM) is microscopic solid or liquid matter suspended in the atmosphere of Earth. The sources of PM can be natural or anthropogenic. The most common used for classification of PM is the size of PM. PM 2.5 influences general health problems. Inhalation of PM 2.5 also causes obstetric complications such as low birth weight, preterm delivery and stillbirth. Thus, pregnant women should avoid the exposure to PM 2.5 for prevention of these obstetric complications.

Correspondence to: Vorapong Phupong, M.D., Placental Related Diseases Research Unit, Department of Obstetrics and Gynecology, Faculty of Medicine, Chulalongkorn University, Rama IV Road, Pathumwan, Bangkok 10330, Thailand, Tel: +66-2-649-2115, Fax: +66-2-256-4275, Email address: vorapong.p@chula.ac.th

Introduction

Particulate matter (PM) is microscopic solid or liquid matter suspended in the atmosphere of Earth^(1, 2). There have been many classifications of PM. The most common used for classification of PM is the size of PM.

PM 10 is defined as particles which pass through a size-selective inlet with a 50% efficiency cut-off at 10 μm aerodynamic diameter. PM10 corresponds to the "thoracic convention"⁽³⁾.

PM 2.5 is defined as particles which pass through a size-selective inlet with a 50% efficiency cut-off at 2.5 μm aerodynamic diameter. PM 2.5 corresponds to the "high-risk respirable convention"⁽³⁾.

The size of PM is very important. Only PM smaller than 10 μm can reach the alveoli. Larger PM are deposited higher up in the respiratory system and removed on the mucocilliary escalator, but may then

be swallowed and subsequently absorbed through the gastrointestinal tract⁽²⁾.

Sources of PM

Sources of PM can be natural or anthropogenic⁽¹⁾. Natural sources include PM originating from volcanoes, dust storms, forest and grassland fires, living vegetation and sea spray. Anthropogenic sources include combustion within car engines, solid-fuel combustion in households, industrial activities (such as building, mining, manufacturing of cement, ceramics and bricks, and smelting), quarrying and mining⁽¹⁾.

Regulation

Due to the highly toxic health effects of particulate matter, most governments have created regulations. Each country has set standards for PM

10 and PM 2.5 concentrations. For example, United States has set standards for PM 2.5 concentrations: 35 µg/m³ for daily average and 12 µg/m³ for yearly average⁽⁴⁾. WHO has set standards for PM 2.5 concentrations: 25 µg/m³ for daily average and 10 µg/m³ for yearly average⁽⁵⁾. In Thailand, the standards for PM 2.5 concentrations are 50 µg/m³ for daily average and 25 µg/m³ for yearly average⁽⁶⁾.

An air quality index (AQI) is used to communicate to the public how polluted the air currently is or how polluted it is forecast to become. As the AQI increases, an increasingly large percentage of the population is likely to experience increasingly severe adverse health effects⁽⁷⁾.

The Air Quality Health Index or (AQHI) is a scale designed to help understand the impact of air quality on health. It has been used in Canada. It is a health protection tool used to make decisions to reduce short-term exposure to air pollution by adjusting activity levels during increased levels of air pollution. The AQHI also provides advice on how to improve air quality by proposing behavioural change to reduce the environmental footprint. The AQHI provides a number from 1 to 10⁺ to indicate the level of health risk associated with local air quality. On occasion, when the amount of air pollution is abnormally high, the number may exceed 10⁽⁸⁾. Table 1 shows the AQHI Health Risk Categories⁽⁸⁾.

Table 1. The AQHI Health Risk Categories⁽⁸⁾.

Health Risk	AQHI	Health Messages	
		At Risk population*	General Population
Low	1–3	Enjoy your usual outdoor activities.	Ideal air quality for outdoor activities
Moderate	4–6	Consider reducing or rescheduling strenuous activities outdoors if you are experiencing symptoms.	No need to modify your usual outdoor activities unless you experience symptoms such as coughing and throat irritation.
High	7–10	Reduce or reschedule strenuous activities outdoors. Children and the elderly should also take it easy.	Consider reducing or rescheduling strenuous activities outdoors if you experience symptoms such as coughing and throat irritation.
Very high	Above 10	Avoid strenuous activities outdoors. Children and the elderly should also avoid outdoor physical exertion.	Reduce or reschedule strenuous activities outdoors, especially if you experience symptoms such as coughing and throat irritation.

* People with heart or breathing problems are at greater risk.

Effect of PM on health problems

The effects of inhaling PM that has been widely studied in humans and animals include respiratory diseases, cardiovascular disease, and obstetric problems. PM has an effect on health problems either acute or chronic effects. Acute health problems include lung inflammatory reactions, respiratory symptoms, adverse effects on the

cardiovascular system, increased in hospital admissions, increased in mortality and increased in medication usage. Chronic health problems include increased in lower respiratory symptoms, reduction in lung function in children, increased in chronic obstructive pulmonary disease, reduction in lung function in adults, and reduction in life expectancy⁽⁹⁾.

Effect of PM 2.5 on obstetric problems

There have been many studies regarding PM 2.5 and obstetric problems. Previous study showed that oxidative stress, deoxyribonucleic acid (DNA) methylation, mitochondrial DNA content alteration, and endocrine disruptions may all play an important role in PM 2.5 induced adverse effects to pregnant women and fetuses. In addition, PM 2.5 exposure can cause male reproductive toxicity, leading to associated adverse pregnancy outcomes⁽¹⁰⁾. Research evidence indicates that PM 2.5 has a potential to induce low birth weight (LBW), preterm birth (PTB), and stillbirth⁽¹⁰⁾.

● **Low birth weight**

Exposure to PM 2.5 during pregnancy may affect the growth and development of infants⁽¹¹⁾. One study found that exposure to specific constituents of PM 2.5, especially traffic-related particles, sulphur constituents, and metals was associated with decreased birth weight. Inhalation of PM can trigger maternal oxidative stress, damage cells, cause inflammation and changes in the blood system, decrease placental blood flow, disrupt transplacental oxygenation, leading to poor growth of the fetus⁽¹²⁾. Previous studies found that exposure to PM 2.5 during pregnancy increased low birth weight infants^(10, 13, 14). Stieb DM, et al demonstrated pooled odds ratios for low birth weight ranged from 1.05 (0.99-1.12) per 10 µg/m³ PM 2.5 to 1.10 (1.05-1.15) per 20 µg/m³ PM 10 based on entire pregnancy exposure⁽¹⁵⁾.

● **Preterm delivery**

Previous studies demonstrated that exposure to PM 2.5 during pregnancy was associated with preterm delivery^(10, 13, 14, 16, 17). DeFranco E et al demonstrated that exposure to high levels of particulate air pollution, PM 2.5, in pregnancy is associated with a 19% increased risk of PTB; with greatest risk with high third trimester exposure⁽¹⁷⁾. Meta-analytic study found that the pooled odds ratio (OR) for PM 2.5 exposure (per 10 µg/m³ increment) during the entire pregnancy on preterm birth was 1.13 (95 % CI 1.03-1.24) in 13 studies with a significant heterogeneity (Q=80.51, p<0.001).

The pooled ORs of PM 2.5 exposure in the first, second and third trimester were 1.08 (95 % CI 0.92-1.26), 1.09 (95 % CI 0.82-1.44) and 1.08 (95 % CI 0.99-1.17), respectively⁽¹⁸⁾.

● **Stillbirth**

Some studies found that exposure to PM 2.5 during pregnancy was associated with stillbirth^(10, 19). DeFranco E et al found that exposure to high levels of PM 2.5 in the third trimester of pregnancy was associated with 42% increased stillbirth risk, aOR 1.42 (95% CI 1.06,1.91)⁽¹⁹⁾.

Prevention of health problems from PM 2.5

The prevention of health problems from PM2.5 include minimizing the releasing of PM 2.5 especially from anthropogenic sources and avoiding the inhalation of PM 2.5.

Risk reduction behaviors of PM 2.5 include minimizing the times for opening windows in order to reduce the PM 2.5 exposure from outside, adjusting time of day or frequency that the physical exercise was done according to AQI, adjusting the physical exercise styles or amounts in relation to the AQI, wearing face masks when going outside in the haze, cleaning the mouths and noses after outdoor activities and using air purifiers when airing the room⁽²⁰⁾.

In conclusion, PM 2.5 can increase obstetric complications such as low birth weight, preterm delivery and stillbirth. Obstetric complications increase when exposure to PM 2.5 during pregnancy, the best way is prevention from PM 2.5 exposure. Pregnant women should avoid the exposure to PM 2.5 for prevention of these obstetric complications

References

1. www.epa.gov/pm-pollution.
2. Lippmann M, Frampton M, Schwartz J, Dockery D, Schlesinger R, Koutrakis P, et al. The US Environmental Protection Agency particulate matter health effects research centers program: A midcourse report of status, progress, and plans. *Environ Health Perspect* 2003;111:1074-92.
3. Air quality - Particle size fraction definitions for health-

- related sampling. International Standards Organization 1995;7708.
4. Cao JJ, Chow JC, Lee FSC, Watson JG. Evolution of PM_{2.5} Measurements and Standards in the US and Future Perspectives for China. *Aerosol Air Qual Res* 2013;13:1197-211.
 5. World Health Organization 2006. WHO Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide.
 6. www.pcd.go.th/info_serv/reg_std_airsnd01.html.
 7. www.gmes-atmosphere.eu/services/raq/raq_nrt/.
 8. www.canada.ca/en/environment-climate-change/services/air-quality-health-index/understanding-messages.html.
 9. World Health Organization 2004. Health aspects of air pollution: Results from the WHO project: Systematic review of health aspects of air pollution in Europe.
 10. Li Z, Tang YQ, Song X, Lazar L, Lie Z, Zhao JS. Impact of ambient PM_{2.5} on adverse birth outcome and potential molecular mechanism. *Ecotoxicol Environ Saf* 2019;169:248-54.
 11. Zhang TL, Zheng XR, Wang X, Zhao H, Wang TT, Zhang HX, et al. Maternal exposure to PM_{2.5} during pregnancy induces impaired development of cerebral cortex in mice offspring. *Int J Mol Sci* 2018;19:pii: E257.
 12. Pedersen M, Gehring U, Beelen R, Wang M, Giorgis-Allemand L, Andersen AMN, et al. Elemental constituents of particulate matter and newborn's size in eight European cohorts. *Environ Health Perspect* 2016;124:141-50.
 13. Shah PS, Balkhair T, P KSGD. Air pollution and birth outcomes: A systematic review. *Environ Int* 2011;37:498-516.
 14. Liu Y, Xu JH, Chen D, Sun P, Ma X. The association between air pollution and preterm birth and low birth weight in Guangdong, China. *BMC Public Health* 2019;19:3.
 15. Stieb DM, Chen L, Eshoul M, Judek S. Ambient air pollution, birth weight and preterm birth: A systematic review and meta-analysis. *Environ Res* 2012;117:100-11.
 16. Laurent O, Hu JL, Li LF, Kleeman MJ, Bartell SM, Cockburn M, et al. A statewide nested case-control study of preterm birth and air pollution by source and composition: California, 2001-2008. *Environ Health Perspect* 2016;124:1479-86.
 17. DeFranco E, Moravec W, Xu F, Hall E, Hossain M, Haynes EN, et al. Exposure to airborne particulate matter during pregnancy is associated with preterm birth: a population-based cohort study. *Environ Health* 2016;15:6.
 18. Sun XL, Luo XP, Zhao CM, Ng RWC, Lim CED, Zhang B, et al. The association between fine particulate matter exposure during pregnancy and preterm birth: a meta-analysis. *BMC Pregnancy Childbirth* 2015;15:300.
 19. DeFranco E, Hall E, Hossain M, Chen A, Haynes EN, Jones D, et al. Air pollution and stillbirth risk: exposure to airborne particulate matter during pregnancy is associated with fetal death. *Plos One* 2015;10:e0120594.
 20. Xiong LL, Li J, Xia T, Hu XY, Wang Y, Sun MN, et al. Risk reduction behaviors regarding PM_{2.5} exposure among outdoor exercisers in the Nanjing metropolitan area, China. *Int J Environ Res Public Health* 2018;15:1728.