

NANO EXPRESS

Open Access



Complex of C₆₀ Fullerene with Doxorubicin as a Promising Agent in Antitumor Therapy

Svitlana V. Prylutska¹, Larysa M. Skivka¹, Gennadiy V. Didenko², Yuriy I. Prylutsky¹, Maxim P. Evstigneev³, Grygoriy P. Potebnya², Rostyslav R. Panchuk^{4*}, Rostyslav S. Stoika⁴, Uwe Ritter⁵ and Peter Scharff⁵

Abstract

The main aim of this work was to evaluate the effect of doxorubicin in complex with C₆₀ fullerene (C₆₀ + Dox) on the growth and metastasis of Lewis lung carcinoma in mice and to perform a primary screening of the potential mechanisms of C₆₀ + Dox complex action. We found that volume of tumor from mice treated with the C₆₀ + Dox complex was 1.4 times less than that in *control* untreated animals. The number of metastatic foci in lungs of animals treated with C₆₀ + Dox complex was two times less than that in *control* untreated animals. Western blot analysis of tumor lysates revealed a significant decrease in the level of heat-shock protein 70 in animals treated with C₆₀ + Dox complex. Moreover, the treatment of tumor-bearing mice was accompanied by the increase of cytotoxic activity of immune cells. Thus, the potential mechanisms of antitumor effect of C₆₀ + Dox complex include both its direct action on tumor cells by inducing cell death and increasing of stress sensitivity and an immunomodulating effect. The obtained results provide a scientific basis for further application of C₆₀ + Dox nanocomplexes as treatment agents in cancer chemotherapy.

Keywords: C₆₀ fullerene, Doxorubicin, Antitumor effect, Antimetastatic effect, Immune response

Background

Suppression of proliferative activity of tumor cells is a basic strategy when using traditional chemotherapeutic drugs [1]. Doxorubicin (Dox) is the anthracycline antibiotic widely used for treatment of cancers of different origin [2]. It uses two main mechanisms in cytotoxic action towards tumor cells: intercalation into DNA helices followed by inhibition of the DNA synthesis and generation of free radicals followed by DNA impairment and cell membrane damage [3]. However, the antitumor effect of traditional chemotherapy is always associated with numerous negative side effects, in particular, the toxicity towards cells of normal organs and tissues. Dox causes potent oxidative stress, mitochondrial dysfunction, and Bcl-2 expression disturbance followed by the apoptotic damage in heart tissue. A number of substances with ability to attenuate the Dox-induced cardiotoxicity have been developed nowadays in order to improve the outcome of the long-term treatment with

Dox [4]. In that regard, C₆₀ fullerene is a promising carbon nanostructure that is characterized by unique physical and chemical properties [5] and biological activity both in vitro and in vivo [6].

It was established that pristine C₆₀ fullerenes at low concentrations are nontoxic [7, 8] and they are able to penetrate through the cytoplasmic membrane of treated cells [9]. One of the biologically most relevant features of C₆₀ fullerene is its antioxidant effect [10]. Our previous results also revealed antioxidant properties of pristine C₆₀ fullerene [11].

C₆₀ fullerene and its derivatives possess potent anticancer activity [12]. It has been reported that C₆₀ fullerene nanocrystal induces certain hallmarks of autophagy in cancer cells [13]. The tumor inhibitory effect of fullerenes is accompanied by the immunomodulatory activity [14].

It is important to emphasize that chemical modification of the surface of C₆₀ molecule for improvement of its water solubility often leads to changes in its physical and chemical properties and to a decrease in specific biological effects. Thus, utilization of pristine C₆₀ fullerene would be more desirable. In our previous study [15], we showed that the water-soluble pristine C₆₀ fullerene

* Correspondence: rpanchuk@ukr.net

⁴Institute of Cell Biology, NAS of Ukraine, 14-16 Drahomanov Str., 79005 Lviv, Ukraine

Full list of author information is available at the end of the article

directly suppresses growth of transplanted malignant tumor. Chen et al. [16] reported that antitumor effect of pristine and functionalized C₆₀ fullerene might be associated with modulation of the oxidative stress and the anti-angiogenic and immunostimulatory activity. Injac et al. [17] demonstrated the ability of fullerenol to decrease the acute Dox pulmototoxicity in rats with malignant neoplasm through inhibition of oxidative stress. In our previous study [18], it was shown that the combination of Dox with C₆₀ fullerene resulted in increase of therapeutic efficacy of the treatment. Taking into account these data, we also suggested [19, 20] that Dox immobilization on C₆₀ fullerene (C₆₀ + Dox complex formation) can reduce negative side effects of this drug towards normal cells as well as enhance its ability to enter target tumor cells.

The main goal of this work was to (1) evaluate the effect of C₆₀ + Dox complex on growth and metastasis of Lewis lung carcinoma (LLC) and (2) perform primary screening of the potential mechanisms of C₆₀ + Dox action.

Methods

Material Preparation and Characterization

The highly stable aqueous colloid solution of purified C₆₀ fullerene (C₆₀FAS; concentration 0.15 mg/ml) was prepared as reviewed in [21, 22]. The method is based on the technology of transferring C₆₀ molecules from toluene to an aqueous phase with the help of ultrasonic treatment.

The atomic force microscopy (AFM) data [21–23] confirm randomly arranged individual C₆₀ molecules with a diameter of ~0.7 nm and their bulk sphere-like aggregates with a height of 2–100 nm in C₆₀FAS.

Dox (“Doxorubicin-TEVA”, Pharmachemie B.V.) was dissolved in saline at initial concentration 0.15 mg/ml and used in all experiments. It was immobilized on the C₆₀ fullerene according to the following protocol: C₆₀FAS (0.15 mg/ml) and Dox (0.15 mg/ml) were mixed in 1:2 volume ratio, and the resulting mixture was treated for 20 min in the ultrasonic disperser. After that, it was subjected to 12-h magnetic stirring at room temperature. Pronounced hypochromic effect observed in spectrophotometric experiment and AFM data clearly indicate a formation of stable C₆₀ + Dox complex [19, 20].

Animals

The male C57Bl/6 mice (20–21 g weight) were kept at 298 ± 1 K on a standard diet in the vivarium of R.E. Kavetsky Institute of Experimental Pathology, Oncology and Radiobiology, NAS of Ukraine (Kyiv). All experiments were conducted in accordance with the international principles of the European Convention for

protection of vertebrate animals under the control of Bio-Ethics Committee of that institution.

Tumor Model, Treatment Regimens, and Study Design

LLC was used as an experimental model. LLC cell line was obtained from the cell line bank of the R.E. Kavetsky Institute of Experimental Pathology, Oncology and Radiobiology, NAS of Ukraine (Kyiv). Tumor cells (5 × 10⁵ in the volume of 100 μl) were transplanted intramuscularly into the mouse limb. After transplantation of tumor cells, the experimental animals were randomized by weight and distributed in four groups with ten animals per group:

Group 1 (C₆₀ fullerene injection). C₆₀FAS was used in 1.5 mg/kg dose (0.2 ml) injected intraperitoneally to mice with transplanted tumor once per day for 5 days with a day interval [15].

Group 2 (Dox injection). Dox was used in 1.5-mg/kg dose (0.2 ml) injected intraperitoneally to mice with transplanted tumor once per day for 5 days with a day interval [24].

Group 3 (C₆₀ + Dox complex injection). C₆₀ + Dox mixture was used in 1.5-mg/kg dose (0.2 ml) injected intraperitoneally to mice with transplanted tumor once per day for 5 days with a day interval.


Control. The mice with transplanted tumor were injected with saline (0.2 ml) once per day for 5 days with a day interval.

Intact animals were used in order to investigate immunological indices (cytotoxic activity of the peritoneal macrophages and mononuclear splenic leucocytes).

The injections of C₆₀ fullerene, Dox, or C₆₀ + Dox complex were started on the 2nd day after tumor cell transplantation. The protocol of injecting C₆₀ fullerenes was based on the fact that C₆₀ fullerenes administered intraperitoneally to rats (500 mg/kg) were subjected to clearance from the organism within 2–4 days [25]. The C₆₀ fullerene dose applied in our experiments was significantly lower than the LD₅₀ value determined for C₆₀ fullerene which, after oral administration to mice, was equivalent to 600 mg/kg of body weight [25].

The kinetics of tumor growth was evaluated as described [15] by linear dimensions of tumor measured every third day with the use of calipers starting from the 9th day after tumor cell inoculation. The euthanasia of experimental animals was performed at the end of the experiment (22nd day), and the number and size of metastases in animal lungs were monitored.

Anticancer effect was also characterized by growth inhibition index, GII, calculated by the formula $GII = (V_c - V_{exp})/V_c \times 100 \%$, where V_c and V_{exp} are the average volumes of tumor of control and experimental



animals, respectively; $V = 1/2 \cdot \left(\frac{a+b}{2}\right)^3$, where a and b are the length and width (in millimeters) of the tumor site, respectively [15].

MTT Assay

To analyze cytotoxic activity of the peritoneal macrophages and mononuclear splenic leukocytes, the modified MTT assay was used as described [26]. Cytotoxic activity of the studied samples was calculated using the formula Cytotoxicity index = $(1 - \epsilon/\epsilon_c) \times 100\%$, where ϵ_c and ϵ are the extinctions of control and test sample, respectively. Measurement of extinction was performed on a digital spectrophotometer (μ Quant, BioTEK, USA) at the wavelength of 540 nm.

The investigation of cytotoxic activity of immunocytes was performed on the 22nd day after tumor cell transplantation. Suspension of tumor cells was prepared from tissue homogenates. Mononuclear splenic leukocytes were obtained from splenocyte suspension by centrifugation (1500 rpm, 40 min) in Ficoll-Hypaque density gradient ($p = 1.077$). Peritoneal macrophages were isolated without preliminary stimulation. Mice were sacrificed, and peritoneal macrophages were harvested using phosphate-buffered saline containing 100 U/ml of heparin. Cells were centrifuged at 300xg for 5 min at 4 °C, washed twice with serum-free DMEM, and re-suspended in DMEM containing 10 % FCS and 40 μ g/ml gentamicin.

To perform cytotoxic assay, LLC cells were placed in 96-well plates (3×10^5 cells/well), and mononuclear splenic leukocytes or peritoneal macrophages were added at 20:1 ratio. Cells were incubated in a RPMI-1640 medium supplemented with gentamicin sulfate (100 μ g/ml) and maintained at 37 °C for 18 h in 5 % CO₂ atmosphere. After incubation, MTT (Sigma) was added to a final

concentration of 0.5 mg/ml followed by culturing for 3 h. After culturing, cells were centrifuged at 4000 rpm (1600xg) for 10 min. Culture medium was removed, and blue formazan crystals were dissolved in 100 μ l DMSO. Optical density was determined at 570 nm.

Western Blot Analysis

The tumors were surgically removed, and cell lysates were prepared by EDTA extraction. After removal of unlysed cell remnants and nuclei by centrifugation in the Eppendorf micro-centrifuge (5 min, 10,000 rpm, 10,200xg); protein concentration was determined by standard method, as described [27]; and 10 μ g of equal amounts of protein was loaded into 15 % polyacrylamide gel. Proteins were resolved and transferred to Immobilon-P membrane (Millipore, Billerica, MA) using semi-dry transfer (Bio-Rad, Hercules, CA). After incubating the membrane in the blocking buffer, the membrane was incubated with heat-shock protein 70 (HSP70) monoclonal antibodies (Santa Cruz Biotechnology, Santa Cruz, CA). For a loading control, the levels of expression of the β -actin were detected in each sample using mouse β -actin monoclonal antibodies (Sigma). Immunoreactive bands were visualized by chemiluminescence using horseradish

Table 1 Tumor growth inhibition index (GIi, %) for each experimental group on the 9th, 13th, 18th, and 22nd day after tumor inoculation

Animal group	Day after tumor cell transplantation			
	9	13	18	22
Group 1 (C ₆₀ fullerene injection), n = 10	26	20	22	12
Group 2 (Dox injection), n = 10	13	22	21	13
Group 3 (C ₆₀ + Dox injection), n = 10	24	22	29	29

Table 2 The effect of C₆₀ fullerene and Dox used alone and in C₆₀ + Dox complex on the LLC metastases

Animal group	The number of tumor nodules in lungs							Total number
	Nodule diameter (mm)							
	<0.5	0.5	1	2	3	4	5	
Control, n = 10	12	9	33	25	3	7	11	100
Group 1(C ₆₀ fullerene injection), n = 10	20	22	23	8	9	1	1	84
Group 2(Dox injection), n = 10	14	5	25	16	7	1	1	69
Group 3(C ₆₀ + Dox injection), n = 10	16	10	6	16	–	–	–	48

peroxidase-conjugated IgG antibodies and ECL Kit (Amersham, Uppsala, Sweden) according to the instructions of the manufacturer.

Statistical Analysis

For statistical analysis of the obtained results, standard variation data within a group was calculated together with a statistical reliability of differences between two groups of data assessed by Student’s *t* test. The level of significance was set to *p* < 0.05.

Results and Discussion

Treatment with C₆₀ + Dox Complex Results in the Inhibition of Tumor Growth and Metastasis and Increases Stress Sensitivity of Tumor Cells In Vivo

Experimental animals tolerated the treatment well and exhibited normal behavior, as determined by the activity level and grooming behavior throughout the study.

LLC was characterized by a significant growth of its size (volume) from the 9th to the 22nd day of the experiment (Fig. 1). One can see that all applied treatments (C₆₀ fullerene, Dox, and C₆₀ + Dox complex) caused a decrease (comparing to *control*, i.e., untreated mice) of tumor volume. Tumor volume in animals of *group 1* (C₆₀ fullerene injection) and *group 2* (Dox injection) differed slightly. The volume of tumor from mice treated with the C₆₀ + Dox complex was significantly lower than that in *control* untreated animals, viz. by 1.4 times.

The effect of C₆₀ fullerene and Dox used alone and in C₆₀ + Dox complex on tumor growth was evaluated by the GII value presented in Table 1.

Thus, on the 13th day after cancer cell inoculation, tumor volume in mice treated with Dox and C₆₀ + Dox

complex was 22 % less than that in the untreated animals. There was no inhibition of tumor growth in mice treated with C₆₀ fullerene alone. On the 18th day after cancer cell transplantation, we observed the most expressed retardation of tumor growth in animals treated with C₆₀ + Dox complex, and the GII value in mice of that group was 29 %. At that time point, the tumor volume in mice treated with C₆₀ fullerene and Dox was ~22 % less than that in the untreated animals. On the 22nd day after cancer cell inoculation, the GII value in mice treated with C₆₀ fullerene and Dox was decreased by ~13 %, but it did not change in mice treated with C₆₀ + Dox complex.

The treatment of tumor-bearing mice with C₆₀ fullerene, Dox, and C₆₀ + Dox complex caused an inhibition of metastasis of the experimental tumor (Table 2). The number of the metastatic foci in lungs of animals of the group treated with C₆₀ + Dox complex was two times less than that in the untreated animals and 1.4 times less than that in mice treated with Dox alone. It should be noted that the metastatic foci in mice treated with C₆₀ fullerene, Dox, and its complex were characterized by different sizes. While in the *control* group, large metastatic foci that infiltrated into the lung parenchyma were observed; the metastatic foci were much smaller and solitary in mice treated with Dox and C₆₀ + Dox complex. In mice treated with C₆₀ + Dox complex, the metastatic foci with the diameter of ≥3 mm were absent. Since only tumor growth beyond the size of 1–2 mm is angiogenesis-dependent [28], we suggested that the small-sized metastatic focus (≤1 mm in diameter) is in a state of dormancy. Therefore, one can suppose that C₆₀ + Dox complex exerts a negative effect towards tumor angiogenesis.





Fig. 2 The effect of treatment with C₆₀ + Dox complex on the level of HSP70 in tumor tissue of animals with LLC (representative Western blot). 1 and 2—control animals (untreated); 3 and 4—C₆₀ + Dox-treated animals




It is known that HSP70 is aberrantly expressed in cancer cells of different origins. The survival of these cells strongly depends upon HSPs due to their role not only in protein refolding and degradation but also in preventing apoptosis [29]. Elevated expression of HSP70 is associated with poor response of tumor cells to chemotherapy, and inhibition of its expression was shown to be an effective strategy against cancer [30]. Therefore, we have measured the level of HSP70 in tumor tissue of animals treated with Dox and C₆₀ + Dox complex. Western blot analysis of tumor cell lysates revealed a significant decrease in HSP70 level only in animals treated with C₆₀ + Dox complex (Fig. 2). The level of HSP70 in tumor lysates from mice treated with Dox alone did not differ from that in the untreated tumor-bearing mice.

The obtained results clearly demonstrate that the anti-cancer activity of Dox is not only well preserved in its complex with C₆₀ fullerene, but it is even enhanced after formation of such complex.

C₆₀ + Dox Complex Modulates Immunological Reactivity of Tumor-Bearing Mice

C₆₀ fullerene and its derivatives were shown to possess the immunomodulating properties [14]. Thus, we supposed that the immunomodulating effect of C₆₀ + Dox complex can be involved in its antitumor action. To testify this hypothesis, the cytotoxic activity of mononuclear splenic leukocytes and macrophages towards autologous tumor cells was evaluated in tumor-bearing mice treated with the C₆₀ + Dox complex.



Growth of experimental tumor was associated with a decrease of macrophage cytotoxicity towards autologous tumor cells *in vitro* (Fig. 3).

Treatment with Dox as well as with C₆₀ fullerene used alone and in C₆₀ + Dox complex resulted in increased cytotoxic activity of the peritoneal macrophages of tumor-bearing mice. Cytotoxic indices of the peritoneal phagocytes in treated animals were comparable with those in the intact mice.

Cytotoxic activity of splenic mononuclear leukocytes in tumor-bearing mice did not differ from that in the intact animals (Fig. 4).

Treatment with Dox resulted in an increase of splenocyte cytotoxicity. A significant individual variability of cytotoxic indices in animals from this group should be noted. Cytotoxicity indices of the mononuclear splenic leukocytes in animals treated with C₆₀ fullerene used alone and in C₆₀ + Dox complex were significantly higher than those in the untreated tumor-bearing mice. The most positive effect was observed in animals receiving C₆₀ + Dox complex. Cytotoxic activity of splenic mononuclear cells towards autologous tumor cells is substantially mediated by splenic NK cells [31]. Turabekova M et al. reported that C₆₀ fullerene might be recognized by Toll-like receptors (TLRs) [32]. TLR-dependent stimulatory effect of the preparation and an increased stress sensitivity of LLC cells associated with a decreased HSP70 expression might be potential reasons of increased cytotoxicity of splenic mononuclear cells in animals receiving C₆₀ + Dox complex.

Conclusions

The results of performed experiments demonstrated that treatment of mice bearing LLC with C₆₀ + Dox nanocomplexes is associated with a significant antitumor effect, namely, (1) the volume of tumor of mice treated with C₆₀ + Dox complex was 1.4 times smaller than that in the *control* untreated animals; (2) the number of metastatic foci in lungs of animals of the group treated with C₆₀ + Dox complex was two times smaller than that in *control* untreated animals; (3) there were no metastatic foci with diameter ≥ 3 mm in mice treated with C₆₀ + Dox complex. Western blot analysis of tumor cell lysates of animals treated with C₆₀ + Dox complex revealed a significant decrease in the HSP70 level. The MTT assay showed that C₆₀ + Dox complex modulates immunological reactivity of tumor-bearing mice. The potential mechanisms of C₆₀ + Dox complex antitumor effect are likely to be based on its direct action on tumor cells with inducing cell death as well as an increasing of stress sensitivity and immunomodulating effect. Thus, the C₆₀ + Dox nanocomplexes might be proposed as new pharmacological agents that are effectively killing tumor cells and simultaneously stimulating immune responses in tumor-bearing mice.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contributions

RS and UR designed the experiments; YP, ME, and PS were involved in the synthesis and characterization of the samples; SP and RP have done *in vitro* studies of C₆₀-Dox complexes (MTT assay, Western blot analysis); and SP, LS, and GD were responsible for *in vivo* studies of C₆₀-Dox complexes. GP, UR, and SP analyzed the data of experiments. SP, YP, and UR wrote the manuscript of the paper. All authors read and approved the final manuscript.

Acknowledgements

The authors thank the Ministry of Education and Science of Ukraine (Yu.K.), DAAD (S.P.), and Russian Science Fund (project no. 14-14-00328) for supporting this work.

Author details

¹Taras Shevchenko National University of Kyiv, 64 Volodymyrska Str., 01601 Kyiv, Ukraine. ²R.E. Kavetsky Institute of Experimental Pathology, Oncology and Radiobiology of NASU, Vasylykivska Str. 45, 03022 Kyiv, Ukraine. ³Belgorod State University, Pobedy Str.85, 308015 Belgorod, Russia. ⁴Institute of Cell Biology, NAS of Ukraine, 14-16 Drahomanov Str., 79005 Lviv, Ukraine. ⁵Technical University of Ilmenau, Institute of Chemistry and Biotechnology, 25 Weimarer Str., 98693 Ilmenau, Germany.

Received: 17 November 2015 Accepted: 16 December 2015

Published online: 29 December 2015

References

- Hirsch J (2006) *JAMA* 296:1518
- Abou EHMA, Verheul HM, Jorna AS, Schalkwijk C, van Bezu J, van der Vijgh WJ, Bast A (2003) *Brit J Cancer* 98:357
- Szulawska A, Czyz M (2006) *Postepy Hig Med Dosw* 60:78
- Jay SM, Murthy AC, Hawkins JF, Wortzel JR, Steinhilber ML, Alvarez LM, Gannon J, Macrae CA, Griffith LG, Lee RT (2013) *Circulation* 128:152
- Fullerenes: Principles and Applications (2nd Ed) (2011). In: Langa De La Puente F, Nierengarten J-F (eds). Cambridge: The Royal Society of Chemistry. p 650. doi:10.1039/9781849732956-PP001.
- Anilkumar P, Lu F, Cao L, Luo P, Liu JH, Sahu S, Tackett K II, Wang Y, Sun YP (2011) *Current Med Chem* 18:2045
- Prylutska SV, Matyshevska OP, Golub AA, Prylutskyi YI, Potebnya GP, Ritter U, Scharff P (2007) Study of C₆₀ fullerenes and C₆₀-containing composites cytotoxicity *in vitro*. *Mater Sci Engineer C* 27:1121–1124
- Johnston HJ, Hutchison GR, Christensen FM, Aschberger K, Stone V (2010) *Toxicol Sci* 114:162
- Prylutska S, Bilyy R, Overchuk M, Bychko A, Andreichenko K, Stoika R, Rybalchenko V, Prylutskyi Y, Tsiherkezos NG, Ritter U (2012) Water-soluble pristine fullerenes C₆₀ increase the specific conductivity and capacity of lipid model membrane and form the channels in cellular plasma membrane. *J Biomed Nanotechnol* 8:522–527
- Markovic Z, Trajkovic V (2008) *Biomaterials* 29:3561
- Prylutska SV, Grynyuk II, Matyshevska OP, Prylutskyi YI, Ritter U, Scharff P (2008) Anti-oxidant properties of C₆₀ fullerenes *in vitro*. *Fullerenes, Nanotubes, Carbon Nanostruct* 16:698–705
- Chen Z, Mao R, Liu Y (2012) *Curr Drug Metabolism* 13:1035
- Zhang Q, Yang W, Man N, Zheng F, Shen Y, Sun K, Li Y, Wen LP (2009) *Autophagy* 5:1107
- Zhu J, Ji Z, Wang J, Sun R, Zhang X, Gao Y, Sun H, Liu Y, Wang Z, Li A, Ma J, Wang T, Jia G, Gu Y (2008) *Small* 4:1168
- Prylutska SV, Burlaka AP, Prylutskyi YI, Ritter U, Scharff P (2011) Pristine C₆₀ fullerenes inhibit the rate of tumor growth and metastasis. *Exp Oncol* 33:162–164
- Chen Z, Ma L, Liu Y, Chen C (2012) *Theranostics* 2:238
- Injac R, Radic N, Govedarica B, Perse M, Cerar A, Djordjevic A, Strukelj B (2009) *Pharmacol Reports* 61:335
- Prylutska S, Grynyuk I, Matyshevska O, Prylutskyi Y, Evstigneev M, Scharff P, Ritter U (2014) C₆₀ fullerene as synergistic agent in tumor-inhibitory doxorubicin treatment. *Drugs R D* 14:333–340
- Evstigneev MP, Buchelnikov AS, Voronin DP, Rubin YV, Belous LF, Prylutskyi YI, Ritter U (2013) Complexation of C₆₀ fullerene with aromatic drugs. *Chem Phys Chem* 14:568–578

20. Prylutskyi YI, Evstigneev MP, Pashkova IS, Wyrzykowski D, Woziwodzka A, Gołuński G, Piosik J, Cherepanov VV, Ritter U (2014) Characterization of C₆₀ fullerene complexation with antibiotic doxorubicin. *Phys Chem Chem Phys* 16:23164–23172
21. Prylutskyi YI, Petrenko VI, Ivankov OI, Kyzyma OA, Bulavin LA, Litsis OO, Evstigneev MP, Cherepanov VV, Naumovets AG, Ritter U (2014) On the origin of C₆₀ fullerene solubility in aqueous solution. *Langmuir* 30:3967–3970
22. Ritter U, Prylutskyi YI, Evstigneev MP, Davidenko NA, Cherepanov VV, Senenko AI, Marchenko OA, Naumovets AG (2015) Structural features of highly stable reproducible C₆₀ fullerene aqueous colloid solution probed by various techniques. *Fullerenes, Nanotubes, Carbon Nanostruct* 23:530–534
23. Prylutskyi YI, Buchelnikov AS, Voronin DP, Kostjukov VV, Ritter U, Parkinson JA, Evstigneev MP (2013) C₆₀ fullerene aggregation in aqueous solution. *Phys Chem Chem Phys* 15:9351–9360
24. Solyanik GI, Todor IN, Kulik GI, Chekhun VF (1999) *Exp Oncol* 21:264
25. Gharbi N, Pressac M, Hadchouel M, Szwarc H, Wilson SR, Moussa F (2005) *Nano Lett* 5:2578
26. Didenko G, Prylutska S, Kichmarenko Y, Potebnya G, Prylutskyi Y, Slobodyanik N, Ritter U, Scharff P (2013) Evaluation of the antitumor immune response to C₆₀ fullerene. *Mat-wiss u Werkstofftech* 44:124–128
27. Skivka LM, Fedorchuk OG, Bezdeneznykh NO, Lykhova OO, Semesiuk NI, Kudryavets YI, Malanchuk OM (2014) *J Exp Integr Med* 4:93
28. Naumov GN, Akslen LA, Folkman J (2006) *Cell Cycle* 5:1779
29. Calderwood SK (2013) *Scientifica* 2013:217513
30. Jego G, Hazoum A, Seigneuric R, Garrido C (2013) *Cancer Lett* 332:275
31. Di Santo JP (2006) *Ann Rev Immunol* 24:257
32. Turabekova M, Rasulev B, Theodore M, Jackman J, Leszczynska D, Leszczynski J (2014) *Nanoscale* 6:3488

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
