

# Second-Order Partial Differentiation of Real Ternary Functions

Takao Inoué  
Inaba 2205, Wing-Minamikan  
Nagano, Nagano, Japan

**Summary.** In this article, we shall extend the result of [17] to discuss second-order partial differentiation of real ternary functions (refer to [7] and [14] for partial differentiation).

MML identifier: PDIFF\_5, version: 7.11.07 4.156.1112

The notation and terminology used here have been introduced in the following papers: [6], [11], [12], [1], [2], [3], [4], [5], [7], [16], [17], [13], [8], [15], [10], and [9].

## 1. SECOND-ORDER PARTIAL DERIVATIVES

For simplicity, we use the following convention:  $x, x_0, y, y_0, z, z_0, r$  denote real numbers,  $u, u_0$  denote elements of  $\mathcal{R}^3$ ,  $f, f_1, f_2$  denote partial functions from  $\mathcal{R}^3$  to  $\mathbb{R}$ ,  $R$  denotes a rest, and  $L$  denotes a linear function.

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . We say that  $f$  is partial differentiable on 1st-1st coordinate in  $u$  if and only if the condition (Def. 1) is satisfied.

(Def. 1) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $x_0$  such that  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 1), u)$  and there exist  $L, R$  such that for every  $x$  such that  $x \in N$  holds  $(\text{SVF1}(1, \text{pdiff1}(f, 1), u))(x) - (\text{SVF1}(1, \text{pdiff1}(f, 1), u))(x_0) = L(x - x_0) + R(x - x_0)$ .

We say that  $f$  is partial differentiable on 1st-2nd coordinate in  $u$  if and only if the condition (Def. 2) is satisfied.

(Def. 2) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $y_0$  such that  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 1), u)$  and there exist  $L, R$  such that for every  $y$  such that  $y \in N$  holds  $(\text{SVF1}(2, \text{pdiff1}(f, 1), u))(y) - (\text{SVF1}(2, \text{pdiff1}(f, 1), u))(y_0) = L(y - y_0) + R(y - y_0)$ .

We say that  $f$  is partial differentiable on 1st-3rd coordinate in  $u$  if and only if the condition (Def. 3) is satisfied.

(Def. 3) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $z_0$  such that  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 1), u)$  and there exist  $L, R$  such that for every  $z$  such that  $z \in N$  holds  $(\text{SVF1}(3, \text{pdiff1}(f, 1), u))(z) - (\text{SVF1}(3, \text{pdiff1}(f, 1), u))(z_0) = L(z - z_0) + R(z - z_0)$ .

We say that  $f$  is partial differentiable on 2nd-1st coordinate in  $u$  if and only if the condition (Def. 4) is satisfied.

(Def. 4) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $x_0$  such that  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 2), u)$  and there exist  $L, R$  such that for every  $x$  such that  $x \in N$  holds  $(\text{SVF1}(1, \text{pdiff1}(f, 2), u))(x) - (\text{SVF1}(1, \text{pdiff1}(f, 2), u))(x_0) = L(x - x_0) + R(x - x_0)$ .

We say that  $f$  is partial differentiable on 2nd-2nd coordinate in  $u$  if and only if the condition (Def. 5) is satisfied.

(Def. 5) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $y_0$  such that  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 2), u)$  and there exist  $L, R$  such that for every  $y$  such that  $y \in N$  holds  $(\text{SVF1}(2, \text{pdiff1}(f, 2), u))(y) - (\text{SVF1}(2, \text{pdiff1}(f, 2), u))(y_0) = L(y - y_0) + R(y - y_0)$ .

We say that  $f$  is partial differentiable on 2nd-3rd coordinate in  $u$  if and only if the condition (Def. 6) is satisfied.

(Def. 6) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $z_0$  such that  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 2), u)$  and there exist  $L, R$  such that for every  $z$  such that  $z \in N$  holds  $(\text{SVF1}(3, \text{pdiff1}(f, 2), u))(z) - (\text{SVF1}(3, \text{pdiff1}(f, 2), u))(z_0) = L(z - z_0) + R(z - z_0)$ .

We say that  $f$  is partial differentiable on 3rd-1st coordinate in  $u$  if and only if the condition (Def. 7) is satisfied.

(Def. 7) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $x_0$  such that  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 3), u)$  and there exist  $L, R$  such that for every  $x$  such that  $x \in N$  holds  $(\text{SVF1}(1, \text{pdiff1}(f, 3), u))(x) - (\text{SVF1}(1, \text{pdiff1}(f, 3), u))(x_0) = L(x - x_0) + R(x - x_0)$ .

We say that  $f$  is partial differentiable on 3rd-2nd coordinate in  $u$  if and only if the condition (Def. 8) is satisfied.

(Def. 8) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $y_0$  such that  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 3), u)$  and there exist  $L, R$  such that for every  $y$  such that  $y \in N$  holds  $(\text{SVF1}(2, \text{pdiff1}(f, 3), u))(y) - (\text{SVF1}(2, \text{pdiff1}(f, 3), u))(y_0) = L(y - y_0) + R(y - y_0)$ .

We say that  $f$  is partial differentiable on 3rd-3rd coordinate in  $u$  if and only if the condition (Def. 9) is satisfied.

(Def. 9) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $z_0$  such that  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 3), u)$  and there exist  $L, R$  such that for every  $z$  such that  $z \in N$  holds  $(\text{SVF1}(3, \text{pdiff1}(f, 3), u))(z) - (\text{SVF1}(3, \text{pdiff1}(f, 3), u))(z_0) = L(z - z_0) + R(z - z_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 1st-1st coordinate in  $u$ . The functor  $\text{hpartdiff11}(f, u)$  yielding a real number is defined by the condition (Def. 10).

(Def. 10) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $x_0$  such that  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 1), u)$  and there exist  $L, R$  such that  $\text{hpartdiff11}(f, u) = L(1)$  and for every  $x$  such that  $x \in N$  holds  $(\text{SVF1}(1, \text{pdiff1}(f, 1), u))(x) - (\text{SVF1}(1, \text{pdiff1}(f, 1), u))(x_0) = L(x - x_0) + R(x - x_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 1st-2nd coordinate in  $u$ . The functor  $\text{hpartdiff12}(f, u)$  yielding a real number is defined by the condition (Def. 11).

(Def. 11) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $y_0$  such that  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 1), u)$  and there exist  $L, R$  such that  $\text{hpartdiff12}(f, u) =$

$L(1)$  and for every  $y$  such that  $y \in N$  holds  $(\text{SVF1}(2, \text{pdiff1}(f, 1), u))(y) - (\text{SVF1}(2, \text{pdiff1}(f, 1), u))(y_0) = L(y - y_0) + R(y - y_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 1st-3rd coordinate in  $u$ . The functor  $\text{hpartdiff13}(f, u)$  yielding a real number is defined by the condition (Def. 12).

(Def. 12) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $z_0$  such that  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 1), u)$  and there exist  $L, R$  such that  $\text{hpartdiff13}(f, u) = L(1)$  and for every  $z$  such that  $z \in N$  holds  $(\text{SVF1}(3, \text{pdiff1}(f, 1), u))(z) - (\text{SVF1}(3, \text{pdiff1}(f, 1), u))(z_0) = L(z - z_0) + R(z - z_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 2nd-1st coordinate in  $u$ . The functor  $\text{hpartdiff21}(f, u)$  yielding a real number is defined by the condition (Def. 13).

(Def. 13) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $x_0$  such that  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 2), u)$  and there exist  $L, R$  such that  $\text{hpartdiff21}(f, u) = L(1)$  and for every  $x$  such that  $x \in N$  holds  $(\text{SVF1}(1, \text{pdiff1}(f, 2), u))(x) - (\text{SVF1}(1, \text{pdiff1}(f, 2), u))(x_0) = L(x - x_0) + R(x - x_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 2nd-2nd coordinate in  $u$ . The functor  $\text{hpartdiff22}(f, u)$  yielding a real number is defined by the condition (Def. 14).

(Def. 14) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $y_0$  such that  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 2), u)$  and there exist  $L, R$  such that  $\text{hpartdiff22}(f, u) = L(1)$  and for every  $y$  such that  $y \in N$  holds  $(\text{SVF1}(2, \text{pdiff1}(f, 2), u))(y) - (\text{SVF1}(2, \text{pdiff1}(f, 2), u))(y_0) = L(y - y_0) + R(y - y_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 2nd-3rd coordinate in  $u$ . The functor  $\text{hpartdiff23}(f, u)$  yielding a real number is defined by the condition (Def. 15).

(Def. 15) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $z_0$  such that  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 2), u)$  and there exist  $L, R$  such that  $\text{hpartdiff23}(f, u) = L(1)$  and for every  $z$  such that  $z \in N$  holds  $(\text{SVF1}(3, \text{pdiff1}(f, 2), u))(z) - (\text{SVF1}(3, \text{pdiff1}(f, 2), u))(z_0) = L(z - z_0) + R(z - z_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 3rd-1st coordinate in  $u$ . The functor

$\text{hpartdiff31}(f, u)$  yields a real number and is defined by the condition (Def. 16).

(Def. 16) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $x_0$  such that  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 3), u)$  and there exist  $L, R$  such that  $\text{hpartdiff31}(f, u) = L(1)$  and for every  $x$  such that  $x \in N$  holds  $(\text{SVF1}(1, \text{pdiff1}(f, 3), u))(x) - (\text{SVF1}(1, \text{pdiff1}(f, 3), u))(x_0) = L(x - x_0) + R(x - x_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 3rd-2nd coordinate in  $u$ . The functor  $\text{hpartdiff32}(f, u)$  yielding a real number is defined by the condition (Def. 17).

(Def. 17) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $y_0$  such that  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 3), u)$  and there exist  $L, R$  such that  $\text{hpartdiff32}(f, u) = L(1)$  and for every  $y$  such that  $y \in N$  holds  $(\text{SVF1}(2, \text{pdiff1}(f, 3), u))(y) - (\text{SVF1}(2, \text{pdiff1}(f, 3), u))(y_0) = L(y - y_0) + R(y - y_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $u$  be an element of  $\mathcal{R}^3$ . Let us assume that  $f$  is partial differentiable on 3rd-3rd coordinate in  $u$ . The functor  $\text{hpartdiff33}(f, u)$  yielding a real number is defined by the condition (Def. 18).

(Def. 18) There exist real numbers  $x_0, y_0, z_0$  such that

- (i)  $u = \langle x_0, y_0, z_0 \rangle$ , and
- (ii) there exists a neighbourhood  $N$  of  $z_0$  such that  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 3), u)$  and there exist  $L, R$  such that  $\text{hpartdiff33}(f, u) = L(1)$  and for every  $z$  such that  $z \in N$  holds  $(\text{SVF1}(3, \text{pdiff1}(f, 3), u))(z) - (\text{SVF1}(3, \text{pdiff1}(f, 3), u))(z_0) = L(z - z_0) + R(z - z_0)$ .

Next we state a number of propositions:

- (1) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 1st-1st coordinate in  $u$ , then  $\text{SVF1}(1, \text{pdiff1}(f, 1), u)$  is differentiable in  $x_0$ .
- (2) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 1st-2nd coordinate in  $u$ , then  $\text{SVF1}(2, \text{pdiff1}(f, 1), u)$  is differentiable in  $y_0$ .
- (3) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 1st-3rd coordinate in  $u$ , then  $\text{SVF1}(3, \text{pdiff1}(f, 1), u)$  is differentiable in  $z_0$ .
- (4) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 2nd-1st coordinate in  $u$ , then  $\text{SVF1}(1, \text{pdiff1}(f, 2), u)$  is differentiable in  $x_0$ .
- (5) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 2nd-2nd coordinate in  $u$ , then  $\text{SVF1}(2, \text{pdiff1}(f, 2), u)$  is differentiable in  $y_0$ .
- (6) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 2nd-3rd coordinate in  $u$ , then  $\text{SVF1}(3, \text{pdiff1}(f, 2), u)$  is differentiable in  $z_0$ .
- (7) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 3rd-1st coordinate in  $u$ , then  $\text{SVF1}(1, \text{pdiff1}(f, 3), u)$  is differentiable in  $x_0$ .

- (8) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 3rd-2nd coordinate in  $u$ , then  $\text{SVF1}(2, \text{pdiff1}(f, 3), u)$  is differentiable in  $y_0$ .
- (9) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 3rd-3rd coordinate in  $u$ , then  $\text{SVF1}(3, \text{pdiff1}(f, 3), u)$  is differentiable in  $z_0$ .
- (10) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 1st-1st coordinate in  $u$ , then  $\text{hpartdiff11}(f, u) = (\text{SVF1}(1, \text{pdiff1}(f, 1), u))'(x_0)$ .
- (11) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 1st-2nd coordinate in  $u$ , then  $\text{hpartdiff12}(f, u) = (\text{SVF1}(2, \text{pdiff1}(f, 1), u))'(y_0)$ .
- (12) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 1st-3rd coordinate in  $u$ , then  $\text{hpartdiff13}(f, u) = (\text{SVF1}(3, \text{pdiff1}(f, 1), u))'(z_0)$ .
- (13) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 2nd-1st coordinate in  $u$ , then  $\text{hpartdiff21}(f, u) = (\text{SVF1}(1, \text{pdiff1}(f, 2), u))'(x_0)$ .
- (14) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 2nd-2nd coordinate in  $u$ , then  $\text{hpartdiff22}(f, u) = (\text{SVF1}(2, \text{pdiff1}(f, 2), u))'(y_0)$ .
- (15) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 2nd-3rd coordinate in  $u$ , then  $\text{hpartdiff23}(f, u) = (\text{SVF1}(3, \text{pdiff1}(f, 2), u))'(z_0)$ .
- (16) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 3rd-1st coordinate in  $u$ , then  $\text{hpartdiff31}(f, u) = (\text{SVF1}(1, \text{pdiff1}(f, 3), u))'(x_0)$ .
- (17) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 3rd-2nd coordinate in  $u$ , then  $\text{hpartdiff32}(f, u) = (\text{SVF1}(2, \text{pdiff1}(f, 3), u))'(y_0)$ .
- (18) If  $u = \langle x_0, y_0, z_0 \rangle$  and  $f$  is partial differentiable on 3rd-3rd coordinate in  $u$ , then  $\text{hpartdiff33}(f, u) = (\text{SVF1}(3, \text{pdiff1}(f, 3), u))'(z_0)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. We say that  $f$  is partial differentiable on 1st-1st coordinate on  $D$  if and only if:

- (Def. 19)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f|_D$  is partial differentiable on 1st-1st coordinate in  $u$ .

We say that  $f$  is partial differentiable on 1st-2nd coordinate on  $D$  if and only if:

- (Def. 20)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f|_D$  is partial differentiable on 1st-2nd coordinate in  $u$ .

We say that  $f$  is partial differentiable on 1st-3rd coordinate on  $D$  if and only if:

- (Def. 21)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f|_D$  is partial differentiable on 1st-3rd coordinate in  $u$ .

We say that  $f$  is partial differentiable on 2nd-1st coordinate on  $D$  if and only if:

- (Def. 22)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f|_D$  is partial differentiable on 2nd-1st coordinate in  $u$ .

We say that  $f$  is partial differentiable on 2nd-2nd coordinate on  $D$  if and only if:

- (Def. 23)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f|_D$  is partial differentiable on 2nd-2nd coordinate in  $u$ .

We say that  $f$  is partial differentiable on 2nd-3rd coordinate on  $D$  if and only if:

(Def. 24)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright D$  is partial differentiable on 2nd-3rd coordinate in  $u$ .

We say that  $f$  is partial differentiable on 3rd-1st coordinate on  $D$  if and only if:

(Def. 25)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright D$  is partial differentiable on 3rd-1st coordinate in  $u$ .

We say that  $f$  is partial differentiable on 3rd-2nd coordinate on  $D$  if and only if:

(Def. 26)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright D$  is partial differentiable on 3rd-2nd coordinate in  $u$ .

We say that  $f$  is partial differentiable on 3rd-3rd coordinate on  $D$  if and only if:

(Def. 27)  $D \subseteq \text{dom } f$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright D$  is partial differentiable on 3rd-3rd coordinate in  $u$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 1st-1st coordinate on  $D$ . The functor  $f \upharpoonright_D^{1\text{st}-1\text{st}}$  yields a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and is defined by:

(Def. 28)  $\text{dom}(f \upharpoonright_D^{1\text{st}-1\text{st}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright_D^{1\text{st}-1\text{st}}(u) = \text{hpartdiff11}(f, u)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 1st-2nd coordinate on  $D$ . The functor  $f \upharpoonright_D^{1\text{st}-2\text{nd}}$  yielding a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  is defined by:

(Def. 29)  $\text{dom}(f \upharpoonright_D^{1\text{st}-2\text{nd}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright_D^{1\text{st}-2\text{nd}}(u) = \text{hpartdiff12}(f, u)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 1st-3rd coordinate on  $D$ . The functor  $f \upharpoonright_D^{1\text{st}-3\text{rd}}$  yields a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and is defined by:

(Def. 30)  $\text{dom}(f \upharpoonright_D^{1\text{st}-3\text{rd}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright_D^{1\text{st}-3\text{rd}}(u) = \text{hpartdiff13}(f, u)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 2nd-1st coordinate on  $D$ . The functor  $f \upharpoonright_D^{2\text{nd}-1\text{st}}$  yielding a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  is defined as follows:

(Def. 31)  $\text{dom}(f \upharpoonright_D^{2\text{nd}-1\text{st}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright_D^{2\text{nd}-1\text{st}}(u) = \text{hpartdiff21}(f, u)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 2nd-2nd coordinate on  $D$ . The functor  $f \upharpoonright_D^{2\text{nd}-2\text{nd}}$  yields a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and is defined by:

(Def. 32)  $\text{dom}(f \upharpoonright_D^{2\text{nd}-2\text{nd}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  $f \upharpoonright_D^{2\text{nd}-2\text{nd}}(u) = \text{hpartdiff22}(f, u)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 2nd-3rd coordinate on  $D$ . The functor  $f \upharpoonright_D^{2\text{nd}-3\text{rd}}$

yields a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and is defined by:

- (Def. 33)  $\text{dom}(f_{\downarrow D}^{2\text{nd}-3\text{rd}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  
 $f_{\downarrow D}^{2\text{nd}-3\text{rd}}(u) = \text{hpartdiff23}(f, u)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 3rd-1st coordinate on  $D$ . The functor  $f_{\downarrow D}^{3\text{rd}-1\text{st}}$  yields a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and is defined as follows:

- (Def. 34)  $\text{dom}(f_{\downarrow D}^{3\text{rd}-1\text{st}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  
 $f_{\downarrow D}^{3\text{rd}-1\text{st}}(u) = \text{hpartdiff31}(f, u)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 3rd-2nd coordinate on  $D$ . The functor  $f_{\downarrow D}^{3\text{rd}-2\text{nd}}$  yields a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and is defined by:

- (Def. 35)  $\text{dom}(f_{\downarrow D}^{3\text{rd}-2\text{nd}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  
 $f_{\downarrow D}^{3\text{rd}-2\text{nd}}(u) = \text{hpartdiff32}(f, u)$ .

Let  $f$  be a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  and let  $D$  be a set. Let us assume that  $f$  is partial differentiable on 3rd-3rd coordinate on  $D$ . The functor  $f_{\downarrow D}^{3\text{rd}-3\text{rd}}$  yielding a partial function from  $\mathcal{R}^3$  to  $\mathbb{R}$  is defined by:

- (Def. 36)  $\text{dom}(f_{\downarrow D}^{3\text{rd}-3\text{rd}}) = D$  and for every element  $u$  of  $\mathcal{R}^3$  such that  $u \in D$  holds  
 $f_{\downarrow D}^{3\text{rd}-3\text{rd}}(u) = \text{hpartdiff33}(f, u)$ .

## 2. MAIN PROPERTIES OF SECOND-ORDER PARTIAL DERIVATIVES

Next we state a number of propositions:

- (19)  $f$  is partial differentiable on 1st-1st coordinate in  $u$  if and only if  $\text{pdiff1}(f, 1)$  is partially differentiable in  $u$  w.r.t. 1.
- (20)  $f$  is partial differentiable on 1st-2nd coordinate in  $u$  if and only if  $\text{pdiff1}(f, 1)$  is partially differentiable in  $u$  w.r.t. 2.
- (21)  $f$  is partial differentiable on 1st-3rd coordinate in  $u$  if and only if  $\text{pdiff1}(f, 1)$  is partially differentiable in  $u$  w.r.t. 3.
- (22)  $f$  is partial differentiable on 2nd-1st coordinate in  $u$  if and only if  $\text{pdiff1}(f, 2)$  is partially differentiable in  $u$  w.r.t. 1.
- (23)  $f$  is partial differentiable on 2nd-2nd coordinate in  $u$  if and only if  $\text{pdiff1}(f, 2)$  is partially differentiable in  $u$  w.r.t. 2.
- (24)  $f$  is partial differentiable on 2nd-3rd coordinate in  $u$  if and only if  $\text{pdiff1}(f, 2)$  is partially differentiable in  $u$  w.r.t. 3.
- (25)  $f$  is partial differentiable on 3rd-1st coordinate in  $u$  if and only if  $\text{pdiff1}(f, 3)$  is partially differentiable in  $u$  w.r.t. 1.
- (26)  $f$  is partial differentiable on 3rd-2nd coordinate in  $u$  if and only if  $\text{pdiff1}(f, 3)$  is partially differentiable in  $u$  w.r.t. 2.



- (27)  $f$  is partial differentiable on 3rd-3rd coordinate in  $u$  if and only if  $\text{pdiff1}(f, 3)$  is partially differentiable in  $u$  w.r.t. 3.
- (28) If  $f$  is partial differentiable on 1st-1st coordinate in  $u$ , then  $\text{hpartdiff11}(f, u) = \text{partdiff}(\text{pdiff1}(f, 1), u, 1)$ .
- (29) If  $f$  is partial differentiable on 1st-2nd coordinate in  $u$ , then  $\text{hpartdiff12}(f, u) = \text{partdiff}(\text{pdiff1}(f, 1), u, 2)$ .
- (30) If  $f$  is partial differentiable on 1st-3rd coordinate in  $u$ , then  $\text{hpartdiff13}(f, u) = \text{partdiff}(\text{pdiff1}(f, 1), u, 3)$ .
- (31) If  $f$  is partial differentiable on 2nd-1st coordinate in  $u$ , then  $\text{hpartdiff21}(f, u) = \text{partdiff}(\text{pdiff1}(f, 2), u, 1)$ .
- (32) If  $f$  is partial differentiable on 2nd-2nd coordinate in  $u$ , then  $\text{hpartdiff22}(f, u) = \text{partdiff}(\text{pdiff1}(f, 2), u, 2)$ .
- (33) If  $f$  is partial differentiable on 2nd-3rd coordinate in  $u$ , then  $\text{hpartdiff23}(f, u) = \text{partdiff}(\text{pdiff1}(f, 2), u, 3)$ .
- (34) If  $f$  is partial differentiable on 3rd-1st coordinate in  $u$ , then  $\text{hpartdiff31}(f, u) = \text{partdiff}(\text{pdiff1}(f, 3), u, 1)$ .
- (35) If  $f$  is partial differentiable on 3rd-2nd coordinate in  $u$ , then  $\text{hpartdiff32}(f, u) = \text{partdiff}(\text{pdiff1}(f, 3), u, 2)$ .
- (36) If  $f$  is partial differentiable on 3rd-3rd coordinate in  $u$ , then  $\text{hpartdiff33}(f, u) = \text{partdiff}(\text{pdiff1}(f, 3), u, 3)$ .
- (37) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(1, 3))(u_0)$ . Suppose  $f$  is partial differentiable on 1st-1st coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 1), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real numbers. Suppose  $\text{rng } c = \{(\text{proj}(1, 3))(u_0)\}$  and  $\text{rng}(h + c) \subseteq N$ . Then  $h^{-1}((\text{SVF1}(1, \text{pdiff1}(f, 1), u_0)_*(h+c)) - (\text{SVF1}(1, \text{pdiff1}(f, 1), u_0)_*c))$  is convergent and  $\text{hpartdiff11}(f, u_0) = \lim(h^{-1}((\text{SVF1}(1, \text{pdiff1}(f, 1), u_0)_*(h+c)) - (\text{SVF1}(1, \text{pdiff1}(f, 1), u_0)_*c)))$ .
- (38) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(2, 3))(u_0)$ . Suppose  $f$  is partial differentiable on 1st-2nd coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 1), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real numbers. Suppose  $\text{rng } c = \{(\text{proj}(2, 3))(u_0)\}$  and  $\text{rng}(h + c) \subseteq N$ . Then  $h^{-1}((\text{SVF1}(2, \text{pdiff1}(f, 1), u_0)_*(h+c)) - (\text{SVF1}(2, \text{pdiff1}(f, 1), u_0)_*c))$  is convergent and  $\text{hpartdiff12}(f, u_0) = \lim(h^{-1}((\text{SVF1}(2, \text{pdiff1}(f, 1), u_0)_*(h+c)) - (\text{SVF1}(2, \text{pdiff1}(f, 1), u_0)_*c)))$ .
- (39) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(3, 3))(u_0)$ . Suppose  $f$  is partial differentiable on 1st-3rd coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 1), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real num-

bers. Suppose  $\text{rng } c = \{(\text{proj}(3,3))(u_0)\}$  and  $\text{rng}(h+c) \subseteq N$ . Then  $h^{-1}((\text{SVF1}(3, \text{pdiff1}(f, 1), u_0)_*(h+c)) - (\text{SVF1}(3, \text{pdiff1}(f, 1), u_0)_*c))$  is convergent and  $\text{hpartdiff13}(f, u_0) = \lim(h^{-1}((\text{SVF1}(3, \text{pdiff1}(f, 1), u_0)_*(h+c)) - (\text{SVF1}(3, \text{pdiff1}(f, 1), u_0)_*c)))$ .

- (40) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(1,3))(u_0)$ . Suppose  $f$  is partial differentiable on 2nd-1st coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 2), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real numbers. Suppose  $\text{rng } c = \{(\text{proj}(1,3))(u_0)\}$  and  $\text{rng}(h+c) \subseteq N$ . Then  $h^{-1}((\text{SVF1}(1, \text{pdiff1}(f, 2), u_0)_*(h+c)) - (\text{SVF1}(1, \text{pdiff1}(f, 2), u_0)_*c))$  is convergent and  $\text{hpartdiff21}(f, u_0) = \lim(h^{-1}((\text{SVF1}(1, \text{pdiff1}(f, 2), u_0)_*(h+c)) - (\text{SVF1}(1, \text{pdiff1}(f, 2), u_0)_*c)))$ .
- (41) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(2,3))(u_0)$ . Suppose  $f$  is partial differentiable on 2nd-2nd coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 2), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real numbers. Suppose  $\text{rng } c = \{(\text{proj}(2,3))(u_0)\}$  and  $\text{rng}(h+c) \subseteq N$ . Then  $h^{-1}((\text{SVF1}(2, \text{pdiff1}(f, 2), u_0)_*(h+c)) - (\text{SVF1}(2, \text{pdiff1}(f, 2), u_0)_*c))$  is convergent and  $\text{hpartdiff22}(f, u_0) = \lim(h^{-1}((\text{SVF1}(2, \text{pdiff1}(f, 2), u_0)_*(h+c)) - (\text{SVF1}(2, \text{pdiff1}(f, 2), u_0)_*c)))$ .
- (42) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(3,3))(u_0)$ . Suppose  $f$  is partial differentiable on 2nd-3rd coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 2), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real numbers. Suppose  $\text{rng } c = \{(\text{proj}(3,3))(u_0)\}$  and  $\text{rng}(h+c) \subseteq N$ . Then  $h^{-1}((\text{SVF1}(3, \text{pdiff1}(f, 2), u_0)_*(h+c)) - (\text{SVF1}(3, \text{pdiff1}(f, 2), u_0)_*c))$  is convergent and  $\text{hpartdiff23}(f, u_0) = \lim(h^{-1}((\text{SVF1}(3, \text{pdiff1}(f, 2), u_0)_*(h+c)) - (\text{SVF1}(3, \text{pdiff1}(f, 2), u_0)_*c)))$ .
- (43) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(1,3))(u_0)$ . Suppose  $f$  is partial differentiable on 3rd-1st coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(1, \text{pdiff1}(f, 3), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real numbers. Suppose  $\text{rng } c = \{(\text{proj}(1,3))(u_0)\}$  and  $\text{rng}(h+c) \subseteq N$ . Then  $h^{-1}((\text{SVF1}(1, \text{pdiff1}(f, 3), u_0)_*(h+c)) - (\text{SVF1}(1, \text{pdiff1}(f, 3), u_0)_*c))$  is convergent and  $\text{hpartdiff31}(f, u_0) = \lim(h^{-1}((\text{SVF1}(1, \text{pdiff1}(f, 3), u_0)_*(h+c)) - (\text{SVF1}(1, \text{pdiff1}(f, 3), u_0)_*c)))$ .
- (44) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(2,3))(u_0)$ . Suppose  $f$  is partial differentiable on 3rd-2nd coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(2, \text{pdiff1}(f, 3), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real numbers. Suppose  $\text{rng } c = \{(\text{proj}(2,3))(u_0)\}$  and  $\text{rng}(h+c) \subseteq N$ . Then

- $h^{-1}((\text{SVF1}(2, \text{pdiff1}(f, 3), u_0)_*(h+c)) - (\text{SVF1}(2, \text{pdiff1}(f, 3), u_0)_*c))$  is convergent and  $\text{hpartdiff32}(f, u_0) = \lim(h^{-1}((\text{SVF1}(2, \text{pdiff1}(f, 3), u_0)_*(h+c)) - (\text{SVF1}(2, \text{pdiff1}(f, 3), u_0)_*c)))$ .
- (45) Let  $u_0$  be an element of  $\mathcal{R}^3$  and  $N$  be a neighbourhood of  $(\text{proj}(3, 3))(u_0)$ . Suppose  $f$  is partial differentiable on 3rd-3rd coordinate in  $u_0$  and  $N \subseteq \text{dom SVF1}(3, \text{pdiff1}(f, 3), u_0)$ . Let  $h$  be a convergent to 0 sequence of real numbers and  $c$  be a constant sequence of real numbers. Suppose  $\text{rng } c = \{(\text{proj}(3, 3))(u_0)\}$  and  $\text{rng}(h+c) \subseteq N$ . Then  $h^{-1}((\text{SVF1}(3, \text{pdiff1}(f, 3), u_0)_*(h+c)) - (\text{SVF1}(3, \text{pdiff1}(f, 3), u_0)_*c))$  is convergent and  $\text{hpartdiff33}(f, u_0) = \lim(h^{-1}((\text{SVF1}(3, \text{pdiff1}(f, 3), u_0)_*(h+c)) - (\text{SVF1}(3, \text{pdiff1}(f, 3), u_0)_*c)))$ .
- (46) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-1st coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-1st coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) + \text{pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 1 and  $\text{partdiff}(\text{pdiff1}(f_1, 1) + \text{pdiff1}(f_2, 1), u_0, 1) = \text{hpartdiff11}(f_1, u_0) + \text{hpartdiff11}(f_2, u_0)$ .
- (47) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-2nd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-2nd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) + \text{pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 2 and  $\text{partdiff}(\text{pdiff1}(f_1, 1) + \text{pdiff1}(f_2, 1), u_0, 2) = \text{hpartdiff12}(f_1, u_0) + \text{hpartdiff12}(f_2, u_0)$ .
- (48) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-3rd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-3rd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) + \text{pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 3 and  $\text{partdiff}(\text{pdiff1}(f_1, 1) + \text{pdiff1}(f_2, 1), u_0, 3) = \text{hpartdiff13}(f_1, u_0) + \text{hpartdiff13}(f_2, u_0)$ .
- (49) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-1st coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-1st coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) + \text{pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 1 and  $\text{partdiff}(\text{pdiff1}(f_1, 2) + \text{pdiff1}(f_2, 2), u_0, 1) = \text{hpartdiff21}(f_1, u_0) + \text{hpartdiff21}(f_2, u_0)$ .
- (50) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-2nd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-2nd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) + \text{pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 2 and  $\text{partdiff}(\text{pdiff1}(f_1, 2) + \text{pdiff1}(f_2, 2), u_0, 2) = \text{hpartdiff22}(f_1, u_0) + \text{hpartdiff22}(f_2, u_0)$ .

- (51) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-3rd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-3rd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) + \text{pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 3 and  $\text{partdiff}(\text{pdiff1}(f_1, 2) + \text{pdiff1}(f_2, 2), u_0, 3) = \text{hpartdiff23}(f_1, u_0) + \text{hpartdiff23}(f_2, u_0)$ .
- (52) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-1st coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-1st coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) - \text{pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 1 and  $\text{partdiff}(\text{pdiff1}(f_1, 1) - \text{pdiff1}(f_2, 1), u_0, 1) = \text{hpartdiff11}(f_1, u_0) - \text{hpartdiff11}(f_2, u_0)$ .
- (53) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-2nd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-2nd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) - \text{pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 2 and  $\text{partdiff}(\text{pdiff1}(f_1, 1) - \text{pdiff1}(f_2, 1), u_0, 2) = \text{hpartdiff12}(f_1, u_0) - \text{hpartdiff12}(f_2, u_0)$ .
- (54) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-3rd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-3rd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) - \text{pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 3 and  $\text{partdiff}(\text{pdiff1}(f_1, 1) - \text{pdiff1}(f_2, 1), u_0, 3) = \text{hpartdiff13}(f_1, u_0) - \text{hpartdiff13}(f_2, u_0)$ .
- (55) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-1st coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-1st coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) - \text{pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 1 and  $\text{partdiff}(\text{pdiff1}(f_1, 2) - \text{pdiff1}(f_2, 2), u_0, 1) = \text{hpartdiff21}(f_1, u_0) - \text{hpartdiff21}(f_2, u_0)$ .
- (56) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-2nd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-2nd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) - \text{pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 2 and  $\text{partdiff}(\text{pdiff1}(f_1, 2) - \text{pdiff1}(f_2, 2), u_0, 2) = \text{hpartdiff22}(f_1, u_0) - \text{hpartdiff22}(f_2, u_0)$ .
- (57) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-3rd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-3rd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) - \text{pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 3 and  $\text{partdiff}(\text{pdiff1}(f_1, 2) - \text{pdiff1}(f_2, 2), u_0, 3) = \text{hpartdiff23}(f_1, u_0) - \text{hpartdiff23}(f_2, u_0)$ .

$\text{hpartdiff23}(f_2, u_0)$ .

- (58) Suppose  $f$  is partial differentiable on 1st-1st coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 1)$  is partially differentiable in  $u_0$  w.r.t. 1 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 1), u_0, 1) = r \cdot \text{hpartdiff11}(f, u_0)$ .
- (59) Suppose  $f$  is partial differentiable on 1st-2nd coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 1)$  is partially differentiable in  $u_0$  w.r.t. 2 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 1), u_0, 2) = r \cdot \text{hpartdiff12}(f, u_0)$ .
- (60) Suppose  $f$  is partial differentiable on 1st-3rd coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 1)$  is partially differentiable in  $u_0$  w.r.t. 3 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 1), u_0, 3) = r \cdot \text{hpartdiff13}(f, u_0)$ .
- (61) Suppose  $f$  is partial differentiable on 2nd-1st coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 2)$  is partially differentiable in  $u_0$  w.r.t. 1 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 2), u_0, 1) = r \cdot \text{hpartdiff21}(f, u_0)$ .
- (62) Suppose  $f$  is partial differentiable on 2nd-2nd coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 2)$  is partially differentiable in  $u_0$  w.r.t. 2 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 2), u_0, 2) = r \cdot \text{hpartdiff22}(f, u_0)$ .
- (63) Suppose  $f$  is partial differentiable on 2nd-3rd coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 2)$  is partially differentiable in  $u_0$  w.r.t. 3 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 2), u_0, 3) = r \cdot \text{hpartdiff23}(f, u_0)$ .
- (64) Suppose  $f$  is partial differentiable on 3rd-1st coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 3)$  is partially differentiable in  $u_0$  w.r.t. 1 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 3), u_0, 1) = r \cdot \text{hpartdiff31}(f, u_0)$ .
- (65) Suppose  $f$  is partial differentiable on 3rd-2nd coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 3)$  is partially differentiable in  $u_0$  w.r.t. 2 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 3), u_0, 2) = r \cdot \text{hpartdiff32}(f, u_0)$ .
- (66) Suppose  $f$  is partial differentiable on 3rd-3rd coordinate in  $u_0$ .  
Then  $r \text{ pdiff1}(f, 3)$  is partially differentiable in  $u_0$  w.r.t. 3 and  
 $\text{partdiff}(r \text{ pdiff1}(f, 3), u_0, 3) = r \cdot \text{hpartdiff33}(f, u_0)$ .
- (67) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-1st coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-1st coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) \text{ pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 1.
- (68) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-2nd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-2nd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) \text{ pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 2.
- (69) Suppose that
- (i)  $f_1$  is partial differentiable on 1st-3rd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 1st-3rd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 1) \text{ pdiff1}(f_2, 1)$  is partially differentiable in  $u_0$  w.r.t. 3.

- (70) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-1st coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-1st coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) \text{ pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 1.
- (71) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-2nd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-2nd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) \text{ pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 2.
- (72) Suppose that
- (i)  $f_1$  is partial differentiable on 2nd-3rd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 2nd-3rd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 2) \text{ pdiff1}(f_2, 2)$  is partially differentiable in  $u_0$  w.r.t. 3.
- (73) Suppose that
- (i)  $f_1$  is partial differentiable on 3rd-1st coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 3rd-1st coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 3) \text{ pdiff1}(f_2, 3)$  is partially differentiable in  $u_0$  w.r.t. 1.
- (74) Suppose that
- (i)  $f_1$  is partial differentiable on 3rd-2nd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 3rd-2nd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 3) \text{ pdiff1}(f_2, 3)$  is partially differentiable in  $u_0$  w.r.t. 2.
- (75) Suppose that
- (i)  $f_1$  is partial differentiable on 3rd-3rd coordinate in  $u_0$ , and
  - (ii)  $f_2$  is partial differentiable on 3rd-3rd coordinate in  $u_0$ .
- Then  $\text{pdiff1}(f_1, 3) \text{ pdiff1}(f_2, 3)$  is partially differentiable in  $u_0$  w.r.t. 3.
- (76) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 1st-1st coordinate in  $u_0$ . Then  $\text{SVF1}(1, \text{pdiff1}(f, 1), u_0)$  is continuous in  $(\text{proj}(1, 3))(u_0)$ .
- (77) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 1st-2nd coordinate in  $u_0$ . Then  $\text{SVF1}(2, \text{pdiff1}(f, 1), u_0)$  is continuous in  $(\text{proj}(2, 3))(u_0)$ .
- (78) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 1st-3rd coordinate in  $u_0$ . Then  $\text{SVF1}(3, \text{pdiff1}(f, 1), u_0)$  is continuous in  $(\text{proj}(3, 3))(u_0)$ .
- (79) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 2nd-1st coordinate in  $u_0$ . Then  $\text{SVF1}(1, \text{pdiff1}(f, 2), u_0)$  is continuous in  $(\text{proj}(1, 3))(u_0)$ .
- (80) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 2nd-2nd coordinate in  $u_0$ . Then  $\text{SVF1}(2, \text{pdiff1}(f, 2), u_0)$  is continuous in  $(\text{proj}(2, 3))(u_0)$ .

- (81) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 2nd-3rd coordinate in  $u_0$ . Then  $\text{SVF1}(3, \text{pdiff1}(f, 2), u_0)$  is continuous in  $(\text{proj}(3, 3))(u_0)$ .
- (82) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 3rd-1st coordinate in  $u_0$ . Then  $\text{SVF1}(1, \text{pdiff1}(f, 3), u_0)$  is continuous in  $(\text{proj}(1, 3))(u_0)$ .
- (83) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 3rd-2nd coordinate in  $u_0$ . Then  $\text{SVF1}(2, \text{pdiff1}(f, 3), u_0)$  is continuous in  $(\text{proj}(2, 3))(u_0)$ .
- (84) Let  $u_0$  be an element of  $\mathcal{R}^3$ . Suppose  $f$  is partial differentiable on 3rd-3rd coordinate in  $u_0$ . Then  $\text{SVF1}(3, \text{pdiff1}(f, 3), u_0)$  is continuous in  $(\text{proj}(3, 3))(u_0)$ .

## REFERENCES

- [1] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [2] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [5] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [6] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [7] Noboru Endou, Yasunari Shidama, and Keiichi Miyajima. Partial differentiation on normed linear spaces  $\mathcal{R}^n$ . *Formalized Mathematics*, 15(2):65–72, 2007, doi:10.2478/v10037-007-0008-5.
- [8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [9] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(2):273–275, 1990.
- [10] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [11] Konrad Raczkowski and Paweł Sadowski. Real function continuity. *Formalized Mathematics*, 1(4):787–791, 1990.
- [12] Konrad Raczkowski and Paweł Sadowski. Real function differentiability. *Formalized Mathematics*, 1(4):797–801, 1990.
- [13] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [14] Walter Rudin. *Principles of Mathematical Analysis*. MacGraw-Hill, 1976.
- [15] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [16] Bing Xie, Xiquan Liang, and Hongwei Li. Partial differentiation of real binary functions. *Formalized Mathematics*, 16(4):333–338, 2008, doi:10.2478/v10037-008-0041-z.
- [17] Bing Xie, Xiquan Liang, and Xiuzhuan Shen. Second-order partial differentiation of real binary functions. *Formalized Mathematics*, 17(2):79–87, 2009, doi: 10.2478/v10037-009-0009-7.

Received January 26, 2010