


Contents lists available at [ScienceDirect](http://ScienceDirect.com)

# Vaccine

journal homepage: [www.elsevier.com/locate/vaccine](http://www.elsevier.com/locate/vaccine)

## Retinoic acid elicits a coordinated expression of gut homing markers on T lymphocytes of Zambian men receiving oral Vivotif, but not Rotarix, Dukoral or OPVERO vaccines

Mpala Mwanza-Lisulo<sup>a,\*</sup>, Mumba S. Chomba<sup>a</sup>, Mubanga Chama<sup>a,b</sup>, Ellen C. Besa<sup>a</sup>, Evelyn Funjika<sup>a,b</sup>, Kanekwa Zyambo<sup>a</sup>, Rose Banda<sup>a</sup>, Mercy Imikendu<sup>a</sup>, Sandie Sianongo<sup>a</sup>, Robert E.W. Hancock<sup>c</sup>, Amy Lee<sup>c</sup>, Roma Chilengi<sup>d</sup>, Andy J. Stagg<sup>e</sup>, Boniface Namangala<sup>f</sup>, Paul M. Kelly<sup>a,e</sup>

<sup>a</sup>Tropical Gastroenterology & Nutrition Group, Department of Medicine, School of Medicine, University of Zambia, Lusaka, Zambia

<sup>b</sup>Department of Chemistry, University of Zambia, Lusaka, Zambia

<sup>c</sup>University of British Columbia, Vancouver, Canada

<sup>d</sup>Centre for Infectious Disease Research in Zambia (CIDRZ), Lusaka, Zambia

<sup>e</sup>Blizard Institute, Barts & The London School of Medicine, Queen Mary University of London, London, UK

<sup>f</sup>Institute of Distance Education, University of Zambia, Lusaka, Zambia

### ARTICLE INFO

#### Article history:

Received 17 January 2018

Received in revised form 29 March 2018

Accepted 30 April 2018

Available online 22 May 2018

#### Keywords:

Retinoic acid

$\alpha 4\beta 7$

CCR9

Gut mucosa

Vaccines

### ABSTRACT

All-trans retinoic acid (ATRA) up-regulates, in laboratory animals, the expression of the gut homing markers  $\alpha 4\beta 7$  integrin and CCR9 on lymphocytes, increasing their gut tropism. Here, we show that, in healthy adult volunteers, ATRA induced an increase of these gut homing markers on T cells *in vivo* in a time dependent manner. The coordinated increase of  $\alpha 4\beta 7$  and CCR9 by ATRA was seen in 57% (12/21) of volunteers and only when given together with an oral Vivotif vaccine. When this coordinated response to ATRA and Vivotif vaccine was present, it was strongly correlated with the gut immunoglobulin A (IgA) specific response to vaccine LPS ( $\rho = 0.82$ ;  $P = 0.02$ ). Using RNA-Seq analysis of whole blood transcription, patients receiving ATRA and Vivotif in conjunction showed transcriptomic changes in immune-related pathways, particularly including interferon  $\alpha/\beta$  signaling pathway, membrane-ECM interactions and immune hubs. These results suggest that exogenous ATRA can be used to manipulate responses to a subclass of oral vaccines, so far limited to a live attenuated Vivotif vaccine.

© 2018 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

### 1. Introduction

There is strong evidence of diminished immunogenicity and efficacy of oral vaccines in developing country populations [1]. For example, oral rotavirus vaccine affords excellent (80–90%) protection against severe rotavirus diarrhea in industrialized and middle income countries but has been less efficacious when tested in infants in the developing world [2,3]. The trivalent oral polio vaccine (OPVERO) has also been found to be less effective in children in the developing world [4] with estimated efficacies of around 21% in India compared to 50% in the United States [5]. In view of this,

*Abbreviations:* ATRA, all-trans retinoic acid; RA, retinoic acid; CCR9, chemokine receptor 9; WGLF, whole gut lavage fluid.

\* Corresponding author at: TROPGAN Office, Department of Internal Medicine, University of Zambia School of Medicine, University Teaching Hospital, Nationalist Road, Lusaka, Zambia.

E-mail address: [mpalamwanza123@gmail.com](mailto:mpalamwanza123@gmail.com) (M. Mwanza-Lisulo).

<https://doi.org/10.1016/j.vaccine.2018.04.083>

0264-410X/© 2018 The Authors. Published by Elsevier Ltd.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

there is an urgent need to identify ways to improve immune responses of these oral vaccines in the settings where they are most needed.

Vitamin A supplementation significantly reduces mortality from diarrheal infections as evidenced by results of several studies [6–9]. The mechanism of protection however is unclear [6,9].

Vitamin A is oxidized to four isoforms of retinoic acid (RA); these include all-trans RA (ATRA), 9-cis RA, 13-cis RA, and 11-cis RA. The RAs have important immunological functions especially on intestinal defense and mucosal immune responses.

Selective migration of the effector T cells to the gut requires the expression of  $\alpha 4\beta 7$ -integrin and the chemokine receptor, CCR9. Integrin  $\alpha 4\beta 7$  is the homing receptor for the small and large intestine, whereas CCR9 is the small intestine-specific homing receptor [10]. Several studies have shown that RA is a key mediator of T cell homing to the gut [11–13]. Iwata and colleagues [11] demonstrated that CD4<sup>+</sup> T cells activated *in vitro* in the presence of ATRA

had enhanced expression of the gut-homing receptors  $\alpha 4\beta 7$ -integrin and CCR9. Hammerschmidt and colleagues [12] demonstrated that RA-assisted subcutaneous immunization resulted in pronounced localization of antigen-specific plasma cells in the small intestine and a robust immunoglobulin A (IgA) response in serum and intestinal washes. In recent preliminary work [14], we found that ATRA could enhance gut IgA responses to Vivotif typhoid vaccine in intestinal washes. Collectively these observations suggest that RA-assisted immunization generates intestinal immunity and confers the benefits of mucosal protection.

Retinoic acid has been known to affect expression of the polymeric immunoglobulin receptor (pIgR) [15] which transports polymeric immunoglobulin A (pIgA) into external secretions as secretory IgA (S-IgA). This suggests that vitamin A might be required for the proper regulation of IgA transport in response to mucosal infection, which would be a desirable characteristic of an orally-active adjuvant [16,17].

The present study investigated whether the ATRA adjuvanticity seen with Vivotif vaccine in African men could be generalized to other oral vaccines. We asked if ATRA given alongside oral vaccines could improve the immune response by enhancing expression of  $\alpha 4\beta 7$ -integrin and CCR9 gut homing molecules, and if so, under what circumstances.

## 2. Methods

### 2.1. Recruitment

We recruited 94 adult male volunteers aged between 18 and 60 years from a high-density township in Lusaka, Zambia, as previously described [14]. A 3-stage consent process was used [20,21]. Nutritional assessment and HIV testing was done. Details in [Supplementary material](#) methods.

### 2.2. Ethical Approval

Approval for the study was obtained from the Biomedical Research Ethics Committee of the University of Zambia (references 012-06-12 and 013-01-14).

### 2.3. Vaccine administration and sample collection

Recruited volunteers ( $n = 94$ ) were randomized to receive one of four oral vaccines: Vivotif (Crucell, Switzerland) [22,23], Dukoral (SBLVaccines), Rotarix (GSK), and OPVERO (Sanofi Pasteur). Each vaccine was given with or without ATRA on day 1. ATRA was administered as 10 mg daily for 8 days and 1 h before vaccination. Vivotif was given on days 1, 3 and 5 while Dukoral, Rotarix and OPVERO were administered as single doses. Whole gut lavage fluid was collected as previously described [14,25] on days 0 and 14 (see [supplementary material](#) methods). Blood was also collected at the same time-points for the serum antibody assays.

### 2.4. Detection of Vivotif and Dukoral-specific IgA and IgG in serum and gut lavages

Vivotif and Dukoral LPS antigen preparations were made using a modified phenol extraction technique as previously described [14,22]. Analysis of vaccine specific antibodies in serum and gut lavages was performed as described previously [14].

### 2.5. Detection of Rotarix-specific IgA and IgG in serum and gut lavages

Analysis of IgA and IgG antibody responses was done using ELISA (see [supplementary material](#)).

### 2.6. Detection of OPVERO-specific IgA and IgG in serum

Analysis of OPVERO-specific antibodies in serum was performed by neutralization.

### 2.7. Analysis of effects of ATRA on polymeric Ig receptor (pIgR)

Biopsies (<20 mg of tissue) were collected from the jejunum of 13 participants and analyzed for the effect of ATRA on expression of pIgR (for details refer to [Supplementary material](#) methods).

### 2.8. Analysis of baseline serum retinol concentrations

Retinol in serum was assayed by high performance liquid chromatography (HPLC) modified version [26].

### 2.9. Transcriptome analysis of effects of ATRA during vaccination

Blood samples were collected from 8 male individuals given both ATRA and Vivotif on day 0 (baseline) and day 8. Each sample comprised 2.5 ml of blood directly drawn into a PAXgene blood RNA tube (Qiagen), inverted 10 times, then kept at room temperature for 2 h before being stored at  $-80^{\circ}\text{C}$  until RNA extraction (see [Supplementary material](#) methods for more details).

### 2.10. Statistical analyses

All graphical and statistical tests were performed using GraphPad Prism 6.0 (La Jolla, CA) and STATA 12 (Stata Corp, College Station, TX, USA). Statistical comparisons were done using nonparametric tests (Mann-Whitney, Kruskal-Wallis and Spearman's rank correlation coefficient). Gut lavage responses are expressed as change in log titer values; P values <0.05 were assumed significant.

## 3. Results

In total, 94 participants were randomized to receive one of 4 vaccines and participant demographics are described in [Table 1](#). The HIV prevalence in this study cohort was 21% with average age being 29 years and nutritional status comparable in each group. We report combined data for HIV infected and non-infected, as we found HIV infection had no effect on the intestinal IgA response to Vivotif ([Supplementary Fig. 1](#)).

### 3.1. ATRA increased gut IgA directed at Vivotif LPS, but not other vaccine antigens

Antibody titers to Vivotif, Dukoral and Rotarix were measured as changes in log titers in response to vaccination with or without daily administration of 10 mg ATRA for 8 days. Antibody titers in whole gut fluid against Vivotif LPS antigen were significantly increased ([Fig. 1A](#)) in the group that received Vivotif vaccine as well as ATRA, consistent with our previous findings [14]. There was no significant change in mucosal responses to Rotarix antigens or Dukoral (whether LPS or CTB) antigens in gut lavage samples ([Fig. 1B-D](#)). Antibody titers in serum showed no significant change in any of the vaccine groups ([Supplementary Fig. 2](#)). This increase in specific IgA was not a consequence of increased pIgR expression in intestinal biopsies ([Supplementary Fig. 3](#)).


### 3.2. Effect of ATRA was vaccine and time dependent

We then analyzed whether expression of gut homing markers was vaccine dependent. We found that there was an increase in

**Table 1**  
Baseline characteristics of study participants.

Number	33	13	7	6	4	5	7	6	13	P
Vaccine	Vivotif	Vivotif	Rotarix	Rotarix	Opvero	Opvero	Dukoral	Dukoral	None	
ATRA	+	–	+	–	+	–	+	–	+	
Age (years)	34 (22–42)	38 (29–40)	21 (19–25)	39 (27–53)	29 (25–33)	20 (19–42)	31 (28–40)	39 (37–40)	28 (22–43)	0.43
HIV Positive (n)	5	2	0	1	1	1	3	4	3	
BMI (kg/m <sup>2</sup> )	20 (19–23)	22 (19–23)	20 (19–21)	20 (19–20)	21 (19–24)	19 (17–25)	20 (18–24)	19 (18–19)	19 (18–24)	0.43

The number of participants randomised to Vivotif includes those in the time course experiments, vaccine comparison experiments and pIgR experiments. Volunteers were all healthy adult men randomised to receive one or none of the vaccines with or without 10 mg ATRA given daily for 8 days. For vaccine comparisons 8 volunteers were recruited, pIgR studies had 13 volunteers while the time course studies had 12 volunteers all given Vivotif + ATRA. These are shown as pooled Vivotif vaccine groups. Continuous variables are shown as median and interquartile range (IQR). ATRA: all-trans retinoic acid, BMI: body mass index.



**Fig. 1.** Change in specific IgA in intestinal lavage fluid in volunteers given one of three vaccines. ‘Change’ is the difference in log titre values of IgA pre- and 14 days post vaccination. Antigen-specific IgA responses to Vivotif and Dukoral LPS, Dukoral CTB and Rotarix antigen were measured in whole gut lavage fluid (WGLF). (A) Specific IgA in WGLF against Vivotif LPS was increased ( $P = 0.01$ ) in Vivotif recipients who received ATRA. No change was seen in responses to the other vaccine antigens: (B) Rotarix ( $P = 0.19$ ), (C) Dukoral LPS ( $P = 0.83$ ) or (D) Dukoral Cholera Toxin B subunit (CTB) ( $P = 0.84$ ). Data were analyzed using Mann-Whitney test. Responses are shown measured as change in log titers.

$\alpha 4\beta 7^+ CD4^+$  (Fig. 2A) and  $\alpha 4\beta 7^+ CD8^+ T$  cells (Fig. 2B) in the Vivotif and Rotarix groups when compared to those that received only ATRA. We only observed slightly enhanced  $CCR9^+ CD4^+ T$  cells (Fig. 2C) but not  $CCR9^+ CD8^+ T$  cells (Fig. 2D) in those given both Vivotif and ATRA. When analyzing only activated (HLA-DR<sup>+</sup>) CD3<sup>+</sup> T cells, it was the CD8<sup>+</sup> cells which showed the greatest change in  $\alpha 4\beta 7$  and CCR9 expression (Supplementary Fig. 4). Conversely, a slight decrease in  $\alpha 4\beta 7^+ DR^+ CD4^+ T$  cells was observed at day 3 ( $P = 0.02$ ) and day 8 ( $P = 0.01$ ) in participants that received both Vivotif and ATRA (Supplementary Fig. 5). These data indicate that the effect of ATRA is seen only in the presence of antigen and on day 14 post vaccination.


### 3.3. ATRA caused a coordinated response of $\alpha 4\beta 7$ integrin and CCR9 only if given simultaneously with Vivotif vaccination

Having established that the co-administration of Vivotif and ATRA resulted in increased  $\alpha 4\beta 7^+ CD4^+$  and  $CCR9^+ CD4^+$  T cells at day 14, we explored the correlation of these two gut-homing


markers. Spearman's rank test revealed a strong correlation ( $\rho = 0.83$ ;  $P < 0.0001$ ) of these gut homing markers on CD4<sup>+</sup> T cells (Fig. 3A), but only when ATRA was given alongside Vivotif. We designated those individuals with a coordinated increase in these two gut homing markers as *positive responders* and those with a coordinated decrease as *negative responders*. A coordinated increase in both gut homing markers was observed in 57% (12/21) of volunteers that received Vivotif and ATRA simultaneously. This effect was not seen with Vivotif alone (Fig. 3B), ATRA alone (Fig. 3C) or Rotarix given with ATRA (Fig. 3D). Our data therefore suggest that the coordinated gut homing response to ATRA is restricted to the situation when co-administered with Vivotif.

### 3.4. Baseline serum retinol was lower in individuals that showed coordinated responses to $\alpha 4\beta 7$ integrin and CCR9

Baseline serum retinol concentrations (Table 2) in the *positive responders* were significantly lower (median of 1.41  $\mu\text{mol/L}$ , interquartile range 1.06–2.48  $\mu\text{mol/L}$ ) than the retinol



**Fig. 2.** Summary of changes on CD4<sup>+</sup> and CD8<sup>+</sup> T cells expressing  $\alpha 4\beta 7$  and CCR9. Volunteers received either one of 3 vaccines (Vivotif, Rotarix or Opvero) plus ATRA or ATRA alone. (A) Change in  $\alpha 4\beta 7$  expression on CD4<sup>+</sup> T cells was significantly higher in participants that received Vivotif with ATRA ( $P < 0.05$ ) or Rotarix with ATRA ( $P = 0.02$ ). (B) Change in  $\alpha 4\beta 7$  expression on CD8<sup>+</sup> T cells was also significantly higher in the group that received Vivotif with ATRA ( $P = 0.03$ ) or Rotarix with ATRA ( $P = 0.002$ ). (C) Change in CCR9 expression on CD4<sup>+</sup> T cells was significantly higher in participants that received Vivotif with ATRA ( $P = 0.02$ ). (D) CCR9 expression on the CD8<sup>+</sup> T cells; no effect of ATRA was observed. The changes were calculated by subtracting the pre- vaccination CD4<sup>+</sup> T cells expressing the gut homing markers of interest from the post vaccination CD4<sup>+</sup> T cells. The effects of ATRA were analyzed using the Kruskal-Wallis test.



**Fig. 3.** ATRA induced coordinated expression of  $\alpha 4\beta 7$ -integrin and CCR9 on CD4<sup>+</sup> T cells in Vivotif recipients. (A) Change in  $\alpha 4\beta 7$ +CD4<sup>+</sup> and CCR9+CD4<sup>+</sup> was closely correlated ( $\rho = 0.83$ ;  $P < 0.0001$ ) in participants that received Vivotif with ATRA. Participants given (B) Vivotif alone ( $\rho = 0.12$ ;  $P = 0.83$ ), (C) ATRA alone ( $\rho = 0.07$ ;  $P = 0.82$ ), or (D) Rotarix with ATRA ( $\rho = 0.39$ ;  $P = 0.38$ ) showed no correlation in gut marker expression respectively. Correlations were analyzed using the Spearman rank correlation test.

concentration in the *negative responders* (median 2.68  $\mu\text{mol/L}$ , interquartile range 1.68–3.29;  $P = 0.03$ ). In the *positive responder* group, 40% (4/10) of the subjects had serum retinol concentrations of  $\leq 1.05 \mu\text{mol/L}$ , consistent with vitamin A deficiency, compared to 0% (0/7) of *negative responders*.

### 3.5. The change in $\alpha 4\beta 7$ on DR<sup>+</sup> CD4<sup>+</sup> T cells correlated strongly with the intestinal IgA response to Vivotif

We then looked for a correlation between the change in gut homing marker expression in blood and the intestinal IgA response to Vivotif (i.e. gut lavage IgA to Vivotif LPS). The change in  $\alpha 4\beta 7$  marker expression on CD4<sup>+</sup>T cells was strongly correlated with

**Table 2**  
Baseline characteristics of positive responders and negative responders.

Characteristic	Positive responders	Negative responders	P
n	12	9	
Sex	Male	Male	
Age (mean, years)	36	29	
HIV seropositive (n)	3	2	
Serum retinol ( $\mu\text{mol/L}$ )	1.4 (1.06–2.48)	2.68 (1.68–3.29)	0.03

A retinol concentration of 1  $\mu\text{mol/L}$  was equivalent to 28.


the change in specific IgA response to Vivotif LPS in the intestine in the *positive responders* only (Fig. 4A). In *negative responders* (Fig. 4B), or those given Vivotif alone (Fig. 4C), there was no correlation.

### 3.6. Transcriptome analysis

To elucidate the transcriptional changes that occurred during co-administration of ATRA and the oral vaccine Vivotif, we performed RNA-Seq on blood samples. Transcriptomic changes in immune-related pathways, particularly including interferon  $\alpha/\beta$  signaling pathway, membrane-ECM interactions and immune hubs were observed (Supplementary Table 1). Key immune hubs identified include fibronectin 1 (FN1), AXL Tyrosine-protein kinase receptor, complement component 1 (C1QB), complement component 4 binding protein (C4BPA) and HLA-DRB1 (Supplementary Fig. 6).

## 4. Discussion

The global burden of diarrheal disease and the reduced efficacy of oral vaccines in populations from developing nations have highlighted the need for new strategies for vaccination against



**Fig. 4.** Correlation of change in gut IgA to change in  $\alpha 4\beta 7^+DR^+CD4^+$  cells in the positive and negative responders. (A) The positive responders showed a strong correlation ( $\rho = 0.82$ ;  $P = 0.02$ ) of gut IgA to gut marker expression. (B) This was not observed in the negative responders ( $\rho = 0.31$ ,  $P = 0.2$ ) or those given (C) Vivotif alone ( $\rho = 0.37$ ;  $P = 0.2$ ). Correlations were analyzed using the Spearman rank correlation test.

diarrheal diseases, for example by using oral adjuvants. This study has revealed four important effects of ATRA on immune responses to oral vaccination: First, we have confirmed earlier findings [14] that ATRA can increase gut mucosal IgA directed at an oral live attenuated typhoid vaccine, and that this increase is not attributable to the increased pIgR expression as had been suggested previously [15]. Secondly, our data suggests that ATRA can increase the number of circulating  $CD4^+$  T cells expressing the gut homing markers  $\alpha 4\beta 7$  and CCR9 in a coordinated and vaccine-specific fashion in a subset of individuals. Thirdly, integrin  $\alpha 4\beta 7$  increase was strongly correlated with the intestinal IgA responses in *positive responders*, i.e. those individuals with a coordinated increase in the gut homing markers. Lastly, we found that ATRA treatment may cause changes in immune related pathways, including interferon  $\alpha/\beta$  signaling pathway (adjusted  $p$ -value = 0.0015), membrane-ECM interactions, and immune hubs including fibronectin 1, AXL kinase, complement pathway components and major histocompatibility complex components such as HLA-DRB1. Since these elements were impacted by the end of the 8 day ATRA treatment, we propose that they are responsible, in part, for the increased responses to Vivotif vaccine.

We have demonstrated previously [14] that, in whole gut fluid, ATRA is able to enhance specific IgA against vaccine antigens suggesting that ATRA could play an adjuvant role when given alongside Vivotif. To determine whether the adjuvanticity of ATRA could be translated to other oral vaccines, we evaluated its effect on three other vaccines; Rotarix, Dukoral and OPVERO vaccines, all orally administered one hour after administration (or not) of 10 mg ATRA given daily for 8 days. Our data revealed that ATRA adjuvanticity in this study only occurred with Vivotif and not the other vaccines, suggesting that the ATRA effect might be vaccine and/or antigen specific and not a generalizable adjuvant property. There may be several contributors to the specificity of ATRA in promoting adjuvanticity with Vivotif and not the other oral vaccines. The most important factor is likely to be the type of immune responses being elicited by the vaccines. Vivotif induces a vigorous T-cell response that favours the production of interferon gamma ( $IFN-\gamma$ ), tumour necrosis factor- $\alpha$  ( $TNF-\alpha$ ), interleukin -1 ( $IL-1$ ) and  $IL-6$ , indicative of a Th1-type response probably through Class II antigen presentation [31,32]. Rotavirus has been shown to drive the secretion of  $TGF-\beta$  resulting in inhibition of Th1 responses by dendritic cells [33]. In one study [34], Rotarix vaccination showed



no discernible effect on IFN- $\gamma$  and TNF- $\alpha$  response in a considerable proportion [76% (16/21)] of vaccinees. Another study [35] showed that oral polio vaccine was associated with down regulation of cytokine (IFN- $\gamma$  and TNF- $\alpha$ ) production when co-administered with BCG. The authors speculated that polio virus might have specific immune modulatory molecules that down-regulate immune responses to antigens to which immune responses are being mounted simultaneously. Finally, the inactivated bacterial vaccine Dukoral tends to elicit diminished or skewed CD4<sup>+</sup> T cell responses towards development of a Th2 T cell phenotype because of the presence of CTB [36]. Of the vaccines studied, Vivotif was the only vaccine that is likely to elicit a Th1-type immune response through Class II antigen presentation. Intriguingly, ATRA led to an increase in several MHC class II components including a major component enabling antigen recognition by T cells, HLADRB1. We hypothesize here that it is the nature of the context of antigen presentation, in this case for a vaccine that generates an IFN- $\gamma$  dominant response, which determines whether ATRA has an effect [37]. Also, in a pro-inflammatory context involving cytokines, such as IL-15, ATRA probably acts through dendritic cells to decrease conversion of naïve T cells into T regs and to enhance Th1 cell polarization. However, this hypothesis will have to be tested empirically by inclusion of another vaccine group that elicits a Th1 type response, and that the experimental protocol would need to include direct demonstration of Th1 and Tregs in comparison with Th2 and Th17 responses.

To our knowledge, this is only the second study to evaluate ATRA effects on immune function in humans *in vivo*, since most other studies [11,13,38–43] have used *in vitro* systems or animal models. Our findings showed an increase in CD4 T cells expressing  $\alpha 4\beta 7$  and CCR9 following oral administration of 10 mg ATRA but only when given with Vivotif but not Rotarix or OPVERO vaccines. These results are consistent with other studies in animal models [11,12,39,40].

There is also evidence that ATRA has effects on other cell types that also possibly migrate to the gut. Pantazi and colleagues [13] demonstrated in a murine model, that retinoic acid signaling in B cells is not essential for their homing to Peyer's Patches but is necessary in generating antigen-specific IgA responses in the gut. ATRA has also been found to enhance the migratory properties of dendritic cells, which is crucial for their antigen presenting function during infection. We must emphasize that while it would have been of interest to extend the scope of our work to include B cells, NK cells and DCs, in a human study one is always constrained by the experimental procedures which we can conduct, acceptable volumes of blood to be drawn, and the invasiveness of endoscopic procedures. Our attention in this study was drawn to T cells because of the mouse work by other groups [11,41,42] and our own previous study [14] suggesting that ATRA can augment responses to Vivotif vaccine.

For safety reasons, one limitation of this study is that it was carried out only in men due to the teratogenicity of ATRA [18,19]; and with no clear safe dose, ATRA could not be used in women who might become pregnant during the study. As other studies [44,45] confirm differences between males and females in response to vaccination and we suspect that responses in women would have been different. A further limitation is that we did not analyse the gut-resident cells to see the effect of ATRA on the gut immune responses, as we did not have ethical approval to perform gut lavage and endoscopic biopsies on the same participants.

When the baseline serum retinol concentration was analyzed, we found that 4/10 (40%) of the *positive responders* had a clear or borderline vitamin A deficiency but none of the *negative responders* had such deficiency. The difference observed might signify that the effect of ATRA is seen predominantly in people with borderline vitamin A deficiency. Also, the data could suggest that the vitamin

A deficiency seen in the positive responders may have resulted in the dendritic cells being altered toward being more inflammatory producing high levels of IL-13 and TNF- $\alpha$  [46] thus creating a favorable environment in which ATRA has an effect. The response to ATRA and Vivotif was also not explained by HIV infection. A high proportion of this variability therefore remains unexplained, as evident from our RNA-Seq studies, so the possibility of an immunogenetic predisposition needs to be considered in future work.

It must be pointed out here that ATRA is also obtained from vitamin A in the diet, and the authors did not control or attempt to control for diet in the cohort because of the recall bias that doing a food frequency questionnaire has. It is also apparent from pharmacological data [14] that oral administration of 10 mg ATRA over a period of 8 days generates a sharp rise in serum concentrations of ATRA required to produce the desired immune effects that were the focus of the current study. Also of note is that ATRA levels were not monitored longitudinally but only evaluated at two time points; baseline and day 14 post vaccination.

### Acknowledgements

We thank Dr. David Allen, Public Health England, Colindale, UK for assistance with analysis of OPVERO specific antibodies in serum. We thank the Tropical Diseases Research Centre, Ndola Central Hospital, Ndola, Zambia (TDRC) for help with analyzing the retinol concentrations in serum samples. We are also grateful to Dr. Yusuf Ahmed, Dr. Patrick Musonda and Dr. Andrew Prendergast for work on the Data and Safety Monitoring Board.

### Funding

This work was supported by the Bill and Melinda Gates Foundation through a Grand Challenges Explorations grant (OPP1043696).

### Disclosure

The authors declare no conflicts of interest.

### Author contributions

P.K. originated the study and is guarantor. M.M-L. and P.K designed the study and the outcome measures. M.S.C, M.C, E.B, E. F and K.Z carried out sample collection and laboratory assays. R. B, M.I and P.K supervised clinical data collection. M.M-L wrote the first draft of the manuscript. All authors contributed to and approved the final manuscript, including the authorship list.

### Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.vaccine.2018.04.083>.

### References

- [1] Levine MM. Immunogenicity and efficacy of oral vaccines in developing countries: lessons from a live cholera vaccine. *BMC Bio* 2010;8:129.
- [2] Vesikari T, Karvonen A, Prymula R, et al. Efficacy of human rotavirus vaccine against rotavirus gastroenteritis in the first two years of life in European infants: randomized double-blind controlled study. *Lancet* 2007;370:1757–63.
- [3] Madhi SA, Cunliffe NA, Steele D, et al. Effect of human Rotavirus vaccine on severe diarrhea in African Infants. *N Engl J Med* 2010;362:289–98.
- [4] Grassly NC, Jafari H, Bahl S, Durrani S, Wenger J, Sutter RW, et al. Asymptomatic wild-type poliovirus infection in India among children with previous oral poliovirus vaccination. *J Infect Dis* 2010;201:1535–43.
- [5] Grassly NC, Fraser C, Wenger J, Deshpande JM, Sutter RW, Heymann DL, et al. New strategies for the elimination of polio from India. *Science* 2006;314:1150–3.

- [6] Sommer A. Vitamin A deficiency and clinical disease: an historical overview. *J Nutr* 2008;138:1835–9.
- [7] Ghana Vitamin A. Supplementation Trials (VAST) Study Team. Vitamin A supplementation in Northern Ghana: effects on clinic attendances, hospital admissions and child mortality. *Lancet* 1994;344:228–31.
- [8] Long K, Rosado JL, DuPont HL, Hertzmark E, Santos JL. Supplementation with vitamin A reduces watery diarrhea and respiratory infections in Mexican children. *Br J Nutr* 2007;97:337–43.
- [9] Hall JA, Cannons JL, Dos Santos LM, et al. Essential role for retinoic acid in the promotion of CD4<sup>+</sup> T cell effector responses via retinoic acid receptor alpha. *Immunity* 2011;34:435–77.
- [10] Kim HM, Taparowsky EJ, Kim CH. Retinoic acid differentially regulates the migration of innate lymphoid cell subsets to the gut. *Immunity* 2015;43:107–19.
- [11] Iwata M, Hirakiyama A, Eshima Y, Kagechika H, Kato C, Song S. Retinoic acid imprints gut-homing specificity on T cells. *Immunity* 2004;21:527–38.
- [12] Hammerschmidt S, Friedrichsen M, Boelter J, Lyszkiewicz M, Kremmer E, Pabst O, et al. Retinoic acid induces homing of protective T and B cells to the gut after subcutaneous immunization of mice. *J Clin Invest* 2011;121:3051–61.
- [13] Pantazi E, Marks E, Stolarczyk E, Lycke N, Noelle R, Elgueta R. Retinoic acid signaling in B-cells is essential for oral immunization and microflora composition. *J Immunol* 2015;195(4):1368–71.
- [14] Lisulo MM, Kapulu MC, Banda R, Sinkala E, Kayamba V, Sianonogo S, et al. Adjuvant potential of low dose all-trans retinoic acid during oral typhoid vaccination in Zambian men. *Clin Exp Immunol* 2014;175:468–75.
- [15] Takenouchi ON, Asano M, Chihaya H, Chung-Hsuing WU, Ishikasa K, Moro I. Retinoic acid enhances the gene expression of human polymeric immunoglobulin receptor (pIgR) by TNF- $\alpha$ . *Clin Exp Immunol* 2004;135:448–54.
- [16] Reed GS, Orr MT, Fox CB. Key role of adjuvants in modern vaccines. *Nat Med* 2013;19:1597–608.
- [17] Pulendran B, Ahmed R. Immunological mechanisms of vaccination. *Nat Immunol* 2011;12:509–17.
- [18] Lee L, Leung CY, Tang WW, et al. A paradoxical teratogenic mechanism for retinoic acid. *Proc Natl Acad Sci USA* 2012;109:13668–73.
- [19] Adams J. The neurobehavioral teratology of retinoids: a 50-year history. *Birth Defects Res A* 2010;88:895–905.
- [20] Kelly P, Todd J, Sianonogo S, Mwansa J, Sinsungwe H, Katubulushi M, et al. Susceptibility to intestinal infection and diarrhea in Zambian adults in relation to HIV status and CD4 count. *Gastroenterology* 2009;9:7.
- [21] Zulu J, Mwanza-Lisulo M, Besa E, et al. Improving validity of informed consent for biomedical research in Zambia using a laboratory exposure intervention. *PLoS One* 2014;9(9):e108305. <https://doi.org/10.1371/journal.pone.0108305>.
- [22] Luderitz O, Risse HJ, Schulte-Holthausen H, Strominger JL, Sutherland IW, Westphal O. Biochemical studies of the smooth-rough mutation in *Salmonella* minnesota. *J Bacteriol* 1965;89:343–54.
- [23] Wahid R, Simon R, Zafar S, Levine M, Szein MB. Live oral typhoid vaccine Ty21a induces cross-reactive humoral immune responses against *Salmonella entericaserovar Paratyphi A* and *S. Paratyphi B* in humans. *Clin Vaccine Immunol* 2012;19:825–34.
- [24] Daley A, Randall R, Darsley M, et al. Genetically modified enterotoxigenic *Escherichia coli* vaccines induce mucosal immune responses without inflammation. *Gut* 2007;56:1550–6.
- [25] Kim YK, Quadro L. Reverse-phase high-performance liquid chromatography (HPLC) analysis of retinol and retinyl esters in mouse serum and tissues. *Methods Mol Biol* 2010;652:263–75.
- [26] Breuer K, Ferooshani AK, Laird MR, et al. InnateDB: systems biology of innate immunity and beyond—recent updates and continuing curation. *Nucl Acids Res (Database issue)* 2013;41:D1228–33.
- [27] Xia J, Gill EE, Hancock REW. NetworkAnalyst for statistical, visual and network-based approaches for meta-analysis of expression data. *Nat Protocols* 2015;10:823–44.
- [28] Xia J, Benner MJ, Hancock REW. NetworkAnalyst – integrative approaches for protein–protein interaction network analysis and visual exploration. *Nucl Acids Res* 2014;42:W167–74.
- [29] Postigo AA, Sanchez-Mateos P, Lazarovits AI, Sanchez-Madrid F, de Landazuri MO.  $\alpha 4\beta 7$  integrin mediates B cell binding to fibronectin and vascular cell adhesion molecule-1. *J Immunol* 1993;151:2471–83.
- [30] Lundin BS, Johansson C, Svennerholm A. Oral immunization with a *Salmonella enterica* serovar Typhi vaccine induces specific circulation mucosa-homing CD4<sup>+</sup> and CD8<sup>+</sup> T cells in humans. *Infect Immun* 2002;70:5622–7.
- [31] Pasetti M, Simon M, Szein M, Levine M. Immunology of gut mucosal vaccines. *Immunol Rev* 2011;239:125–48.
- [32] Clarke E, Desselberger U. Correlates of protection against human rotavirus disease and the factors influencing protection in low-income settings. *Mucosal Immunol* 2015;8:1–17.
- [33] Banda R, Yambayamba V, Lalusha BD, Sinkala E, Kapulu MC, Kelly P. Safety of live, attenuated oral vaccines in HIV- infected Zambian adults. *Oral vaccine in HIV*. *Vaccine* 2012;30:5656–60.
- [34] Sartono E, Lisse IM, Terveer EM, et al. Oral polio vaccine influences the immune response to BCG vaccination. A natural experiment. *PLoS ONE* 2010;5(5):e10328. <https://doi.org/10.1371/journal.pone.0010328>.
- [35] Jensen KJ, Karkov HS, Lund N, et al. The immunological effects of oral polio vaccine provided with BCG vaccine at birth: a randomized trial. *Vaccine* 2014;32:5949–56.
- [36] Kuchta A, Rahman T, Sennott ET, et al. *Vibrio cholerae* O1 infection induces proinflammatory CD4<sup>+</sup> T cell responses in blood and intestinal mucosa of infected humans. *Clin Vaccine Immunol* 2011;18:1371–7.
- [37] Brown CC, Esterhazy D, Sarde A, et al. Retinoic acid is essential for Th1 cell lineage stability and prevents transition to a Th17 cell program. *Immunity* 2015;42:499–511.
- [38] Kang S, Lim H, Andrisani O, Broxmeyer H, Kim C. Vitamin A metabolites induce gut-homing FoxP3<sup>+</sup> regulatory T cells. *J Immunol* 2007;179:3724–33.
- [39] Eksteen B, Mora JR, Houghton EL, et al. Gut homing receptors on CD8 T cells are retinoic acid dependent and not maintained by liver dendritic or stellate cells. *Gastroenterology* 2009;137:320–9.
- [40] Tang X, Gudas LJ. Retinoids, retinoic acid receptors and cancer. *Annu Rev Pathol Mech Dis* 2011;6:345–64.
- [41] Kaufman DR, De Calisto J, Simmons NL, Cruz AN, Villablanca EJ, Mora JR, et al. Vitamin A deficiency impairs vaccine-elicited gastrointestinal immunity. *J Immunol* 2011;187:1877–83.
- [42] Mora JR, Bono MR, Manjunath N, et al. Selective imprinting of gut-homing T cells by peyer's patch dendritic cells. *Nature* 2003;424:88–93.
- [43] Aaby P, Jensen H, Gomes J, Fernandes M, Lisse IM. The introduction of diphtheria-tetanus-pertussis vaccine and child mortality in rural Guinea-Bissau: an observational study. *Int J Epidemiol* 2004;33:374–80.
- [44] Klein S, Marriot I, Fish E. Sex-based differences in immune function and responses to vaccination. *Trans R Soc Trop Med Hyg* 2015;109:9–15.
- [45] Yokota-Nakatsuma A, Takeuchi H, Ohoka Y, et al. Retinoic acid prevents mesenteric lymph node dendritic cells from inducing IL-13 producing inflammatory Th2 cells. *Mucosal Immunol* 2014;7(4):786–801. <https://doi.org/10.1038/mi.2013.96>. Epub 2013 Nov 13.
- [46] Kim D, Perteau G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol* 2013;14(4):R36.
- [47] Li H, Handsaker B, Wysoker A, et al. The Sequence alignment/map (SAM) format and SAMtools. *Bioinformatics* 2009;25:2078–9.
- [48] Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* 2015;31(2):166–9.
- [49] Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014;15:550.

### Further reading

- [24] Lucas MES, Dean JL, von Seidlein L, et al. Effectiveness of mass oral cholera vaccination in Beira, Mozambique. *N Engl J Med* 2005;352:757–67.