

White-winged Vampire Bat, *Diaemus youngi* (Jentink, 1893) (Mammalia, Chiroptera): range extension in the Cerrado biome and new locality in Mato Grosso do Sul, southwestern Brazil

Gustavo Lima Urbietta,^{1,3} Thawane Yvin Sanches de Siqueira² & Gustavo Gracioli¹

¹Programa de Pós-Graduação em Biologia Animal, Laboratório de Sistemática, Ecologia e Evolução (LSEE), Centro de Ciências Biológicas e da Saúde, Universidade Federal de Mato Grosso do Sul (UFMS), 79090-900, Campo Grande, MS, Brazil

²Ciências Biológicas, Laboratório de Ecologia de Fragmentos (EcoFrag), Universidade Católica Dom Bosco, CEP 79117-900, Campo Grande, MS, Brazil

³Corresponding author. E-mail: gustavo.cx@hotmail.com

Abstract. We document a new locality of *Diaemus youngi* in the Cerrado biome, extend the distribution in Mato Grosso do Sul state (southwestern Brazil), and present the first record of a vampire bat species from the state's capital, Campo Grande. We recorded one male adult in an urban remnant of Cerrado. There are few locality data for this species in the Cerrado biome and our report significantly adds to the knowledge of this species' distribution within the state and throughout Brazil.

Key words. Bats; distribution; urban area; secondary rainforest; Campo Grande

Brazil presents the second largest diversity of bat species in South America, and up to 9 families, 68 genera, and 178 species have been recorded (NOGUEIRA et al. 2014), but information on the occurrence and distribution of some species is still lacking. This is especially so for areas within the Cerrado domain, such as in the state of Mato Grosso do Sul, which is under anthropogenic threats.

In the most recent checklist of bats from Mato Grosso do Sul, FISCHER et al. (2015) recorded 74 species of bats belonging to 42 genera and 7 families. However, for some subfamilies, such as Desmodontinae, occurrences were scarce or recorded from isolated localities, making it difficult to understand the distribution of bat species in Mato Grosso do Sul and in the Cerrado. The subfamily Desmodontinae includes 3 species (*Desmodus rotundus* (Geoffroy, 1810), *Diphylla ecaudata* Spix, 1823, and *Diaemus youngi* (Jentink, 1893), with hematophagous feeding behavior. (GREENHALL & SCHUTT 1996).

Diaemus youngi (Jentink, 1893) preferably feeds on the blood of birds (UIEDA 1993). It is rare throughout Brazil (e.g., AGUIAR et al. 2006; GREENHALL & SCHUTT 1996), and therefore, information on its conservation, biology, and ecology is important (COSTA et al. 2008). Here, we report a new locality record of *D. youngi* in the Cerrado and extend its distribution. This represents the first record of *D. youngi* and a hematophagous bat species from Campo Grande, the state's capital.

We captured the specimen at the São Vicente Institute (20°23'08.5" S, 054°36'28.8" W), in the municipality of Campo Grande, capital of Mato Grosso do Sul state. The institute has an area of 191 ha, of which 20 ha are designated as legal reserve and about 30 ha are used for farming of crops such as corn, beans, and soy (CEREZOLI et al. 2014). Additionally, at the site there is a fish pond for fish farming and livestock (e.g., cattle, goats, sheep, and horses) are raised (CEREZOLI et al. 2014). The study area is characterized as a remnant of the Cerrado biome within an urban area.

In November 2016 we captured the specimen with a permanent license (number 10.566) issued by the Instituto Brasileiro de Meio Ambiente (IBAMA). We used 6 mist nets (9.0 m × 3.0 m) at a height of 0.5 m above the ground and maximum height of 3.5 m. These were distributed on trails, in clearings, and at the edges of and inside the remnant. The individual captured



Figure 1. The individual *Diaemus youngi* (Jentink, 1893) captured in the present work. (Photographed by Allan Corral).



Figure 2. The arrows show the main external morphological features of *Diaemus youngi* (Jentink, 1893): the ear edges and wings are white, as well as the membrane between the second and third fingers. (Photographed by Allan Corral).

was deposited in the zoological reference collection of Universidade Federal de Mato Grosso do Sul (ZUFMS) with the voucher code ZUFMS-CHI02310.

To identify the specimen we followed diagnoses of REIS et al. (2007) and the key by GARDNER (2008). The male adult specimen captured was identified as *D. youngi* (Figure 1). According to REIS et al. (2007), *D. youngi* resembles *Desmodus rotundus* (E. Geoffroy 1810) and *Diphylla ecaudata* Spix, 1823, but can be easily distinguished from these other vampire bats because it lacks a calcar and has no evident tail. The thumb of *D. youngi* has a single pad while *D. rotundus* has 2 pads (AGUIAR et al. 2006). In *D. youngi*, both sexes possess cup-shaped oral scent glands located bilaterally inside the mouth. These scent glands are only seen when the bat is disturbed and directs the glands forward and emits an offensive odor (GREENHALL & SCHUTT 1996). Other features that clearly distinguish *D. youngi* from other vampire bats are the white edges of its ears and wings, as well as the white membrane between the second and third fingers (GREENHALL & SCHUTT 1996) (Fig. 2).

Up to 2006 only 3 records of *D. youngi* were recorded in the Cerrado (AGUIAR et al. 2006); ours is the tenth record from the Cerrado (Fig. 3). Within Mato Grosso do Sul, our record is 143 km away from the closest locality recorded for this species in state. Most of the known records of *D. youngi* are distributed in the Pantanal biome (7 localities), with only 2 localities in the Cerrado of the Mato Grosso do Sul (FISCHER et al. 2015), and ours is the third record from the Cerrado of the state.

Campo Grande has an area of 8,092,951 (km²) and 863,982 inhabitants (IBGE 2016). The capture location is considered an area of transition between urban and periurban perimeters, with some housing complexes and neighborhoods nearby.

The capture of this species in an urban area should alert us to the possibility of potential contact with humans, which was emphasized by ITO et al. (2016) who reported human blood in the diet of *Diphylla ecaudata*. However, species like *D. youngi*

are considered specialists for bird blood, so while the habitat fragmentation and loss can negatively affect the availability of natural prey, conversion of natural landscape to livestock farming can greatly increase the availability of introduced prey (BOBROWIEC et al. 2015). Hence, vampire bats may favor some degree of environmental disturbance, especially when such disturbances lead to an increase in the availability of domesticated animals, such as chickens, goats, and cows (FENTON et al. 1992, GOMES et al. 2010).

The process of habitat fragmentation in Mato Grosso do Sul can have varying consequences for the regional biodiversity, by changing species abundance and increasing the abundance of species adapted to anthropic environments (WILLIG et al. 2007). We also understand that the presence of *D. youngi* in an urban area of Cerrado may be due to the destruction of caves, habitat, and deforestation in nearby fragmented areas, which may cause these bats to inhabit new urban environments (PACHECO et al. 2010).

ACKNOWLEDGEMENTS

We thank Instituto de Pesquisa São Vicente and the Universidade Católica Dom Bosco (UCDB) for accommodating the researchers, CAPES (process number 1591062), CNPq (process number 304616/2015-0) and Fundect for scholarships. We also thank Luciana Mendes Valério for making available to us her laboratory, Elaine Anjos for help with English translation, and Allan Corral for the photographs.

LITERATURE CITED

AGUIAR, L.M.S., W.R. CARMAGO & A.S. PORTELA. 2006. Occurrence of White-winged Vampire Bat, *Diaemus youngi* (Mammalia, Chiroptera), in the Cerrado of Distrito Federal, Brazil. *Revista Brasileira de Zoologia* 23: 893–896. <http://doi.org/c45mb6>




Figure 3. Distribution of *Diaemus youngi* (Jentink, 1893) in the Cerrado biome. Red star: new locality. Black circles 1–9: other localities, in Table 1. Geodetic datum used: WGS84.

Table 1. Localities (numbers correspond to Figure 3) and specimens of *Diaemus youngi* in Cerrado, Brazil. State abbreviations: MS, Mato Grosso do Sul; DF, Distrito Federal; MG, Minas Gerais; TO, Tocantins; MA, Maranhão.

| Number | Locality | State | Source | Latitude (S) | Longitude (W) |
|--------|------------------------------|-------|-------------------------|--------------|---------------|
| 1 | Caimã Farm and resort | MS | ALHO et al. (2011) | 19°18'13.3" | 056°01'24.5" |
| 2 | Coxim city | MS | FISCHER et al. (2015) | 18°33'45.04" | 054°47'01.70" |
| 3 | Brazlândia city | DF | AGUIAR et al. (2006) | 15°30' | 048°10' |
| 4 | Uberlândia city | MG | UIEDA & ARAÚJO (1987) | 18°55'18.06" | 048°16'30.60" |
| 5 | Uberaba city | MG | UTUTZ et al. (2004) | 19°44'59.76" | 047°56'20.20" |
| 6 | Esmeraldas city | MG | UIEDA (1993) | 19°45'56.70" | 044°17'08.80" |
| 7 | São José da Lapa city | MG | TORQUETTI et al. (2013) | 19°42'44.72" | 043°56'44.60" |
| 8 | Mateiros city | TO | GREGORIN et al. (2011) | 10°40'47.36" | 046°52'46.90" |
| 9 | APA Inhamum | MA | OLÍMPIO et al. (2016) | 04°53'29.81" | 043°24'48.70" |
| 10 | Campo Grande city (red star) | MS | This study | 20°23'08.5" | 054°36'28.8" |

ALHO, C. JR., E. FISCHER, L.F. OLIVEIRA-PISSINI & C.F. SANTOS. 2011. Bat-species richness in the Pantanal floodplain and its surrounding uplands. *Brazilian Journal of Biology* 71: 311–320. <http://doi.org/b7ns>

BOBROWIEC, P.E.D., M.R. LEMES & R. GRIBEL. 2015. Prey preference of the common vampire bat (*Desmodus rotundus*, Chiroptera) using molecular analysis. *Journal of Mammalogy* 96: 54–63. <http://doi.org/b7nt>

CEREZOLI, C.C.J., J.M. TORRES & E.A.C ANJOS. 2014. Plants consumed by bats at Lagoa da Cruz, Campo Grande, MS. *Multitemas* 45: 123–136.

COSTA, L.M., D.M. OLIVEIRA, A.F.P.D. FERNANDES & C.E.L. ESBERARD. 2008. Ocorrência de *Diaemus youngi* (Jentink, 1893)

Chiroptera, no Estado do Rio de Janeiro. *Biota Neotropica* 8: 2017–220. <http://doi.org/dhxhn2>

FENTON, M.B. 1992. Wounds and the origin of blood-feeding in bats. *Biological Journal of the Linnean Society* 47: 161–171. <http://doi.org/bwbs5g>

FISCHER, E., C.F. SANTOS, L.F.A.C. CARVALHO, G. CAMARGO, N.L. CUNHA, M. SILVEIRA, M.O. BORGINON & C.L. SILVA. 2015. Bat fauna of Mato Grosso do Sul, southwestern Brazil. *Biota Neotropica* 15: 1–16. <http://doi.org/f7fm96>

GARDNER, A.L. 2008. Subfamily Desmodontidae, pp. 218–224, in: A.L. GARDNER (ed.). *Mammals of South America. Volume 1, marsupials, xenarthrans, shrews, and bats*. Chicago: University of Chicago Press.

- GOMES, M.N., A.M.V. MONTEIRO, N. LEWIS, C.A. GONÇALVES & V.S. NOGUEIRA-FILHO. 2010. Landscape risk factors for attacks of vampire bats on cattle in São Paulo, Brazil. *Preventive Veterinary Medicine* 93: 139–146. <https://doi.org/10.1016/j.prevetmed.2009.10.006>
- GREENHALL, A.M. & W.A. SCHUTT JR. 1996. *Diaemus youngi*. *Mammalian Species* 533: 1–7.
- GREGORIN, R., E. GONÇALVES, C.C. AIRES & A.A.P. CARMIGNOTTO. 2011. Morcegos (Mammalia: Chiroptera) da Estação Ecológica Serra Geral do Tocantins: composição específica e considerações taxonômicas. *Biota Neotropica* 11: 299–312.
- IBGE. 2016. Cidades. Instituto Brasileiro de Geografia e Estatística. Accessed at <http://cod.ibge.gov.br/CMC>, 4 April 2016.
- ITO, F., E. BERNARD & R.A. TORRES. 2016. What is for dinner? First report of human blood in the diet of the hairy-legged vampire bat *Diphylla ecaudata*. *Acta Chiropterologica* 18: 509–515. <https://doi.org/10.3161/15081109ACC2016.18.2.017>
- NOGUEIRA, M. R., I.P. LIMA, R. MORATELLI, V.C. TAVARES, R. GREGORIN & A.L. PERRACCHI. 2014. Checklist of Brazilian bats, with comments on original records. *Check List* 10: 808–821. <https://doi.org/10.15560/10.4.808>
- OLÍMPIO, A.P.M., M.C.S. VENTURA, M.J.O. MASCARENHAS, D.C. NASCIMENTO, F.A. DE ANDRADE, E.C. FRAGA & M.C. BARROS. 2016. Bat fauna of the Cerrado savanna of eastern Maranhão, Brazil, with new species occurrences. *Biota Neotropica* 16: 1–6. <https://doi.org/10.1590/1676-0611-BN-2015-0089>
- PACHECO, S.M., M. SODRÉ, A.R. GAMA, A. BREDT, E.M. CAVALLINI-SANCHES, R.V. MARQUES, M.M. GUIMARÃES & G. BIANCONI. 2010. Morcegos urbanos: status do conhecimento e plano de ação para a conservação no Brasil. *Chiroptera Neotropical* 16: 630–647.
- REIS, N.R., A.L. PERRACCHI, W.A. PEDRO & I.P. LIMA. 2007. Morcegos do Brasil. Londrina: Editora da Universidade Estadual de Londrina. 253 pp.
- STUTZ, W.H., M.C. ALBUQUERQUE, W. UIEDA, E.M. MACEDO & C.B. FRANCA. 2004. Updated list of Uberlândia bats (Minas Gerais state, southeastern Brazil). *Chiroptera Neotropical* 10: 188–190.
- TORQUETTI, C.G., M.A.C. ASSIS, S.S.A. CARMO & S.A. TALAMONI. 2013. Observations of a diurnal roost of the White-winged Vampire Bat *Diaemus youngi* in a karstic area of southeastern Brazil. *Chiroptera Neotropical* 19: 1147–1150.
- UIEDA W. & V.F. DE ARAÚJO. 1987. Manutenção dos morcegos hematófagos *Diaemus youngii* e *Diphylla ecaudata* (Chiroptera, Phyllostomidae), em Cataveiro [unpublished report]. Uberaba, ON: Seminário de Ciências Fiurbe. 30 + [42] pp.
- UIEDA, W. 1993. Comportamento alimentar do morcego hematófago *Diaemus youngi*, em aves domésticas. *Revista Brasileira de Biologia* 53: 529–538. <https://doi.org/10.1590/S1676-06032008000100023>
- WILLIG M.R., S.J. PRESLEY, C.P. BLOCH, C.L. HICE & S.P. YANOVIK. 2007. Phyllostomid bats of lowland Amazonia: effects of habitat alteration on abundance. *Biotropica* 39: 737–746. <http://doi.org/dvpx6k>

Authors' contributions. GLU collected the data. GLU, TYSS and GG wrote the text, and GLU made the map and table.

Received: 8 February 2017

Accepted: 21 April 2017

Academic editor: Annia Rodriguez-San Pedro