



New records of spiders (Arachnida, Araneae) from the state of Roraima, northern Brazil

Leonardo S. Carvalho^{1,2,3,7}, Pedro H. Martins³, Marielle C. Schneider⁴ & Jimmy J. Cabra-Garcia^{5,6}

¹ Universidade Federal do Piauí, Campus Amílcar Ferreira Sobral, BR 343, KM 3.5, Bairro Meladão, CEP 64800-000, Floriano, PI, Brazil

² Programa de Pós-Graduação em Zoologia, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

³ Universidade Federal de Minas Gerais, Instituto de Ciências Biológicas, Departamento de Zoologia, Av. Antônio Carlos, 6627, Bairro Pampulha, Caixa Postal 486, CEP 31270-901, Belo Horizonte, MG, Brazil

⁴ Universidade Federal de São Paulo, Departamento de Ciências Biológicas, Av. Prof. Artur Riedel, 275, CEP 09972-270, Diadema, São Paulo, Brazil

⁵ Universidade de São Paulo, Instituto de Biociências, Departamento de Zoologia, São Paulo, SP, 05508-090, Brazil

⁶ Universidad del Valle, Departamento de Biología, Cali, AA 25360, Colombia

⁷ Corresponding author. E-mail: carvalho@ufpi.edu.br

Abstract: The Brazilian spider fauna comprises thousands of described species, mostly known by only one or two records, and there are large sampling gaps. The spider fauna of the state of Roraima is enigmatic in Brazil and remains largely unknown. Herein, we present a list of spider species recently collected during an expedition in Roraima. Species-level identifications were possible for 229 adult individuals of 54 species. Five species are newly recorded from Brazil, and 30 species are presented for the first time from Roraima. Most of these new records are represented by widespread species, representing the huge and historical deficiency in the spider sampling throughout Roraima.

Key words: Amazon Forest; Cerrado; geographical distribution; Neotropical Region

INTRODUCTION

Brazilian spiders are suitable for the study of diversity patterns and geographical distribution owing to the existence of a large database that includes all records in taxonomic publications, among other records (i.e., BRESCOVIT et al. 2011; OLIVEIRA et al. 2015). As of 2013, there were 3,425 spider species recorded from Brazil, but about 40% of these species are known only by single records, while about 45% have between two and 15 records, 10% have between 16 and 60 records, and only 2% of the species are represented by more than 100 records (OLIVEIRA et al. 2015). This heterogeneous knowledge is highly influenced by where the main national research institutions, which harbor the most significant collections, and researchers on spider systematics are located. Collections are mostly composed of species gathered nearby (BRESCOVIT et al. 2011), significantly associated to access routes, creating larger knowledge shortfalls in localities far from the main

research centers (OLIVEIRA et al. 2016).

For example, the Amazon has the lowest record density for invertebrates, vertebrates and angiosperms among all Brazilian biomes (OLIVEIRA et al. 2016). Regarding spiders, there is a historical concentration of sampling effort mostly in small regions (OLIVEIRA et al. 2015). The states of Pará (e.g., RICETTI & BONALDO 2008; BONALDO et al. 2009b) and Amazonas (e.g., HÖFER & BRESCOVIT 2001; ADIS et al. 2002; BONALDO et al. 2009a; DIAS & BONALDO 2012) concentrate most of the published spider inventories, collaborating for such heterogeneous knowledge. Additionally, in the last 15 years, knowledge of spiders in the states of Roraima, Rondônia, Acre, and Amapá has increased mainly through taxonomic papers (e.g., POLOTOW & BRESCOVIT 2008; ABRAHIM et al. 2012; RUIZ & BRESCOVIT 2013; RODRIGUES 2013; PAULA et al. 2014; COSTA & RUIZ 2014; BERTANI et al. 2016), but no lists of spider species have been published or updated.

For Roraima, there is a single spider species inventory carried out in *terra firme* forests in the Island of Maracá, in the municipality of Uraricoera; it is almost two decades old (LISE 1998a, 1998b). In that paper, LISE (1998a, 1998b) recorded 145 spider species (92 determined and 55 undetermined species), a species richness presently considered low, if compared to well-sampled Amazon Forest localities such as the Serra do Cachimbo (427 spp., RICETTI & BONALDO 2008), the Reserva Florestal Adolpho Ducke (506 spp., HÖFER & BRESCOVIT 2001; ADIS et al. 2002), the Floresta Nacional de Caxiuanã (591 spp., BONALDO et al. 2009b), or the Porto Urucu River basin (623 spp., DIAS & BONALDO 2012).

As a consequence of the general deficiency of sampling in Roraima, any expedition focusing on sampling arachnids in the region is expected to result in a large number of undescribed or, at least, under-recorded species (even




Figure 1. Location of the sampling points in the municipalities of Amajari (squares), Boa Vista (circles), Bonfim (triangles) and Cantá (pentagons), state of Roraima, northern Brazil.

for widespread species). Recently, we performed a short survey in four municipalities (Amajari, Boa Vista, Bonfim, and Cantá) at Roraima, and here we present a species list of these spiders and update their geographical distribution.

MATERIAL AND METHODS

The sampling was conducted in the state of Roraima in northern Brazil. The state borders Venezuela on the north and west and Guyana on the east. To the south is the state of Amazonas, and to the southeast, the state of Pará. Additionally, the region represents a large ecotone area; flanked by the Amazon Forest to the south and west, by savannas to the east, and by mountainous regions, with altitudes up to 3,000 m, to the north (EDEN & MCGREGOR 1998). The climate is characterized by three climate categories in Köppen's classification: Af (constantly humid regions with low annual variations in rainfall and temperature), Am (humid summer and short term dry winter seasons), and Aw (with four months of a real drought) (BARBOSA et al.

1997; FALCÃO & COSTA 2012). In general, lowland Roraima has annual mean temperature ranging between 26 and 27°C, which decreases considerably with elevation (EDEN & MCGREGOR 1998). The mean annual rainfall varies from 1,100–1,400 mm/year in the northeast to 2,000–2,300 mm/year in the southwest (BARBOSA et al. 1997). Such heterogeneous climate and topography allow for the existence of different habitats, such as *terra firme* forest, seasonally flooded forest (*várzea* and *igapó*), white-sand forest (*campina* and *campinarana*), savanna, gallery, and dry forests, and various types of montane forests (Naka et al. 2006).

In July 2014, a 10-day sampling campaign was carried out in Roraima (Figure 1), in the municipalities of Amajari (Figures 2–6), Boa Vista, Bonfim (Figure 7), and Cantá. Collecting permits were issued by the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) through the Sistema de Autorização e Informação em Biodiversidade (SISBIO #44225-1). Arachnids were collected through diurnal and nocturnal manual searches that did not follow any



Figures 2–7. Aspects of the vegetation in sampled localities throughout Roraima. **2 and 3.** Forested areas in the Estância Ecológica SESC Tepequém, Vila Tepequém, Amajari. **4.** Gallery forest near the Cachoeira do Paiva, Vila Tepequém, Amajari. **5 and 6.** Savannas and rocky outcrops in the uphill of the Serra do Tepequém, Vila Tepequém, Amajari. **7.** Seasonally flooded lowland savanna in the surroundings of Bonfim. Photos by L.S. Carvalho.

standardized protocol.

Specimens were sorted then identified using relevant taxonomic literature (i.e., CHICKERING 1943, 1966; LEVI, 1963, 1968, 1971, 1975, 1976, 1985, 1991a, b, 1992, 1993, 1994, 1995, 1996, 1997, 2004; EXLINE & LEVI 1965; PLATNICK 1975, 1986; SHEAR 1970; OPELL, 1979; DEE-LEMAN-REINHOLD & PRINSEN 1987; HARROD et al. 1991;

BRESCOVIT & BONALDO 1993; HÖFER et al. 1994; SAA-GER 1994; GLUECK 1994; COYLE 1995; BRESCOVIT 1996; CORRONCA 1998; BRESCOVIT & RHEIMS 2000; HÖFER & BRESCOVIT 2000; HUBER 2000, 2005; SANTOS & BRESCO-VIT 2001; DEBLEMAN-REINHOLD & VAN HARTEN 2001; AGNARSSON 2003; RHEIMS & BRESCOVIT 2004; SANTOS & RHEIMS 2005; BERTANI & ARAÚJO 2006; GUADANUCCI

et al. 2007; SANTOS 2007; POLOTOW & BRESOVIT 2008, 2009; RHEIMS et al. 2008; RUDLOFF & WEINMANN 2010; CREWS 2011; PIACENTINI 2011; SILVA & CARICO 2012; COSTA & RUIZ 2014; BERTANI et al. 2016; CABRA-GARCÍA & BRESOVIT 2016), all available on-line (WORLD SPIDER CATALOG 2017). Only specimens identified to the species level are listed here. Thus, undetermined or undescribed specimens are not discussed. Names of species follow the WORLD SPIDER CATALOG (2017).

All specimens were deposited in the following arachnid collections: Coleções Taxonômicas da Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais (UFMG; curator A. J. Santos); Coleção de História Natural da Universidade Federal do Piauí, Floriano, Piauí (CHNUFPI; curator E. F. B. Lima); and Museu de Zoologia da Universidade de São Paulo, São Paulo, São Paulo (MZSP; curator R. Pinto-da-Rocha). Photographs were made using a Leica M205C stereomicroscope equipped with a Leica DFC295 digital camera. Images in different focal planes were assembled in multi-focus images using the software Leica Application Suite ver. 3.3.0 (Leica Inc.). The map of the sampling points was constructed in ArcMap 10.0 (ESRI), using WGS84 datum (the same used to obtain the geographical coordinates during the sampling).

RESULTS

Species identification was achieved for 229 adult individuals (86 males and 143 females) belonging to 54 species and 19 families. Only six species presented more than 10 individuals: *Metagonia mariguitarensis* (González-Sponga, 1998) ($n = 24$), *Wagneriana atuna* Levi, 1991 ($n = 24$), *Trochanteria gomezi* Canals, 1933 ($n = 23$), and *Carapoia paraguaensis* González-Sponga, 1998 ($n = 19$), *Architis tenuis* Simon, 1898 ($n = 18$) and *Metazygia gregalis* (O. Pickard-Cambridge, 1889) ($n = 12$). Eighteen species were doubletons and 18 species were singletons.

The list of spiders presented herein is mainly composed by widely distributed species (50 spp.; 92% of the total), and only four species are known to have a restricted **Distribution**: the theraphosids *Guyruita waikoshiemi* (Bertani & Araújo, 2006), *Psalmopoeus irminia* Saager, 1994 and *Theraphosa stirmi* Rudloff & Weinmann, 2010, and the orb-weaver araneid *Hypognatha cacau* Levi, 1996. The theraphosid species are distributed only in the Guiana shield, while *H. cacau* is known from the northern Amazon Forest.

Four species are recorded in Brazil for the first time: the theraphosids *G. waikoshiemi* and *T. stirmi*, the selenopid *Selenops geraldinae* Corronca, 1996, and the orb-weaver araneid *Wagneriana taboga* Levi, 1991. Additionally, 30 species are recorded from Roraima for the first time. No species of medical importance was recorded. Additionally, five recorded species are widely distributed and/or can be found in synanthropic environments: *Argiope argentata* (Fabricius, 1775) (Araneidae), *Micropholcus fauroti* (Simon, 1887) (Pholcidae), *Philoponella vittata* (Keyserling, 1881)

(Uloboridae), *Oecobius concinnus* Simon, 1893 (Oecobiidae), and *Scytodes fusca* Walckenaer, 1837 (Scytodidae). Below, we present a brief discussion on each species collected.

Infraorder Mygalomorphae

Family Dipluridae Simon, 1889

Ischnothele guianensis (Walckenaer, 1837)

Mygale guianensis WALCKENAER (1837).

Pezionyx guianensis — TACZANOWSKI (1874).

Entomothele guianensis — SIMON (1889).

Thelechoris guianensis — SIMON (1891).

Ischnothele siemensis F. O. PICKARD-CAMBRIDGE (1896).

Ischnothele guianensis — F. O. PICKARD-CAMBRIDGE (1896); COYLE (1995).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l. [above sea level]), 1 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17230); (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♂, 16.VII.2014, same collectors (UFMG 17085).

Distribution: Bolivia, Brazil (Acre, Amapá, Amazonas, Ceará, Pará, Piauí and Roraima [**new record**]), Colombia, French Guiana, Guyana, Peru, and Surinam (Coyle 1995; Carvalho et al. 2014).

Family Theraphosidae Thorell, 1869

Subfamily Schismatothelinae Guadanucci, 2014

Guyruita waikoshiemi (Bertani & Araújo, 2006)

(Figures 8, 12–15)

Holothele waikoshiemi BERTANI & ARAÚJO (2006).


Guyruita waikoshiemi — GUADANUCCI et al. (2007).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♀, 16.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 16690).

Distribution: *Guyruita waikoshiemi* was previously known only from its type locality, the Yanomami community Motorema (02°30'27.0" N, 065°9'39.9" W) in the Reserva de Biosfera Alto Orinoco-Casiquiare, 411 km from Puerto Ayacucho, Estado Amazonas, Venezuela (BERTANI & ARAÚJO 2006). The new record increases the geographical distribution by at least 410 km to the northeast.

Natural history: The specimen was found on the ground inside a burrow, which was lined internally with threads (as also described by BERTANI & ARAÚJO 2006). It was captured by digging out the soil. The surrounding area consisted of secondary *terra firme* vegetation.

Note: The spider genus *Guyruita* Guadanucci was described in 2007 and comprises three described species. It is distinguished from the remaining Schismatothelinae theraphosid genera by the labium densely occupied by more than 100 cusps, absence of intercheliceral intumescence, posterior sternal sigilla remote from margin, absence of teeth in tarsal claws, undivided tarsal scopula



Figures 8–11. Representatives of the spiders collected in the State of Roraima. **8.** Adult female of *Guyruita waikoshiemi* (UFMG 16690), collected at Amajari, **9.** Adult male of *Theraphosa stirmi* (UFMG 17210), collected at Amajari. **10.** Adult female of *Psalmopoeus irminia* (UFMG 17888), collected at Amajari. **11.** Adult female of *Eriophora edax* (UFMG 17073), collected at Bonfim. Photos by L.S. Carvalho (8, 9, 11) and P.H. Martins (10).

I–II, and divided tarsal scopula III–IV (GUADANUCCI et al. 2007). The females of *G. waikoshiemi* (Figures 8 and 12–15) can be distinguished from congeners by the incrassate tibia I (GUADANUCCI et al. 2007; see Figure 14) and by the spermathecae being multilobulated at the apical portion (GUADANUCCI et al. 2007; see Figure 15). The female specimen presented the diagnostic features for the genus (Figures 12 and 13) and species (Figures 14 and 15), but its spermathecae presented sclerotized receptacles positioned more apically than the holotype specimen drawn by BERTANI & ARAÚJO (2006). However, we consider this to be intraspecific variation.

Subfamily Selenocosmiinae Simon, 1889

Psalmopoeus irminia Saager, 1994 (Figure 10)

Psalmopoeus irminia SAAGER (1994) — PETERS (2000, 2003); SCHMIDT (2003); SCHMIDT et al. (2006); MENDOZA (2014).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♂, 16.VII.2014, L.S.

Carvalho & M.C. Schneider *leg.* (UFMG 17888).


Distribution: This species is known from its type locality in northeastern Venezuela (Gran Sabana, Estado Guyana) (SAAGER 1994) and the municipality of Pacaraima (Roraima) near the Brazil–Venezuela border (BERTANI et al. 2016). The new record extends the geographical distribution of this species by about 102 km toward the southwest.

Subfamily Theraphosinae Thorell, 1869

Theraphosa stirmi Rudloff & Weinmann, 2010 (Figure 9)
Theraphosa stirmi RUDLOFF & WEINMANN (2010).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♂, 16.VII.2014, L.S. Carvalho & M.C. Schneider (UFMG 17210).

Distribution: This species was known only from its type locality in southeastern Guyana (Essequibo, Takutu River). The new record is the first from Brazil (RUDLOFF & WEINMANN 2010) and extends the species' geographical distribution by about 420 km toward the northwest.



Figures 12–15. Diagnostic features of *Guyruita waikoshiemi* (UFMG 16690), collected at Amajari, Roraima, Brazil. **12.** Sternum. **13.** Labium and endites. **14.** Tibia I. **15.** Spermathecae. Photos by P.H. Martins.

Infraorder Araneomorphae
Family Araneidae Clerck, 1757

Acacesia hamata (Hentz, 1847)

Epeira hamata HENTZ (1847).
Epeira foliata HENTZ (1847).
Epeira folifera — MARX (1890).
Acacesia foliata — SIMON (1895a).
Araneus hallucinator PETRUNKEVITCH (1911).
Acacesia lanceolata BADCOCK (1932).
Acacesia hamata — BRYANT (1945); LEVI (1976, 2002); GLUECK (1994).
Araneus nigrolineatus CAPORACCO (1955).
Acacesia nigrolineata — LEVI (1991a).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém; 03°45'54.1" N, 061°43'14" W; 665 m a.s.l.), 1 ♀, 18.VI.2014, J. Cabra et al. *leg.* (MZSP 69709); (Vila Tepequém, near Pousada PSJ; 03°46'55.4" N, 061°43'19.0" W; 587 m a.s.l.), 1 ♂, 15.vi.2014, J. Cabra et al. *leg.* (MZSP 69714).

Distribution: This species is widely distributed in the New World and is known from Argentina, Bahamas, Bolivia, Brazil (Amapá, Espírito Santo, Goiás, Mato Grosso, Minas Gerais, Pará, Paraíba, Paraná, Pernambuco, Rio de Janeiro, Rio Grande do Norte, Rio Grande do Sul, Roraima [new

record] and São Paulo), British Virgin Islands, Colombia, Costa Rica, Cuba, Ecuador, Guatemala, El Salvador, Guyana, Honduras, Jamaica, Mexico, Panama, Nicaragua, Paraguay, Peru, Trinidad, and United States (LEVI 1976; GLUECK 1994).

Aculepeira travassosi (Soares & Camargo, 1948)

Neosconella travassosi SOARES & CAMARGO (1948).
Neosconella cutucensis KRAUS (1955).
Araneus cutucensis — BRIGNOLI (1983).
Araneus travassosi BRIGNOLI (1983).
Aculepeira travassosi — LEVI (1991a; 2002); DIERKENS (2012).

Material examined: BRAZIL, RORAIMA: Bonfim (03°19'35.9" N, 059°56'41.7" W; 132 m a.s.l.), 1 ♀, 20.VI.2014, J. Cabra et al. *leg.* (MZSP 69707).

Distribution: This species, widely distributed in the New World, is known from Argentina, Bolivia, Brazil (Amazonas, Bahia, Mato Grosso, Pará, Roraima and São Paulo), Mexico, Nicaragua, Panamá, and Paraguay (LEVI 1991a).

Amazonopeira masaka Levi, 1994

Amazonopeira masaka LEVI (1994, 2002)

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ; 03°47'10.4" N, 061°43'15.3"

W; 640 m a.s.l.), 3 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17227); (Vila Tepequém, Pousada PSJ, 03°46'55.4"N, 061°43'19.0" W; 587 m a.s.l.), 1 ♀, 15.VI.2014, J. Cabra et al. *leg.* (MZSP 69708).

Distribution: Brazil (Amazonas, Roraima and Pará) and Ecuador (LEVI 1994; BONALDO et al. 2009b).

Araneus guttatus (Keyserling, 1865)

Epeira guttata KEYSERLING (1865).

Epeira similis TACZANOWSKI (1873).

Aranea glabrata F. O. PICKARD-CAMBRIDGE (1904).

Araneus glabratus — PETRUNKEVITCH (1911).

Araneus guttatus — PETRUNKEVITCH (1911).

Araneus similis — PETRUNKEVITCH (1911).

Aranea similella — ROEWER (1942).

Neosconella bipunctata MELLO-LEITÃO (1948).

Araneus leitaoi BRIGNOLI (1983).

Araneus guttatus — LEVI (1991a, 2002).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, Cachoeira do Paiva, 03°45'18.4" N, 061°42'58.9" W; 523 m a.s.l.), 1 ♀, 15.VI.2014, J. Cabra et al. *leg.* (MZSP 69712); Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♀, 23.VII.2014, L.S. Carvalho et al. *leg.* (UFMG 17072); (02°38'8.2"N, 060°34'15.6" W; 96 m a.s.l.), 1 ♀, 21.VI.2014, J. Cabra et al. *leg.* (MZSP 69713).

Distribution: This species, widely distributed in the New World, is known from Argentina, Brazil (Amazonas, Goiás, Mato Grosso, Minas Gerais, Paraná, Rio de Janeiro, Rio Grande do Sul, Rondônia, Roraima [new record], and São Paulo), Colombia, Costa Rica, Ecuador, Panama, Paraguay, Peru, and Venezuela (LEVI 1991a).

Argiope argentata (Fabricius, 1775)

Aranea argentata FABRICIUS (1775) — OLIVIER (1789).

Aranea mammeata DE GEER (1778) — OLIVIER (1789).

Epeira mammata WALCKENAER (1805).

Argyopes argentatus — C. L. KOCH (1838).

Argyopes fenestrinus C. L. KOCH (1838).

Epeira argentata — WALCKENAER (1841).

Epeira amictoria — WALCKENAER (1841).

Plectana sloanii WALCKENAER (1841).

Epeira gracilis KEYSERLING (1865).

Argiope carinata L. KOCH (1871).

Argyopes maronicus TACZANOWSKI (1873).

Argyopes subtilis TACZANOWSKI (1873).

Acrosoma sloanii — BUTLER (1873).

Argyopes hirtus TACZANOWSKI (1879).

Epeira gracilis KEYSERLING (1893).

Argiope argentata — MCCOOK (1894).

Argiope waughii SIMON (1896).

Araneus gracilis — PETRUNKEVITCH (1911).

Micrathena sloanei — PETRUNKEVITCH (1911).

Gea panamensis CHAMBERLIN (1917).

Argiope argyrea BADCOCK (1932).

Argiope cuyunii HINGSTON (1932).

Argiope filiargentina HINGSTON (1932).

Argiope filinfracta HINGSTON (1932).

Singa gracilis — MELLO-LEITÃO (1941).

Aranea gracilentata ROEWER (1942).

Argiope indistincta MELLO-LEITÃO (1944).

Argiope hirta ARCHER (1963).

Argiope argentata — LEVI (1968, 1983, 1991a, 1993, 2004); AGNARSSON et al. (2016).

Material examined: BRAZIL, RORAIMA, Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 2 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17198).

Distribution: This species is one of the most widespread and widely distributed spider species in the New World (LEVI 2004: map 2A) and is sometimes associated with synanthropic environments (LEVI 1968; TAUCARE-RÍOS 2012).

Eriophora edax (Blackwall, 1863) (Figure 11)

Epeira edax BLACKWALL (1863).

Eriophora edax — BANKS (1909); PETRUNKEVITCH (1930); BRYANT (1948); LEVI (1971, 1991, 2002).

Araneus edax — PETRUNKEVITCH (1911).

Araneus argyronotus — MELLO-LEITÃO (1939c).

Araniella geayi CAPORACCO (1954).

Araneus geayi — LEVI (1974).

Material examined: BRAZIL, RORAIMA: Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♀, 23.VII.2014, L.S. Carvalho et al. *leg.* (UFMG 17073).

Distribution: This species is widely distributed in the New World (LEVI 1971: map 3), with published records from United States to Brazil (LEVI 1971; BONALDO et al. 2009b; WORLD SPIDER CATALOG, 2017). This is the first record of *E. edax* from Roraima and fills a gap in its known geographical distribution.

Hypognatha cacau Levi, 1996

Hypognatha cacau LEVI (1996).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, Cachoeira do Paiva, 03°45'18.4" N, 061°42'58.9" W; 523 m a.s.l.), 1 ♀, 17.VI.2014, J. Cabra et al. *leg.* (MZSP).

Distribution: This species is known only from Peru (San Martín) and Brazil (Rondônia) (LEVI 1996), and the new record from Roraima extends its distribution by at least 1,500 km toward the northeast.

Larinia directa (Hentz, 1847)

Epeira directa HENTZ (1847).

Epeira rubella HENTZ (1847).

Epeira tetragnathoides O. PICKARD-CAMBRIDGE (1889).

Epeira intercisa O. PICKARD-CAMBRIDGE (1889).

Drexelia directa — MCCOOK (1892).

Epeira deludens KEYSERLING (1893).

Larinia directa — BANKS (1894); LEVI (1975, 2002); HARROD et al. (1991); DIERKENS (2012).

Larinia bellona BANKS (1898).

Drexelia bellona — F. O. PICKARD-CAMBRIDGE (1903).

Larinia cymotypha CHAMBERLIN (1924).

Larinia albonigra FRANGANILLO (1931).

Metazygia albonigra — BRYANT (1940).

Larinia nigrovittata MELLO-LEITÃO (1947).

Drexelia octopunctata CAPORACCO (1955).

Material examined: BRAZIL, RORAIMA: Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♀, 23.VII.2014, L.S. Carvalho et al. *leg.* (UFMG 17076); 1 ♀, 24.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1604).

Distribution: This species is widely distributed in the New World (HARROD et al. 1991: map 1). There are published records from the following countries: Bahamas, Brazil (Amapá, Amazonas, Minas Gerais, Roraima, and São Paulo), Colombia, Costa Rica, Cuba, Ecuador, Guatemala, Honduras, Jamaica, Mexico, Nicaragua, Panama, Paraguay, Surinam, United States (Alabama, Arizona, Arkansas, California, Georgia, Louisiana, Mississippi, North Carolina, Tennessee, Texas, Virginia, and South Carolina), and Venezuela (LEVI 1975; HARROD et al. 1991; DIERKENS 2012).

Manoega porracea (C. L. Koch, 1838)

Miranda porracea C. L. KOCH (1838).

Epeira porracea — WALCKENAER (1841).

Zilla guyanensis KEYSERLING (1881b, 1893).

Cyrtophora porracea — SIMON (1895a).

Cyrtophora grammica SIMON (1895b).

Araneus poraceus — PETRUNKEVITCH (1911).

Araneus guyanensis PETRUNKEVITCH (1911).

Zygiella guyanensis — ROEWER (1942).

Mangora octolineata CAPORACCO (1947).

Mecynogea guianensis — MELLO-LEITÃO (1948); LEVI (1980).

Meta brasílica SOARES & CAMARGO (1948).

Meta berlandi CAPORACCO (1954).

Meta espiritosantensis SOARES & CAMARGO (1955).

Cyrtophora guyanensis — LEVI (1986).

Manoega porracea — LEVI (1997, 2002).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♂, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17228).

Distribution: This species is widely distributed over the South America (LEVI 1997: map 3). There are published records from the following countries: Argentina, Brazil (Amapá, Amazonas, Espírito Santo, Mato Grosso, Pará, Paraíba, Paraná, Pernambuco, Rio Grande do Sul, Roraima, São Paulo), Colombia, Ecuador, French Guiana, Guyana, Panama, Paraguay, Peru, and Surinam (LEVI 1997).

Metazygia gregalis (O. Pickard-Cambridge, 1889)

Epeira gregalis O. PICKARD-CAMBRIDGE (1889).

Metazygia gregalis — F. O. PICKARD-CAMBRIDGE (1904); LEVI (1995); DIERKENS (2012).

Aranea gregalis — Strand (1907a).

Araneus gregalis — Petrunkevitch (1911).

Eustala tuceps CHAMBERLIN (1925).

Metazygia manni BRYANT (1945).

Metazygia similis CAPORACCO (1947).

Material examined: BRAZIL, RORAIMA: Bonfim (03°21'56.3" N, 059°50'23.7" W; 132 m a.s.l.), 1 ♀, 19.VI.2014, J. Cabra et al. *leg.* (MZSP 69703); Boa Vista (Campus de Cauamé, Universidade Federal de Roraima, 02°52'38.4" N, 060°43'13.1" W; 91 m a.s.l.), 1 ♀, 22.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17074); (Campus de Cauamé, Universidade Federal de Roraima, 02°52'36.9" N,

060°43'9.3" W; 82 m a.s.l.), 6 ♀, 4m, 22.VI.2014, J. Cabra et al. *leg.* (MZSP 69701).

Distribution: This species is widely distributed over the Central and South Americas (LEVI 1995: map 3e). There are published records from the following countries: Argentina, Bolivia, Brasil (Acre, Amazonas, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Paraná, Rio de Janeiro, Rio Grande do Sul, Rondônia, Santa Catarina, and São Paulo), Costa Rica, Cuba, Dominican Republic, Ecuador, French Guiana, Guyana, Haiti, Nicaragua, Panama, Paraguay, Peru, Puerto Rico, Tobago, Uruguay, and Venezuela (LEVI 1995; DIERKENS 2012).

Metazygia lopez Levi, 1995

Metazygia lopez LEVI (1995).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, 03°45'54.1" N, 061°43'14.2" W; 665 m a.s.l.), 1 ♂, 18.VI.2014, J. Cabra et al. *leg.* (MZSP 69715).

Distribution: This species is known to Brazil (Amazonas, and Roraima [new record]), Colombia, Peru, and Venezuela (LEVI 1995).

Micrathena acuta (Walckenaer, 1841)

Plectana acuta WALCKENAER (1841).

Acrosoma acutum — KEYSERLING (1864).

Micrathena acuta — SIMON (1895a); LEVI (1985).

Micrathena alpha CAPORACCO (1947).

Ildibaha inermis SCHENKEL (1953).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♀, 16.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1603).

Distribution: This species is widely distributed over South America (LEVI 1985: map 7). There are published records from the following countries: Argentina, Brazil (Amapá, Amazonas, Bahia, Goiás, Mato Grosso, Pará, Rio de Janeiro, Rondônia, and Roraima [new record]), Colombia, Ecuador, Guyana, Peru, Trinidad, and Venezuela (LEVI 1985; BONALDO et al. 2009b).

Micrathena aureola (C. L. Koch, 1836)

Acrosoma aureolum C. L. KOCH (1836).

Acrosoma affine C. L. KOCH (1839).

Plectana affinis — WALCKENAER (1841).

Plectana aureola — WALCKENAER (1841).

Chaetacis affinis — SIMON (1895a).

Micrathena aureola — PETRUNKEVITCH (1911); MAGALHÃES & SANTOS (2012).

Chaetacis aureola — REIMOSER (1917); LEVI (1985); DIERKENS (2011).

Chaetacis hirsuta MELLO-LEITÃO (1932).

Chaetacis aculeata CHICKERING (1960).

Chaetacis dentata CHICKERING (1960).

Material examined: BRAZIL, RORAIMA: Cantá (near the road BR432, about 10 km from Cantá, 02°38'9.3" N, 060°34'17.1" W; 106 m a.s.l.), 1 ♀, 21.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17292); 1 ♀ (CHNUFPI 1322).

Distribution: This species is widely distributed over the South America (LEVI 1985: map 15), with records from Bra-

zil (Amazonas, Bahia, Mato Grosso, Minas Gerais, Pará, Rio de Janeiro, Rondônia, and Roraima [**new record**]), French Guiana, Paraguay, and Surinam (LEVI 1985; BONALDO et al. 2009b).

Micrathena coca Levi, 1985

Micrathena coca LEVI (1985).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♂, 16.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1606).

Distribution: This species is distributed over the northern South America, in the Amazon Basin (LEVI 1995: map 11). There are published records from the following countries: Brazil (Amazonas, and Roraima [**new record**]), Colombia, Ecuador, Peru, and Venezuela (LEVI 1995).

Micrathena evansi Chickering, 1960

Micrathena evansi CHICKERING (1960). — LEVI (1985); DIERKENS (2011); MAGALHÃES & SANTOS (2012).

Micrathena insolita CHICKERING (1961).

Micrathena lepida CHICKERING (1964).

Micrathena levii CHICKERING (1964).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, Cachoeira do Paiva, 03°45'45.6" N, 061°45'18" W; 582 m a.s.l.), 1 ♂, 1 ♀, 17.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17290); same data, 1 ♂, 1 ♀, (CHNUFPI 1318); Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♀, 23.VII.2014, L.S. Carvalho et al. *leg.* (UFMG 17293).

Distribution: This species is widely distributed over the South America (LEVI 1995: map 7). There are records to the following countries: Brazil (Amazonas, Bahia, Ceará, Espírito Santo, Mato Grosso, Pará, Pernambuco, and Roraima [**new record**]), Bolivia, Colombia, French Guiana, Guyana, Peru, Surinam, Trinidad, and Venezuela (LEVI 1995; BONALDO et al. 2009b).

Micrathena picta (C. L. Koch, 1836)

Acrosoma pictum C. L. KOCH (1836).

Plectana picta WALCKENAER (1841).

Micrathena picta — PETRUNKEVITCH (1911); MAGALHÃES & SANTOS (2012).

Chaetacis picta — REIMOSER (1917); LEVI (1985, 2002).

Micrathena conspicuum MELLO-LEITÃO (1929a).

Chaetacis evansi CHICKERING (1960)

Chaetacis rugosa CHICKERING (1960).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17288); Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♀, 23.VII.2014, L.S. Carvalho et al. (UFMG 17294); 1 ♀, 24.VII.2014 (UFMG 17296) 1 ♀, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1317); same data, 1 ♀ (CHNUFPI 1605).

Distribution: This species is widely distributed over the South America (LEVI 1995: map 15). There are published records from the following countries: Brazil (Amapá, Maranhão, Mato Grosso, Pará, and Roraima [**new record**]), Bolivia, Guyana, Paraguay, and Peru (LEVI 1995).

Micrathena plana (C. L. Koch, 1836)

Acrosoma planum C. L. KOCH (1836).

Plectana degeerii WALCKENAER (1841).

Plectana plana — WALCKENAER (1841).

Plectana alata WALCKENAER (1841).

Acrosoma maronica TACZANOWSKI (1873).

Acrosoma alatum — BUTLER (1873).

Acrosoma degeeri BUTLER (1873).

Micrathena plana — SIMON (1895); REIMOSER (1917); LEVI (1985);

GONZAGA & SANTOS (2004); MAGALHÃES & SANTOS (2011, 2012).

Micrathena alata — SIMON (1895).

Micrathena maronica — SIMON (1895).

Micrathena degeeri — PETRUNKEVITCH (1911).

Micrathena ornata MELLO-LEITÃO (1932) — LEVI (1985).

Micrathena nitida CHICKERING (1964).

Material examined: BRAZIL, RORAIMA: Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♀, 24.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17295); 1 ♀ (CHNUFPI 1320).

Distribution: This species is widely distributed over the South America (LEVI 1995: map 6). There are published records from the following countries: Argentina, Bolivia, Bolivia, Brazil (Amazonas, Bahia, Espírito Santo, Goiás, Mato Grosso, Minas Gerais, Pará, Rio de Janeiro, Rio Grande do Sul, Rondônia, Roraima [**new record**]), and São Paulo), Colombia, Ecuador, French Guiana, Guyana, Panama, Paraguay, Peru, Surinam, Trinidad, Venezuela, and Virgin Islands (LEVI 1995; CARVALHO et al. 2014).

Micrathena pungens (Walckenaer, 1841)

Plectana pungens WALCKENAER (1841).

Acrosoma pungens — KEYSERLING (1864, 1892).

Acrosoma luctuosa TACZANOWSKI (1873).

Micrathena pungens SIMON (1895a).

Micrathena luctuosa — PETRUNKEVITCH (1911).

Micrathenapungens — REIMOSER (1917); CHICKERING (1960); LEVI (1985).

Micrathena carvalhoi MELLO-LEITÃO (1944).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♂, 16.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17289); 1 ♂ (CHNUFPI 1319).

Distribution: This species is widely distributed over northern South America (LEVI 1995: map 14). There are published records from the following countries: Bolivia, Brazil (Amazonas, Maranhão, Mato Grosso, Pará, Rondônia, and Roraima [**new record**]), Colombia, Ecuador, Guyana, Peru, Surinam, and Venezuela (LEVI 1995; Bonaldo et al. 2009b).

Micrathena schreibersi (Perty, 1833)

Acrosoma schreibersii PERTY (1833).

Acrosoma spinosum C. L. KOCH (1836).

Plectana macrocantha WALCKENAER (1841).
Plectana duplicata WALCKENAER (1841).
Plectana vespoides WALCKENAER (1841).
Acrosoma macrocantha TACZANOWSKI (1873).
Acrosoma tenuis TACZANOWSKI (1873).
Acrosoma subtilis TACZANOWSKI (1873).
Acrosoma myrmeciaeformis TACZANOWSKI (1873).
Acrosoma macracanthum — BUTLER (1873).
Acrosoma vespoides — BUTLER (1873).
Acrosoma duplicatum BUTLER (1873).
Micrathena schreibersi — SIMON (1895a); CHICKERING (1961); LEVI (1985); DIERKENS (2011); MAGALHÃES & SANTOS (2012).
Micrathena tenuis — SIMON (1895a).
Micrathena subtilis — SIMON (1895a).
Ildibaha myrmeciaeformis — SIMON (1895a).
Micrathena duplicata — PETRUNKEVITCH (1911).
Micrathena vespoides — PETRUNKEVITCH (1911).
Micrathena coleophora CHAMBERLIN & IVIE (1936).
Micrathena lesserti MELLO-LEITÃO (1939b).
Ildibaha subtilis — MELLO-LEITÃO (1949).

Material examined: BRAZIL, RORAIMA: Amajari (Cachoeira do Paiva, Vila Tepequém, 03°45'45.6" N, 061°45'18" W; 582 m a.s.l.), 1 ♀, 17.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17291); 1 ♀ (CHNUFPI 1321).

Distribution: This species is widely distributed over the South America (LEVI 1995: map 9). There are published records from the following countries: Brazil (Amapá, Amazonas, Bahia, Espírito Santo, Minas Gerais, Pará, Pernambuco, Rondônia, Roraima, and Santa Catarina), Colombia, Costa Rica, Ecuador, French Guiana, Guyana, Nicaragua, Panama, Trinidad, and Venezuela (LEVI 1995; BONALDO et al. 2009b; CARVALHO et al. 2014).

Neoscona moreli (Vinson, 1863)

Epeira morelii VINSON (1863).
Epeira lanuginosa LENZ (1886).
Epeira moreli — SIMON (1886).
Epeira theisii KEYSERLING (1893) — MCCOOK (1894).
Araneus neotheis PETRUNKEVITCH (1911).
Cubanella nidicola FRANGANILLO (1926a, b).
Cubanella recta FRANGANILLO (1930).
Neoscona neotheis — PETRUNKEVITCH (1930); GERTSCH & MULAİK (1936); BERMAN & LEVI (1971).
Neoscona nidicola — FRANGANILLO (1936); BRYANT (1940).
Neoscona recta — FRANGANILLO (1936).
Neoscona moreli — GRASSHOFF (1980, 1986); LEVI (1993); SARISTO (2010).
Neoscona seca ROBERTS (1983).
Neoscona neotheis — BARRION et al. (1988).

Material examined: BRAZIL, RORAIMA: Bonfim (03°19'35.9" N, 059°56'41.7" W; 132 m a.s.l.), 1 ♀, 20.VI.2014, J. Cabra et al. (MZSP 69704); Boa Vista (Campus de Cauamé, Universidade Federal de Roraima, 02°52'36.9" N, 060°43'9.3" W; 82 m a.s.l.), 1 ♂, 22.VI.2014, J. Cabra et al. *leg.* (MZSP 69705).

Distribution: This species is widespread in Subsaharan Africa, Madagascar, Réunion, Mauritius and Seychelles, West Indies, and South America (LEVI 1993: map 1). In Brazil, the species is recorded from the states of Amapá, Bahia, Mato Grosso, Minas Gerais, Rio de Janeiro, Rio Grande do Sul, Roraima (**new record**), and São Paulo (LEVI 1993).

Parawixia audax (Blackwall, 1863)

Epeira audax BLACKWALL (1863) — KEYSERLING (1892).
Epeira meridionalis KEYSERLING (1865).
Epeira amaurophila HOLMBERG (1876).
Epeira coronigera TACZANOWSKI (1878).
Epeira duodecimtuberculata BERTKAU (1880).
Araneus audax SIMON (1897); FRANGANILLO (1936).
Araneus coronigerus — PETRUNKEVITCH (1911).
Aranea audax — STRAND (1916b).
Aranea eumeniphila STRAND (1916b).
Araneus rugosus BADCOCK (1932).
Verrucosa audax — MELLO-LEITÃO (1933).
Parawixia audax — MELLO-LEITÃO (1940a); LEVI (1992).
Parawixia rugosa — LEVI (1991a).
Parawixia coronigera — LEVI (1991a).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1308); 1 ♀, 17.VII.2014 (UFMG 17075); Vila Tepequém (03°45'54.1" N, 061°43'14.2" W; 665 m.a.s.l.), 1 ♂, 1 ♀, 18.VI.2014, J. Cabra et al. *leg.* (MZSP 69716).

Distribution: This species is widely distributed over the South America (LEVI 1992: map 3). There are published records from the following countries: Argentina, Bolivia, Brasil (Acre, Bahia, Espírito Santo, Goiás, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Pará, Paraná, Rio de Janeiro, Rio Grande do Sul, Roraima [**new record**], Santa Catarina, and São Paulo), Colombia, Ecuador, Guyana, Peru, and Uruguay (LEVI 1992).

Parawixia velutina (Taczanowski, 1878)

Epeira velutina TACZANOWSKI (1878).
Araneus velutinus — PETRUNKEVITCH (1911).
Araneus eriophoroides CAPORACCO (1954).
Parawixia eriophoroides — LEVI (1991a).
Parawixia velutina — LEVI (1991a, 1992).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, 03°45'54.1" N, 061°43'14.2" W; 665 m a.s.l.), 1 ♂, 1 ♀, 18.vi.2014, J. Cabra et al. *leg.* (MZSP).

Distribution: This species is widely distributed over the South America (LEVI 1992: map 3). There are published records from the following countries: Argentina, Bolivia, Brazil (Goiás, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Paraná, Rio Grande do Sul, Roraima [**new record**], and São Paulo), Colombia, Guyana, Paraguay, and Peru (LEVI 1992).

Wagneriana atuna Levi, 1991

Wagneriana atuna LEVI (1991b).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, Pousada PSJ, 03°46'55.4" N, 061°43'19.0" W; 587 m a.s.l.), 1 ♂ (MZSP 69720), 3 ♀ (MZSP 69719), 15.VI.2014, J. Cabra et al. *leg.*; 1 ♂ (MZSP 69718), 3 ♀ (MZSP 69717), 17.VI.2014; (Vila Tepequém, 03°45'54.1" N, 061°43'14.2" W; 665 m a.s.l.), 1 ♂, 1 ♀, 18.VI.2014, J. Cabra et al. *leg.* (MZSP 69721); 8 ♂, 6 ♀, (MZSP 69722); Bonfim (03°21'56.3" N, 059°50'23.7" W; 132 m a.s.l.), 1 ♂, 19.VI.2014, J. Cabra et al. *leg.* (MZSP 69724); (03°19'35.9"

N, 059°56'41.7" W; 132 m a.s.l.), 1 ♂, 1 ♀, 20.VI.2014, J. Cabra et al. *leg.* (MZSP 69723); Cantá (02°38'8.2" N, 060°34'15.6" W; 96 m a.s.l.), 1 ♀, 21.VI.2014, J. Cabra et al. *leg.* (MZSP 69727).

Distribution: This species is widely distributed in the Neotropical Region (LEVI 1991b: map 3), with known records in Brazil (Mato Grosso, Pará, Rio de Janeiro, Rio Grande do Sul, Roraima [**new record**]), Colombia, Costa Rica, Guyana, Paraguay and Peru (LEVI 1991b).

Wagneriana jelskii (Taczanowski, 1873)

Epeira jelskii TACZANOWSKI (1873).

Wagneriana jelskii — CAPORACCO (1954); LEVI (1991b).

Material examined: BRAZIL, RORAIMA: Cantá (02°38'8.2" N, 060°34'15.6" W; 96 m a.s.l.), 1 ♂, 21.VI.2014, J. Cabra, et al. *leg.* (MZSP 69728).

Distribution: This species is widely distributed in the Amazon basin (LEVI 1991b: map 3) and known from Bolivia, Brazil (Amapá, Amazonas, and Roraima), Ecuador, Guyana, Peru, Surinam, Trinidad, and Venezuela (LEVI 1991b).

Wagneriana taboga Levi, 1991

Wagneriana taboga LEVI (1991b).

Material examined: BRAZIL, RORAIMA: Cantá (Serra Grande, 02°35'16.6" N, 060°38'23.8" W; 111 m a.s.l.), 2m, 23.VI.2014, J. Cabra et al. *leg.* (MZSP 69725); (02°38'8.2" N, 060°34'15.6" W; 96 m a.s.l.), 1 ♀, 21.VI.2014, J. Cabra et al. *leg.* (MZSP 69726).

Distribution: This species was recorded only from Panama, Venezuela, and Colombia (LEVI 1991b: map 3). The new Roraima record is the first from Brazil, which extends the species' distribution by at least 900 km to the southeast.

Family Ctenidae Keyserling, 1877

Ancylometes bogotensis (Keyserling, 1877)

Ctenus bogotensis KEYSERLING (1877).

Lycotenus bogotensis F. O. PICKARD-CAMBRIDGE (1897, 1901).

Lycotenus colombianus F. O. PICKARD-CAMBRIDGE (1897).

Ancylometes palustris SIMON (1898c).

Ancylometes orinocensis SIMON (1898b).

Lycotenus palustris F. O. PICKARD-CAMBRIDGE (1899a).

Ancylometes bogotensis — STRAND (1907b); LUCAS (1964); SCHIAPELLI & GERSCHMAN (1970, 1973); MERRETT (1988); HÖFER & BRESCOVIT (2000).

Lycotenus caracasensis STRAND (1909a).

Lycotenus venezuelensis STRAND (1909b).

Ancylometes acostae SCHENKEL (1953).

Ctenus nasutus KRAUS (1955).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♂, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17081); Bonfim, (Near the road RR401, about 26km from Bonfim, 03°16'20.5" N, 060°3'9.3" W; 140 m a.s.l.), 1 ♀, 20.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17080).

Distribution: This species is widely distributed over the Neotropical Region (HÖFER & BRESCOVIT 2000: fig. 62).

There are published records from the following countries: Bolivia, Brazil (Amazonas, Pernambuco, and Roraima), Colombia, Costa Rica, Ecuador, Guyana, Honduras, Nicaragua, Panama, Trinidad, and Venezuela (HÖFER & BRESCOVIT 2000).

Ancylometes rufus (Walckenaer, 1837)

Ctenus rufus WALCKENAER (1837).

Ctenus fuscus WALCKENAER (1837).

Ctenus giganteus TACZANOWSKI (1874).

Leptoctenus tenkatei HASSELT (1888).

Lycotenus brunneus F. O. PICKARD-CAMBRIDGE (1897).

Lycotenus gigas F. O. PICKARD-CAMBRIDGE (1897).

Lycotenus demerarensis F. O. PICKARD-CAMBRIDGE (1897).

Ancylometes gigas — SIMON (1898c).

Lycotenus saraensis STRAND (1909b).

Ancylometes vulpes Petrunkevitch (1910).

Ctenus tenkatei — PETRUNKEVITCH (1911).

Lycotenus paraensis STRAND (1916a).

Ancylometes pindarensis MELLO-LEITÃO (1921, 1924).

Ctenus juruensis MELLO-LEITÃO (1922a).

Ctenus striolatus MELLO-LEITÃO (1922a).

Ctenus xerophilus MELLO-LEITÃO (1936).

Lycotenus titanus CAPORACCO (1947, 1948).

Ancylometes rufus HÖFER & BRESCOVIT (2000).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♀, 17.VII.2014, L.S. Carvalho et al. *leg.* (UFMG 17201).

Distribution: This species is widely distributed over the South America (HÖFER & BRESCOVIT 2000: fig. 61). There are published records from the following countries: Bolivia, Brazil (Amazonas, Ceará, Goiás, Pará, Paraná, Pernambuco, Piauí, Rio de Janeiro, Rio Grande do Sul, Rondônia, Roraima, and São Paulo), Ecuador, French Guiana, Guyana, Peru, Surinam, and Venezuela (HÖFER & BRESCOVIT 2000; BONALDO et al. 2009b; CARVALHO et al. 2014).

Centroctenus auberti (Caporiacco, 1954)

Ctenus auberti CAPORACCO (1954).

Acanthoctenus penicillatus CAPORACCO (1955).

Ctenus tapereba HÖFER et al. (1994).

Centroctenus auberti BRESCOVIT (1996).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17200).

Distribution: This species is distributed over the northern Amazon Basin, with published records in Brazil (Amapá, Amazonas, Pará, and Roraima [**new record**]), French Guiana, and Venezuela (HÖFER et al. 1994; BRESCOVIT 1996).

Parabatinga brevipes (Keyserling, 1891)

Ctenus brevipes KEYSERLING (1891); MELLO-LEITÃO (1936).

Ctenus taeniatus KEYSERLING (1891); MELLO-LEITÃO (1941); EICKSTEDT (1978).

Ctenus thomasi F. O. PICKARD-CAMBRIDGE (1902b).

Ctenus tatarendensis TULLGREN (1905).

Ctenus brevibris — STRAND (1909a).

Ctenus anisitsi STRAND (1909c).

Ctenus atrivulva STRAND (1909c).
Ctenus mentor STRAND (1909c).
Ctenus binotatus MELLO-LEITÃO (1936).
Ctenus gynheraldicus MELLO-LEITÃO (1936).
Isoctenus masculus MELLO-LEITÃO (1939a).
Ctenus albovittatus MELLO-LEITÃO (1939b).
Oligoctenus taeniatus — LEHTINEN (1967).
Parabatinga brevipes — POLOTOW & BRESCOVIT (2009, 2014).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♂, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17082); 1 ♀ (UFMG 17229).

Distribution: This species is widely distributed over the South America (POLOTOW & BRESCOVIT 2009). There are published records from the following countries: Argentina, Bolivia, Brazil (Bahia, Bahia, Ceará, Distrito Federal, Espírito Santo, Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Pará, Paraná, Piauí, Rio de Janeiro, Rio Grande do Sul, Roraima [**new record**], Santa Catarina, São Paulo, São Paulo, Sergipe, and Tocantins), Colombia, Paraguay, and Uruguay (POLOTOW & BRESCOVIT 2009; CARVALHO et al. 2014).

Family Hersiliidae Thorell, 1870

Neotama mexicana (O. Pickard-Cambridge, 1893)

Hersilia mexicana O. PICKARD-CAMBRIDGE (1893).
Tama mexicana — BANKS (1898); F. O. PICKARD-CAMBRIDGE (1902a); WUNDERLICH (1988).
Tama guyanensis MELLO-LEITÃO (1948).
Neotama mexicana — RHEIMS & BRESCOVIT (2004).

Material examined: BRAZIL, RORAIMA: Bonfim (near the road RR401, about 26 km from Bonfim, 03°16'20.5" N, 060°3'9.3" W; 140 m a.s.l.), 1 ♂, 20.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1309).

Distribution: This species is widely distributed for the Central America and northern South America (RHEIMS & BRESCOVIT 2004: fig. 126). There are published records from the following countries: Brazil (Acre, Amazonas, Pará, Roraima [**new record**], Rondônia), Colombia, Costa Rica, Ecuador, El Salvador, Mexico, Panama, Peru, and United States (Texas) (RHEIMS & BRESCOVIT 2004; BONALDO et al. 2009b; REGO et al. 2009).

Family Lycosidae Sundevall, 1833

Aglaoctenus lagotis (Holmberg, 1876)

Ocyale lagotis HOLMBERG (1876).
Diapontia freiburgensis KEYSERLING (1877).
Diapontia granadensis KEYSERLING (1877).
Podophthalma diversa O. PICKARD-CAMBRIDGE (1877).
Tetragonophthalma granadensis — KEYSERLING (1891).
Tetragonophthalma obscura KEYSERLING (1891).
Porrina diversa — SIMON (1898c).
Aglaoctenus bifasciatus TULLGREN (1905).
Tetragonophthalma freiburgensis — PETRUNKEVITCH (1911).
Porrina harknessi CHAMBERLIN (1916); ROEWER (1960); BRADY (1962); CAPOCASALE (1982, 1991).
Architis paulistana MELLO-LEITÃO (1917).

Porrina glieschi MELLO-LEITÃO (1926); ROEWER (1960); CAPOCASALE (1982).

Porrina callipoda MELLO-LEITÃO (1934).
Porrina granadensis — MELLO-LEITÃO (1941b).
Porrina lagotis — MELLO-LEITÃO (1941c), ROEWER (1960); CAPOCASALE (1982, 1991).
Porrina freiburgensis — MELLO-LEITÃO (1943b).
Porrina obscura — MELLO-LEITÃO (1943b).
Porrinosa diversa — Capocasale (1982).
Porrinosa callipoda CAPOCASALE (1982).
Aglaoctenus lagotis — SANTOS & BRESCOVIT (2001); SANTOS (2007); PIACENTINI (2011).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♀, 16.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17086).

Distribution: This species is widely distributed over the South America (SANTOS & BRESCOVIT 2001: fig. 8; Piacentini 2011: fig. 21). There are published records from the following countries: Bolivia, Brazil (Bahia, Ceará, Distrito Federal, Goiás, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Paraíba, Paraná, Piauí, Rio de Janeiro, Rio Grande do Norte, Rio Grande do Sul, Roraima [**new record**], Santa Catarina, São Paulo, and Tocantins), Colombia, Peru, Uruguay, and Venezuela (SANTOS & BRESCOVIT 2001; PIACENTINI 2011; CARVALHO et al. 2014).

Family Oecobiidae Blackwall, 1862

Oecobius concinnus Simon, 1893

Oecobius concinnus SIMON (1892, 1893a) — SHEAR (1970); SANTOS & GONZAGA (2003); ONO (2011); JÄGER & PRAXAYSOMBATH (2011).
Oecobius nieborowskii KULCZYŃSKI (1909).
Oecobius benneri PETRUNKEVITCH (1929).
Thalamia nieborowskii — BANKS (1930).
Oecobius vokesi GERTSCH & DAVIS (1942).
Oecobius audanti BRYANT (1948).
Tarapaca nieborowskii — LEHTINEN (1967).
Tarapaca concinnus — SAARISTO (2010).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♂, 1 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17308).

Distribution: This is pantropical synanthropical species (SHEAR 1970; SANTOS & GONZAGA 2003). Nevertheless, this is the first record of this species from Roraima (SHEAR 1970; SANTOS & GONZAGA 2003; ONO 2011; JÄGER & PRAXAYSOMBATH 2011; SANTOS et al. 2009).

Family Oxyopidae Thorell, 1870

Oxyopes incertus Mello-Leitão, 1929

Oxyopes incertus MELLO-LEITÃO (1929b).

Material examined: BRAZIL, RORAIMA: Cantá (near the road BR432, about 10 km from Cantá, 02°38'9.3" N, 060°34'17.1" W; 106 m a.s.l.), 2 ♀, 21.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17102).

Distribution: This species is recorded only from Brazil (Amazonas, Goiás, Pará, and Roraima [**new record**]) and Peru (MELLO-LEITÃO 1942b; BONALDO et al. 2009b).

Peucetia flava Keyserling, 1877

Peucetia flava KEYSERLING (1877) — MELLO-LEITÃO (1929b); SANTOS & BRESOVIT (2003).

Peucetia tranquilini MELLO-LEITÃO (1922b, 1929b).

Peucetia villosa MELLO-LEITÃO (1929b).

Peucetia meridionalis MELLO-LEITÃO (1929b).

Peucetia rubrigastra MELLO-LEITÃO (1929b, 1943b).

Peucetia viridisternis MELLO-LEITÃO (1945).

Material examined: BRAZIL, RORAIMA: Amajari (Cachoeira do Paiva, Vila Tepequém, 03°45'45.6" N, 061°45'18.0" W; 582 m a.s.l.), 2 ♂, 3 ♀, 17.VII.2014, L.S. Carvalho & M.C. Schneider leg. (UFMG 17313).

Distribution: This species is widely distributed over the South America (SANTOS & BRESOVIT 2003: fig. 45). There are published records from the following countries: Argentina, Bolivia, Brazil (Bahia, Ceará, Distrito Federal, Espírito Santo, Goiás, Mato Grosso do Sul, Minas Gerais, Pará, Paraíba, Paraná, Pernambuco, Piauí, Rio de Janeiro, Rio Grande do Norte, Rio Grande do Sul, Roraima [**new record**], Santa Catarina, São Paulo, and Sergipe), Guyana, Surinam, and Venezuela. (SANTOS & BRESOVIT 2003; CARVALHO et al. 2014).

Peucetia rubrolineata Keyserling, 1877

Peucetia rubrolineata KEYSERLING (1877) — SANTOS & BRESOVIT (2003).

Peucetia similis KEYSERLING (1877); MELLO-LEITÃO (1929b).

Peucetia heterochroma MELLO-LEITÃO (1929a).

Peucetia amazonica MELLO-LEITÃO (1929b).

Peucetia maculipedes PIZA (1938).

Peucetia trivittata MELLO-LEITÃO (1940b).

Peucetia duplovittata MELLO-LEITÃO (1941a).

Tapinillus argentinus MELLO-LEITÃO (1941c).

Peucetia roseonigra MELLO-LEITÃO (1943a).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 8 ♀, 16.VII.2014, L.S. Carvalho & M.C. Schneider leg. (UFMG 17103).

Distribution: This species is widely distributed over the Neotropical Region (SANTOS & BRESOVIT 2003: fig. 46). There are published records from the following countries: Argentina, Brazil (Acre, Amazonas, Bahia, Ceará, Espírito Santo, Mato Grosso, Minas Gerais, Paraíba, Paraná, Pernambuco, Piauí, Rio de Janeiro, Rio Grande do Norte, Roraima [**new record**], São Paulo, and Sergipe), Chile, Colombia, Ecuador, Panama, Peru, and Venezuela (SANTOS & BRESOVIT 2003; CARVALHO et al. 2014).

Family Palpimanidae Thorell, 1870

Otiotrops oblongus Simon, 1891

Otiotrops oblongus SIMON (1891).

Otiotrops lapidicola SIMON (1893a) — MELLO-LEITÃO (1927).

Otiotrops carpenteri CHICKERING (1966)

Otiotrops oblongus PLATNICK (1975).

Material examined: BRAZIL, RORAIMA: Boa Vista (Campus de Cauamé, Universidade Federal de Roraima, 02°52'38.4" N, 060°43'13.1" W; 91 m a.s.l.), 1 ♂, 22.VII.2014, M.C. Schneider leg. (UFMG 17090); 3m, 1 ♀, L.S. Carvalho & M.C. Schneider leg. (CHNUFPI 1306); Bonfim (near the road RR401, about 7.5 km from Bonfim, 03°21'1.8" N, 059°54'15.8" W; 89 m a.s.l.), 1 ♂, 20.VII.2014, L.S. Carvalho & M.C. Schneider leg. (UFMG 17089).

Distribution: This species is recorded only for northern South America (PLATNICK 1975: map 2). There are published records for the following countries: Brazil (Amazonas, and Roraima), Guyana, St. Vincent, Trinidad, and Venezuela (CHICKERING 1966; PLATNICK 1975; BRESOVIT & BONALDO 1993).

Family Pholcidae C. L. Koch, 1850

Carapoia paraguensis González-Sponga, 1998

Carapoia paraguensis GONZÁLEZ-SPONGA (1998) — HUBER (2000, 2005).

Material examined: BRAZIL, RORAIMA: Amajari (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°45'8.4" N, 061°42'49.3" W; 647 m a.s.l.), 1 ♀, 16.VII.2014, L.S. Carvalho & M.C. Schneider leg. (CHNUFPI 1296); 1 ♂ (CHNUFPI 1288); 2m, 1 ♀ (UFMG 17109); Boa Vista (Campus de Cauamé, Universidade Federal de Roraima, 02°52'38.4" N, 060°43'13.1" W; 91 m a.s.l.), 1 ♀, 22.VII.2014, M.C. Schneider leg. (UFMG 17110); 2 ♀ (CHNUFPI 1273); 5 ♂ (UFMG 17111); Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♀, 23.VII.2014, L.S. Carvalho et al. leg. (UFMG 17113); 1 ♀ (UFMG 17114); 1 ♂ (CHNUFPI 1272); 1 ♂ (CHNUFPI 1274); 1 ♂ (UFMG 17112); 1 ♂ (UFMG 17115).

Distribution: This species is recorded only for northern South America (HUBER 2005: fig. 1). There are published records for Brazil (Amazonas, and Roraima), Guyana, and Venezuela (HUBER 2000, 2005).

Metagonia mariguitarensis (González-Sponga, 1998)

Anomalaia mariguitarensis GONZÁLEZ-SPONGA (1998).

Metagonia mariguitarensis HUBER (2000, 2004).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider leg. (CHNUFPI 1265); 1 ♀ (UFMG 17117); 3m (UFMG 17116); (Cachoeira do Paiva, Vila Tepequém, 03°45'45.6" N, 061°45'18.0" W; 582 m a.s.l.), 1 ♀, 17.VII.2014, L.S. Carvalho & M.C. Schneider leg. (CHNUFPI 1277); 1 ♀ (UFMG 17118); 1 ♂, 2 ♀ (UFMG 17119); 1 ♂, 3 ♀ (UFMG 17120); Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♀, 23.VII.2014, L.S. Carvalho et al. leg. (UFMG 17122); 5 ♀ (UFMG 17121); 2 ♀, 24.VII.2014, L.S. Carvalho & M.C. Schneider leg. (UFMG 17123); 1 ♂ (CHNUFPI 1263); 1 ♂ (CHNUFPI 1275).

Distribution: This species is recorded only from northern South America and published records exist only for Brazil (Roraima), Peru, and Venezuela (HUBER 2000).

Micropholcus fauroti (Simon, 1887)*Pholcus fauroti* SIMON (1887).*Pholcus infirmus* THORELL (1895).*Pholcus unicolor* PETRUNKEVITCH (1929).*Leptopholcus occidentalis* MELLO-LEITÃO (1929a).*Pholcus senegalensis* MILLOT (1941).*Pholcus chavaneii* MILLOT (1946).*Micromerys occidentalis* — MELLO-LEITÃO (1946).*Micropholcus fauroti* — DEELEMEN-REINHOLD & PRINSEN (1987);

HUBER (2000, 2011); BEATTY et al. (2008).

Mariguitaia divergentis GONZÁLEZ-SPONGA (2004).*Mariguitaia museorum* GONZÁLEZ-SPONGA (2004).*Mariguitaia neoespartana* GONZÁLEZ-SPONGA (2004).*Mariguitaia sucrensii* GONZÁLEZ-SPONGA (2004).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, Pousada PSJ, 03°46'56.2" N, 061°43'20.8" W; 640 m.a.s.l.), 2 ♂, 18.VII.2014, L.S. Carvalho *leg.* (UFMG 17125); Bonfim, (Pousada Takutu, 03°21'48.7" N, 059°50'15.7" W; 92 m a.s.l.), 1 ♀, 20.VII.2014, L.S. Carvalho *leg.* (UFMG 17124).

Distribution: This is a widespread pantropical synanthropic species (DEELEMEN-REINHOLD & PRINSEN 1987; DEELEMEN-REINHOLD & VAN HARTEN 2001; IRIE 2000; SAARISTO 2001; CARVALHO et al. 2014). In Brazil, it is known from only Pernambuco and Roraima (**new record**) (CARVALHO et al. 2014).

Family Pisauridae Simon, 1890

Architis tenuipes (Simon, 1898)*Drances tenuipes* SIMON (1898a).*Architis vilhena* CARICO (1981, 1989).*Architis tenuipes* — CARICO (1993); SANTOS (2007).

Material examined: BRAZIL, RORAIMA: Cantá (Near the road BR432, about 10 km from Cantá, 02°38'9.3" N, 060°34'17.1" W; 106 m a.s.l.), 1 ♂, 21.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1313); 1 ♂, 1 ♀ (UFMG 17319).

Distribution: This species is widely distributed in the northern Brazil (SANTOS 2007: fig. 7; SANTOS & NOGUEIRA 2008: fig. 12). There are published records for the following Brazilian states: Acre, Amazonas, Maranhão, Mato Grosso, Pará, Rondônia, and Roraima (SANTOS 2007; SANTOS & NOGUEIRA 2008; BONALDO et al. 2009b).

Architis tenuis Simon, 1898*Architis tenuis* SIMON (1898a) — CARICO (1981); SANTOS (2007).*Architis nitidoplumosa* SIMON (1898a); CARICO (1981); SIERWALD (1989).*Thanatidius proximus* MELLO-LEITÃO (1921).*Thanatidius parahybensis* MELLO-LEITÃO (1924).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1269); 1 ♂ (UFMG 17317); (Cachoeira do Paiva, Vila Tepequém, 03°45'45.6" N, 061°45'18" W; 582 m a.s.l.), 10 ♀, 17.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17095); Bonfim, (near the road RR401, surroundings of Bonfim, 03°22'9.0" N, 059°50'18.4" W, 95 m a.s.l.),

1 ♀, 20.VII.2014, L.S. Carvalho *leg.* (UFMG 17096); Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♂, 23.VII.2014, L.S. Carvalho et al. *leg.* (UFMG 17098), 3 ♀ (UFMG 17320); (near the road BR432, about 10 km from Cantá, 02°38'09.3" N, 060°34'17.1" W; 106 m a.s.l.), 1 ♀, 21.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17097).

Distribution: This species is widely distributed in the Neotropical Region (SANTOS 2007: fig. 6). There are published records from the following countries: Brazil (Alagoas, Amazonas, Bahia, Ceará, Espírito Santo, Mato Grosso, Minas Gerais, Pará, Pernambuco, Piauí, Rio Grande do Sul, Rondônia, Roraima, São Paulo, and Sergipe), Colombia, French Guiana, Guyana, Panama, Peru, Surinam, and Venezuela (LISE 1998B; SANTOS 2007; BONALDO et al. 2009b; CARVALHO et al. 2014).

Family Scytodidae Blackwall, 1864

Scytodes fusca Walckenaer, 1837*Scytodes fusca* WALCKENAER (1837) — BRESCOVIT & RHEIMS (2000); ONO (2011); ŠESTÁKOVÁ et al. (2014).*Scytodes domestica* DOLESCHALL (1859) — KULCZYŃSKI (1911).*Scytodes guianensis* TACZANOWSKI (1872).*Dictis fumida* THORELL (1891).*Scytodes hebraica* SIMON (1891) — F. O. PICKARD-CAMBRIDGE (1899b).*Scytodes bajula* SIMON (1891).*Dictis domestica* — THORELL (1895).*Scytodes atrofusca* STRAND (1916b).*Scytodes campinensis* MELLO-LEITÃO (1918).*Scytodes discolor* MELLO-LEITÃO (1918).*Scytodes iguassuensis* MELLO-LEITÃO (1918).*Scytodes nannipes* CHAMBERLIN & IVIE (1936).*Scytodes velutina* MILLOT (1941).*Scytodes torquatus* KRAUS (1955).*Scytodes torquata* — BRIGNOLI (1983).

Material examined: BRAZIL, RORAIMA: Bonfim, (near the road RR401, surroundings of Bonfim, 03°22'9.0" N, 059°50'18.4" W; 95 m a.s.l.), 1 ♂, 1 ♀, 20.VII.2014, L.S. Carvalho *leg.* (UFMG 17105).

Distribution: This species has a pantropical distribution and is introduced to Europe (ŠESTÁKOVÁ et al. 2014). It is commonly associated with human habitations throughout Central and South America (BRESCOVIT & RHEIMS 2000).

Family Selenopidae Simon, 1897

Selenops geraldinae Corronca, 1996*Selenops geraldinae* CORRONCA (1996, 1998) — CREWS (2011); GALVIS & FLÓREZ (2015).*Selenops willinki* CORRONCA (1998).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 17092); 1 ♀ (CHNUFPI 1315).

Distribution: This species is distributed over northern South America (CORRONCA 1998: map 1; CREYS 2011: map 2). There are published records from the departments of Mara and Bolívar, Venezuela (CORRONCA 1998), and Trini-

dad and Tobago (CREWS 2011). The new Roraima record is the first for *S. geraldinae* in Brazil and extends the known distribution of the species by at least 450 km southwards as far as Amajari.

Family Sparassidae Bertkau, 1872

Quemedice piracuruca Rheims, Labarque & Ramírez, 2008

Quemedice piracuruca RHEIMS et al. (2008).

Material examined: BRAZIL, RORAIMA: Boa Vista, (Campus de Cauamé, Universidade Federal de Roraima, 02°52'38.4" N, 060°43'13.1" W; 91 m a.s.l.), 1 ♂, 22.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (UFMG 15791).

Distribution: This species was previously recorded only from Colombia and Brazil (Minas Gerais and Piauí) (Rheims et al. 2008), and the new record from Roraima extends the known geographical distribution by at least 1,400 km toward the northeast.

Family Synotaxidae Simon, 1894

Synotaxus monoceros (Caporiacco, 1947)

Argyrodina monoceros CAPORIAMCO (1947, 1948).

Synotaxus pupularum EXLINE & LEVI (1965).

Synotaxus monoceros EXLINE & LEVI (1965).

Synotaxus monoceros AGNARSSON (2003, 2004); AGNARSSON et al. (2007).

Material examined: BRAZIL, RORAIMA: Boa Vista, (Campus de Cauamé, Universidade Federal de Roraima, 02°52'38.4" N, 060°43'13.1" W; 91 m a.s.l.), 2 ♀, 22.VII.2014, M.C. Schneider (UFMG 17091); 2 ♀ (CHNUFPI 1276).

Distribution: This species is known only for northern Amazon basin, with published records for Guyana, Trinidad and Brazil (Amazonas) (EXLINE & LEVI 1965; AGNARSSON 2003; SANTOS & RHEIMS 2005). This is the first record of *S. monoceros* from Roraima and fills a gap in its known geographical distribution.

Family Tetragnathidae Menge, 1866

Glenognatha gaujoni Simon, 1895

Glenognatha gaujoni SIMON (1895b) — CABRA-GARCÍA et al. (2014); CABRA-GARCÍA & BRESCOVIT (2016).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, Pousada PSJ, 03°46'55.4" N, 061°43'19.0" W; 587 m a.s.l.), 1 ♂, 17.VI.2014, J. Cabra et al. (MZSP 69710); 2 ♂, 1 ♀, 15.VI.2014 (MZSP 69711); (Estância Ecológica SESC Tepequém, Vila Tepequém, 03°46'55.2" N, 061°43'19.3" W; 655 m a.s.l.), 1 ♀, 16.VI.2014, J. Cabra et al. *leg.* (MZSP 69706).

Distribution: This species is widely distributed over South America, with known records in Brazil (Amazonas, Rondônia, Roraima, Mato Grosso and São Paulo), Colombia, Ecuador, Peru and Venezuela (CABRA-GARCÍA & BRESCOVIT 2016).

Family Theridiidae Sundevall, 1833

Dipoena atlantica Chickering, 1943

Dipoena atlantica CHICKERING (1943) — LEVI (1963).

Material examined: BRAZIL, RORAIMA: Cantá (near the road BR432, about 10 km from Cantá, 02°35'15.3" N, 060°38'27.6" W; 105 m a.s.l.), 1 ♂, 23.VII.2014, L.S. Carvalho et al. *leg.* (UFMG 16878).

Distribution: This species is widely distributed in the Neotropical Region, with published records in the following countries: Brazil (Mato Grosso, Minas Gerais, Rio de Janeiro and Roraima [new record]), Panama, Paraguay, Peru, and Venezuela (CHICKERING 1943; LEVI 1963).

Dipoena duodecimpunctata Chickering, 1943

Dipoena 12-punctata CHICKERING (1943).

Dipoena duodecimpunctata LEVI (1963).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♂, 17.VII.2014, L.S. Carvalho et al. *leg.* (UFMG 16877).

Distribution: This species is distributed in through Brazil (Amazonas, Roraima, Pará, Acre, Tocantins, São Paulo, Paraná, and Santa Catarina), Panama, and Venezuela (CHICKERING 1943; LEVI 1963; RODRIGUES 2013).

Family Trochanteriidae Karsch, 1879

Trochanteria gomezi Canals, 1933

Trochanteria gomezi CANALS (1933); PLATNICK (1986).

Oltaclaea major MELLO-LEITÃO (1942a).

Material examined: BRAZIL, RORAIMA: Boa Vista (Campus de Cauamé, Universidade Federal de Roraima, 02°52'38.4" N, 060°43'13.1" W; 91 m a.s.l.), 2 ♂, 6 ♀, 22.VII.2014, L.S. Carvalho & M.C. Schneider *leg.* (CHNUFPI 1307); 1 ♂ (CHNUFPI 1571); 1 ♂ (UFMG 15584); 1 ♀ (UFMG 15954); 2 ♂, 8 ♀ (UFMG 17087); 1 ♀, 23.VII.2014 (UFMG 15955); 1 ♀, 24.VII.2014 (UFMG 15956).

Distribution: The distribution pattern of this species was unclear for many years. It was known only from Argentina and Paraguay (PLATNICK 1986), until recently found in the state of Piauí, northeastern Brazil (Carvalho et al. 2014). The new record from Boa Vista, Roraima, represents a range extension of 2,300 km toward the northwest. This species may be widely distributed over the Cerrado biome, but the scarcity of sampling in appropriate microhabitats for the species hinders the validation of this hypothesis.

Natural history: All specimens of *T. gomezi* were collected from under bark of *Curatella americana* L. (Dilleniaceae) by carefully removing pieces of bark with a forceps. These spiders usually run laterally on the tree trunks to hide in small spaces in, or under, the bark. The association between *T. gomezi* and *C. americana* was also reported in the state of Piauí by Carvalho et al. (2014). As *C. americana* is widely distributed from Central America to Bolivia,

and in almost all of Brazil, especially in the Cerrado biome (Sarmiento and Monasterio 1983; Vilar et al. 2009), further sampling effort from the barks of *C. americana* should be made. This may provide more precise information on the natural history of *T. gomezi*.

Family Uloboridae Thorell, 1869

Philoponella vittata (Keyserling, 1881)

Uloborus vittatus KEYSERLING (1881a).

Uloborus servulus SIMON (1893a).

Uloborus semiargenteus SIMON (1893b).

Uloborus amazonicus MELLO-LEITÃO (1949).

Philoponella semiargentea — LEHTINEN (1967).

Philoponella servula — LEHTINEN (1967).

Philoponella vittata — OPELL (1979).

Material examined: BRAZIL, RORAIMA: Amajari (Vila Tepequém, near Pousada PSJ, 03°47'10.4" N, 061°43'15.3" W; 640 m a.s.l.), 1 ♂, 6 ♀, 15.VII.2014, L.S. Carvalho & M.C. Schneider leg. (UFMG 17106).

Distribution: This species is widely distributed in the Neotropical Region (OPELL 1979: map 5). There are published records from the following countries: Bolivia, Brazil (Amazonas, Pará and Roraima [**new record**]), Colombia, Guyana, Panama, Paraguay, Peru, and Venezuela (OPELL 1979; ALVES-COSTA & GONZAGA 2001).

DISCUSSION

The data presented here were opportunistically collected and may be taxonomically, temporally, and spatially biased. That is, relatively conspicuous (widespread) species were more often collected, the sampling period was short, and sampling sites were close to main roads. Most of the species recorded, including 32 of the 36 species newly recorded from Brazil and/or Roraima, are widespread species, but the new data are not unimportant. On the contrary, these data reflect the historical paucity of spider sampling throughout Roraima. Thus, there is the need to expand spider surveys in this region. The sampling heterogeneity in Roraima is not different from the overall situation throughout Brazil (BRESCOVIT et al. 2011: fig. 5), and corresponds to a general rule for Brazilian invertebrates, vertebrates and angiosperms (LEWINSOHN et al. 2005; OLIVEIRA et al. 2016). Even within the relatively better-studied animal groups and biomes, such as the Atlantic and the Amazon Forests, geographic coverage is very restricted and often just a few localities have been sampled adequately (Lewinsohn et al. 2005). As a result, there is a strong spatial bias regarding the species richness, species composition and endemism knowledge in the Brazilian biodiversity (OLIVEIRA et al. 2016).

Information from natural history collections, such these presented here, are important as they are easily usable and present a high precision in georeferencing (GRAHAM et al. 2004). This type of data can be used in species distribution models for many purposes, such as conservation (FERRAZ et al. 2012). These collection data are also crucial because

they are a permanent record of a species at a given place and time (FUNK & RICHARDSON 2002).

Our sampled localities are within the areas of endemism of Roraima, supported by a relatively low number (at most 12 spp.) of synendemic spider species (OLIVEIRA et al. 2015). However, our results suggest that these areas harbor a higher number of synendemic species than the survey by OLIVEIRA et al. (2015), which therefore had clearly sampling bias. Additionally, the relevancy of our results relies on the use of a high number of species associated with high endemism levels for conservation purposes. The basis for environmental management and conservation policies is, at a bare minimum, the distribution and abundance of every species that requires conservation (POSSINGHAM et al. 2007).

The spider fauna of Roraima is one of the most poorly known in Brazil, with only a few published papers and mostly related to the spiders of the Maracá Island (where there are 92 recorded; number updated using the current spider nomenclature; see WORLD SPIDER CATALOG 2017), or individual species descriptions (LISE 1998a, 1998b; BUCKUP & MARQUES 1989, 1991, 1992; MARQUES & BUCKUP 1992). Of these, only seven species (about 7.5% of the total) were recorded in our samples: the araneid orb-weavers *Aculepeira travassosi*, *Amazonpeira masaka*, *Araneus guttatus*, and *Argiope argentata*, the pisaurid *Architis tenuis*, the theridiid *Dipoena atlantica*, and the palpimanid *Otiotrops oblongus*.

As observed, we expected a large number of previously unrecorded species as result of sampling anywhere in the state. However, future surveys should focus on remote and mountainous regions (e.g., Mount Roraima), or unexplored niches (e.g., soil and leaf litter). Such surveys may provide a wider overview of the whole spider fauna. Additionally, long-term or ecological studies on arachnids of Roraima have never been done. Such research would be a very interesting because the spider dynamics in the Cerrado (savanna) patches within the Amazon Forest is still completely unknown.

ACKNOWLEDGEMENTS

We thank Ricardo Pinto da Rocha (USP), Rafael Boldrini (UFRR), and Ranyse Querino Barbosa da Silva (EMBRAPA Meio Norte) for their suggestions of sampling places and/or logistic support in Roraima; Adalberto J. Santos (UFMG), Bárbara T. Faleiro (UFMG) and José Paulo Leite Guadanucci (UNESP) for assistance with species identification; and Regiane Saturnino (MPEG) and two anonymous reviewers for the revision of an early version of the manuscript. The fieldwork was supported by the project “Meiose em escorpiões (Arachnida): modelo para compreender a evolução em espécies com cromossomos aquiasmáticos”, funded by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, Process #2011/21643-1). JCG is financially supported from FAPESP, grant #2013/20262-0. The surveys in Roraima were part of LSC’s Ph.D. project.

LITERATURE CITED

- ABRAHIM, N., A.D. BRESCOVIT, C.A. RHEIMS, A.J. SANTOS, R. OTT & A.B. BONALDO. 2012. A revision of the Neotropical goblin spider genus *Neoxyphinus* Birabén, 1953 (Araneae, Oonopidae). *American Museum Novitates* 3743: 1–75. <http://hdl.handle.net/2246/6172>
- Adis, J., A.B. Bonaldo, A.D. Brescovit, R. Bertani, A.B. Kury, et al. 2002. Arachnida at Reserva Ducke, Central Amazonia/Brazil. *Amazoniana* (Kiel) 17: 1–14.
- AGNARSSON, I. 2003. The phylogenetic placement and circumscription of the genus *Synotaxus* (Araneae: Synotaxidae), a new species from Guyana, and notes on theridioid phylogeny. *Invertebrate Systematics* 17: 719–734. doi: [10.1071/IS03002](https://doi.org/10.1071/IS03002)
- AGNARSSON, I. 2004. Morphological phylogeny of cobweb spiders and their relatives (Araneae, Araneioidea, Theridiidae). *Zoological Journal of the Linnean Society* 141: 447–626. [10.1111/j.1096-3642.2004.00120.x](https://doi.org/10.1111/j.1096-3642.2004.00120.x)
- AGNARSSON, I., J.A. CODDINGTON & B. KNOFLACH. 2007. Morphology and evolution of cobweb spider male genitalia (Araneae, Theridiidae). *Journal of Arachnology* 35: 334–395. [10.1636/SH-06-36.1](https://doi.org/10.1636/SH-06-36.1)
- AGNARSSON, I., S.M. LEQUIER, M. KUNTNER, R.-C. CHENG, J.A. CODDINGTON & G. BINFORD. 2016. Phylogeography of a good Caribbean disperser: *Argiope argentata* (Araneae, Araneidae) and a new ‘cryptic’ species from Cuba. *ZooKeys* 625: 25–44. [10.3897/zookeys.625.8729](https://doi.org/10.3897/zookeys.625.8729)
- ALVES-COSTA, C.P. & M.O. GONZAGA. 2001. Prey capture and spatial distribution of *Philoponella vittata* (Araneae Uloboridae) in host webs. *Ethology Ecology and Evolution* 13(3): 239–246. doi: [10.1080/08927014.2001.9522773](https://doi.org/10.1080/08927014.2001.9522773)
- ARCHER, A.F. 1963. Catalogo de las arañas chilenas de las familias de la division Metarachnae. *Publicaciones Ocasionales del Museo Nacional de Historia Natural, Santiago* 1: 1–32.
- BADCOCK, A.D. 1932. Reports of an expedition to Paraguay and Brazil in 1926–1927 supported by the Trustes of the Percy Sladen Memorial Fund and the Executive Committee of the Carnegie Trust for the Universities of Scotland. Arachnida from the Paraguayan Chaco. *Journal of the Linnean Society of London, Zoology* 38: 1–48.
- BANKS, N. 1894. Notes on Larinia and Cercidia. *Entomological News* 5: 8–9.
- BANKS, N. 1898. Arachnida from Baja California and other parts of Mexico. *Proceedings of the California Academy of Sciences* (3) 1: 205–308.
- BANKS, N. 1909. Arachnida from Costa Rica. *Proceedings of the Academy of Natural Sciences of Philadelphia* 61: 194–234.
- BANKS, N. 1930. Arachnida. In: *The Norwegian Zoological Expedition of the Galapagos Islands 1925 conducted by Alf Wollebaek*. *Nyt Magazin for Naturvidenskaberne* 68: 271–278.
- BARBOSA, R. I., E.J.G. FERREIRA & E.G. CASTELLÓN (eds.). 1997. *Homem, ambiente e ecologia no estado de Roraima*. Manaus: INPA. 613 pp.
- BARRION, A.A., A.T. BARRION, C.V. CASAL, L.D. TAYLO & D.M. AMALIN. 1988. The orb-weaving spiders genus *Neoscona* (Araneae: Araneidae) in the Philippines. *The Philippine Agricultural Scientist* 69: 385–409.
- BEATTY, J.A., J.W. BERRY & B.A. HUBER. 2008. The pholcid spiders of Micronesia and Polynesia (Araneae, Pholcidae). *Journal of Arachnology* 36: 1–25. doi: [10.1636/H05-66.1](https://doi.org/10.1636/H05-66.1)
- BERMAN, J. D. & H.W. LEVI. 1971. The orb weaver genus *Neoscona* in North America (Araneae: Araneidae). *Bulletin of the Museum of Comparative Zoology at Harvard College* 141: 465–500.
- BERTANI, R., A.S. SANTOS, A.D. ABEGG, F.R. ORTIZ & M.A. FREITAS. 2016. First record of the genus *Psalmopoeus* (Araneae: Theraphosidae) in Brazil. *Check List* 12(2): 1860. doi: [10.15560/12.2.1860](https://doi.org/10.15560/12.2.1860)
- BERTANI, R. & Y. ARAÚJO. 2006. A new species of *Holothele* Karsch, 1879 (Theraphosidae, Ischnocolinae) from Venezuela. *Revista Ibérica de Aracnología* 12: 13–16. http://www.sea-entomologia.org/Publicaciones/RevistaIbericaAracnologia/RIA12/013_016Holothele.pdf
- BERTKAU, P. 1880. Verzeichniss der von Prof. Ed. van Beneden auf seiner im Auftrage der Belgischen Regierung unternommen wissenschaftlichen Reise nach Brasilien und La Plata im Jahren 1872–73 gesammelten Arachniden. *Mémoires Couronnés et Mémoires des Savants Étrangers de l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique* 43: 1–120.
- BLACKWALL, J. 1863. Description of newly discovered spiders captured in Rio de Janeiro, by John Gray and the Rev. Hamlet Clark [continued]. *Annals and Magazine of Natural History* (3) 11: 29–45.
- BONALDO, A.B., A.D. BRESCOVIT, H. HÖFER, T. GASNIER & A.A. LISE. 2009a. A araneofauna (Arachnida, Araneae) da Reserva Florestal Adolfo Ducke, Manaus, Amazonas, Brasil; pp. 201–222, in: E.C. FRANKLIN & C. MAGALHÃES. (org.). *A fauna de artrópodos da Reserva Florestal Adolpho Ducke: estado atual do conhecimento taxonômico e biológico*, 2ed. Manaus: Instituto Nacional de Pesquisa da Amazonia.
- BONALDO, A.B., L.S. CARVALHO, R. PINTO-DA-ROCHA, A.L. TOURINHO, L.T. TOURINHO, et al. 2009b. Inventário e história natural dos aracnídeos da Floresta Nacional de Caxiuanã; pp. 577–622, in: P.L.B. LISBOA (org.). *Caxiuanã: desafios para a conservação de uma Floresta na Amazônia*, 1ed. Belém: MPEG.
- BRADY, A.R. 1962. The spider genus *Sosippus* in North America, Mexico, and Central America (Araneae, Lycosidae). *Psyche*, Cambridge 69: 129–164.
- BRESCOVIT, A.D. 1996. Revisão do gênero *Centroctenus* Mello-Leitão (Araneae, Ctenidae, Cteninae). *Revista Brasileira de Entomologia* 40(2): 301–313.
- BRESCOVIT, A.D. & A.B. BONALDO. 1993. On the genus *Otiotrops* in Brazil (Araneae, Palpimanidae). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique (Entomologie)* 63: 47–50.
- BRESCOVIT, A.D. & C.A. RHEIMS. 2000. On the synanthropic species of the genus *Scytodes* Latreille (Araneae, Scytodidae) of Brazil, with synonymies and records of these species in other Neotropical countries. *Bulletin of the British Arachnological Society* 11: 320–330.
- BRESCOVIT, A.D., U. OLIVEIRA & A.J. SANTOS. 2011. Aranhas (Araneae, Arachnida) do Estado de São Paulo, Brasil: diversidade, esforço amostral e estado do conhecimento. *Biota Neotropica* 11: 1–32. <http://www.biotaneotropica.org.br/v11n1a/en/fullpaper?bn0381101a2011+pt>
- BRIGNOLI, P.M. 1983. *A catalogue of the Araneae described between 1940 and 1981*. Manchester: Manchester University Press. 755 pp.
- BRYANT, E.B. 1940. Cuban spiders in the Museum of Comparative Zoology. *Bulletin of the Museum of Comparative Zoology at Harvard College* 86: 247–532.
- BRYANT, E.B. 1945. The Argiopidae of Hispaniola. *Bulletin of the Museum of Comparative Zoology at Harvard College* 95: 357–422.
- BRYANT, E.B. 1948. The spiders of Hispaniola. *Bulletin of the Museum of Comparative Zoology at Harvard College* 100: 329–447.
- BUCKUP, E.H. & M.A.L. MARQUES. 1989. Aranhas Theridiidae da Ilha de Maracá, Roraima, Brasil. I. Nova espécie de *Echinotheridion* e descrição da fêmea de *Phoroncidia moyobamba* (Araneae). *Iheringia (Zoology)* 69: 123–129.
- Buckup, E.H. and M.A.L. Marques. 1991. Aranhas Theridiidae da Ilha de Maracá, Roraima, Brasil. II. Gênero *Achaearanea* (Araneae). *Iheringia* 71: 81–89.
- BUCKUP, E.H. & M.A.L. MARQUES. 1992. Aranhas Theridiidae da

- Ilha de Maracá, Roraima, Brasil. III. Gêneros Chrysso e Episinus (Araneae). *Iheringia (Zoology)* 72: 121–125.
- BUTLER, A.G. 1873. A list of the spiders of the genus *Acrosoma*. *Proceedings of the Zoological Society of London* 1873: 420–429.
- CABRA-GARCÍA, J. & A.D. BRESCOVIT. 2016. Revision and phylogenetic analysis of the orb-weaving spider genus *Glenognatha* Simon, 1887 (Araneae, Tetragnathidae). *Zootaxa* 4069: 1–183. doi: [10.11646/zootaxa.4069.1.1](https://doi.org/10.11646/zootaxa.4069.1.1)
- CABRA-GARCÍA, J., G. HORMIGA, & A.D. BRESCOVIT. 2014. Female genital morphology in the secondarily haplogyne spider genus *Glenognatha* Simon, 1887 (Araneae, Tetragnathidae), with comments on its phylogenetic significance. *Journal of Morphology* 275: 1027–1040. [10.1002/jmor.20280](https://doi.org/10.1002/jmor.20280)
- CANALS, J. 1933a. Curiosa araña argentina del genero *Trochantheria* Karsch. *Physis* 11: 233–237.
- CAPOCASALE, R.M. 1982. Las especies del genero *Porrimosa* Roewer, 1959 (Araneae, Hippasinae). *Journal of Arachnology* 10: 145–156.
- CAPOCASALE, R.M. 1991. Nuevos aportes al género *Porrimosa* Roewer (Araneae, Lycosidae). *Journal of Arachnology* 19: 93–96.
- CAPORIACCO, L. DI 1947. Diagnosi preliminari de specie nuove di aracnidi della Guiana Britannica raccolte dai professori Beccari e Romiti. *Monitore Zoologico Italiano* 56: 20–34.
- CAPORIACCO, L. DI 1948. Arachnida of British Guiana collected in 1931 and 1936 by Professors Beccari and Romiti. *Proceedings of the Zoological Society of London* 118: 607–747.
- CAPORIACCO, L. DI 1954. Araignées de la Guyane Française du Muséum d'Histoire Naturelle de Paris. *Commentationes Pontificia Academia Scientiarum* 16: 45–193
- CAPORIACCO, L. DI 1955. Estudios sobre los aracnidos de Venezuela. 2a parte: Araneae. *Acta Biologica Venezuelica* 1: 265–448.
- CARICO, J.E. 1981. The Neotropical spider genera *Architis* and *Staberius* (Pisauridae). *Bulletin of the American Museum of Natural History* 170: 140–153.
- CARICO, J.E. 1989. Descriptions of two new species of the genus *Architis* (Araneae, Pisauridae) and the female of *A. vilhena*. *Journal of Arachnology* 17: 221–224.
- Carico, J.E. 1993. Taxonomic notes on the genus *Architis* (Araneae, Pisauridae) and status of the genus *Sisenna* Simon. *Journal of Arachnology* 21: 202–204.
- Carvalho, L.S., A.D. Brescovit, A.J. Santos, U. Oliveira and J.P.L. Guadanucci. 2014. Aranhas da Caatinga; pp. 15–32, in: Bravo, F. and Calor, A. (orgs.). *Artrópodes do semiárido: biodiversidade e conservação*, 1ed. Feira de Santana: Printmídia.
- CHAMBERLIN, R.V. 1916. Results of the Yale Peruvian expedition of 1911: The Arachnida. *Bulletin of the Museum of Comparative Zoology at Harvard College* 60: 177–299.
- CHAMBERLIN, R.V. 1917. New spiders of the family Aviculariidae. *Bulletin of the Museum of Comparative Zoology at Harvard College* 61: 25–75.
- CHAMBERLIN, R.V. 1924. The spider fauna of the shores and islands of the Gulf of California. *Proceedings of the California Academy of Sciences* 12: 561–694.
- CHAMBERLIN, R.V. 1925. Diagnoses of new American Arachnida. *Bulletin of the Museum of Comparative Zoology at Harvard College* 67: 209–248.
- CHAMBERLIN, R.V. & W. IVIE. 1936. New spiders from Mexico and Panama. *Bulletin of the University of Utah* 27(5): 1–103.
- CHICKERING, A.M. 1943. Twenty-one new species of *Dipoena* (Theridiidae) from Panama. *Transactions of the American Microscopical Society* 62: 329–378.
- CHICKERING, A.M. 1960. New species of *Chaetacis* (Araneae, Argiopidae) from South America. *Annals and Magazine of Natural History* (13) 2: 465–475.
- CHICKERING, A.M. 1960. Six new species of *Micrathena* (Araneae, Argiopidae) from South America with notes on known species. *Proceedings of the Zoological Society of London* 135: 65–89.
- CHICKERING, A.M. 1961. The genus *Micrathena* (Araneae, Argiopidae) in Central America. *Bulletin of the Museum of Comparative Zoology at Harvard College* 125: 391–470.
- CHICKERING, A.M. 1964. The genus *Micrathena* (Araneae, Araneidae) in the West Indies. *Bulletin of the Museum of Comparative Zoology at Harvard College* 131: 251–281.
- CHICKERING, A.M. 1966. New species of Palpimanidae (Araneae) from the West Indies. *Psyche, Cambridge* 73: 208–216.
- CORRONCA, J.A. 1996. Tres nuevas especies de *Selenops* Latreille, 1819 (Araneae: Selenopidae) en América del Sur. *Neotropica* 42: 91–96.
- CORRONCA, J.A. 1998. The South American spiders of the genus *Selenops* (Araneae, Selenopidae) with a description of three new species. *Studies on Neotropical Fauna and Environment* 33: 124–148.
- COSTA, E.L.S. & G.R.S. RUIZ. 2014. Taxonomic revision of *Scopocira* Simon, 1900 (Araneae: Salticidae). *Zootaxa* 3893(2): 151–195. doi: [10.11646/zootaxa.3893.2.1](https://doi.org/10.11646/zootaxa.3893.2.1)
- COYLE, F.A. 1995. A revision of the funnelweb mygalomorph spider subfamily Ischnothelinae (Araneae, Dipluridae). *Bulletin of the American Museum of Natural History* 226: 1–133. <http://hdl.handle.net/2246/1665>
- CREWS, S.C. 2011. A revision of the spider genus *Selenops* Latreille, 1819 (Arachnida, Araneae, Selenopidae) in North America, Central America and the Caribbean. *ZooKeys* 105: 1–182. doi: [10.3897/zookeys.105.724](https://doi.org/10.3897/zookeys.105.724)
- DE GEER, C. 1778. *Mémoires pour servir à l'histoire des insectes*. Stockholm 7(3–4): 176–324.
- DEELEMEN-REINHOLD, C.L. & A. VAN HARTEN. 2001. Description of some interesting, new or little known Pholcidae (Araneae) from Yemen; pp. 193–207, in: I. PRAKASH (ed.). *Ecology of desert environments*. Jodhpur: Scientific Publishers.
- DEELEMEN-REINHOLD, C.L. & J.D. PRINSEN. 1987. *Micropholcus fauroti* (Simon) n. comb., a pantropical, synanthropic spider (Araneae: Pholcidae). *Entomologische Berichten, Amsterdam* 47: 73–77.
- DIAS, S.C. & A.B. BONALDO. 2012. Abundância relativa e riqueza de espécies de aranhas (Arachnida, Araneae) em clareiras originadas da exploração de petróleo na Bacia do Rio Urucu (Coari, Amazonas, Brasil). *Boletim do Museu Paraense Emílio Goeldi. Ciências Naturais* 7: 123–152. <http://scielo.iec.pa.gov.br/pdf/bmpegcn/v7n2/v7n2a02.pdf>
- DIERKENS, M. 2011. Contribution à l'étude des Araneidae de la Guyane française 1: Gasteracanthinae. *Cahiers Musée des Confluences – Études Scientifiques* 2: 99–108.
- DIERKENS, M. 2012. Contribution à l'étude de divers genres d'Araneidae (Araneae) de Guyane française. *Bulletin Mensuel de la Société Linnéenne de Lyon* 81: 23–33.
- DOLESCHALL, L. 1859. Tweede Bijdrage tot de Kennis der Arachniden van den Indischen Archipel. *Acta Societatis Scientiarum Indica-Neerlandica* 5: 1–60.
- EDEN, M.J. & D.F.M. MCGREGOR. 1998. Ilha de Maraca and the Roraima Region; pp. 1–11, in: W. MILLIKEN & J.A. RATTER (eds.). *Maraca: ecology of an amazonian rainforest*. London: John Wiley and Sons.
- EICKSTEDT, V.R. VON. 1978. Estudo sobre a sistemática de *Ctenus taeniatus* (Araneae; Labidognatha). *Memórias do Instituto Butantan* 40/41: 211–219.
- EXLINE, H. & H.W. LEVI. 1965. The spider genus *Synotaxus* (Araneae, Theridiidae). *Transactions of the American Microscopical Society* 84: 177–184.
- FABRICIUS, J.C. 1775. *Systema entomologiae, sistens insectorum classes, ordines, genera, species, adiectis, synonymis, locis descriptionibus observationibus*. Flensburg and Lipsiae, 832 pp. (Araneae), pp. 431–441.
- FALCÃO, M.T. & J.A.V. COSTA. 2012. Geomorfologia da região centro-norte de Roraima utilizando técnicas de tratamento e

- interpretação de imagens raster da missão subtle radar topography mission (SRTM). *Revista Geonorte* 2: 1484–1496.
- FERRAZ, K.M.P.M.B., S.F.B. FERRAZ, R.C. PAULA, B. BEISIEGEL & C. BREITENMOSER. 2012. Species distribution modeling for conservation purposes. *Natureza & Conservação* 10: 214–220. doi: [10.4322/natcon.2012.032](https://doi.org/10.4322/natcon.2012.032)
- FRANGANILLO, B.P. 1926a. Arácnidos de Cuba: Arácnidos nuevos o poco conocidos de la Isla de Cuba. *Habana*, pp. 1–24. (possibly reprinted from “Cuba Contemporanea 41(161)”)
- FRANGANILLO, B.P. 1926b. Arácnidos nuevos o poco conocidos de la Isla de Cuba. *Boletín de la Sociedad Entomológica de España* 9: 42–68.
- FRANGANILLO, B.P. 1930. Arácnidos de Cuba: Mas arácnidos nuevos de la Isla de Cuba. *Memorias del Instituto Nacional de Investigaciones Científicas* 1: 47–99.
- FRANGANILLO, B.P. 1931. Excursiones arachnológicas, durante el mes de agosto de 1930. *Estudios de “Belen”* 1931(29): 44–49.
- FRANGANILLO, B.P. 1936. Los arácnidos de Cuba hasta 1936. *Cultural*, La Habana, 183 pp.
- FUNK, V.A. & K.S. RICHARDSON. 2002. Systematic data in biodiversity studies: use it or lose it. *Systematic Biology* 51: 303–316. doi: [10.1080/10635150252899789](https://doi.org/10.1080/10635150252899789)
- GALVIS, W. & E.D. FLÓREZ. 2015. New records of the flattie spider genus *Selenops* Latreille, 1819 (Araneae: Selenopidae) from Colombia. *Revista Ibérica de Aracnología* 27: 139–144.
- GERTSCH, W.J. AND L.I. DAVIS. 1942. Report on a collection of spiders from Mexico. IV. *American Museum Novitates* 1158: 1–19.
- GERTSCH, W. J. & S. MULAİK. 1936. New spiders from Texas. *American Museum Novitates* 863: 1–22.
- GLUECK, S. 1994. A taxonomic revision of the orb weaver genus *Acacesia* (Araneae: Araneidae). *Psyche* 101: 59–84.
- GONZAGA, M.O. & A.J. SANTOS. 2004. A new species and a new synonymy in the spiny orb-weaver spider genus *Micrathena* (Araneae, Araneidae). *Journal of Arachnology* 32: 332–335. [10.1636/H03-35](https://doi.org/10.1636/H03-35)
- GONZÁLEZ-SPONGA, M.A. 1998. Arácnidos de Venezuela. Cuatro nuevos géneros y cuatro nuevas especies de la familia Pholcidae Koch, 1850 (Araneae). *Memorias de la Sociedad de Ciencias Naturales La Salle* 57: 17–31.
- GONZÁLEZ-SPONGA, M.A. 2004. Arácnidos de Venezuela. Un nuevo género y nuevas especies de la familia Pholcidae (Araneae). *Aula y Ambiente* 4: 63–76.
- GRAHAM, C.H., C.H. GRAHAM, C. MORITZ, S. FERRIER, F. HUETTMAN & A.T. HUETTMAN. 2004. New developments in museum-based informatics and applications in biodiversity analysis. *Trends in Ecology and Evolution* 19: 497–503.
- GRASSHOFF, M. 1980. Contributions à l'étude de la faune terrestre des îles granitiques de l'archipel des Séchelles (Mission P.L.G. Benoit – J.J. Van Mol 1972). Araneidae-Argiopinae, Araneidae-Araneinae (Araneae). *Revue Zoologique Africaine* 94: 387–409.
- GRASSHOFF, M. 1986. Die Radnetzspinnen-Gattung *Neoscona* in Afrika (Arachnida: Araneae). *Annalen Zoologische Wetenschapen* 250: 1–123.
- Guadanucci, J.P.L. 2014. Theraphosidae phylogeny: relationships of the ‘Ischnocolinae’ genera (Araneae, Mygalomorphae). *Zoologica Scripta* 43(5): 508–518. doi: [10.1111/zsc.12065](https://doi.org/10.1111/zsc.12065)
- Guadanucci, J.P.L., S.M. Lucas, R.P. Indicatti and F.U. Yamamoto. 2007. Description of *Guyruita* gen. nov. and two new species (Ischnocolinae, Theraphosidae). *Revista Brasileira de Zoologia* 24: 991–996. doi: <http://doi.org/b7s9nz>
- HARROD, J.C., H.W. LEVI & L.B. LEIBENSPERGER. 1991. The Neotropical orbweavers of the genus *Larinia* (Araneae: Araneidae). *Psyche*, Cambridge 97: 241–265.
- HASSELT, A.W.M. VAN. 1888. Araneae exoticae quas collegit, pro Museo Lugdunensi, Dr H. Ten Kate Jr. in Guyanâ Hollandicâ (Suriname). *Tijdschrift voor Entomologie* 31: 165–200.
- HENTZ, N.M. 1847. Descriptions and figures of the araneides of the United States. *Boston Journal of Natural History* 5: 443–478.
- HINGSTON, R.W.G. 1932. A naturalist in the Guiana forest. London: Edward Arnold & Co. 384 pp. [Araneae, pp. 363–377].
- HÖFER, H. & A.D. BRESCOVIT. 2000. A revision of the Neotropical spider genus *Ancylometes* Bertkau (Araneae: Pisauridae). *Insect Systematics and Evolution* 31: 323–360. doi: [10.1163/187631200X00075](https://doi.org/10.1163/187631200X00075)
- HÖFER, H. & A.D. BRESCOVIT. 2001. Species and guild structure of a Neotropical spider assemblage (Araneae) (Reserva Florestal Adolpho Ducke, Manaus, Amazonas, Brazil). *Andrias* (Karlsruhe) 15: 99–120.
- HÖFER, H., A.D. BRESCOVIT & T. GASNIER. 1994. The wandering spiders of the genus *Ctenus* (Ctenidae, Araneae) of Reserva Ducke, a rainforest reserve in central Amazonia. *Andrias* 13: 81–98.
- HOLMBERG, E.L. 1876. Arácnidos argentinos. *Anales de Agricultura de la República Argentina* 4: 1–30.
- HUBER, B.A. 2000. New World pholcid spiders (Araneae: Pholcidae): A revision at generic level. *Bulletin of the American Museum of Natural History* 254: 1–348. <http://hdl.handle.net/2246/1601>
- HUBER, B.A. 2004. Evidence for functional segregation in the directionally asymmetric male genitalia of the spider *Metagonia mariguitarensis* (González-Sponga) (Pholcidae: Araneae). *Journal of Zoology, London* 262: 317–326. [10.1017/S0952836903004709](https://doi.org/10.1017/S0952836903004709)
- HUBER, B.A. 2005. Revision and cladistic analysis of the spider genus *Carapoia* González-Sponga (Araneae: Pholcidae), with descriptions of new species from Brazil's Atlantic forest. *Invertebrate Systematics* 19: 541–556. doi: [10.1071/IS05038](https://doi.org/10.1071/IS05038)
- HUBER, B.A. 2011. Revision and cladistic analysis of *Pholcus* and closely related taxa (Araneae, Pholcidae). *Bonner Zoologische Monographien* 58: 1–509. http://pholcidae.de/PDFs/Huber_2011_BZM_low.pdf
- IRIE, T. 2000. A newly recorded spider from Japan, *Micropholcus fauroti* (Simon 1887) (Araneae: Pholcidae). *Acta Arachnologica* 49: 215–217. doi: [10.2476/asjaa.49.215](https://doi.org/10.2476/asjaa.49.215)
- JÄGER, P. & B. PRAXAYSOMBATH. 2011. Spiders from Laos with forty-three new records and first results from the provinces Bolikhamxay and Champasak (Arachnida: Araneae). *Acta Arachnologica* 61: 9–31. doi: [10.2476/asjaa.60.9](https://doi.org/10.2476/asjaa.60.9)
- KEYSERLING, E. 1864. Beschreibungen neuer und wenig bekannter Arten aus der Familie Orbitelae Latr. oder Epeiridae Sund. *Sitzungsberichte und Abhandlungen der Naturwissenschaftlichen Gesellschaft Isis in Dresden* 1863: 63–98, 119–154.
- KEYSERLING, E. 1865. Beiträge zur Kenntniss der Orbitelae Latr. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 15: 799–856.
- KEYSERLING, E. 1877. Ueber amerikanische Spinnenarten der Unterordnung Citigradae. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 26: 609–708.
- KEYSERLING, E. 1881a. Neue Spinnen aus Amerika. III. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 31: 269–314.
- KEYSERLING, E. 1881b. Neue Spinnen aus Amerika. II. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 30: 547–582.
- KEYSERLING, E. 1891. Die Spinnen Amerikas. *Brasilianische Spinnen*. Nürnberg. 3: 1–278.
- KEYSERLING, E. 1892. Die Spinnen Amerikas. *Epeiridae*. Nürnberg. 4: 1–208.
- Keyserling, E. 1893. Die Spinnen Amerikas. *Epeiridae*. Nürnberg. 4: 209–377.
- KOCH, C. L. 1836. Die Arachniden. Nürnberg. 3: 1–104.
- KOCH, C. L. 1838. Die Arachniden. Nürnberg. 4: 109–144; 5: 1–124.
- KOCH, C. L. 1839. Die Arachniden. Nürnberg. 5: 125–158; 6: 1–156; 7: 1–106.
- KOCH, L. 1871. Die Arachniden Australiens, nach der Natur beschrieben und abgebildet. Nürnberg 1: 1–104.
- KRAUS, O. 1955. Spinnen aus El Salvador (Arachnoidea, Araneae).

- Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft 493: 1–112.
- KULCZYŃSKI, W. 1909. Fragmenta Arachnologica. XI–XIII. Bulletin International de l'Academie des Sciences de Cracovie 1909: 427–472.
- KULCZYŃSKI, W. 1911. Symbola ad faunam Araneorum Javae et Sumatrae cognoscendam. II. Sicariidae, Dysderidae, Drassodidae, Zodariidae. Bulletin International de l'Academie des Sciences de Cracovie 1911: 451–496.
- LEHTINEN, P.T. 1967. Classification of the cribellate spiders and some allied families, with notes on the evolution of the suborder Araneomorpha. Annales Zoologici Fennici 4: 199–468.
- LENZ, H. 1886. Beiträge zur Kenntniss der Spinnenfauna Madagascars. Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere 1: 379–408.
- LEVI, H.W. 1963. American spiders of the genera *Audifia*, *Euryopsis* and *Dipoena* (Araneae: Theridiidae). Bulletin of the Museum of Comparative Zoology at Harvard College 129: 121–185.
- LEVI, H.W. 1968. The spider genera *Gea* and *Argiope* in America (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 136: 319–352.
- LEVI, H.W. 1971. The *ravilla* group of the orbweaver genus *Eriophora* in North America (Araneae: Araneidae). Psyche 77: 280–302.
- LEVI, H.W. 1974. The orb-weaver genera *Araniella* and *Nuctenea* (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 146: 291–316.
- LEVI, H.W. 1975. The American orb-weaver genera *Larinia*, *Cercidia* and *Mangora* north of Mexico (Araneae, Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 147: 101–135.
- LEVI, H.W. 1976. The orb-weaver genera *Verrucosa*, *Acanthepeira*, *Wagneriana*, *Acacesia*, *Wixia*, *Scoloderus* and *Alpaida* north of Mexico. Bulletin of the Museum of Comparative Zoology at Harvard College 147: 351–391.
- LEVI, H.W. 1980. The orb-weaver genus *Mecynogea*, the subfamily Metinae and the genera *Pachygnatha*, *Glenognatha* and *Azilia* of the subfamily Tetragnathinae north of Mexico (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 149: 1–74.
- LEVI, H.W. 1985. The spiny orb-weaver genera *Micrathena* and *Chaetacis* (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 150: 429–618.
- LEVI, H.W. 1986. The Neotropical orb-weaver genera *Chrysometa* and *Homalometa* (Araneae: Tetragnathidae). Bulletin of the Museum of Comparative Zoology at Harvard College 151: 91–215.
- LEVI, H.W. 1991a. The Neotropical and Mexican species of the orb-weaver genera *Araneus*, *Dubiepeira*, and *Aculepeira* (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 152: 167–315.
- LEVI, H.W. 1991b. The Neotropical orb-weaver genera *Edricus* and *Wagneriana* (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 152: 363–415.
- LEVI, H.W. 1992. Spiders of the orb-weaver genus *Parawixia* in America (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 153: 1–46.
- LEVI, H.W. 1993. American *Neoscona* and corrections to previous revisions of Neotropical orb-weavers (Araneae: Araneidae). Psyche, Cambridge 99: 221–239.
- LEVI, H.W. 1994. New species of *Bertrana* and *Amazonpeira*, orb-weaving spiders from the Neotropics (Araneae: Araneidae). Transactions of the American Microscopical Society 113: 229–241.
- LEVI, H.W. 1995. The Neotropical orb-weaver genus *Metazygia* (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 154: 63–151.
- LEVI, H.W. 1996. The American orb weavers *Hypognatha*, *Encyosaccus*, *Xylethrus*, *Gasteracantha*, and *Enacrosoma* (Araneae, Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 155: 89–157.
- LEVI, H.W. 1997. The American orb weavers of the genera *Mecynogea*, *Manogea*, *Kapogea* and *Cyrtophora* (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 155: 215–255.
- LEVI, H.W. 2002. Keys to the genera of araneid orbweavers (Araneae, Araneidae) of the Americas. Journal of Arachnology 30: 527–562.
- LEVI, H.W. 2004. Comments and new records for the American genera *Gea* and *Argiope* with the description of new species (Araneae: Araneidae). Bulletin of the Museum of Comparative Zoology at Harvard College 158: 47–65.
- LEWINSOHN, T.M., A.V.L. FREITAS & P.I. PRADO. 2005. Conservation of terrestrial invertebrates and their habitats in Brazil. Conservation Biology 19: 640–645. doi: [10.1111/j.1523-1739.2005.00682.x](https://doi.org/10.1111/j.1523-1739.2005.00682.x)
- LISE, A.A. 1998a. Arachnids of the Ilha de Maracá — Notes on the spiders of the Ilha de Maracá; pp. 377–380, in: W. MILLIKEN & J. RATTER (orgs.). Maracá: The biodiversity and environment of an Amazonian Rainforest. Chichester: John Wiley and Sons.
- LISE, A.A. 1998b. Preliminary list of spiders collected on the Ilha de Maracá; pp. 479–481, in: W. MILLIKEN & J. RATTER (orgs.). Maracá: The biodiversity and environment of an Amazonian Rainforest. Chichester: John Wiley and Sons.
- LUCAS, S. 1964. Sobre a posição sistemática de algumas espécies de aranhas verdadeiras do gênero *Cupiennius*, Simon 1891, da família Ctenidae, em relação ao gênero *Ancylometes*, Bertkau 1880, da família Pisauridae. Memórias do Instituto Butantan 31: 127–134.
- MAGALHÃES, I.L.F. & A.J. SANTOS. 2011. Two new species and taxonomic notes on the Neotropical spiny orb-weaving spiders *Micrathena* and *Chaetacis* (Araneae: Araneidae), with remarks on the development of *Micrathena excavata*. Zootaxa 2983: 39–56.
- MAGALHÃES, I.L.F. & A.J. SANTOS. 2012. Phylogenetic analysis of *Micrathena* and *Chaetacis* spiders (Araneae: Araneidae) reveals multiple origins of extreme sexual size dimorphism and long abdominal spines. Zoological Journal of the Linnean Society 166: 14–53. [10.1111/j.1096-3642.2012.00831.x](https://doi.org/10.1111/j.1096-3642.2012.00831.x)
- MARQUES, M.A.L. & E.H. BUCKUP. 1992. Aranhas Theridiidae da Ilha de Maracá, Roraima, Brasil. IV. Gênero *Thymoites* (Araneae). Iheringia (Série Zoologia) 73: 55–58.
- MARX, G. 1890. Catalogue of the described Araneae of temperate North America. Proceedings of the United States National Museum 12: 497–594.
- MCCOOK, H.C. 1892. *Drexelia*, a new genus of spiders. Proceedings of the Academy of Natural Sciences of Philadelphia 1892: 127.
- MCCOOK, H.C. 1894. American spiders and their spinningwork. Proceedings of the Academy of Natural Sciences of Philadelphia 1893: 1–285.
- MELLO-LEITÃO, C.F. 1917. Notas arachnologicas. 5, Espécies novas ou pouco conhecidas do Brasil. Brotéria (Serie Zool.) 15: 74–102.
- MELLO-LEITÃO, C.F. 1918. Scytodidas e pholcidas do Brasil. Revista do Museu Paulista 10: 83–144.
- MELLO-LEITÃO, C.F. 1921. Um genero e quatro especies novas de aranhas do Brasil. Revista de Ciencias 5: 179–180.
- MELLO-LEITÃO, C.F. 1922a. Novas clubionidas do Brasil. Archivos da Escola Superior de Agricultura e Medicina Veterinaria, Rio de Janeiro 6: 17–56.
- MELLO-LEITÃO, C.F. 1922b. Quelques araignées nouvelles ou peu connues du Bresil. Annales de la Société Entomologique de France 91: 209–228.
- MELLO-LEITÃO, C.F. 1924. Algumas aranhas novas do Brasil. Boletim do Museu Nacional do Rio de Janeiro 1(4): 275–281.
- MELLO-LEITÃO, C.F. 1926. Algumas aranhas do Brasil meridional. Boletim do Museu Nacional do Rio de Janeiro 2: 1–18.
- MELLO-LEITÃO, C.F. 1927. Notes sur quelques araignées bresiliennes de la collection E. Simon. 1. Les palpimanides de l'Amerique du Sud. Bulletin du Muséum National d'Histoire Naturelle de Paris

- 1927: 86–92.
- MELLO-LEITÃO, C.F. 1929a. Aranhas do Pernambuco colhidas por D. Bento Pickel. *Anais da Academia Brasileira de Ciências* 1: 91–112.
- MELLO-LEITÃO, C.F. 1929b. Oxyopideos do Brasil. *Revista do Museu Paulista* 16: 489–536.
- MELLO-LEITÃO, C.F. 1932. Notas sobre ao Micratheneas do Brasil. *Anais da Academia Brasileira de Ciências* 4: 73–97.
- MELLO-LEITÃO, C.F. 1933. Catalogo das aranhas argentinas. *Archivos da Escola Superior de Agricultura e Medicina Veterinaria, Rio de Janeiro* 10: 3–63.
- MELLO-LEITÃO, C.F. 1934. Tres aranhas novas nas collecoes do Instituto Butantan. *Memórias do Instituto Butantan* 8: 401–407.
- MELLO-LEITÃO, C.F. 1936. Contribution à l'etude des Ctenides du Bresil. *Festschrift Embrik Strand* 1: 1–31.
- MELLO-LEITÃO, C.F. 1939a. Algumas aranhas de S.-Paulo e Santa Catarina. *Memórias do Instituto Butantan* 12: 523–531.
- MELLO-LEITÃO, C.F. 1939b. Araignées américaines du Musee d'histoire naturelle de Bâle. *Revue Suisse de Zoologie* 46: 43–93.
- MELLO-LEITÃO, C.F. 1939c. Some new argiopid spiders of British Guiana taken by Mr C. W. Richards from the nests of solitary wasps. *Anais da Academia Brasileira de Ciências* 11: 105–112.
- MELLO-LEITÃO, C.F. 1940a. Arañas de la provincia de Buenos Aires y de las gobernaciones de La Pampa, Neuquén, Río Negro y Chubut. *Revista del Museo de La Plata* 2: 3–62.
- MELLO-LEITÃO, C.F. 1940b. Aranhas do Espírito Santo coligidas por Mario Rosa, em 1936 e 1937. *Arquivos de Zoologia do Estado de Sao Paulo* 2: 199–214.
- MELLO-LEITÃO, C.F. 1941a. Notas sobre a sistemática das aranhas com descrição de algumas novas especies Sul Ameicanas. *Anais da Academia Brasileira de Ciências* 13: 103–127.
- MELLO-LEITÃO, C.F. 1941b. Catalogo das aranhas da Colombia. *Anais da Academia Brasileira de Ciências* 13: 233–300.
- MELLO-LEITÃO, C.F. 1941c. Las arañas de Córdoba, La Rioja, Catamarca, Tucumán, Salta y Jujuy colectadas por los Profesores Birabén. *Revista del Museo de La Plata (N.S., Zool.)* 2: 99–198.
- MELLO-LEITÃO, C.F. 1942a. Arañas del Chaco y Santiago del Estero. *Revista del Museo de La Plata* 2: 381–426.
- MELLO-LEITÃO, C.F. 1942b. Cinco aranhas novas do Perú. *Revista Brasileira de Biologia* 2: 429–434.
- MELLO-LEITÃO, C.F. 1943a. Aranhas do Chile coligidas pelo Dr. J. C. Carvalho. *Revista Brasileira de Biologia* 3: 403–409.
- MELLO-LEITÃO, C.F. 1943b. Catálogo das aranhas do Rio Grande do Sul. *Arquivos do Museu Nacional do Rio de Janeiro* 37: 147–245.
- MELLO-LEITÃO, C.F. 1944. Arañas de la provincia de Buenos Aires. *Revista del Museo de La Plata* 3: 311–393.
- MELLO-LEITÃO, C.F. 1945. Arañas de Misiones, Corrientes y Entre Ríos. *Revista del Museo de La Plata* 4: 213–302.
- MELLO-LEITÃO, C.F. 1946. Notas sobre os Filistatidae e Pholcidae. *Anais da Academia Brasileira de Ciências* 18: 39–83.
- MELLO-LEITÃO, C.F. 1947. Aranhas de Carmo do Rio Claro (Minas Gerais) coligidas pelo naturalista José C. M. Carvalho. *Boletim do Museu Nacional do Rio de Janeiro* 80: 1–34.
- MELLO-LEITÃO, C.F. 1948. Contribuição ao conhecimento da fauna araneológica das Guianas. *Anais da Academia Brasileira de Ciências* 20: 151–196.
- MELLO-LEITÃO, C.F. 1949. Aranhas da Foz do Kuluene. *Boletim do Museu Nacional do Rio de Janeiro* 92: 1–19.
- MENDOZA, J.I.M. 2014. *Psalmopoeus victori*, the first arboreal theraphosid spider described for Mexico (Araneae: Theraphosidae: Aviculariinae). *Revista Mexicana de Biodiversidad* 85: 728–735. [10.7550/rmb.44597](https://doi.org/10.7550/rmb.44597)
- MERRETT, P. 1988. Notes on the biology of the neotropical pisaurid, *Ancylometes bogotensis* (Keyserling) (Araneae: Pisauridae). *Bulletin of the British Arachnological Society* 7: 197–201.
- MILLOT, J. 1941. Les araignées de l'Afrique Occidentale Française – sicariides et pholcides. *Mémoires de l'Académie des Sciences de l'Institut de France* 64: 1–53.
- MILLOT, J. 1946. Les pholcides de Madagascar (Aranéides). *Mémoires du Muséum National d'Histoire Naturelle de Paris (N.S.)* 22: 127–158.
- MILLOT, J. 1946. Les Scytodes d'Afrique Noire française (Araneae). *Revue Française d'Entomologie* 13: 156–168.
- NAKA, L.N., M. COHN-HAFT, F. MALLETT-RODRIGUES, M.P.D. SANTOS & M.F. TORRES. 2006. The avifauna of the Brazilian state of Roraima: bird distribution and biogeography in the Rio Branco basin. *Revista Brasileira de Ornitologia* 14: 197–238. http://www4.museu-goeldi.br/revistabornito/revista/index.php/BJO/article/view/2502/pdf_369
- OLIVEIRA, U., A.D. BRESCOVIT & A.J. SANTOS. 2015. Delimiting Areas of Endemism through Kernel Interpolation. *Plos One* 10: e0116673-18. doi: [10.1371/journal.pone.0116673](https://doi.org/10.1371/journal.pone.0116673)
- Oliveira, U., A.P. Paglia, A.D. Brescovit, C.J.B. Carvalho, D.P. Silva, et al. 2016. The strong influence of collection bias on biodiversity knowledge shortfalls of Brazilian terrestrial biodiversity. *Diversity and Distributions*: 1–13. [10.1111/ddi.12489](https://doi.org/10.1111/ddi.12489)
- OLIVIER, G.A. 1789. Araignée, Aranea. *Encyclopédie Méthodique, Histoire Naturelle, Insectes, Paris* 4: 173–240.
- ONO, H. 2011. Spiders (Arachnida, Araneae) of the Ogasawara Islands, Japan. *Memoirs of the National Museum of Nature and Science Tokyo* 47: 435–470.
- OPELL, B.D. 1979. Revision of the genera and tropical American species of the spider family Uloboridae. *Bulletin of the Museum of Comparative Zoology at Harvard College* 148: 443–549.
- PAULA, F.S., R. GABRIEL, R.P. INDICATTI, A.D. BRESCOVIT & S.M. LUCAS. 2014. On the Brazilian Amazonian species of *Acanthoscuria* (Araneae: Theraphosidae). *Zoologia* 31: 63–80. doi: [10.1590/S1984-46702014000100008](https://doi.org/10.1590/S1984-46702014000100008)
- PERTY, M. 1833. Arachnides Brasilienses. In: J.B. DE SPIX AND F.P. MARTIUS (eds.). *Delectus animalium articulorum quae in itinere per Braziliam ann. 1817 et 1820 colligerunt. Monachii*, pp. 191–209, pls. 38–39.
- PETERS, H.-J. 2000. Tarantulas of the world: Kleiner Atlas der Vogelspinnen – Band 2. Wegberg: published by the author, 162 pp.
- PETERS, H.-J. 2003. Tarantulas of the world: Amerika's Vogelspinnen. Wegberg: published by the author. 328 pp.
- PETRUNKOVITCH, A. 1910. Some new or little known American Spiders. *Annals of the New York Academy of Science* 19: 205–224.
- PETRUNKOVITCH, A. 1911. A synonymic index–catalogue of spiders of North, Central and South America with all adjacent islands, Greenland, Bermuda, West Indies, Terra del Fuego, Galapagos, etc. *Bulletin of the American Museum of Natural History* 29: 1–791.
- PETRUNKOVITCH, A. 1929. The spiders of Porto Rico. Part I. *Transactions of the Connecticut Academy of Arts and Sciences* 30: 1–158.
- PETRUNKOVITCH, A. 1930. The spiders of Porto Rico. Part II. *Transactions of the Connecticut Academy of Arts and Sciences* 30: 159–356.
- PIACENTINI, L.N. 2011. Three new species and new records in the wolf spider subfamily Sosippinae from Argentina (Araneae: Lycosidae). *Zootaxa* 3018: 27–49. doi: [10.11646/zootaxa.3790.1.1](https://doi.org/10.11646/zootaxa.3790.1.1)
- PICKARD-CAMBRIDGE, F.O. 1896. On the Theraphosidae of the lower Amazons: being an account of the new genera and species of this group of spiders discovered during the expedition of the steamship “Faraday” up the river Amazons. *Proceedings of the Zoological Society of London* 1896: 716–766.
- PICKARD-CAMBRIDGE, F.O. 1897. On cteniform spiders from the lower Amazons and other regions of North and South America, with a list of all known species of these groups hitherto recorded from the New World. *Annals and Magazine of Natural History* (6) 19: 52–106.
- PICKARD-CAMBRIDGE, F.O. 1899a. On new species of spiders from Trinidad, West Indies. *Proceedings of the Zoological Society of London* 1898: 890–900. [10.1111/j.1096-3642.1898.tb03191.x](https://doi.org/10.1111/j.1096-3642.1898.tb03191.x)

- PICKARD-CAMBRIDGE, F.O. 1899b. Arachnida – Araneida and Opiliones. In: *Biologia Centrali-Americana, Zoology*. London 2: 41–88.
- PICKARD-CAMBRIDGE, F.O. 1901. Arachnida – Araneida and Opiliones. In: *Biologia Centrali-Americana, Zoology*. London 2: 193–312.
- PICKARD-CAMBRIDGE, F.O. 1902a. Arachnida – Araneida and Opiliones. In: *Biologia Centrali-Americana, Zoology*. London 2: 313–424.
- PICKARD-CAMBRIDGE, F.O. 1902b. New species of spiders belonging to the genus *Ctenus*, with supplementary notes. *Annals and Magazine of Natural History* (7) 9: 401–415.
- PICKARD-CAMBRIDGE, F.O. 1903. Arachnida – Araneida and Opiliones. In: *Biologia Centrali-Americana, Zoology*. London 2, 425–464.
- PICKARD-CAMBRIDGE, F.O. 1904. Arachnida – Araneida and Opiliones. In: *Biologia Centrali-Americana, Zoology*. London 2: 465–560.
- PICKARD-CAMBRIDGE, O. 1877. On some new species of Araneida, with characters of two new genera and some remarks on the families Podophthalmides and Dinopides. *Proceedings of the Zoological Society of London* 45(3): 557–578, Pl. LVI–LVII.
- PICKARD-CAMBRIDGE, O. 1889. Arachnida. Araneida. In: *Biologia Centrali-Americana, Zoology*. London 1: 1–56.
- PICKARD-CAMBRIDGE, O. 1893. Arachnida. Araneida. In: *Biologia Centrali-Americana, Zoology*. London 1: 105–120.
- PIZA JR., S. DE T. 1938. Duas novas aranhas oxyópidas do Brasil. *Boletim Biológico Sao Paulo* 3: 47–48.
- PLATNICK, N.I. 1975. A revision of the palpimanid spiders of the new subfamily Otiiothopinae (Araneae, Palpimanidae). *American Museum Novitates* 2562: 1–32. <http://hdl.handle.net/2246/2755>
- PLATNICK, N.I. 1986. A revision of the spider genus *Trochanteria* (Araneae: Gnaphosoidea). *Bulletin of the British Arachnological Society* 7: 29–33.
- POLOTOW, D. & A.D. BRESCOVIT. 2008. Revision of the Neotropical spider genus *Gephyroctenus* (Araneae: Ctenidae: Calocteninae). *Revista Brasileira de Zoologia* 25: 705–715. doi: [10.1590/S0101-81752008000400016](https://doi.org/10.1590/S0101-81752008000400016)
- POLOTOW, D. & A.D. BRESCOVIT. 2009. Revision and cladistic analysis of *Isoctenus* and description of a new Neotropical genus (Araneae, Ctenidae, Cteninae). *Zoological Journal of the Linnean Society* 155: 583–614. doi: [10.1111/j.1096-3642.2008.00452.x](https://doi.org/10.1111/j.1096-3642.2008.00452.x)
- POLOTOW, D. & A.D. BRESCOVIT. 2014. Phylogenetic analysis of the tropical wolf spider subfamily Cteninae (Arachnida, Araneae, Ctenidae). *Zoological Journal of the Linnean Society* 170: 333–361. [10.1111/zoj.12101T](https://doi.org/10.1111/zoj.12101T)
- POSSINGHAM, H.P., H. GRANTHAM & C. RONDININI. 2007. How can you conserve species that haven't been found? *Journal of Biogeography* 34: 758–759. doi: [10.1111/j.1365-2699.2007.01717.x](https://doi.org/10.1111/j.1365-2699.2007.01717.x)
- REGO, F.N.A.A., E.M. VENTICINQUE, A.D. BRESCOVIT, C.A. RHEIMS & A.L.M.K. ALBERNAZ. 2009. A contribution to the knowledge of the spider fauna (Arachnida: Araneae) of the floodplain forests of the main Amazon River channel. *Revista de Aracnologia* 17: 85–96.
- REIMOSER, E. 1917. Die Spinnengattung *Micrathena* Sundevall. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 67: 73–160.
- RHEIMS, C.A. & A.D. BRESCOVIT. 2004. Revision and cladistic analysis the family Hersiliidae (Arachnida, Araneae) with emphasis on Neotropical and Nearctic species. *Insect Systematics and Evolution* 35: 189–239.
- RHEIMS, C.A., F.M. LABARQUE & M.J. RAMÍREZ. 2008. The South American genus *Quemedice* Mello-Leitão (Araneae: Sparassidae): familial placement and taxonomic revision. *Zootaxa* 1813: 60–68.
- RICETTI, J. & A.B. BONALDO. 2008. Diversidade e estimativas de riqueza de aranhas em quatro fitofisionomias na Serra do Cachimbo, Pará, Brasil. *Theringia, Série Zoologia* 98: 88–99. doi: [10.1590/S0073-47212008000100013](https://doi.org/10.1590/S0073-47212008000100013)
- ROBERTS, M.J. 1983. Spiders of the families Theridiidae, Tetragnathidae and Araneidae (Arachnida: Araneae) from Aldabra atoll. *Zoological Journal of the Linnean Society* 77: 217–291.
- RODRIGUES, E.N.L. 2013. Six new species, complementary descriptions and new records from the Neotropical Region of the spider genus *Dipoena* (Araneae: Theridiidae). *Zootaxa* 3750: 1–25. doi: [10.11646/zootaxa.3750.1.1](https://doi.org/10.11646/zootaxa.3750.1.1)
- ROEWER, C.F. 1942. *Katalog der Araneae von 1758 bis 1940*. Bremen. 1: 1–1040.
- ROEWER, C.F. 1960. Araneae Lycosaeformia II (Lycosidae) (Fortsetzung und Schluss. Exploration du Parc National de l'Upemba, Mission G. F. de Witte 55: 519–1040.
- RUDLOFF, J.-P. AND D. WEINMANN. 2010. A new giant tarantula from Guyana. *Arthropoda Scientia* 1: 21–40.
- RUIZ, G.R.S. & A.D. BRESCOVIT. 2013. Revision of *Breda* and proposal of a new genus (Araneae: Salticidae). *Zootaxa* 3664: 401–433. doi: [10.11646/zootaxa.3664.4.1](https://doi.org/10.11646/zootaxa.3664.4.1)
- SAAGER, F. 1994. *Psalmopoeus irminia* sp. n., Beschreibung einer neuen Aviculariinae (Theraphosidae, Aviculariinae, genus *Psalmopoeus*) inclusive einem Vergleich mit *Psalmopoeus cambridgei*. *Arthropoda* 2: 59–71.
- SAARISTO, M.I. 2001. Pholcid spiders of the granitic Seychelles (Araneae, Pholcidae). *Phelsuma* 9: 9–28.
- SAARISTO, M.I. 2010. Araneae; pp. 8–306, in: J. GERLACH & Y. MARUSIK (eds.). *Arachnida and Myriapoda of the Seychelles islands*. Manchester: Siri Scientific Press.
- SANTOS, A. J. & A.D. BRESCOVIT. 2003. A revision of the Neotropical species of the lynx spider genus *Peucetia* Thorell 1869 (Araneae: Oxyopidae). *Insect Systematics and Evolution* 34: 95–116.
- SANTOS, A.J. 2007. A revision of the Neotropical nursery-web spider genus *Architis* (Araneae: Pisauridae). *Zootaxa* 1578: 1–40.
- SANTOS, A.J. & A.A. NOGUEIRA. 2008. Three new species, new records and notes on the nursery-web spider genus *Architis* in Brazil (Araneae: Pisauridae). *Zootaxa* 1815: 51–61.
- SANTOS, A.J. & A.D. BRESCOVIT. 2001. A revision of the South American spider genus *Aglaoctenus* Tullgren, 1905 (Araneae, Lycosidae, Sosippinae). *Andrias* 15: 75–90.
- SANTOS, A.J. & A.D. BRESCOVIT. 2003. A revision of the Neotropical species of the lynx spider genus *Peucetia* Thorell 1869 (Araneae: Oxyopidae). *Insect Systematics and Evolution* 34: 95–116. doi: [10.1163/187631203788964863](https://doi.org/10.1163/187631203788964863)
- SANTOS, A.J. & C.A. RHEIMS. 2005. Four new species and new records for the spider genus *Synotaxus* Simon, 1895 (Araneae: Synotaxidae) from Brazil. *Zootaxa* 937: 1–12.
- SANTOS, A.J. & M.O. GONZAGA. 2003. On the spider genus *Oecobius* Lucas, 1846 in South America (Araneae, Oecobiidae). *Journal of Natural History* 37: 239–252. doi: [10.1080/713834668](https://doi.org/10.1080/713834668)
- SANTOS, A.J., M.O. GONZAGA & G. HORMIGA. 2009. Notes on two problematic eastern Asian species of the spider genus *Oecobius* (Araneae, Oecobiidae, Linyphiidae). *Journal of Arachnology* 37: 101–102. doi: [10.1636/A08-23.1](https://doi.org/10.1636/A08-23.1)
- SARMIENTO, G. & M. MONASTERIO. 1983. Life forms and phenology; pp. 79–108, in: F. BOURLIÈRE (ed.) *Ecosystems of the world. XIII, tropical savannas*. Amsterdam: Elsevier.
- SCHENKEL, E. 1953. Bericht über einige Spinnentiere aus Venezuela. *Verhandlungen der Naturforschenden Gesellschaft in Basel* 64: 1–57.
- SCHIAPPELLI, R.D. & P.B.S. GERSCHMAN. 1970. Consideraciones sobre el género *Ancylometes* Bertkau 1880 (Araneae: Pisauridae). *Acta Zoologica Lilloana* 27: 155–179.
- SCHIAPPELLI, R.D. & P.B.S. GERSCHMAN. 1973. Diagnosis de *Phoneutria reidyi* (F. O. Pickard-Cambridge, 1897) y de *Phoneutria boliviensis* (F. O. Pickard-Cambridge, 1897) (Araneae, Ctenidae). *Revista de la Sociedad Entomológica Argentina* 34: 31–38.

- SCHMIDT, G. 2003. Die Vogelspinnen: Eine weltweite Übersicht. Neue Brehm-Bücherei, Hohenwarsleben, 383 pp.
- SCHMIDT, G., M. BULLMER & M. THIERER-LUTZ. 2006. Eine neue *Psalmopoeus*-Art aus Venezuela, *Psalmopoeus langensbucheri* sp. n. (Araneae: Theraphosidae: Aviculariinae). *Tarantulas of the World* 121/122: 3–17.
- ŠESTÁKOVÁ, A., L. ČERNECKÁ, J. NEUMANN & N. REISER. 2014. First record of the exotic spitting spider *Scytodes fusca* (Araneae, Scytodidae) in Central Europe from Germany and Slovakia. *Arachnologische Mitteilungen* 47: 1–6. doi: 10.5431/aramit4701
- SHEAR, W.A. 1970. The spider family Oecobiidae in North America, Mexico, and the West Indies. *Bulletin of the Museum of Comparative Zoology at Harvard College* 140: 129–164.
- SIERWALD, P. 1989. Morphology and ontogeny of female copulatory organs in American Pisauridae, with special reference to homologous features (Arachnida: Araneae). *Smithsonian Contributions to Zoology* 484: 1–24.
- SILVA, E.L.C. & J.E. CARICO. 2012. Revision of the Neotropical nursery-web spider genus *Thaumasia* Perty, 1833 (Araneae: Lycosoidea: Pisauridae: Thaumasiinae). *Zootaxa* 3567: 1–64.
- SIMON, E. 1886. Etudes arachnologiques. 18e Mémoire. XXVI. Matériaux pour servir à la faune des Arachnides du Sénégal. (Suivi d'une appendice intitulé: descriptions de plusieurs espèces africaines nouvelles. *Annales de la Société Entomologique de France* (6) 5: 345–396.
- SIMON, E. 1887. Arachnides recueillis à Obock en 1886, par M. le Dr L. Faurot. *Bulletin de la Société Zoologique de France* 12: 452–455.
- SIMON, E. 1889. Arachnides. In: *Voyage de M. E. Simon au Venezuela (décembre 1887–avril 1888. 4e Mémoire. Annales de la Société Entomologique de France* 6 9, 169–220.
- SIMON, E. 1891. On the spiders of the island of St. Vincent. Part 1. *Proceedings of the Zoological Society of London* 1891: 549–575.
- SIMON, E. 1892. Histoire naturelle des araignées. Paris. 1: 1–256.
- SIMON, E. 1893a. Arachnides; pp. 423–462, in: *Voyage de M.E. Simon au Venezuela (décembre 1887 – avril 1888. 21e Mémoire. Annales de la Société Entomologique de France* 61.
- SIMON, E. 1893b. Études arachnologiques. 25e Mémoire. XL. Descriptions d'espèces et de genres nouveaux de l'ordre des Araneae. *Annales de la Société Entomologique de France* 62: 299–330.
- SIMON, E. 1895a. Histoire naturelle des araignées. Paris 1, 761–1084.
- SIMON, E. 1895b. Etudes arachnologiques. 26e. XLI. Descriptions d'espèces et de genres nouveaux de l'ordre des Araneae. *Annales de la Société Entomologique de France* 64: 131–160.
- SIMON, E. 1896. Étude sur les Arachnides du Chili. Premier mémoire. *Actes de la Société Scientifique du Chili* 6: 63–70, civ–cvii.
- SIMON, E. 1897. Liste de arachides recueillis aux îles du Cap Vert, dans la République Argentine et le Paraguay et descriptions d'espèces nouvelles. In: *Viaggio del Dott. A. Borelli nella Repubblica Argentina e nel Paraguay. Bollettino dei Musei di Zoologia ed Anatomia Comparata della Reale Università di Torino* 12(270): 1–8.
- SIMON, E. 1898a. Descriptions d'arachnides nouveaux des familles des Agelenidae, Pisauridae, Lycosidae et Oxyopidae. *Annales de la Société Entomologique de Belgique* 42: 5–34.
- SIMON, E. 1898b. Descriptions d'arachnides nouveaux des familles des Agelenidae, Pisauridae, Lycosidae et Oxyopidae. *Annales de la Société Entomologique de Belgique* 42: 5–34.
- SIMON, E. 1898c. Histoire naturelle des araignées. Paris. 2: 193–380.
- SOARES, B.A.M. & H.F.A. CAMARGO. 1948. Aranhas coligadas pela Fundação Brasil-Central (Arachnida-Araneae). *Boletim do Museu Paraense Emílio Goeldi* 10: 355–409.
- SOARES, B.A.M. & H.F.A. CAMARGO. 1955. Algumas novas espécies de aranhas brasileiras (Araneae, Anyphaenidae, Argiopidae, Eusparassidae, Theridiidae). *Arquivos do Museu Nacional do Rio de Janeiro* 42: 577–580.
- STRAND, E. 1907a. Eine neue Avicularia nebst Bemerkungen über andere südamerikanische Spinnen. *Jahrbücher des Nassauischen Vereins für Naturkunde* 60: 220–227.
- STRAND, E. 1907b. Über drei Clubioniden und eine Pisauride vom Sorata in den Cordilleren (Günther leg.), *Museum Lübeck. Zeitschrift für Naturwissenschaften* 79: 422–431.
- STRAND, E. 1909a. Neue oder wenig bekannte neotropische cteniforme Spinnen des Berliner Museums. *Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere* 28: 401–428.
- STRAND, E. 1909b. Neue oder wenig bekannte *Lycotenus*-Arten des Berliner Museums. *Zoologischer Anzeiger* 34: 329–337.
- STRAND, E. 1909c. Neue oder wenig bekannte südamerikanische *Cupiennius*- und *Ctenus*-Arten. *Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere* 28: 293–328.
- STRAND, E. 1916a. *Arachnologica varia*, I–IX. *Archiv für Naturgeschichte* 81(A11): 112–123.
- STRAND, E. 1916b. Systematische-faunistische Studien über paläarktische, afrikanische und amerikanische Spinnen des Senckenbergischen Museums. *Archiv für Naturgeschichte* 81: 1–153.
- TACZANOWSKI, L. 1872. Les aranéides de la Guyane française. *Horae Societatis Entomologicae Rossicae* 9: 64–112.
- TACZANOWSKI, L. 1873. Les aranéides de la Guyane française. *Horae Societatis Entomologicae Rossicae* 9: 113–150, 261–286.
- TACZANOWSKI, L. 1874. Les aranéides de la Guyane française. *Horae Societatis Entomologicae Rossicae* 10: 56–115.
- TACZANOWSKI, L. 1878. Les Aranéides du Pérou central. *Horae Societatis Entomologicae Rossicae* 14: 140–175.
- TACZANOWSKI, L. 1879. Les aranéides du Pérou central (suite). *Horae Societatis Entomologicae Rossicae* 15: 102–136.
- TAUCARE-RÍOS, A.O. 2012. Notas acerca de la ecología de *Argiope argentata* (Fabricius, 1775) (Araneidae) en Chile. *Boletín de Biodiversidad de Chile* 7: 42–47.
- THORELL, T. 1891. Spindlar från Nikobarerna och andra delar af södra Asien. *Bihang till Kongliga Svenska Vetenskaps-Akademiens Handlingar* 24(2): 1–149.
- THORELL, T. 1895. *Descriptive catalogue of the spiders of Burma*. London, pp. 1–406.
- TULLGREN, A. 1905. Aranea from the Swedish expedition through the Gran Chaco and the Cordilleras. *Arkiv för Zoologi* 219: 1–81.
- VILAR, J.B., L.S. DE-ANDRADE, K.R. LEITE, H.D. FERREIRA & L.C. CHEN. 2009. Assessment of genotoxicity and cytotoxicity of “lixreira” (*Curatella americana* L.) using the prophage lambda induction test (SOS inductest). *Brazilian Journal of Pharmacological Science* 45: 491–496. doi: 10.1590/S1984-82502009000300015
- VINSON, A. 1863. Aranéides des îles de la Réunion, Maurice et Madagascar. Paris, i–cxx, 1–337.
- WALCKENAER, C.A. 1805. Tableau des aranéides ou caractères essentiels des tribus, genres, familles et races que renferme le genre Aranea de Linné, avec la désignation des espèces comprises dans chacune de ces divisions. Paris. 88 pp.
- WALCKENAER, C.A. 1837. Histoire naturelle des insectes. Aptères. Paris. 1: 1–682.
- WALCKENAER, C.A. 1841. Histoire naturelle des insectes. Aptères. Paris. 2: 1–549.
- WORLD SPIDER CATALOG. 2017. World spider catalog, version 17.5. Bern: Natural History Museum. Accessed at <http://wsc.nmbe.ch>, 13 January 2017.
- WUNDERLICH, J. 1988. Die Fossilen Spinnen im Dominikanischen Bernstein. *Straubenhardt: published by the author*. 378 pp.

Authors' contributions: LSC, MCS and JJCC collected the data; LSC, PHM and JJCC identified the specimens; LSC, PHM, MCS and JJCC wrote the text.

Received: 2 April 2016

Accepted: 2 September 2016

Academic editor: Regiane Saturnino