



Floristic survey of the Brazilian Ages Memorial: a Cerrado *sensu stricto* area with an educational relevance

Thaís N. C. Vasconcelos^{1,2,3*}, Juliana S. Silva¹, Marcelo L. Ianhez¹ and Carolyn E. B. Proença¹

1 Departamento de Botânica, Instituto de Ciências Biológicas, Universidade de Brasília, Brasília, Distrito Federal, Brazil CEP 70919-970

2 Research Department of Genetics, Evolution and Environment, University College London, Gower Street, London WC1E6BT United Kingdom

3 Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, United Kingdom

Corresponding author. E-mail: T.vasconcelos@kew.org

Abstract: The Cerrado Biome is currently classified as one of the most diverse savannas in the world. The anthropic destruction of this biome led the Cerrado to be recognized as one of the world's hot spots for bioconservation. Cerrado *sensu stricto* phytophysiology represents 70% of the original Cerrado biome surface and floristic surveys are basic and important studies for conservation initiatives in these areas. Our survey area is a Cerrado *sensu stricto* of 6 ha attached to the Brazilian Ages Memorial, an open air museum located in Distrito Federal, in the center of the Cerrado biome. Our list counted 222 species of angiosperms, with the families Asteraceae, Fabaceae, Malpighiaceae and Myrtaceae being the richest ones. Our work contributes towards a better understanding of the Cerrado biome and also may help future educational programs in the museum.

Key words: Distrito Federal, museum, cerrado

INTRODUCTION

The Cerrado biome in Brazil is currently classified among the savannas with highest biodiversity in the world, with an estimate in excess of 12,000 angiosperms species (Mendonça et al. 2008). Furthermore, it is also significant in terms of its large surface area, having originally covered 21% of Brazilian territory (Aguar et al. 2004) and for including parts of the three largest hydrological systems of South America in its area (Alho and Martins 1995; Klink 1996).

One of the most remarkable characteristics of the Cerrado vegetation is the great variety of phytophysiology, which are represented by a diversity of forests, savannas and meadows (Ribeiro and Walter 2008). Among the savannas phytophysiology, the Cerrado *sensu stricto* is the most common, occurring in almost 70% of the biome (Eiten 1972).

According to Ribeiro and Walter (2008), the main

characteristics of Cerrado *sensu stricto* are a defined layer of trees and shrubs, covering between 10% to 60%, and an herbaceous layer approximately 60 cm high. The high species number of this phytophysiology is represented by a great diversity of herbs, subshrubs and shrubs mainly represented by the families Fabaceae, Rubiaceae, Myrtaceae and Asteraceae (Ratter 1997). The destruction of Cerrado *sensu stricto* areas, which has seen more than 50% of its area lost or degraded, has resulted in the Brazilian cerrado being recognized as one of the world hotspots for bioconservation (Felfili and Silva Junior 2001).

The Distrito Federal of Brazil (DF) is located in the center of the Cerrado biome (Figure 1) and has suffered from increasing exploitation of its natural resources.




Figure 1. The original area of Cerrado Biome in Brazil (based on Ratter et al. 1997) and the study site location within the biome. DF: Distrito Federal, GO: Goiás state.

After 44 years of occupation, 73.8% of its original Cerrado has already been lost (Felfili 2000). Conservation areas represent 42% of the DF area, but many of these have already been invaded by illegal occupation which has led to biodiversity loss and to the contamination and siltation of rivers (Felfili 2000).

In general, the Cerrado fragmentation has reinforced the importance of creating new conservation areas and, particularly, in preserving the areas that already exist. Knowledge of the Cerrado flora in the Distrito Federal is a significant step in planning and selecting new representative areas of the biome that will be prioritized for conservation and recovery (Felfili et al. 1993; Mendonça et al. 2008). Therefore, works like this are very useful towards a better understanding of areas where limited information is available. In addition to that, floristic surveys are a tool to improve herbarium collections, which then will constitute the basis of studies in taxonomy, biogeography and ecology.

MATERIALS AND METHODS

Study site

The Brazilian Ages Memorial (in Portuguese “Memorial das Idades do Brasil”) is a privately owned open air museum located at 15°77' S, 047°80' W, on the boundary of Paranoá city, Distrito Federal, Brazil. The area is a memorial about the ancient people who inhabited the region of the central Brazilian highlands and is frequently visited by school groups.

According to the official website (<http://www.paulobertran.com.br>), the area was given this name because it represents the “three ages” of the central highlands formation. The first age is the “geological age”, represented by the rocks and stones, some of them up to 1 billion years old and originally under the Pangean sea. The second is the “anthropological age”, represented by reproductions of ancient cave paintings made by local artists (Figure 2). The third age is the “plant age”, represented by the trails through the area of Cerrado *sensu stricto* in which the survey was done.



Figure 2. Cave painting reproductions from the site.



Figure 3. Aerial view of the Brazilian Ages Memorial and region on Google Earth™. a) Area in which our floristic survey was done; b) Brazilian Ages Memorial museum; c) *Pinus* sp. plantation.; d) Paranoá Lake.

The site of study is a Cerrado *sensu stricto* located at an altitude of 1,050 m with approximately 6 ha attached to the Brazilian Ages Memorial Museum. The site is close to a large plantation of *Pinus* sp. which has already started to invade the area’s native vegetation via seed dispersal (Figure 3).

Data collection

Our species list was mostly defined by weekly collections over a period of a year, from May 2009 to May 2010. Collections made by the Ethnobotany course of the University of Brasilia, conducted by Prof. Carolyn E. B. Proença in 2009, were also included in our survey. Some sporadic collections made in the following years were also considered. The collections were made mainly next to the trails and fertile plants with buds, flowers and fruits were prioritized. Some characteristics of the plants, such as color, scent and latex were also recorded and are available in BRAHMS at the University of Brasilia Herbarium (UB).

The species identification was mostly made by comparison with dry specimens in the UB herbarium. In some cases, species were also identified using specialized keys, literature, and consulting taxonomists. The collected material was incorporated into the UB collection.

Synonyms follow the classification listed on the Flora do Brasil website (Lista de Espécies da Flora do Brasil, consulted in 2015), for it presents the most up to date classification of plants in Brazil. The family classification system follows the Angiosperm Phylogeny Group III (APG III 2009). Classification of the species as native or invasive was the same as used by Mendonça et al. (2008). Information of habit (i.e., shrub, tree) was also taken from the “Espécies da Flora do Brasil” website (consulted in 2015) and from Mendonça et al. (2008).

RESULTS

Our floristic survey counted 222 species of angiosperms, representing 148 genera and 59 families,




Figure 4. Families with the number of species in the Brazilian Ages Memorial. Aca = Acanthaceae; Ast = Asteraceae; Fab = Fabaceae; Lam = Lamiaceae; Lyt = Lythraceae; Mal = Malpighiaceae; Mel = Melastomataceae; Mlv = Malvaceae; Myr = Myrtaceae; Poa = Poaceae; Rub = Rubiaceae; Sol = Solanaceae.

of which 50 are eudicots, 8 are monocots, and one is a magnoliid (Annonaceae). The complete species list, with collection number, habit, category (if endemic or invasive) is presented in Table 1.

Family with the largest species number in the area was Fabaceae (represented by 32 species), followed by Asteraceae (27 species), Malpighiaceae (16 species), Myrtaceae (13 species), Rubiaceae (9 species), Melastomataceae and Lamiaceae (8 species each), Acanthaceae (7 species), Poaceae and Malvaceae (6 species each), Lythraceae and Solanaceae (5 species each) (Figure 4).

Among the genera, the richest in species number were *Banisteriopsis* (Malpighiaceae) with seven species, *Chamaecrista* (Fabaceae), *Eugenia* (Myrtaceae) and *Miconia* (Melastomataceae) with six species each, *Byrsonima* (Malpighiaceae), *Hyptis* (Lamiaceae) and *Myrcia* (Myrtaceae) with five species each, and *Mimosa* (Fabaceae) represented by four species.

DISCUSSION

The high diversity of Asteraceae and Fabaceae species was expected since these are among those families considered hyperdiverse in Brazil (Rapini et al. 2009). They also appear as the two most diverse species of eudicots in other surveys in similar areas of Cerrado (Carvalho et al. 2010; Ishara and Rodela 2012).

The families Poaceae and Cyperaceae usually are represented by a high species number in Cerrado *sensu stricto* environments (Ratter et al. 1997). In our study, however, we found only three species of Cyperaceae and six of Poaceae, a number abnormally low. This result can be explained by either (1) the families were poorly sampled or (2) because the site is small and well-cared for, there was no sign of recent fire in the area. Amaral et

al. 2013, by analyzing an area of Cerrado also in Distrito Federal, found that the absence of fire in Cerrado leads to an increasing in the number of woody species and a decreasing in the number of herbaceous plants, such as grasses, what could explain the little diversity of these families in the area.

Myrtaceae and Malpighiaceae families, the third and fourth most diverse families in our survey, also appear in similar studies between the most diverse families in Cerrado *sensu stricto* areas (Carvalho et al. 2010; Ishara and Rodela 2012) confirming the great ecological importance of these two families in the biome.

Among the listed species, *Anemopaegma arvense* (Bignoniaceae) is cited as endangered by Martinelli and Moraes (2013). *Caryocar brasiliense* (Caryocaraceae) is cited in the same literature as a species with commercial value undergoing distribution decline. These threatened species are concerning as the area has already been invaded by individuals of *Pinus* sp. Furthermore, six species are not cited in the books of the Flora do Distrito Federal (Cavalcanti et al. Vols. 1 to 10, from 2001 to 2012) nor in the Flora do Brasil website for Distrito Federal (Lista de Espécies da Flora do Brasil, consulted in 2015) (Table 1), highlighting that the flora of DF, although confined to small preserved areas, requires additional study.

Our study concludes that the Brazilian Ages Memorial, while not a conservation area, is an area of ecological relevance to the Distrito Federal. The area is rich in native species of Cerrado *sensu stricto* vegetation, which in turn attracts species of fauna associated with this kind of vegetation. Furthermore, it is an area suitable for new studies that will contribute to its management and conservation. This list can also be used for ethnobotanical studies in order to make the link between the “anthropological age” and “plant age” at the museum. Additionally, it will improve the educational purposes of the Memorial of Brazilian Ages.

ACKNOWLEDGEMENTS

We would like to acknowledge André Bertran, for help and making the study site available, and the Foundation for Research Support of Federal District (FAP-DF), the body responsible for funding this project. We would like also to thank Petra Broddle for comments on the manuscript and Regina Célia de Oliveira for helping with identification.

LITERATURE CITED

- Aguiar, L.M.S., B.M. Machado and J. Marinho-Filho. 2004. A diversidade biológica do Cerrado; pp. 17–40, in: L.M.S. Aguiar and A.J.A. Camargo (eds.) Cerrado: ecologia e caracterização. 0. Embrapa Cerrados, Planaltina.
- Amaral, G.A., C.B.R. Munhoz, C.U.O. Eugênio and J.M. Felfili. 2013. Vascular flora in dry-shrub and wet grassland Cerrado seven years after a fire, Federal District, Brazil. Check List 9(3): 487–503. <http://www.checklist.org.br/getpdf?SL121-12>

- Alho, C.J.R. and E.S. Martins. 1995. De Grão em Grão, o Cerrado Perde Espaço. Cerrado — Impactos do Processo de Ocupação. Brasília: WWF-Fundo Mundial para a Natureza. 66 pp.
- APG (Angiosperm Phylogeny Group). 2009. The Linear Angiosperm Phylogeny Group (APG) III: a linear sequence of the families in APG III. Botanical Journal of the Linnean Society 161: 128–131. doi: [10.1111/j.1095-8339.2009.01000.x](https://doi.org/10.1111/j.1095-8339.2009.01000.x)
- Carvalho, M.B., K.L. Ishara and R.C.S. Maimoni-Rodella. 2010. Vascular flora of a Cerrado *sensu stricto* remnant in Pratânia, state of São Paulo, southeastern Brazil. Check List 6(3): 350–357. <http://www.checklist.org.br/getpdf?SLo12-10>
- Cavalcanti T.B. and A.E. Ramos. 2001. Flora do Distrito Federal, Brasil. Vol. 1. Brasília: Embrapa recursos genéticos e tecnologia. 359 pp.
- Cavalcanti T.B. and A.E. Ramos. 2002. Flora do Distrito Federal, Brasil. Vol. 2. Brasília: Embrapa recursos genéticos e tecnologia 184 pp.
- Cavalcanti T.B. and A.E. Ramos. 2003. Flora do Distrito Federal, Brasil. Vol. 3. Brasília: Embrapa recursos genéticos e tecnologia. 240 pp.
- Cavalcanti T.B. and A.E. Ramos. 2005. Flora do Distrito Federal, Brasil. Vol. 4. Brasília: Embrapa recursos genéticos e tecnologia. 312 pp.
- Cavalcanti T.B. 2006. Flora do Distrito Federal, Brasil. Vol. 5. Brasília: Embrapa recursos genéticos e tecnologia 204 pp.
- Cavalcanti T.B. 2007. Flora do Distrito Federal, Brasil. Vol. 6. Brasília: Embrapa recursos genéticos e tecnologia 192 pp.
- Cavalcanti T.B. and M.F. Batista. 2009. Flora do Distrito Federal, Brasil. Vol. 7. Brasília: Embrapa recursos genéticos e tecnologia 328 pp.
- Cavalcanti T.B. and M.F. Batista. 2010. Flora do Distrito Federal, Brasil. Vol. 8. Brasília: Embrapa recursos genéticos e tecnologia 184 pp.
- Cavalcanti T.B. and A.P. Silva. 2011. Flora do Distrito Federal, Brasil. Vol. 9. Brasília: Embrapa recursos genéticos e tecnologia 224 pp.
- Cavalcanti T.B. and E.B.A. Dias. 2012. Flora do Distrito Federal, Brasil. Vol. 10. Brasília: Embrapa recursos genéticos e tecnologia 340 pp.
- Eiten, G. 1972. The Cerrado vegetation of Brazil. Botanical Review 38(2): 201–341.
- Felfili, J.M., M.C. Silva Júnior, A.V. Rezende, J.W.B. Machado, B.M.T. Walter, P.E.N. Silva and J.D. Hay. 1993. Análise comparativa da florística e fitossociologia da vegetação arbórea do cerrado *sensu stricto* na Chapada da Pratinha, DF- Brasil. Acta Botanica Brasílica 6(2): 27–46.
- Felfili, J.M. 2000. Vegetação no Distrito Federal: tempo e espaço. Brasília: UNESCO. 80 pp.
- Felfili, J.M. and M.C. Silva Júnior. 2001. Biogeografia do Bioma Cerrado: estudo fitofisionômico da Chapada do Espigão Mestre do São Francisco. Brasília: Universidade de Brasília, Faculdade de Tecnologia, Departamento de Engenharia Florestal 152 pp.
- IPNI. 2014. International Plant Names Index. Accessed at <http://www.ipni.org>, July 2014.
- Ishara, K.L. and R.C.S. Maimoni-Rodella. 2012. Richness and similarity of the Cerrado vascular flora in the central west region of São Paulo state, Brazil. Check List 8(1): 32–42. <http://www.checklist.org.br/getpdf?SLo52-11>
- Klink, C.A. 1996. Relação entre o desenvolvimento agrícola e a biodiversidade; pp. 25–27, in: R.C. Pereira and L.C.B. Nasser (eds.). Anais VIII Simpósio sobre o Cerrado, 1st International Symposium on Tropical Savanas — Biodiversidade e Produção Sustentável de Alimentos e fibras nos Cerrados. Brasília: Embrapa CPAC..
- Lista de Espécies da Flora do Brasil. 2014. Jardim Botânico do Rio de Janeiro. Accessed at <http://floradobrasil.jbrj.gov.br>, July 2014.
- Martinelli, G. and Moraes M.A. (orgs.) 2013. Livro Vermelho da Flora do Brasil. Centro Nacional de Conservação da Flora (CNCFLORA). Rio de Janeiro: Centro Nacional de Conservação da Flora. 1102 pp.
- Mendonça, R.C., J.M. Felfili, B.M.T. Walter, M.C. Silva Júnior, A.V. Rezende, T.S. Filgueiras and P.E. Nogueira. 1998. Flora vascular do cerrado; pp. 289–556, in: S.M. Sano, S.P. Almeida and J.F. Ribeiro (eds.). Cerrado: ecologia e flora. Planaltina: Embrapa Cerrados.
- Mendonça, R.C., J.M. Felfili, B.M.T. Walter, M.C. Silva Júnior, A.V. Rezende, T.S. Filgueiras, P.E. Nogueira and C.W. Fagg. 2008. Flora vascular do bioma Cerrado: checklist com 12.356 espécies; pp. 421–1279, in: S.M. Sano, S.P. Almeida and J.F. Ribeiro (orgs.). Cerrado: ecologia e flora. Brasília: Embrapa.
- Missouri Botanical Gardens. 2014. Tropicos. Accessed at <http://www.tropicos.org/>, July 2014
- Rapini, A., M.J.G. Andrade, A.M. Giulietti, L.P. Queiroz and J.M.C. Silva. 2009. Introdução; pp. 23–35, in: A.M. Giulietti, A. Rapini, M.J.G. Andrade, L.P. Queiroz and J.M.C. Silva (eds.). Plantas raras do Brasil. Belo Horizonte: Conservação Internacional.
- Ratter, J.A., J.F. Ribeiro and S. Bridgewater. 1997. The Brazilian Cerrado vegetation and threats to its biodiversity. Annals of Botany 80: 223–230. doi: [10.1006/anbo.1997.0469](https://doi.org/10.1006/anbo.1997.0469)
- Ribeiro, J.F. and B.M.T. Walter. 1998. Fitofisionomias do bioma cerrado; pp. 89–166, in: S.M. Sano, S.P. Almeida and J.F. Ribeiro (eds.). Cerrado: ecologia e flora Planaltina: Embrapa Cerrados.
- Ribeiro, J.F. and B.M.T. Walter. 2008. As principais fitofisionomias do Bioma Cerrado, pp. 151–212 in: Sano, S.M., S.P. Almeida and J.F. Ribeiro (eds.). Cerrado: ecologia e flora Planaltina: Embrapa Cerrados.
- The Plants List. 2014. The Plant List. Accessed at <http://www.theplantlist.org>, July 2014

Authors' contribution statement: TNCV, MLI, JSS and CEBP collected the data; TNCV, JSS and CEBP wrote the text. TNCV and MLI both had a scholarship from FAP-DF to produce this study and were supervised by CEBP.

Received: 3 December 2014

Accepted: 26 May 2015

Academic editor: Rubens Luiz Gayoso Coelho

Table 1. List of families and species of angiosperms from Memorial das Idades do Brasil, Federal District, Brazil, with their vouchers (UB), collectors (name and number), habit, categories and citation on the Flora do Distrito Federal. Collector: TN = T.N.C.Vasconcelos; MLI= M.L. Ianhez; CP= C.E.B. Proença. Habit: Tre: tree; Smt: small tree; Shr: Shrub; Ssh: subshrub; Her: herb; Lia: liana; Cre: creeper. Category. ECe: endemic to the Cerrado; eGO: endemic to Goiás state. Thr: threatened; Inv: invasive. N = species not cited in the Flora do Distrito Federal books. * species observed in the study site, but not collected.

Family/Species, Author	Voucher	Collector	Habit	Category	Flora DF
Acanthaceae					
<i>Justicia chrysotrichoma</i> (Nees) Benth.	109229; 121092	MLI 97; CP 3611	Her — Ssh	ECe, eGO	
<i>Justicia lanstykii</i> Rizzini	106714; 119026	MLI 23; TN 37	Ssh	ECe	
<i>Justicia sarothroides</i> Lindau	106686	TN 81	Ssh	ECe, eGO	
<i>Lepidagathis cyanea</i> (Leonard) Kameyama	84580; 109225; 121088	TN 10; MLI 93; CP 3607a	Her	ECe	
<i>Ruellia hapalotricha</i> Lindau	119017; 109239; 109274	TN 28; MLI 110; TN 102	Her	ECe, eGO	
<i>Ruellia incomta</i> (Nees) Lindau	126544	MRL 21	Ssh	ECe	
<i>Ruellia nitens</i> (Nees) Washh.	126545	MRL 22	Ssh	—	
Amaranthaceae					
<i>Alternanthera brasiliana</i> (L.) Kuntze	106674	TN 69	Her	—	
Anacardiaceae					
<i>Anacardium humile</i> A.St.-Hil.	106676; 106741	TN 71; MLI 50	Shr	—	
Annonaceae					
<i>Annona crassiflora</i> Mart.	165336	TN 126	Tre	—	
<i>Annona monticola</i> Mart.	165342	TN 132	Shr	—	
<i>Duguetia furfuracea</i> (A.St.-Hil.) Saff.	82020; 119028	CP 3571; TN 39	Shr	—	
<i>Duguetia marcgraviana</i> Mart.	106726	MLI 35	Tre	—	N
Apocynaceae					
<i>Aspidosperma macrocarpon</i> Mart.	82019	CP 3570	Tre	—	
<i>Hancornia speciosa</i> Gomes	82025; 106690	CP 3576; TN 85	Tre	—	
<i>Himatanthus obovatus</i> (Müll.Arg.) Woodson	82043	CP 3594	Tre	ECe	
<i>Odontadenia lutea</i> (Vell.) Markgr.	106734; 109259	MLI 43;	Cre	—	
Araliaceae					
<i>Schefflera macrocarpa</i> (Cham. & Schltdl.) Frodin	109255	MLI 126	Tre	—	
Areaceae					
<i>Syagrus flexuosa</i> (Mart.) Becc.	109262	TN 90	Shr	—	
Asteraceae					
<i>Achyrocline satureioides</i> (Lam.) DC.	119055	TN 66	Her	—	
<i>Aldama robusta</i> (Gardner) E.E. Schill & Panero	121084	TN 116	Ssh	—	
<i>Ageratum conyzoides</i> L.	119056	MLI 58	Her	Inv	
<i>Aspilia foliacea</i> (Spreng.) Baker	82027	CP 3578	Her	—	
<i>Baccharis reticularia</i> DC.	119031	TN 42	Shr	—	
<i>Bidens pilosa</i> L.	109276	TN 104	Her	Inv	
<i>Calea quadrifolia</i> Pruski & Urbatsch	109266	TN 94	Ssh	ECe, eGO	
<i>Calea sickii</i> (G.M.Barroso) Urbatsch et al.	84597	TN 27	Ssh	ECe, eGO	
<i>Chresta curumbensis</i> (Philipson) H.Rob.	106736	MLI 45	Her	ECe	
<i>Chresta exsucca</i> DC.	119020; 119035	TN 31; TN 46	Her	ECe	
<i>Chromolaena chaseae</i> (B.L.Rob.) R.M.King & H.Rob.	109251	MLI 122	Ssh	ECe	
<i>Dimerostemma vestitum</i> (Baker) S.F.Blake	109204	MLI 72	Ssh	ECe	
<i>Echinocoryne holosericea</i> (Mart. ex DC.) H.Rob.	106708; 118997; 119012	MLI 18; TN 8; TN 23	Shr	ECe	
<i>Elephantopus biflorus</i> (Less.) Sch.Bip.	119042	TN 53	Ssh	—	
<i>Eremanthus glomeratus</i> Less.	119123	MLI 108	Tre	ECe	
<i>Eremanthus goyazensis</i> (Gardner) Sch.Bip.	84594; 109279	TN 24; TN 107	Smt	ECe	
<i>Eremanthus mollis</i> Sch.Bip.	109246	MLI 117	—	ECe	
<i>Ichthyothere latifolia</i> Baker	106748	MLI 57	Ssh	ECe	
<i>Lepidaploa aurea</i> (Mart. ex DC.) H.Rob.	106683; 106729; 106737	TN 78; MLI 38; MLI 46	Ssh	—	
<i>Lepidaploa rufogrisea</i> (A.St.-Hil.) H.Rob.	119048	TN 59	Shr	ECe	
<i>Lessingianthus durus</i> (Mart. ex DC.) H.Rob.	119047	TN 58	Ssh	ECe	
<i>Lessingianthus floccosus</i> (Gardner) H.Rob.	119054	TN 65	Shr	ECe	
<i>Piptocarpha</i> sp.	—	TN 96	NA	—	
<i>Strophopappus glomerulatus</i> (Gardner) R.Esteves	121100	CP 3619	Ssh	?	
<i>Trichogonia salviifolia</i> Gardner	109222	MLI 90	Her	—	
<i>Vernonia cf. rubriramea</i>	84585	TN 15	Shr	—	
<i>Vernonia rubriramea</i> Mart. ex DC.	121093	CP 3612	Shr	?	
Bignoniaceae					
<i>Anemopaegma arvense</i> (Vell.) Stellfeld ex de Souza	106701	MLI 11	Ssh	—	
<i>Cuspidaria sceptrum</i> (Cham.) L.G.Lohman	84579; 119046; 109258	TN 9; TN 57; TN 86	Shr	ECe	

Continued

Table 1. Continued.

Family/Species, Author	Voucher	Collector	Habit	Category	Flora DF
<i>Jacaranda ulei</i> Bureau & K.Schum.	82033; 106707; 119041	CP 3584; MLI 17; TN 52	Shr	ECe	
<i>Zeyheria montana</i> Mart.	119043; 109280	TN 54; TN 108	Shr — Smt	—	
Bixaceae					
<i>Cochlospermum regium</i> (Mart. ex Schrank) Pilg.	106715	MLI 24	Shr	—	
Boraginaceae					
<i>Euploca salicoides</i> (Cham.) J.I.M.Melo & Semir	106718	MLI 27	Her — Ssh	?	
Bromeliaceae					
<i>Dyckia brasiliiana</i> L.B.Sm.	84582	TN 12	Her	ECe, eGO	
Burseraceae					
<i>Protium ovatum</i> Engl.	82034; 106675; 106702	CP 3585; TN 70; MLI 12	Shr	ECe	
Calophyllaceae					
<i>Kielmeyera coriacea</i> Mart. & Zucc.	106724	MLI 33	Tre	—	
<i>Kielmeyera rubriflora</i> Cambess.	84600	TN 30	Shr	ECe	
<i>Kielmeyera speciosa</i> A.St.-Hil.	109264; 109284; 121094	TN 92; TN 112; CP 3613	Tre	ECe	
Caryocaraceae					
<i>Caryocar brasiliense</i> Cambess.	82041	CP 3592	Tre	—	
Celastraceae					
<i>Plenckia populnea</i> Reissek	165340	TN 130	Tre	—	
<i>Salacia crassifolia</i> (Mart. ex Schult.) G.Don	82031	CP 3582	Shr – Smt	—	
Combretaceae					
<i>Terminalia fagifolia</i> Mart.	109200	MLI 68	Tre	—	
Connaraceae					
<i>Connarus suberosus</i> Planch.	106684; 106727; 109206	TN 79; MLI 36; MLI 74	Shr	ECe	
Convolvulaceae					
<i>Merremia tomentosa</i> (Choisy) Hallier f.	84573; 84596; 109195	TN 3; TN 26; MLI 63	Ssh	ECe	
Cucurbitaceae					
<i>Melothria campestris</i> (Naudin) H. Schaef. & S.S. Renner	—	*	NA	—	
<i>Perianthopodus</i> sp.	82038	CP 3589	NA	—	
Cyperaceae					
<i>Bulbostylis jacobinae</i> (Steud.) Lindm.	109215	MLI 83	Her	—	
<i>Rhynchospora consanguinea</i> (Kunth) Boeckeler	106709	MLI 19	Her	—	
<i>Rhynchospora exaltata</i> Kunth	84587	TN 17	Her	—	
Dilleniaceae					
<i>Davilla elliptica</i> A.St.-Hil.	106746; 109287	MLI 55; TN 115	Shr	—	
Ebenaceae					
<i>Diospyros hispida</i> A.DC.	82026; 109286; 121015; 165337165335	CP 3577; T 114; MLI 78; TN 127; TN 125	Tre	—	
Eriocaulaceae					
<i>Paepalanthus</i> sp.	—	MLI 78	NA	—	
Erythroxylaceae					
<i>Erythroxylum deciduum</i> A.St.-Hil.	82044	CP 3595	Ter – Smt	—	
<i>Erythroxylum suberosum</i> A.St.-Hil.	109193; 109197	MLI 61; MLI 65	Smt	—	
Euphorbiaceae					
<i>Croton campestris</i> A.St.-Hil.	109205	MLI 73	Shr	—	
<i>Dalechampia caperonioides</i> Baill.	106700	MLI 10	Her	ECe	
<i>Manihot anomala</i> Pohl	106740	MLI 49	Shr	—	
<i>Maprounea guianensis</i> Aubl.	106747; 109263	MLI 56; TN 91	Tre	ECe	
Fabaceae					
<i>Aeschynomene paucifolia</i> Vogel	106688	TN 83	Her — Ssh	ECe	
<i>Anadenanthera colubrina</i> (Vell.) Brenan	165353	TN 143	Tre	—	
<i>Ancistrotropis firmula</i> (Mart. ex Benth.) A.Delgado	109249; 109283	MLI 120; TN 111	Shr — Cre	—	
<i>Bauhinia dumosa</i> Benth.	165348	TN 138	Ssh	ECe	
<i>Bauhinia rufa</i> (Bong.) Steud.	119029; 119038	TN 40; TN 49	Ssh	ECe	
<i>Calliandra dysantha</i> Benth.	82037; 119027	CP 3588; TN 38	Ssh — Shr	—	
<i>Chamaecrista basifolia</i> (Vogel) H.S.Irwin & Barneby	109226	MLI 94	Ssh	ECe	
<i>Chamaecrista benthamiana</i> (Harms) H.S.Irwin & Barneby	106679	TN74	Ssh — Shr	ECe, eGO	
<i>Chamaecrista clauseni</i> (Benth.) H.S.Irwin & Barneby	11936; 106680	TN 47; TN 75	Ssh	ECe	
<i>Chamaecrista conferta</i> (Benth.) H.S.Irwin & Barneby	106696	MLI 6	Ssh	—	
<i>Chamaecrista desvauxii</i> (Collad.) Killip	109227; 109233; 109267	MLI 95; MLI 102; TN 95	Shr	—	
<i>Chamaecrista filicifolia</i> (Benth.) H.S.Irwin & Barneby	109245	MLI 116	Ssh	—	
<i>Crotalaria flavicoma</i> Benth.	109236	MLI 106	Her — Ssh	—	

Continued

Table 1. Continued.

Family/Species, Author	Voucher	Collector	Habit	Category	Flora DF
<i>Dalbergia miscolobium</i> Benth.	109269	TN 97	Tre	—	
<i>Dimorphandra mollis</i> Benth.	109209; 119039	MLI 77; TN 50	Tre	—	
<i>Enterolobium gummiferum</i> (Mart.) J.F.Macbr.	165343	TN 133	Tre	ECe	
<i>Eriosema glaziovii</i> Harms	109196	MLI 64	Her	ECe, eGO	
<i>Galactia heringeri</i> Burkart	109234	MLI 103	Her	—	
<i>Galactia peduncularis</i> (Benth.) Taub.	82017; 106692; 109252	CP 3568; MLI 2b; MLI 123	Her	ECe	
<i>Hymenaea stigonocarpa</i> Mart. ex Hayne	165341	TN 131	Tre	—	
<i>Leptolobium dasycarpum</i> Vogel	82023	CP 3574	Tre – Smt	—	
<i>Macroptilium</i> sp.	119135	MLI 120b	NA	—	
<i>Mimosa albolanata</i> Taub.	84599; 119030	TN 29; TN 41	Shr	ECe	
<i>Mimosa debilis</i> Humb. & Bonpl. ex Willd.	106730	MLI 39	Her — Ssh	—	
<i>Mimosa lanuginosa</i> Glaz. ex Burkart	119021; 119034	TN 32; TN 45	Ssh	—	
<i>Mimosa somnians</i> Humb. & Bonpl. ex Willd.	121097	CP 3616	Shr	—	
<i>Periandra mediterranea</i> (Vell.) Taub.	84571; 109221; 121102	TN 1; MLI 89; CP 3621	Ssh — Shr	—	
<i>Pterodon emarginatus</i> Vogel	165352	TN 142	Tre	—	
<i>Senna rugosa</i> (G.Don) H.S.Irwin & Barneby	84576; 109275; 121091	TN 6; TN 103; CP 3610a	Shr	—	
<i>Stryphnodendron adstringens</i> (Mart.) Coville	165338	TN 128	Tre	—	
<i>Strychantes guianensis</i> (Aubl.) Sw.	109238	MLI 109	Her	ECe	
<i>Tachigali vulgaris</i> L.G.Silva & H.C.Lima	119040	TN 51	Tre	—	
Gentianaceae					
<i>Deianira chiquitana</i> Herzog	165349	TN 139	Ssh	—	
Iridaceae					
<i>Alophia drummondii</i> (Grah.) R.C.Foster	109194	MLI 62	?	?	N
<i>Sisyrinchium vaginatum</i> Spreng.	121101	CP 3620	Her	—	
<i>Trimezia juncifolia</i> (Klatt) Benth. & Hook.	109231	MLI 99	Her	—	
Lamiaceae					
<i>Aegiphila verticillata</i> Vell.	82045; 165344	CP 3596; TN 134	Shr — Tre	—	
<i>Hypenia macrantha</i> (A.St.-Hil. ex Benth.) Harley	109253; 119051	MLI 124; TN 62	Ssh	—	
<i>Hypptis conferta</i> Pohl ex Benth.	109232	MLI 100	Shr	—	
<i>Hypptis rubiginosa</i> Benth.	106704; 119050	MLI 14; TN 61	Ssh	ECe	
<i>Hypptis villosa</i> Pohl ex Benth.	119033	TN 44	Her	ECe	
<i>Medusantha crinita</i> (Benth.) Harley & J.F.B.Pastore	119049	TN 60	Ssh	ECe	
<i>Oocephalus lytroides</i> (Pohl ex Benth.) Harley & J.F.B.Pastore	84592	TN 22	Ssh	ECe, eGO	
<i>Salvia tomentella</i> Pohl	109203	MLI 71	Her	—	
Loganiaceae					
<i>Strychnos pseudoquina</i> A.St.-Hil.	82022	CP 3573	Tre	—	
Lythraceae					
<i>Cuphea ferruginea</i> Pohl ex Koehne	109201	MLI 69	Ssh	ECe, eGO	
<i>Cuphea spermacoce</i> A.St.-Hil.	106711	MLI 21	Ssh	ECe, eGO	
<i>Diplusodon rosmarinifolius</i> A.St.-Hil.	119037; 119044	TN 48; TN 55	Ssh	ECe, eGO	
<i>Diplusodon sessiliflorus</i> Koehne	121099	CP 3618	Ssh	ECe, eGO	
<i>Lafoensia pacari</i> A.St.-Hil.	109242	MLI 113	Tre	—	
Malpighiaceae					
<i>Banisteriopsis argyrophylla</i> (A.Juss.) B.Gates	109268	TN 96	Shr — Cre	ECe	
<i>Banisteriopsis campestris</i> (A.Juss.) Little	106732; 109235	MLI 41; MLI 104	Shr	—	
<i>Banisteriopsis laevifolia</i> (A.Juss.) B.Gates	109243	MLI 114	Shr — Cre	ECe	
<i>Banisteriopsis latifolia</i> (A.Juss.) B.Gates	121096	CP 3615	Smt	ECe, eGO	
<i>Banisteriopsis malifolia</i> (Nees & Mart.) B.Gates	84591; 121098	TN 21; CP 3617	Shr	—	
<i>Banisteriopsis megaphylla</i> (A.Juss.) B.Gates	106681; 109265; 109248	TN 76; TN 93; MLI 119	Shr	ECe	
<i>Banisteriopsis</i> sp.	84595; 109247	TN 25; MLI 118	NA	—	
<i>Byrsonima basiloba</i> A.Juss.	109223; 119032	MLI 91; TN 43	Ssh	ECe	
<i>Byrsonima coccolobifolia</i> Kunth	82018; 106728	CP 3569; MLI 37	Tre	—	
<i>Byrsonima guilleminiana</i> A.Juss.	82028; 119022; 106689	CP 3579; TN 33; TN 84	Shr	ECe	
<i>Byrsonima rotunda</i> Griseb.	109248	MLI 119	—	—	
<i>Byrsonima</i> sp.	119045; 109277	TN 56; TN 105	NA	—	
<i>Camarea affinis</i> A.St.-Hil.	109220	MLI 88	Ssh	ECe	
<i>Peixotoa cordistipula</i> A.Juss.	109250	MLI 121	Ssh	ECe	N
<i>Peixotoa goiana</i> C.E.Anderson	84586	TN 16	Ssh	ECe, eGO	
<i>Pterandra pyroidea</i> A.Juss.	106698	MLI 8	Ssh	ECe	

Continued

Table 1. Continued.

Family/Species, Author	Voucher	Collector	Habit	Category	Flora DF
Malvaceae					
<i>Eriotheca pubescens</i> (Mart. & Zucc.) Schott & Endl.	109257	MLI 128	Shr — Tre	ECe	
<i>Pavonia pohlii</i> Gürke	84577; 84584; 109260	TN 7; TN 14; TN 88	Ssh	ECe, eGO	
<i>Sida cerradoensis</i> Krap.	109207	MLI 75	Ssh — Shr	—	
<i>Sida linifolia</i> Cav.	106723	MLI 32	Her	Inv	
<i>Sida santaremensis</i> Mont.	109278	TN 106	Her — Ssh	Inv	
<i>Waltheria indica</i> L.	109230	MLI 98	Ssh	Inv	
Melastomataceae					
<i>Miconia albicans</i> (Sw.) Triana	—	*	Smt	—	
<i>Miconia burchellii</i> Triana	—	*	Tre	ECe	
<i>Miconia fallax</i> DC.	84590; 106703	TN 20; MLI 13	Smt	—	
<i>Miconia ferruginata</i> DC.	119025; 119052	TN 36; TN 63	Tre	—	
<i>Miconia pohliana</i> Cogn.	106743	MLI 52	Tre	—	
<i>Miconia rubiginosa</i> (Bonpl.) DC.	106714	MLI 23	Shr	—	
<i>Tibouchina aegopogon</i> Cogn.	109240	MLI 111	Ssh	ECe	
<i>Tibouchina laevicaulis</i> Wurdack	84572; 109282; 106745	TN 2; TN 110; MLI 54	Shr	ECe, eGO	
Menispermaceae					
<i>Cissampelos ovalifolia</i> DC.	106706	MLI 16	Ssh	—	
Moraceae					
<i>Brosimum gaudichaudii</i> Trécul	82040; 106721	CP 3591; MLI 30	Shr — Tre	—	
Myrtaceae					
<i>Campomanesia pubescens</i> (Mart. ex DC.) O.Berg	82032	CP 3583	Shr	—	
<i>Eugenia angustissima</i> O.Berg	82015; 106725	CP 3566; MLI 34	Ssh	—	
<i>Eugenia bimarginata</i> DC.	106678	TN 73	Tre	—	
<i>Eugenia dysenterica</i> (Mart.) DC.	82039	CP 3590	Tre	—	
<i>Eugenia involucrata</i> DC.	82036	CP 3587	Shr — Tre	—	
<i>Eugenia paranahybensis</i> O.Berg	106722	MLI 31	Ssh	ECe	N
<i>Eugenia puniceifolia</i> (Kunth) DC.	106699	MLI 9	Shr	—	
<i>Myrcia goyazensis</i> Cambess.	106687	TN 82	Shr	ECe	
<i>Myrcia guianensis</i> (Aubl.) DC.	106673; 106695; 109192	MLI 5, MLI 60, TN 68	Shr	—	
<i>Myrcia nivea</i> Cambess.	106685	TN 80	Shr	ECe	
<i>Myrcia variabilis</i> Mart. ex DC.	106731	MLI 40	Ssh	—	
<i>Psidium firmum</i> O.Berg	106682; 106720	TN 77; MLI 29	Shr	—	
<i>Psidium laruotteanum</i> Cambess.	109285	TN 113	Shr	—	
Nyctaginaceae					
<i>Guapira noxia</i> (Netto) Lundell	165350	TN 140	Tre	ECe	
<i>Guapira psammophila</i> (Mart. ex J.A.Schmidt) Angely.	109241	MLI 112	?	?	N
Ochnaceae					
<i>Ouratea floribunda</i> (A.St.-Hil.) Engl.	165345	TN 135	Ssh — Shr	ECe	
<i>Ouratea hexasperma</i> (A.St.-Hil.) Baill.	84589; 109256	TN 19; MLI 127	Shr – small tree	ECe	
Orchidaceae					
<i>Cyrtopodium</i> sp.	165356	TN 146	NA	—	
Orobanchaceae					
<i>Buchnera lavandulacea</i> Cham. & Schltdl.	106677	TN 72	Her	—	
<i>Buchnera rosea</i> Kunth	109254	MLI 125	Her	—	
Oxalidaceae					
<i>Oxalis barrelieri</i> L.	109191	MLI 59	Her — Ssh	Inv	
Poaceae					
<i>Aristida setifolia</i> Kunth	119053	TN 64	Her	—	
<i>Axonopus aureus</i> P.Beauv.	119024	TN 35	Her	—	
<i>Axonopus pressus</i> (Sw.) P.Beauv.	152130	MLI 89b	Her	—	
<i>Ichnanthus inconstans</i> (Trin. ex Nees) Döll	106738	MLI 47	Her	—	
<i>Paspalum eucomum</i> Nees ex Trin.	121090	CP 3609a	Her	ECe	
<i>Urochloa decumbens</i> (Stapf) R.D.Webster	152131	MLI 93b	Her	Inv	
Primulaceae					
<i>Cybianthus densiflorus</i> Miq.	109244	MLI 115	Shr	—	
<i>Cybianthus detergens</i> Mart.	82013; 106697; 106712	CP 3564; MLI 7; MLI 22	Shr	—	
<i>Myrsine guianensis</i> (Aubl.) Kuntze	109202	MLI 70	Tre	—	
Proteaceae					
<i>Roupala montana</i> Aubl.	106733	MLI 42	Tre	—	

Continued

Table 1. *Continued.*

Family/Species, Author	Voucher	Collector	Habit	Category	Flora DF
Rubiaceae					
<i>Borreria capitata</i> (Ruiz & Pav.) DC.	121089	CP 3608a	Her — Ssh	—	
<i>Borreria tenella</i> (Kunth) Cham. & Schltldl.	165351	TN 141	Ssh	—	
<i>Chomelia ribesoides</i> Benth. ex A.Gray	106693	MLI 3	Shr	ECe	
<i>Declieuxia fruticosa</i> (Willd. ex Roem. & Schult.) Kuntze	106744; 109261	MLI 53; TN 89	Ssh — Shr	—	
<i>Palicourea officinalis</i> Mart.	109289	TN 117	Ssh	ECe	
<i>Palicourea rigida</i> Kunth	82042; 106717	CP 3593; MLI 26	Shr – small tree	—	
<i>Planaltina capitata</i> (K.Schum.) R.M.Salas & E.L.Cabral	84588; 109281	TN 18; TN 109	Ssh	ECe, eGO	
<i>Sabicea brasiliensis</i> Wernham	84574; 106739	TN 4; MLI 48	Ssh	—	
<i>Tocoyena formosa</i> (Cham. & Schltldl.) K.Schum.	82029; 106710; 109211	CP 3580; MLI 20; MLI 79	Smt	—	
Salicaceae					
<i>Casearia sylvestris</i> Sw.	106672; 106691	TN 67; MLI 1	Shr – Smt	—	
Sapindaceae					
<i>Serjania cf. obtusidentata</i> Radlk.	165346	TN 136	Cre	—	
<i>Serjania</i> sp.	82035	CP 3586	NA	—	
Smilacaceae					
<i>Smilax goyazana</i> A.DC.	82021; 121095; 106694	CP 3572; CP 3614; MLI 4	Shr	—	
Solanaceae					
<i>Cestrum cf. obovatum</i> Sendtn.	106735	MLI 44	Ssh	—	
<i>Schwenckia americana</i> Rooyen ex L.	109273	TN 101	Ssh	—	
<i>Solanum lycocarpum</i> A. St.-Hil.	82016; 106705	CP 3567; MLI 15	Shr – Smt	—	
<i>Solanum subumbellatum</i> Vell.	106716	MLI 25	Ssh	—	
<i>Solanum viarum</i> Dunal	109208	MLI 76	Shr	—	
Styracaceae					
<i>Styrax ferrugineus</i> Nees & Mart.	84581	TN 11	Tre — Smt	ECe	
Turneraceae					
<i>Turnera lamiifolia</i> Cambess.	109224	MLI 92	Her	ECe	
<i>Turnera longiflora</i> Cambess.	109237	MLI 107	Ssh	ECe	
Urticaceae					
<i>Cecropia pachystachya</i> Trécul	—	*	Tre	—	
Velloziaceae					
<i>Vellozia squamata</i> Pohl	121087	CP 3606a	Ssh — Shr	ECe	
Verbenaceae					
<i>Lippia origanoides</i> Kunt	84575	TN 5	Shr	—	
Violaceae					
<i>Pombalia lanata</i> (A.St.-Hill.) Paula-Souza	82014	CP 3565	Her	ECe	
Vitaceae					
<i>Cissus erosa</i> Rich.	109199	MLI 67	Cre	—	
Vochysiaceae					
<i>Qualea grandiflora</i> Mart.	165339	TN 129	Tre	—	
<i>Qualea multiflora</i> Mart.	82024; 84583	CP 3575; TN 13	Tre	—	
<i>Vochysia elliptica</i> Mart.	106742; 119023	MLI 51; TN 34	Tre	—	
<i>Vochysia thyrsoidea</i> Pohl	165347	TN 137	Tre	—	