

# *Eratyrus mucronatus* Stål, 1859 (Hemiptera: Reduviidae: Triatominae): First report in the State of Acre, Brazil, and updated geographic distribution in South America

Marcos Takashi Obara<sup>1</sup>, Acigelda da Silva Cardoso<sup>2</sup>, Maria Carmelinda Gonçalves Pinto<sup>3</sup>, Cláudio Rodrigues de Souza<sup>4</sup>, Rafaella Albuquerque e Silva<sup>5</sup> and Rodrigo Gurgel-Gonçalves<sup>6\*</sup>

<sup>1</sup> Universidade de Brasília, Faculdade de Ceilândia. QNN 14, Área Especial, Ceilândia Sul. CEP 72220-140. Brasília, DF, Brasil.

<sup>2</sup> Secretaria Municipal de Saúde, Área Técnica de Entomologia Municipal. Av. Ceará, Nº 3188, Bairro Abraão Alab. CEP 69960-160. Rio Branco, AC, Brasil.

<sup>3</sup> Secretaria de Estado de Saúde, Área Técnica de Doença de Chagas. Rua Benjamin Constant, 830- 1º Andar, Centro. CEP 69908-420. Rio Branco, AC, Brasil.

<sup>4</sup> Secretaria de Estado de Saúde, Área Técnica de Entomologia Estadual. Av. Antônio da Rocha Viana, Nº 1584, Bairro Vila Igonete. CEP: 69980-560. Rio Branco, AC, Brasil.

<sup>5</sup> Ministério da Saúde, Secretaria de Vigilância em Saúde, Unidade Técnica de Vigilância das Doenças de Transmissão Vetorial. Setor Comercial Sul, Quadra 4, Bloco A, 2º andar, Edifício Principal. CEP 70304-000. Brasília, DF, Brasil.

<sup>6</sup> Universidade de Brasília, Faculdade de Medicina, Área de Patologia, Laboratório de Parasitologia Médica e Biologia de Vetores. Campus Universitário Darcy Ribeiro, Asa Norte. CEP 70904-970. Brasília, DF, Brasil.

\* Corresponding author. E-mail: [rurgel@unb.br](mailto:rurgel@unb.br)

**ABSTRACT:** A new occurrence record of *Eratyrus mucronatus* (Hemiptera, Reduviidae, Triatominae) is reported from Rio Branco, Acre, Brazil. The distribution of this triatomine species was updated and a potential geographic distribution map in South America was presented based on ecological niche modeling.

Currently, 145 species are recognized in the subfamily Triatominae (Costa *et al.* 2006; Costa and Felix 2007; Schofield and Galvão 2009; Frías-Lasserre 2010; Rosa *et al.* 2012; Gonçalves *et al.* 2013; Jurberg *et al.* 2013) of which 63 occurs in Brazil (Gurgel-Gonçalves *et al.* 2012; Rosa *et al.* 2012). The genus *Eratyrus* Stål, 1859 is represented by two species: *Eratyrus cuspidatus* Stål, 1859 and *Eratyrus mucronatus* Stål, 1859.

*Eratyrus mucronatus* occurs in Bolivia, Colombia, Ecuador, Guatemala, Guyana, French Guyana, Panama, Peru, Suriname, Trinidad, Venezuela and Brazil (Carcavallo *et al.* 1999). In Brazil, this species was registered in the states of Amazonas, Tocantins, Maranhão, Mato Grosso, Rondônia and Pará, in humid forest ecoregions (Abad-Franch and Monteiro 2007; Gurgel-Gonçalves *et al.* 2012).

Tree holes, palm trees and rocks are known habitats of *E. mucronatus* in forests (Carcavallo *et al.* 1998b). In nature, nymphs can feed on the haemolymph of arachnids (Amblypygi) inside hollow trees and adults are usually associated with mammals, such as *Coendou prehensilis*, *Didelphis marsupialis* and bats (Carcavallo *et al.* 1998a; Gaunt and Miles 2000).


In the Amazon, invasion of houses by triatomines, apparently related to artificial light sources, should favor transmission of *Trypanosoma cruzi* to humans either by direct contact or by food contamination, especially through *açaí* fruit (Coura *et al.* 1999; Valente *et al.* 1999; Nobrega *et al.* 2009; Pereira *et al.* 2009).

According to Morocoima *et al.* (2010) *E. mucronatus* specimens are highly attracted to artificial light sources. In Brazil, *E. mucronatus* has been registered in houses in Manaus (Castro *et al.* 2010) and Rondônia (Meneguetti

*et al.* 2011). Furthermore, detection of infected colonies of *E. mucronatus* in houses and peridomestic has been observed in Bolivia (Noireau *et al.* 1995; Depickère *et al.* 2012), Colombia (Guhl 2007), Peru (Torres and Cabrera 2010) and Venezuela (Rojas *et al.* 2008) indicating the potential risk of transmission of *T. cruzi* to humans. Thus, it is important to record the locations where *E. mucronatus* has been detected in domiciles, as well as update the geographic distribution of this species and predict areas of potential occurrence. In this study we registered a new occurrence of *E. mucronatus* in Brazil, updated its distribution and analyzed its potential geographic distribution in South America.

In November 2011, a triatomine was detected in a house located in the Vila Irineu Serra, Rio Branco, state of Acre, Brazil. This state is located in northern Brazil, bordering the states of Amazonas and Rondônia. It has an area of 152,581.4 km<sup>2</sup> equivalent to about 2% of the total area of the country. The climate is hot and humid with average monthly temperatures ranging between 24°C and 27°C. The annual rainfall is 2.100 mm, with a dry season during the months of June, July and August (IBGE 2012).


The residence was located in a periurban area with the presence of secondary forests, palm trees and livestock. The insect was sent to the Laboratory of Entomology of Rio Branco, Acre, and identified as *E. mucronatus* (Figure 1). The identification was confirmed by the Laboratory of Medical Entomology of General Coordination of Public Health Laboratories of the Health Surveillance Secretariat of the Ministry of Health. The specimen is deposited at the Laboratory of Entomology of the Municipal Health Secretariat of Rio Branco, Acre, Brazil.



**FIGURE 1.** *Eratyrus mucronatus* specimen (male) from municipality of Rio Branco, state of Acre, Brazil. A. dorsal view. B. lateral view. C. *Attalea* palm trees found in the peridomestic of the house where the specimen was detected, probably the natural ecotope of *E. mucronatus*.

Data on the occurrence of *E. mucronatus* were obtained from Lent and Wygodzinsky (1979), Silveira et al. (1984), Rebelo et al. (1998), Soto Vivas et al. (2001), Chávez (2006), Guhl et al. (2007), Cortez et al. (2007), Rojas et al. (2008), Serrano et al. (2008), Castro et al. (2010), Morocoima et al. (2010), Torres and Cabrera (2010), Meneguetti et al. (2011), Depickère et al. (2012). Records of *E. mucronatus* in collections of Rodolfo Carcavallo and Herman Lent at the National and International Reference Laboratory on Triatomine Taxonomy at the Oswaldo Cruz Institute, FIOCRUZ, were analyzed. The distributional data for *E. mucronatus* available in other museums provided by the Global Biodiversity Information Facility - GBIF (<http://data.gbif.org>), species link (<http://splink.cria.org.br/>) and VertNet (<http://vertnet.org/index.php>) were also analyzed.

In order to develop ecological niche models for *E. mucronatus*, data on climate were obtained from the WorldClim project (<http://www.worldclim.org>). The environmental data used in the analyses cover South America, at a spatial resolution of 2.5' (5 km). To avoid the confounding effects of calibrating models in an overly dimensional environmental space, only a subset of the 19 variables in the climatic data were chosen: annual mean temperature, mean diurnal range (mean of monthly (max temp - min temp)), maximum temperature of warmest month, minimum temperature of coldest month, annual precipitation, precipitation of wettest month, and precipitation of driest month (Gurgel-Gonçalves et al. 2012).



**FIGURE 2.** Geographic distribution of *Eratyrus mucronatus* in South America. The squares represent the known records for this species in Venezuela, Colombia, Ecuador, Peru, Bolivia and Brazil. The triangle represents the new record from the state of Acre, Brazil described in the present article. Areas identified as suitable based on climatic variables using ecological niche modeling (Maxent method) are shown in redscale: light red (low suitability) to dark red (high suitability).

Data on a total of 58 occurrences of *E. mucronatus* were assembled. These occurrences were referenced to geographic coordinates with an uncertainty of  $\leq 5$  km and precision of  $0.01^\circ$ . The records were georeferenced based on data found at <http://www.fallingrain.com/world>. Occurrence data were separated into two sets: one for model calibration (75% of points) and the other for model evaluation (25% of points). Maps of potential distribution of *E. mucronatus* were produced using Maxent version 3.2.1 (Phillips *et al.* 2006), applying the default parameters with 10 replicates using Bootstrap replication method. Then, the averaged model was imported into ARC VIEW (version 3.2) to create and edit the distribution map. Predicted areas in semi-arid caatinga ecoregion were eliminated in the final edited map.

The quality of the models was evaluated using the ROC (receiver operating characteristic) curve, which correlates the sensitivity and specificity of model predictions across multiple thresholds (Phillips *et al.* 2006). Model accuracy was also assessed by examining the proportion of test points falling into regions of predicted presence (Anderson *et al.* 2002). Using Maxent's internal Jackknife test, the variables that most influence the distribution of triatomine species were identified.

The updated geographic distribution map of *E. mucronatus* (Figure 2) presents the known records in South America and the new record of the state of Acre, Brazil. The ecological niche model indicates that *E. mucronatus* should occur widely in Amazon. The species also showed a potential geographic distribution in drier ecoregions outside the Amazon region, in northern Venezuela and Colombia, such as Maracaibo dry forests, Lara-Falcón dry forests, Paraguana xeric scrub and La Costa xeric shrublands. All of the models derived from the analysis performed well ( $AUC > 0.92$ ), summarizing the necessary ecological conditions for the occurrence of *E. mucronatus*. All test points fell in the area of predicted presence of *E. mucronatus*. The Jackknife tests showed that precipitation of the wettest month was the variable that best explained the models.

The importance of *E. mucronatus* as a potential vector of *T. cruzi* to humans has been highlighted in Bolivia, Peru and Venezuela (Rojas *et al.* 2008; Torres and Cabrera 2010; Depickère *et al.* 2012). The new record of *E. mucronatus* in domicile in Brazil is further evidence of the synanthropic potential of this species. Ecological niche modeling predicts that *E. mucronatus* is widespread in Amazon, suggesting that the invasion of this species in households must be very common in humid ecoregions of South America.

**ACKNOWLEDGMENTS:** The authors are grateful to Gilberto Gil da Silva Serato and Janis Lunier de Souza, entomology technicians from the Municipal Health of Rio Branco, for their support in identifying the species. We also thank Patricia Gurgel Gonçalves for reviewing the English version of the manuscript.

#### LITERATURE CITED

- Abad-Franch, F. and F.A. Monteiro. 2007. Biogeography and evolution of Amazonian triatomines (Heteroptera: Reduviidae): implications for Chagas disease surveillance in humid forest ecoregions. *Memórias do Instituto Oswaldo Cruz* 102: 57-69.
- Anderson, R.P., M. Gómez-Laverde and A.T. Peterson. 2002. Geographical distributions of spiny pocket mice in South America: insights from predictive models. *Global Ecology and Biogeography* 11: 131-141.
- Carcavallo, R.U., D.S. Rocha, I. Galíndez Girón, I. Sherlock, C. Galvão, A. Martínez, R.J. Tonn and E. Cortón. 1998a. Feeding sources and patterns; p. 537-560 In R.U. Carcavallo, I. Galíndez Girón, J. Jurberg and H. Lent. (ed.) *Atlas of Chagas Disease Vectors in Americas*. Volume II. Rio de Janeiro: Fiocruz.
- Carcavallo, R.U., M. Rodríguez, R. Salvatella, S.I. Curto de Casas, I. Sherlock and C. Galvão. 1998b. Habitats and related fauna; p. 561-619 In R.U. Carcavallo, I. Galíndez Girón, J. Jurberg and H. Lent (ed.). *Atlas of Chagas Disease Vectors in Americas*, Volume II. Rio de Janeiro: Fiocruz.
- Carcavallo, R.U., S.I. Curto de Casas, I. Sherlock, I. Galíndez Girón, J. Jurberg, C. Galvão, C.A. Mena Segura and F. Noireau. 1999. Geographical distribution and alti-latitude dispersion; p. 747-792 In R.U. Carcavallo, I. Galíndez Girón, J. Jurberg and H. Lent. (ed.) *Atlas of Chagas disease vectors in Americas*, Volume III. Rio de Janeiro: Fiocruz.
- Castro, M.C.M., T.V. Barrett, W.S. Santos, F. Abad-Franch and J.A. Rafael. 2010. Attraction of Chagas disease vectors (Triatominae) to artificial light sources in the canopy of primary Amazon rainforest. *Memórias do Instituto Oswaldo Cruz* 105: 1061-1064.
- Chávez, J. 2006. Contribución al estudio de los triatominos del Perú: Distribución geográfica, nomenclatura y notas taxonómicas. *Anales de la Facultad de Medicina* 67(1): 65-76.
- Cortez, M.R., M. Avalos, V. Rocha and D. Gorla. 2007. Distribución biogeográfica de los triatominos en Bolivia: discriminación de la distribución de las especies en relación a variables ambientales; p. 67-137 In: Cortez, M.R. (ed.) *Triatominos de Bolivia y la enfermedad de Chagas*. La Paz: Ministerio de Salud y Deportes. Programa Nacional de Chagas.
- Costa, J. and M. Felix. 2007. *Triatomata juazeirensis* sp. nov. from the state of Bahia, northeastern Brazil (Hemiptera: Reduviidae: Triatominae). *Memórias do Instituto Oswaldo Cruz* 102: 87-90.
- Costa, J., A.M. Argolo and M. Felix. 2006. Redescription of *Triatomata melanica* Neiva & Lent, 1941, new status (Hemiptera: Reduviidae: Triatominae). *Zootaxa* 1385: 47-52.
- Coura, J.R., A.C.V. Junqueira, M.N. Bóia, O. Fernandes. 1999. Chagas disease: from bush to husts and houses. Is it the case of the Brazilian Amazon? *Memórias do Instituto Oswaldo Cruz* 94 (1): 379-384.
- Depickère, S., P. Durán, R. López, E. Martínez and T. Chávez. 2012. After five years of chemical control: colonies of the triatomine *Eratyrus mucronatus* are still present in Bolivia. *Acta Tropica* 123: 234-238.
- Frías-Lasserre, D. 2010. A new species and karyotype variation in the bordering distribution of *Mepraia spinolai* (Porter) and *Mepraia gajardoi* Frias et al (Hemiptera: Reduviidae: Triatominae) in Chile and its parapatric model of speciation. *Neotropical Entomology* 39: 572-583.
- Gaunt, M. and M. Miles. 2000. The ecotopes and evolution of triatomine bugs (Triatominae) and their associated trypanosomes. *Memórias do Instituto Oswaldo Cruz* 95: 557-565.
- Gonçalves, T.C.M., S.C. Neves-Neves, J.R.S. Mallet, A.L. Carbajal-de-la-Fuente and C.M. Lopes. 2013. *Triatomata jatai* sp. nov. in the state of Tocantins, Brazil (Hemiptera: Reduviidae: Triatominae). *Memórias do Instituto Oswaldo Cruz* 108: 429-437.
- Guhl, F. 2007. Chagas disease in Anden countries. *Memórias do Instituto Oswaldo Cruz* 102(1): 29-37.
- Guhl, F., G. Aguilera, N. Pinto and D. Vergara. 2007. Actualización de la distribución geográfica y ecoepidemiología de la fauna de triatominos (Reduviidae: Triatominae) en Colombia. *Biomédica* 27: 143-162.
- Gurgel-Gonçalves, R., C. Galvão, J. Costa and A.T. Peterson. 2012. Geographic distribution of Chagas disease vectors in Brazil based on ecological niche modeling. *Journal of Tropical Medicine* 2012: 1-15.
- IBGE 2012. Instituto Brasileiro de Geografia e Estatística. *Geografia e demografia do Estado do Acre*. Database accessible at <http://www.ibge.gov.br/>. Captured on 29 October 2012.
- Jurberg, J., V. Cunha, S. Cailleaux, R. Raigorodski, M.S. Lima, D.S. Rocha and F.F.F. Moreira. 2013. *Triatomata pintodiiasi* sp. nov. do subcomplexo *T. rubrovaria* (Hemiptera: Reduviidae, Triatominae). *Revista Pan-Amazônica de Saúde* 4: 43-56.
- Lent, H. and P. Wygodzinsky. 1979. Revision on the Triatominae (Hemiptera, Reduviidae) and their significance as vectors of Chagas'disease. *Bulletin of the American Museum of Natural History* 163: 123-520.
- Meneguetti, D.U.O., O. Trevisan, R.M. Rosa and L.M. Camargo. 2011. First report *Eratyrus mucronatus*, Stal, 1859, (Hemiptera, Reduviidae, Triatominae), in the State of Rondonia, Brazil. *Revista da Sociedade Brasileira de Medicina Tropical* 44: 511-512.
- Morocoima, A., J. Chique, L. Herrera and S.U. Morales. 2010. *Eratyrus mucronatus* (Stal, 1859) (Hemiptera, Reduviidae, Triatominae): primer registro para el estado Anzoátegui (Venezuela). *Boletín de Marariología y Salud Ambiental* 2: 307-310.
- Nóbrega, A.A., M.H. Garcia, E. Tatto, M.T. Obara, E. Costa, J. Sobel and W.N. Araujo. 2009. Oral transmission of Chagas Disease by consumption of açaí Palm Fruit, Brasil. *Emerging Infectious Diseases* 15: 653-655.
- Noireau, F., M.F. Bosseno, R. Carrasco, J. Telleria, F. Vargas, C.

- Camacho, N. Yaksi and S.F. Breniere. 1995. Sylvatic triatomines (Hemiptera: Reduviidae) in Bolivia: trends towards domesticity and possible infection with *Trypanosoma cruzi* (Kinetoplastida: Trypanosomatidae). *Journal of Medical Entomology* 32: 594-598.
- Pereira, K.S., F.L. Schmidt, A.M.A. Guaraldo, R.M.B. Franco, V.L. Dias and L.A.C. Passos. 2009. Chagas disease as a foodborne illness. *Journal Food Protection* 72: 441-446.
- Phillips, S.J., R.P. Anderson and R.E. Schapire. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190: 231-259.
- Rebelo, J.M.M., V.L.L. Barros and W.A. Mendes. 1998. Espécies de Triatominae (Hemiptera: Reduviidae) do Estado do Maranhão, Brasil. *Cadernos de Saúde Pública* 14: 187-192.
- Rojas, M.E., P. Várquez, M.F. Villarreal, C. Velandia, L. Vergara, Y.H. Móran-Borges, J. Ontiveros, M.Y. Calderón, M.A. Chiurillo-Siervo, C.C. Rodríguez-Bonfate, E. Aldana, J.L. Concepción and R.A. Bonfate-Cabarcas. 2008. Estudio soroepidemiológico y entomológico sobre la enfermedad de Chagas en un área infestada por *Triatoma maculata* (Erichson 1848) en el centro-occidente de Venezuela. *Cadernos de Saúde Pública* 24: 2323-2333.
- Rosa, J.A., C.S. Rocha, S. Gardim, M.C. Pinto, V.J. Mendonça, J.C.R.F. Filho, E.O.C. Carvalho, L.M.A. Camargo, J. Oliveira, J.D. Nascimento, M. Cilense and C.E. Almeida. 2012. Description of *Rhodnius montenegrensis* sp.n. (Hemiptera: Reduviidae: Triatominae) from the state of Rondônia, Brazil. *Zootaxa* 3478: 62-76.
- Schofield, C.J. and C. Galvão. 2009. Classification, evolution, and species groups within the Triatominae. *Acta Tropica* 110: 88-100.
- Silveira, A.C., V.R. Feitosa and R. Borges. 1984. Distribuição de triatomíneos capturados no ambiente domiciliar, no período de 1975/83, Brasil. *Revista Brasileira de Malariologia e Doenças Tropicais* 39: 15-312.
- Serrano, O., F. Mendoza, B. Suárez and A. Soto. 2008. Seroepidemiology of Chagas disease in two rural populations in the municipality of Costa de Oro, at Aragua State, northern Venezuela. *Biomedica* 28: 108-15.
- Soto Vivas, A., H. Barazarte and D.M. Fernández. 2001. Primer registro de *Eratyrus mucronatus* Stål, 1959 (Hemiptera: Reduviidae) en el ambiente domiciliario en Venezuela. *Entomotropica* 16: 215-217.
- Torres, D.B. and R. Cabrera. 2010. Geographical distribution and intra-domiciliary capture of sylvatic triatomines in La Convención Province, Cusco, Peru. *Revista do Instituto de Medicina Tropical de São Paulo* 52: 157-160.
- Valente, S.A.S., V.C. Valente and H.F. Neto. 1999. Considerations on the epidemiology and transmission of Chagas disease in the Brazilian Amazon. *Memórias do Instituto Oswaldo Cruz* 94 (Suppl 1): 395-398.

RECEIVED: December 2012

ACCEPTED: February 2013

PUBLISHED ONLINE: September 2013

EDITORIAL RESPONSIBILITY: Lívia Aguiar Coelho