

First record of *Scinax tripui* Lourenço, Nascimento and Pires, 2010 (Amphibia: Anura: Hylidae) from Espírito Santo state, Brazil

Guilherme Ramos da Silva^{1*}, Cyro de Luna-Dias^{2,3}, Fabio Silva Fernandes dos Santos Hepp⁴ and Sergio Potsch de Carvalho e Silva³

- 1 Universidade Estadual do Piauí, Campus Professor Alexandre Alves Oliveira. Avenida Nossa Senhora de Fátima s/nº. CEP 64202-220. Parnaíba, PI, Brazil.
 - 2 Programa de Pós Graduação em Zoologia, Universidade Federal do Rio de Janeiro, Museu Nacional. Quinta da Boa Vista. CEP 20940-040. Rio de Janeiro, RJ, Brazil.
 - 3 Universidade Federal do Rio de Janeiro, Instituto de Biologia, Departamento de Zoologia. Av. Brigadeiro Trompowsky, s/nº, CCS, Bl. A, Ilha do Fundão. CEP 21941-590. Rio de Janeiro, RJ, Brazil.
 - 4 Universidade Federal do Rio de Janeiro, Museu Nacional, Departamento de Vertebrados. Quinta da Boa Vista, São Cristóvão. CEP 20940-040. Rio de Janeiro, RJ, Brasil.
- * Corresponding author. E-mail: grscinax@gmail.com

ABSTRACT: *Scinax tripui* is a medium sized frog that belongs to the *Scinax catharinae* clade. We provide a new record of *Scinax tripui* from Serra de Torres, Espírito Santo state, southeastern Brazil. This is the first record outside the specie's type locality (Ouro Preto, Minas Gerais) and the first record from the state of Espírito Santo.

The genus *Scinax* Wagler ranges from eastern and southern Mexico to Argentina and Uruguay; Trinidad and Tobago; St. Lucia (Frost 2013). The genus is composed by two large clades: the *S. ruber* and the *S. catharinae* clades (Faivovich 2002; Faivovich *et al.* 2005). The *S. ruber* clade contains the *S. rostratus* group, *S. uruguayus* group, and a large number of species currently unassigned to any group. The *S. catharinae* clade contains the groups *catharinae* and *perpusillus* (Faivovich *et al.* 2005). *Scinax tripui* is a medium sized frog diagnosed by: snout subovoid in dorsal view; canthus rostralis slightly concave and well marked; dorsum surface dark brown with scattered dark brown and green blotches; nuptial pads developed, but nonhypertrophied; interorbital region with a W-shaped mark posteriorly; light green coloration on hidden regions of flanks and thighs of adults; and belly and throat cream with great density of dark brown spots (Lourenço *et al.* 2009).

During field work in September 2010 at Serra de Torres (21°02'60"S, 41°15'0.8"W), between the municipalities of Atilio Vivácqua, Mimoso do Sul, and Muqui, Espírito Santo state, we collected four adults of *Scinax tripui* (Figure 1). Specimens were collected in streams in altitude between 900 and 950 m, in three different types of Atlantic Forest physiognomies: seasonal semideciduous, submontane dense ombrophilous, and montane dense ombrophilous. Specimens were deposited in the Amphibian Collection of the Departamento de Zoologia, Universidade Federal do Rio de Janeiro (ZUF RJ 12307-12310). The permit for collecting zoological material was provided by Instituto Chico Mendes (ICMBio) – Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais e Renováveis (IBAMA; number 12164-2). Before our record, the species was known only from its type locality, at the Estação Ecológica do Tripuí, municipality of Ouro Preto (20°23'22"S,




FIGURE 1. *Scinax tripui* from the municipality of Mimoso do Sul, Espírito Santo state, Brazil. A: dorsal view of an adult male (ZUF RJ 12308, SVL 28.8 mm). B: Amplectant pair (male: ZUF RJ 12310, SVL 24.4 mm; female: ZUF RJ 12309, SVL 47.0 mm). C: detail of light green coloration on hidden regions of flanks and thighs (ZUF RJ 12307). Photos: C. Luna-Dias.

43°32'32"W, elev. 1070m), Minas Gerais state, Brazil, in the Serra Espinhaço. This record of *Scinax tripui* in Serra das Torres extends its distribution southwestwards at about 252 km in a straight line from the type locality (Figure 2).

Several species of amphibians had its boundaries recently extended to Serra das Torres (ES) (Oliveira et al. 2009; Oliveira et al. 2012; Silva et al. 2012), increasingly revealing the biogeographic relevance of the region for this taxonomic group. We suggest that future intensified studies in the region will likely reveal new records of species. Our record contributes to the knowledge of the geographical distribution of *Scinax tripui* and enables a better assessment of their conservation status.




FIGURE 2. Geographic distribution of *Scinax tripui*. Red circle: new record in Serra de Torres, municipality of Mimoso do Sul, Espírito Santo state. White circle: type locality in Estação Ecológica do Tripuí, municipality of Ouro Preto, Minas Gerais state.

ACKNOWLEDGMENTS: We are grateful to Maria das Graças de Castro Sedano e Valdeir de Castro Sedano for hospitality and help in field samplings.

LITERATURE CITED

- Faivovich, J. 2002. A cladistic analysis of *Scinax* (Anura: Hylidae). *Cladistics* 18(4): 367-393.
- Faivovich, J., C.F.B. Haddad, P.C.A. Garcia, D.R. Frost, J.A. Campbell and W.C. Wheeler. 2005. Systematic review of the frog family Hylidae, with special reference to Hylinae: Phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 294: 1-240.
- Frost, D. R. 2013. *Amphibian Species of the World: an online reference. Version 5.6 (9 January, 2013)*. Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA. Captured on 22 april 2012.
- Lourenço, A.C.C, L.B. Nascimento and M.R.S. Pires. 2009. A new species of the *Scinax catharinae* species group (Anura: Hylidae) from Minas Gerais, southeastern Brazil. *Herpetologica* 65(4): 468-479.
- Oliveira, J.C.F, L. Coco, F.F. de Deus, R. Pagotto, E.S. Silva, C.F.D. Rocha and D. Vrcibradic. 2009. Geographic Distribution. *Phasmahyla guttata*. *Herpetological Review* 40(4): 446.
- Oliveira, J.C.F, L. Coco, R. V. Pagotto, E. Pralon, D. Vrcibradic, J. P. Pombal Jr and C.F.D. Rocha. 2012. Amphibia, Anura, *Brachycephalus didactylus* (Izecksohn, 1971) and *Zachaeus parvulus* (Girard, 1853): Distribution extension. *Check List* 8(2): 242-244.
- Silva, G. R., C. Luna-Dias, F. S. F. S. Hepp, A. M. P. T. Carvalho-e-Silva and S. P. Carvalho-e-Silva. 2012. New record of *Aplastodiscus weygoldti* (Cruz & Peixoto, 1987) in the municipality of Mimoso do Sul, Espírito Santo State, southeastern Brazil (Anura, Hylidae). *Herpetology Notes* 5: 371-373.

RECEIVED: November 2012

ACCEPTED: May 2013

PUBLISHED ONLINE: June 2013

EDITORIAL RESPONSIBILITY: Fernanda Werneck