

# First record of *Flabellina dana* Millen and Hamann, 2006 (Mollusca: Nudibranchia) in the South Atlantic Ocean

Licia Sales<sup>1\*</sup>, Marlon Delgado<sup>2</sup>, Vinicius Queiroz<sup>1</sup>, Vinicius Padula<sup>3</sup>, Cláudio L. S. Sampaio<sup>4</sup> and Rodrigo Johnsson<sup>1</sup>

1 Universidade Federal da Bahia, Instituto de Biologia, Departamento de Zoologia, Avenida Adhemar de Barros s/nº, Campus Ondina, CEP 40170-290, Salvador, BA, Brazil.

2 Universidade Federal do Rio Grande do Norte, Centro de Biociências, Departamento de Botânica, Ecologia e Zoologia, Laboratório de Invertebrados Bentônicos, Avenida Senador Salgado Filho, CEP 59072-970, Natal, RN, Brazil.

3 Zoologische Staatssammlung München, Mollusca Sektion, Münchhausenstr. 21, 81247, München, Germany.

4 Universidade Federal de Alagoas, Unidade de Ensino Penedo, Avenida Beira Rio, s/nº, Centro Histórico, CEP 57200-000, Penedo, AL, Brazil.

\* Corresponding author: E-mail: [licia\\_so@yahoo.com.br](mailto:licia_so@yahoo.com.br)

**ABSTRACT:** The nudibranch *Flabellina dana* Millen and Hamann, 2006 is reported from two localities in the northeastern Brazilian coast. These are the first records of this species, previously recorded from localities in the Bahamas and the Caribbean Sea, in South Atlantic Ocean, extending its known geographic distribution more than 3500 km southward.

Nudibranchia comprehends around 3000 marine gastropod species highly diverse in morphological, ecological and biological aspects. Nudibranchs lack the shell in adult forms, are hermaphrodite and most species are carnivorous, feeding on other marine invertebrates, mainly sponges. It is the richest group within Opisthobranchia, with approximately half of the total number of species (Behrens 2005; Wagële and Klussman-Kolb 2005).

So far, around 100 nudibranch species were reported to Brazil (DaCosta *et al.* 2010), a small number if compared to more studied places and even the Caribbean Sea (191 species, Valdés *et al.* 2006). In Brazil, most studies were carried out during three decades, ranging from 1955 to 1983 by the German couple Ernst and Eveline Marcus (Marcus 1955; Marcus 1970; Marcus 1983). Recently, new species were discovered and described from Brazil (DaCosta *et al.* 2010; Padula and Delgado 2010; Alvim *et al.* 2011) indicating that the diversity is clearly underestimated.

Among nudibranchs, *Flabellina* Voigt, 1834 is characterized by a triseriate radula with denticulate rachidian teeth; strong jaws; presence of corners on the foot (propodial tentacles); grouped cerata and pleuroproctus anus (Millen and Hermosillo 2007). The genus has over 54 species distributed in temperate and tropical seas (Millen and Hamann 2006). In Brazil, only four species are known: *Flabellina dushia* (Marcus and Marcus, 1963) from the region of Búzios, Rio de Janeiro state (García García *et al.* 2008); *Flabellina engeli* Marcus and Marcus, 1968 known from Cabo Frio and Arraial do Cabo, Rio de Janeiro state to Arvoredo, Santa Catarina state (DaCosta *et al.* 2007); *Flabellina verta* (Marcus, 1970) from Cananéia, São Paulo state (García García *et al.* 2008; Rios 2009) and *Flabellina marcusorum* Gosliner and Kuzirian, 1990 from Santos Bay, São Paulo state (García García *et al.* 2008; Rios 2009).

This work reports for the first time the occurrence of *Flabellina dana* Millen and Hamann, 2006 in the Brazilian

coast, South Atlantic Ocean. This species was previously recorded in the Caribbean Sea, more exactly in St. Lucia (type locality), Curaçao, Costa Rica and also in the island of Abaco, Bahamas (Figure 1) (Redfern, 2001; Millen and Hamann 2006; Valdés *et al.* 2006).

The collected and examined material is deposited in the malacological collection of the Museu de Zoologia of the Universidade de São Paulo (MZSP). Specimens were collected through snorkeling in Santa Rita beach, Extremoz, Rio Grande do Norte state (RN) (05°41'41" S, 35°11'58" W) on August 2009 (MZSP 99636) (Figure 2B) and in Barra beach, Salvador, Bahia state (BA) (13°00'37" S, 38°31'49" W) on January 2010 (MZSP 96480) (Figure 2A and 2C). The specimen from Bahia (MZSP 96480) was photographed *in situ* (Figure 2A).

**Order Nudibranchia Blainville, 1814**

**Family Flabellinidae Bergh, 1889**

***Flabellina dana* Millen and Hamann, 2006**

**Description:** long and extremely slender body, specimens length 6,0 mm (MZSP 99636) and 7,0 mm (MZSP 96480). Bilabiate foot with long propodial tentacles and tapered posterior end. Oval head as wide as foot. Mouth with vertical aperture. Oral tentacles smooth, length corresponding to approximately 1/5 of the total body length. Annulate and wide rhinophores with pointed tip, each rhinophore with up to eight wide and spaced rings. Eyes situated laterally on the head just below the basis of each rhinophore. Cerata latero-dorsally positioned, arranged in five to six paired groups along the body length. Pair of pre-cardiac groups with five to six ceratas, post-cardiac groups decreasing in number of cerata towards distal end. Translucid white body with a longitudinal dorsal white line. Body sides opaque white. Cerata red with white cnidosacs. Rhinophores opaque white except for their translucid basis (Figure 2).

The specimens found in Brazil are consistent with the


original description of *F. dana* (Millen and Hamann 2006) and additional works (Redfern, 2001, as *Flabellina* sp. A; Valdés *et al.* 2006, as *Flabellina* sp. 3), based on material from the Bahamas and the Caribbean Sea. As the species is easily recognized by its external morphology, with specific characteristics such as the annulate rhinophores, the small specimens herein studied were not dissected to preserve its morphology and were deposited intact in the collection. Our records from Brazil extend the known geographic distribution of the species more than 3500 km southward, in the South Atlantic Ocean.

Among the Flabellinidae the species that most resembles *F. dana* is *F. dushia*. Some common characteristics are the opaque white color on the head, oral tentacles and most of the dorsum; the foot bilabiate and notched with long propodial tentacles; the pleuroproctic anus, posterior to inter-hepatic space; and the renal and genital opening location. However, the two species can be easily morphologically differentiated due to characteristics in the head, mouth and rhinophores. *Flabellina dushia* presents rounded head, terminal and triangular mouth and smooth rhinophores, while *F. dana* has oval head, vertical and sub-terminal mouth and annulate rhinophores (Millen and Hamann 2006; Valdés *et al.* 2006). The rhinophore is an important diagnostic characteristic of *F. dana*. It is the only known western Atlantic Flabellinidae with annulate rhinophores (Figure 2C) (Millen and Hamann, 2006).



**FIGURE 1.** Geographic distribution of *Flabellina dana*. Previous records (circles): 1. Bahamas; 2. St. Lucia (type locality); 3. Curaçao and 4. Costa Rica (Valdés *et al.* 2006). New records (squares): 5. Rio Grande do Norte and 6. Bahia, Brazilian northeastern coast.

Species distribution ranging from the Caribbean Sea as far as the Brazilian coast is not novelty among marine organisms, with examples in cnidarians (Neves *et al.* 2006; Neves *et al.* 2008; Neves *et al.* 2010) and reef fishes (Floeter *et al.* 2001; Joyeux *et al.* 2001; Rocha *et al.* 2005; Floeter *et al.* 2008; Olavo *et al.* 2011). Concerning nudibranchs, the similarity between the Brazilian and Caribbean fauna has been reported since 1970's (Marcus 1970; Marcus and Marcus 1970; Padula and Santos 2006). *Flabellina dana* is one more species in common between these two regions, being the present records important additions to the little known nudibranch biodiversity in northeastern Brazil.



**FIGURE 2.** *Flabellina dana*, specimens from Rio Grande do Norte and Bahia, northeastern Brazil. A. Specimen *in situ* from Salvador, Bahia State (7 mm long); B. Specimen from Extremoz, Rio Grande do Norte State (6 mm long); C. Detail of the annulate rhinophores of the specimen from Salvador.

**ACKNOWLEDGMENTS:** The authors thank Cristiane Xerez Barroso (UFC) and one anonymous referee for valuable comments on the manuscript. We also thank Sandra Millen (University of British Columbia) for her incentive and opinion on the identity of the studied material, and Luiz Simone (Museu de Zoologia of the Universidade de São Paulo) for providing registration numbers for specimens. V.P. is a scholarship holder from the Council for Scientific and Technological Development (CNPq-Brazil) and the German Academic Exchange Service (DAAD-Germany). R.J. would like to thank CNPq and FAPESB for the financial support.

#### LITERATURE CITED

- Alvim, J., V. Padula and A.D. Pimenta. 2011. First record of the genus *Onchidoris* (Gastropoda: Nudibranchia: Onchidorididae) from the South Atlantic Ocean, with the description of a new species from Brazil. *Journal of the Marine Biological Association of the United Kingdom* 91(2): 505-511.
- Behrens D.W. 2005. *Nudibranch Behaviour*. Jacksonville: New World Publications, Inc. 176 p.
- DaCosta, S., C.M. Cunha, L.R.L. Simone and M. Schrödl. 2007. Computerbased 3-dimensional reconstruction of major organ systems of a new aeolid nudibranch subspecies, *Flabellina engeli lucianae*, from Brazil (Gastropoda, Opisthobranchia). *Journal of Molluscan Studies* 73: 339-353.
- DaCosta, S., V. Padula and M. Schrödl. 2010. A new species of *Hypselodoris* and a redescription of *Hypselodoris picta lajensis* (Nudibranchia: Chromodorididae) from Brazil. *The Veliger* 51(1): 15-25.

- Floeter, S.R., R.Z.P. Guimarães, L.A. Rocha, C.E.L. Ferreira, C.A. Rangel and J.L. Gasparini. 2001. Geographic variation in reef-fish assemblages along the Brazilian coast. *Global Ecology & Biogeography* 10: 423-431.
- Floeter, S.R., L.A. Rocha, D.R. Robertson, J.C. Joyeux, W.F. Smith-Vaniz, P. Wirtz, A.J. Edwards, J.P. Barreiros, C.E.L. Ferreira, J.L. Gasparini, A. Brito, J.M. Falcón, B.W. Bowen and G. Bernardi. 2008. Atlantic reef fish biogeography and evolution. *Journal of Biogeography* 35: 22-47.
- García García, F.J., M. Domínguez Álvarez and J.S. Troncoso. 2008. *Opisthobranchios de Brasil: Descripción y distribución de opisthobranchios del litoral de Brasil y del Archipiélago Fernando de Noronha*. Vigo: Feito, S.L. 215 p.
- Joyeux, J.C., S.R. Floeter, C.E.L. Ferreira and J.L. Gasparini. 2001. Biogeography of tropical reef fishes: the South Atlantic puzzle. *Journal of Biogeography* 28: 831-841.
- Marcus, Er. 1955. Opisthobranchia from Brazil. *Boletim da Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo, Zoologia* 20: 89-262.
- Marcus, Er. and Ev. Marcus. 1970. Opisthobranchs from Curaçao and faunistically related regions. *Studies on the Fauna of Curaçao and other Caribbean Islands* 122: 1-129.
- Marcus, Ev. 1970. Opisthobranchs from Northern Brazil. *Bulletin of Marine Science* 20: 922-951.
- Marcus, Ev. 1983. The Western Atlantic Tritoniidae. *Boletim de Zoologia da Universidade de São Paulo* 6: 177-214.
- Millen, S. and J.C. Hamann. 2006. A new nudibranch species, genus *Flabellina* (Opisthobranchia: Aeolidacea) from the Caribbean with redescription of *F. verta* (Marcus 1970), and *F. dushia* (Marcus and Marcus, 1963). *Proceedings of the California Academy of Sciences* 57 (30): 928-936.
- Millen, S. and A. Hermsillo. 2007. The *Flabellina* Voight, 1834 (Mollusca: Opisthobranchia) from Bahía de Banderas (Pacific Coast of Mexico) with ecological observations, the description of a new species, and redescription of *Flabellina cynara*. *Proceedings of the California Academy of Sciences* 58 (26): 543-559.
- Neves, E.G., S.C. Andrade, F.L. Silveira and V.N. Solferini. 2008. Genetic variation and population structuring in two brooding coral species (*Siderastrea stellata* and *Siderastrea radians*) from Brazil. *Genetica* (The Hague) 132: 243-254.
- Neves, E.G., R. Johnsson, C.L.S. Sampaio and M. Pichon. 2006. The occurrence of *Scolymia cubensis* in Brazil: revising the problem of the Caribbean solitary mussids. *Zootaxa* 1366: 45-54.
- Neves, E.G., F.L. Silveira, M. Pichon, and R. Johnsson. 2010. Cnidaria, Scleractinia, Siderastreaidae, *Siderastrea siderea* (Ellis and Solander, 1786): Hartt Expedition and the first record of a Caribbean siderastroid in tropical Southwestern Atlantic. *Check List* 6: 505-510.
- Olavo, G., P.A.S. Costa, A.S. Martins and B.P. Ferreira. 2011. Shelf-edge reefs as priority areas for conservation of reef fish diversity in the tropical Atlantic. *Aquatic Conservation: Marine and Freshwater Ecosystems*. DOI: 10.1002/aqc.1174
- Padula, V. and M. Delgado. 2010. A new species of *Cerberilla* (Gastropoda: Nudibranchia: Aeolidiidae) from northeastern Brazil. *The Nautilus* 124 (4):175-180.
- Padula, V. and F.N. Santos. 2006. Three new records of Nudibranchia (Mollusca, Gastropoda) – additions on the Brazilian biodiversity. *Biociências* 14: 214-220.
- Redfern, C. 2001. *Bahamian Seashells: a Thousand Species from Abaco, Bahamas*. Florida: Bahamianseashells.com, Inc., 261 p.
- Rios, E.C. 2009. *Compendium of Brazilian Sea Shells*. Rio Grande: Evangraf. 668 p.
- Rocha, L.A., D.R. Robertson, J. Roman and B.W. Bowen. 2005. Ecological speciation in tropical reef fishes. *Proceedings of the Royal Society B* 272: 573-579.
- Valdés, A., J. Hamann, D.W. Behrens and A. Dupont. 2006. *Caribbean Sea Slugs: A field guide to the the opisthobranch mollusks from the tropical northwestern Atlantic*. Washington: A Sea Challengers Natural History Books Publication. 289 p.
- Wägele, H. and A. Klussmann-Kolb. 2005. Opisthobranchia (Mollusca, Gastropoda) - more than just slimy slugs. Shell reduction and its implications on defense and foraging. *Frontiers in Zoology* 2(3): 1-18.

RECEIVED: July 2011

LAST REVISED: August 2011

ACCEPTED: September 2011

PUBLISHED ONLINE: December 2011

EDITORIAL RESPONSIBILITY: Luis E. Arruda Bezerra