



Corrigendum: Role of Hippocampal Lipocalin-2 in Experimental Diabetic Encephalopathy

OPEN ACCESS

Approved by:

Frontiers in Endocrinology Editorial Office, Frontiers Media SA, Switzerland

*Correspondence:

Kyoungho Suk ksuk@knu.ac.kr

[†]These authors have contributed equally to this work

Specialty section:

This article was submitted to Experimental Endocrinology, a section of the journal Frontiers in Endocrinology

Received: 25 March 2019 Accepted: 26 March 2019 Published: 09 April 2019

Citation:

Bhusal A, Rahman MH, Lee I-K and Suk K (2019) Corrigendum: Role of Hippocampal Lipocalin-2 in Experimental Diabetic Encephalopathy. Front. Endocrinol. 10:239. doi: 10.3389/fendo.2019.00239 Anup Bhusal^{1†}, Md Habibur Rahman^{1†}, In-Kyu Lee² and Kyoungho Suk^{1,3*}

¹ BK21 Plus KNU Biomedical Convergence Program, Departments of Biomedical Science and Pharmacology, School of Medicine, Kyungpook National University, Daegu, South Korea, ² Division of Endocrinology and Metabolism, Department of Internal Medicine, School of Medicine, Kyungpook National University, Daegu, South Korea, ³ Brain Science and Engineering Institute, Kyungpook National University, Daegu, South Korea

Keywords: Lipocalin-2, diabetic encephalopathy, hippocampus, glia, neuroinflammation, cognitive dysfunction

A Corrigendum on

Role of Hippocampal Lipocalin-2 in Experimental Diabetic Encephalopathy

by Bhusal, A., Rahman, M. H., Lee, I.-K., and Suk, K. (2019). Front. Endocrinol. 10:25. doi: 10.3389/fendo.2019.00025

In the original article, there was a mistake in **Figure 1** as published. The *Gapdh* band image used in **Figure 1C** was from the pilot experiment. The corrected **Figure 1** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2019 Bhusal, Rahman, Lee and Suk. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 1 Expression of LCN2 in the hippocampus of diabetic mice. The expression of *Lcn2* mRNA in the hippocampus at 8 w post MLDS and 4 w post HDS injection was assessed by conventional PCR **(A,C)**. Further, the expression level of LCN2 protein in the hippocampus of STZ-induced diabetic mice was estimated by ELISA assay **(B,D)**. Similar upregulation of *Lcn2* mRNA and LCN2 protein was detected in the hippocampus at 24 w post HFD feeding **(E,F)**. **p* < 0.05 vs. the vehicle-treated control animals; Student's *t*-test; *n* = 3 for each group; data are represented as mean \pm SEM. STZ, streptozotocin; MLDS, multiple low dose of STZ; HDS, high dose of STZ; HFD, high fat diet; LCN2, Lipocalin-2; w, weeks; SEM, standard error of the mean.