
Surveys in Mathematics and its Applications

ISSN 1842-6298 (electronic), 1843-7265 (print)
Volume 14 (2019), 109 – 140

A SURVEY ON THE TERNARY PURELY EXPONENTIAL DIOPHANTINE EQUATION

$$a^x + b^y = c^z$$

Maohua Le, Reese Scott and Robert Styer

Abstract. Let a, b, c be fixed coprime positive integers with $\min\{a, b, c\} > 1$. In this survey, we consider some unsolved problems and related works concerning the positive integer solutions (x, y, z) of the ternary purely exponential diophantine equation $a^x + b^y = c^z$.

1 Introduction

Let \mathbb{Z}, \mathbb{N} be the sets of all integers and positive integers, respectively. A great deal of number theory arises from the discussion of the integer or rational solutions of a polynomial equation with integer coefficients. Such equations are called diophantine equations. Using the definition given by T. N. Shorey and R. Tijdeman [158], those diophantine equations with fixed bases and variable exponents are called purely exponential equations. Let a, b, c be fixed coprime positive integers with $\min\{a, b, c\} > 1$. In this survey we investigate the ternary purely exponential equations of the form

$$a^x + b^y = c^z, x, y, z \in \mathbb{N}. \quad (1.1)$$

We give an outline of the contents of this survey.

First, in this Introduction (Section 1), after some explanatory comments to help orient the reader, we give the current bounds on the variable exponents x, y , and z . We then give the current bounds on the number of solutions (x, y, z) to (1.1).

Section 2 deals with the Jeśmanowicz conjecture, which states that, if (a, b, c) is a Pythagorean triple, then the only solution to (1.1) is $(x, y, z) = (2, 2, 2)$. Work on this conjecture is characterized by many different highly specific results, making it difficult for those working on this conjecture to know if they are making full use of what is available and not reproving old results. Here the first author presents the existing work in an organized manner, which we hope will greatly facilitate future work on this problem.

2010 Mathematics Subject Classification: 11D61.

Keywords: ternary purely exponential diophantine equation; Jeśmanowicz conjecture, Terai-Jeśmanowicz conjecture.

<http://www.utgjiu.ro/math/sma>

Section 3 introduces the Terai-Jeśmanowicz conjecture, which states that (1.1) has at most one solution with $\min\{x, y, z\} > 1$. Here the first author clarifies the various oversights which have occurred in previous formulations of the conjecture. Then a third conjecture, which includes both the Jeśmanowicz and Terai-Jeśmanowicz conjectures, is stated and briefly discussed. The remainder of Section 3 is largely concerned with various cases for which it can be shown that (1.1) has at most one solution.

Before proceeding, it might be helpful to point out what is *not* included in this survey.

First, when one of x or y is fixed, (1.1) becomes the familiar Pillai equation, which has already been thoroughly handled in an excellent survey by M. Waldschmidt [189]. So here we treat only cases in which all of x , y , and z are variable. Second, (1.1) can be viewed as a specific case of an S -unit equation. The many results of this type are usually not included here. Third, we do not reference any work from the vast and familiar literature dealing with cases in which at least one of the bases a , b , c is variable. Finally, we deal largely with fairly recent results; we do not attempt to give a complete history of the problem.

We note that the first author conceived the project and drafted the bulk of this paper and its extensive bibliography, the second author contributed some sections showing connections to more general work on exponential equations as well as contributing multiple improvements, and the third author edited previous versions and prepared the final version as well as handling various technical matters.

We now begin our discussion of (1.1).

In 1933, K. Mahler [128] used his p -adic analogue of the diophantine approximation method of Thue-Siegel to prove that (1.1) has only finitely many solutions (x, y, z) . His method is ineffective. An effective result for solutions of (1.1) was given by A. O. Gel'fond [47]. In 1999, M.-H. Le [89] used some elementary methods to prove that if $c \equiv 1 \pmod{2}$, then the solutions (x, y, z) of (1.1) satisfy $z < \frac{2}{\pi} ab \log(2eab)$. This gives a bound on z independent of c . Recently, R. Scott and R. Styer [157] proved that if $c \equiv 1 \pmod{2}$ then $z < \frac{1}{2} ab$. In some cases, finding a bound on z independent of c can be taken quite a bit further: when every prime dividing ab is in a given finite set of primes S , it is often possible to show that (1.1) implies $z = 1$ except for a finite list of specified exceptions; to do this, elementary methods sometimes suffice (e. g., $S = \{2, 3, 5\}$), but in general they do not, particularly when $2 \notin S$. M. A. Bennett and N. Billerey [3] deal with the case $S = \{3, 5, 7\}$, completely handling not only (1.1) but also the more general case in which A and B are S -units such that $A + B = c^z$, $z > 1$.

We now consider finding bounds on $\max\{x, y, z\}$. N. Hirata-Kohno [65] showed that if $c \equiv 1 \pmod{2}$ then $\max\{x, y, z\} < 2^{288} \sqrt{abc} (\log(abc))^3$. For general a , b , and c , combining a lower bound for linear forms in two logarithms and an upper bound

for the p -adic logarithms, Y.-Z. Hu and M.-H. Le [69] proved that

$$\max\{x, y, z\} < 155000(\log(\max\{a, b, c\}))^3. \quad (1.2)$$

Very recently they improved that to

$$\max\{x, y, z\} < 6500(\log(\max\{a, b, c\}))^3. \quad (1.3)$$

(see Y.-Z. Hu and M.-H. Le [71]).

Let $N(a, b, c)$ denote the number of solutions (x, y, z) of (1.1). As a straightforward consequence of an upper bound for the number of solutions of binary S -unit equations due to F. Beukers and H. P. Schlickewei [6], we have $N(a, b, c) \leq 2^{36}$. Considering the parity of x and y , all solutions (x, y, z) of (1.1) can be put into the following four classes:

- Class I: $x \equiv y \equiv 0 \pmod{2}$.
- Class II: $x \equiv 1 \pmod{2}$ and $y \equiv 0 \pmod{2}$.
- Class III: $x \equiv 0 \pmod{2}$ and $y \equiv 1 \pmod{2}$.
- Class IV: $x \equiv y \equiv 1 \pmod{2}$.

Under this approach, M.-H. Le [89] proved that if $a < b$ and $c \equiv 1 \pmod{2}$, then each class has at most $2^{\omega(c)-1}$ solutions (x, y, z) , where $\omega(c)$ is the number of distinct prime divisors of c , except for $(a, b, c) = (3, 10, 13)$. This implies that $N(a, b, c) \leq 2^{\omega(c)+1}$ if $c \equiv 1 \pmod{2}$. Recently, R. Scott and R. Styer [157] improved this: if $c \equiv 1 \pmod{2}$, all solutions (x, y, z) of (1.1) occur in at most two distinct parity classes and each class has at most one solution when solutions exist in two distinct classes, and at most two solutions otherwise. Therefore, we have $N(a, b, c) \leq 2$ if $c \equiv 1 \pmod{2}$. Very recently, combining the upper bound 1.3 with some elementary methods, Y.-Z. Hu and M.-H. Le [70, 71] proved that if $\max\{a, b, c\} > 5 \times 10^{27}$ then $N(a, b, c) \leq 3$. Moreover, if $2 \mid c$ and $\max\{a, b, c\} \geq 10^{62}$, then $N(a, b, c) \leq 2$.

Notice that, for any positive integer k with $k \geq 2$, if $(a, b, c) = (2, 2^k - 1, 2^k + 1)$, then (1.1) has two solutions $(x, y, z) = (1, 1, 1)$ and $(k + 2, 2, 2)$. Hence, as observed by R. Scott and R. Styer [157], there exist infinitely many triples (a, b, c) with $N(a, b, c) = 2$. Therefore, in general, $N(a, b, c) \leq 2$ may be the best upper bound for $N(a, b, c)$.

By combining the results in [71] and [157], one reduces the problem of finding $N(a, b, c) > 2$ to a finite search.

Because computer searches suggest that $N(a, b, c) \leq 1$ except for the known cases of double solutions, there have been a series of conjectures concerning exact upper bounds for $N(a, b, c)$. In the next two sections, we shall introduce these conjectures and related works.

2 Jeśmanowicz conjecture

A positive integer triple (A, B, C) is called a Pythagorean triple if

$$A^2 + B^2 = C^2. \quad (2.1)$$

In particular, if $\gcd(A, B) = 1$, then (A, B, C) is called a primitive Pythagorean triple. By (2.1), every Pythagorean triple (A, B, C) can be expressed as

$$A = A_1n, B = B_1n, C = C_1n, n \in \mathbb{N}, \quad (2.2)$$

where (A_1, B_1, C_1) is a primitive Pythagorean triple. Notice that every primitive Pythagorean triple (A_1, B_1, C_1) satisfies $C_1 \equiv 1 \pmod{2}$, also A_1 and B_1 have opposite parity. We may therefore assume without loss of generality that $A_1 \equiv 1 \pmod{2}$ and $B_1 \equiv 0 \pmod{2}$. So we have

$$A_1 = f^2 - g^2, B_1 = 2fg, C_1 = f^2 + g^2, f, g \in \mathbb{N}, f > g, \gcd(f, g) = 1, fg \equiv 0 \pmod{2} \quad (2.3)$$

(see L. J. Mordell [148, Chapter 3]).

In 1956, L. Jeśmanowicz [75] conjectured that if (A, B, C) is a Pythagorean triple, then the equation

$$A^x + B^y = C^z, x, y, z \in \mathbb{N} \quad (2.4)$$

has only one solution $(x, y, z) = (2, 2, 2)$. By (2.2), (2.3), and (2.4), Jeśmanowicz' conjecture can be rewritten as follows:

Conjecture 2.1. *For any positive integer n , the equation*

$$((f^2 - g^2)n)^x + (2fgn)^y = ((f^2 + g^2)n)^z, x, y, z \in \mathbb{N} \quad (2.5)$$

has only the single solution $(x, y, z) = (2, 2, 2)$, where f, g are positive integers satisfying (2.3).

Jeśmanowicz' conjecture has been proven to be true in many special cases. But, in general, the problem is not solved as of yet.

2.1 Primitive cases

In this subsection we consider Conjecture 2.1 for $n = 1$. Many of the early works on Conjecture 2.1 deal with (2.5) for the case

$$n = 1, f = g + 1. \quad (2.6)$$

When (2.6) holds, Conjecture 2.1 is true if one of the following conditions is satisfied:

- (i) (W. Sierpiński [159]) $g = 1$.

- (ii) (L. Jeśmanowicz [75]) $2 \leq g \leq 5$.
- (iii) (Z. Ke [77], [76]) $g \equiv 1, 3, 4, 5, 7, 9, 10, 11 \pmod{12}$, $g \equiv 2 \pmod{5}$, $g \equiv 3 \pmod{7}$, $g \equiv 4 \pmod{9}$, $g \equiv 5 \pmod{11}$, $g \equiv 6 \pmod{13}$, or $g \equiv 7 \pmod{15}$.
- (iv) (D.-M. Rao [153]) $g \equiv 2, 6 \pmod{12}$.
- (v) (Z. Ke [78], [79], [80]); Z. Ke and Q. Sun [81]) $g \leq 6144$.

In 1965, V. A. Dem'janenko [30] completely solved the case (2.6). Four decades later, Y.-Z. Hu and P.-Z. Yuan [74] gave a new proof of Dem'janenko's result. Moreover, Z. Ke, T. Józefiak, J.-R. Chen, V. D. Podyspanin, Z.-F. Cao, W.-J. Chen, A. Grytczuk and A. Grelak successively discussed some rather special cases of Conjecture 2.1 for $n = 1$. The proofs of the above results are elementary. A detailed record can be found in Z. Ke and Q. Sun [82, Section 7.1], Z.-F. Cao [14, Section 9.2], G. Soydan, M. Demirci, I. N. Cangul, and A. Togbé [160].

For any positive integer t , let $P(t)$ denote the product of the distinct prime divisors of t .

Except for the above mentioned works, the existing results on Conjecture 2.1 for $n = 1$ can be divided into three cases as follows.

Case I. A solution (x, y, z) of (2.5) with $(x, y, z) \neq (2, 2, 2)$ is called exceptional. If $n = 1$, then the exceptional solutions of (2.5) have the following properties:

- (i) (Z. Li [108]) If $x \equiv y \equiv z \equiv 0 \pmod{2}$, then $x \equiv y \equiv z \equiv 2 \pmod{4}$.
- (ii) (T. Miyazaki [134]) If $y \equiv z \equiv 0 \pmod{2}$, then $y \equiv z \equiv 2 \pmod{4}$.
- (iii) (M.-J. Deng and D.-M. Huang [38]) If $f \equiv 2 \pmod{4}$, $g \equiv 3 \pmod{4}$, and $f + g \equiv 1 \pmod{16}$, then $y = 1$.
- (iv) (M.-M. Ma and Y.-G. Chen [125]) If $fg \equiv 2 \pmod{4}$, then $y = 1$.

Case II. If f and g take the following values, then Conjecture 2.1 is true for $n = 1$.

- (i) (W.-D. Lu [117]) $g = 1$.
- (ii) (H.-L. Liu [115]); X.-F. An [1]) $g = 2$ and f is an odd prime power.
- (iii) (N. Terai [183]) $g = 2$.
- (iv) (C.-Y. Fu [42]) $g = 6$ and f is an odd prime power.
- (v) (T. Miyazaki [137]) $f^2 - 2fg - g^2 = \pm 1$.

Case III. If f and g satisfy the following congruence and divisibility conditions, then Conjecture 2.1 is true for $n = 1$.

- (i) (K. Takakuwa [168]) $f \equiv 2 \pmod{4}$ and $g \in \{3, 7, 11, 15\}$.
- (ii) (M.-H. Le [85]) $fg \equiv 2 \pmod{4}$ and $f^2 + g^2$ is an odd prime power.
- (iii) (Y.-D. Guo and M.-H. Le [57]; M.-H. Le [86]) $f \equiv 2 \pmod{4}$, $g \equiv 3 \pmod{4}$, and $f > 81g$.
- (iv) (K Takakuwa and Y. Asaeda [169], [170], [171]) $f \equiv 2 \pmod{4}$, g is an odd prime with $g \equiv 3 \pmod{4}$, and the divisors of $f^2 - g^2$ satisfy certain conditions.
- (v) (M.-J. Deng [31], [32]; J.-H. Wang and M.-J. Deng [190]; M.-J. Deng and G. L. Cohen [36]; T. Miyazaki [131]; S.-Z. Li [106]; Y. An [2]; J.-J. Xing [196]; S.-S. Gou

[49]; C.-Y. Zheng [208]) The divisors of $2fg$ and $f^2 + g^2$ satisfy certain congruence properties.

- (vi) (W.-J. Guan [51]) $(f, g) = (1, 6), (2, 5), (5, 2), (6, 1) \pmod 8$.
- (vii) (T. Miyazaki [133]) $f^2 - g^2 \equiv \pm 1 \pmod{2fg}$ or $f^2 + g^2 \equiv 1 \pmod{2fg}$.
- (viii) (T. Miyazaki, P.-Z. Yuan, and D.-Y. Wu [147]) $f^2 - g^2 \equiv \pm 1 \pmod{P(2fg)}$ or $f^2 + g^2 \equiv \pm 1 \pmod{P(2fg)}$.
- (ix) (Y. Fujita and T. Miyazaki [45]) $2fg \equiv 0 \pmod{2^k}$ and $2fg \equiv \pm 2^k \pmod{(f^2 - g^2)}$, where k is a positive integer with $k \geq 2$.
- (x) (Y. Fujita and T. Miyazaki [46]) $2fg$ has a divisor d with $d \equiv \pm 1 \pmod{(f^2 - g^2)}$.
- (xi) (Q. Han and P.-Z. Yuan [60]) $fg \equiv 2 \pmod 4$ and $f + g$ has a prime divisor p with $p \not\equiv 1 \pmod{16}$.
- (xii) (M.-J. Deng and J. Guo [37]) $g \equiv 2 \pmod 4$ and $n < 600$.
- (xiii) (T. Miyazaki and N. Terai [144]) $f > 72g$, $g \equiv 2 \pmod 4$, and $g/2$ is an odd prime power or a square.
- (xiv) (P.-Z. Yuan and Q. Han [206]) $fg \equiv 2 \pmod 4$, $f > 72g$, and the divisors of f, g satisfy some conditions.
- (xv) (M.-H. Le [102]) $fg \equiv 2 \pmod 4$ and $f > 30.8g$.
- (xvi) (M.-H. Le [100]) $f^2 + g^2 > 4 \times 10^9$ and $\gcd(f^2 + g^2, ((f^2 - g^2)^l - \lambda)/(f^2 + g^2)) = 1$, where l is the least positive integer with $(f^2 - g^2)^l \equiv \lambda \pmod{(f^2 + g^2)}$, $\lambda \in \{1, -1\}$.

2.2 Non-primitive cases

In this subsection we consider Conjecture 2.1 for $n > 1$. There was no study of this problem until 1998. In 1998, M.-J. Deng and G. L. Cohen [35] proved that if $n > 1$, $f^2 - g^2$ is an odd prime power, and either $n \equiv 0 \pmod{P(2fg)}$ or $2fg \not\equiv 0 \pmod{P(n)}$, then (2.5) has only one solution $(x, y, z) = (2, 2, 2)$. One year later, M.-H. Le [88] gave a more general result on Conjecture 2.1 for $n > 1$. He proved that if $n > 1$, then every exceptional solution (x, y, z) of (2.5) satisfies one of the following conditions:

- (i) $\max\{x, y\} > \min\{x, y\} > z$, $f^2 + g^2 \equiv 0 \pmod{P(n)}$, and $P(n) < P(f^2 + g^2)$.
- (ii) $x > z > y$ and $2fg \equiv 0 \pmod{P(n)}$.
- (iii) $y > z > x$ and $f^2 - g^2 \equiv 0 \pmod{P(n)}$.

Sixteen years later, H. Yang and R.-Q. Fu [198] simplified Le's result, showing that the condition (i) can be removed. The above two results are invariably used for solving (2.5) with $n > 1$.

Many works investigated (2.5) in the case

$$n > 1, f = 2^k, g = 1, k \in \mathbb{N}. \quad (2.7)$$

When (2.7) holds, if k takes the following values, then Conjecture 2.1 is true.

- (i) (M.-J. Deng and G. L. Cohen [35]) $k = 1$.
- (ii) (Z.-J. Yang and M. Tang [203]) $k = 2$.

- (iii) (S.-S. Gou and H. Zhang [50]) $k = 3$.
- (iv) (M. Tang and Z.-J. Yang [175]) $k \in \{1, 2, 4, 8\}$.
- (v) (M.-J. Deng [34]) $k \in \{2, 3, 4, 5\}$.
- (vi) (M. Tang and J.-X. Weng [173]) $k = 2^s$, where s is a nonnegative integer.

In about 2014, X.-W. Zhang and W.-P. Zhang [207], and T. Miyazaki [140] independently solved the case (2.7), namely, they proved that if n , f , and g satisfy (2.7), then Conjecture 2.1 is true.

Conjecture 2.1 for $n > 1$ has also been verified in the following cases.

- (i) (Z.-J. Yang and J.-X. Weng [204]) $(f, g) = (6, 1)$.
- (ii) (G. Soydan, M. Demirci, I. N. Cangul, and A. Togbé [160]) $(f, g) = (10, 1)$.
- (iii) (B.-L. Liu [114]) $(f, g) = (12, 1)$.
- (iv) (D.-R. Ling and J.-X. Weng [111]) $(f, g) = (14, 1)$.
- (v) (M.-M. Ma and J.-D. Wu [127]; C.-F. Sun and Z. Cheng [164]) $g = 1$, either $n \equiv 0 \pmod{P(f^2 - 1)}$ or $f = 2p^k$ and $4p^{2k} - 1 \not\equiv 0 \pmod{P(n)}$, where p is an odd prime with $p \equiv 3 \pmod{4}$, k a positive integer.
- (vi) (Z. Cheng, C.-F. Sun, and X.-N. Du [28]) $(f, g) = (5, 2)$.
- (vii) (C.-F. Sun and Z. Cheng [162]; G. Tang [172]) $(f, g) = (7, 2)$.
- (viii) (C.-Y. Fu and M.-J. Deng [43]) $(f, g) = (7, 2)$ or $(49, 2)$.
- (ix) (C.-F. Sun and Z. Cheng [163]) $(f, g) = (9, 2)$.
- (x) (W.-Y. Lu, L. Gao, and H.-F. Hao [118]) $(f, g) = (11, 2)$.
- (xi) (H. Yang, R.-Z. Ren, and R.-Q. Fu [200]; F.-J. Chen [24]) If f is an odd prime and $g = 2$, then (2.5) has no solutions (x, y, z) with $x > z > y$.
- (xii) (Y.-M. Li [107]) $(f, g) = (8, 3)$.
- (xiii) (M.-M. Ma [124]; M.-M. Ma and J.-D. Wu [126]) $(f, g) = (9, 4)$.
- (xiv) (M.-J. Deng and G. L. Cohen [35]) $(f, g) = (3, 2), (4, 3), (5, 4), (6, 5)$.
- (xv) (M.-J. Deng [33]) $(f, g) = (7, 6), (8, 7)$.
- (xvi) (H. Che [23]) $(f, g) = (11, 10)$.
- (xvii) (H.-N. Sun [166]) $(f, g) = (18, 17)$.
- (xviii) (L.-L. Wang [192]) $(f, g) = (20, 19)$.
- (xix) (C.-N. Lin [109]) $(f, g) = (26, 25)$.
- (xx) (J. Ma [123]) $(f, g) = (29, 28)$.
- (xxi) (W.-Y. Lu, L. Gao, X.-H. Wang, and H.-F. Hao [119]) $(f, g) = (46, 45)$.
- (xxii) (H. Yang and R.-Q. Fu [198]) $(f, g) = (2^k, 2^k - 1)$ and $2^k - 1$ is an odd prime, where k is a positive integer.
- (xxiii) (H. Yang and R.-Q. Fu [199]) $(f, g) = (2^k + 1, 2^k)$ and $2^k + 1$ is an odd prime, where k is a positive integer.
- (xxiv) (T.-T. Wang, X.-H. Wang, and Y.-Z. Jiang [193]) If $f = g + 1$ and $c > 500000$, then (2.5) has no solutions (x, y, z) with $y > z > x$.

2.3 A shuffle variant of the Jeśmanowicz conjecture

Let (A_1, B_1, C_1) be a primitive Pythagorean triple with $B_1 \equiv 0 \pmod{2}$. Then A_1 , B_1 , and C_1 can be expressed as in (2.3). In 2011, T. Miyazaki [135] proposed the following conjecture:

Conjecture 2.2. *If $f = g + 1$, then the equation*

$$(f^2 + g^2)^x + (2fg)^y = (f^2 - g^2)^z, x, y, z \in \mathbb{N} \quad (2.8)$$

has only one solution $(x, y, z) = (1, 1, 2)$, otherwise (2.8) has no solutions (x, y, z) .

This is an unsolved problem as well. Up to now, it has been verified in the following cases.

- (i) (T. Miyazaki [135]) $f^2 + g^2 \equiv 1 \pmod{2fg}$.
- (ii) (Z. Rabai [152]) $f^2 + g^2 \equiv 1 \pmod{d}$, where d is the greatest odd divisor of $2fg$.
- (iii) (B.-L. Liu [112]) $(f, g) \equiv (0, 1), (0, 5), (1, 2), (2, 3), (3, 4), (4, 1), (4, 5), (5, 6), (6, 7)$, or $((7, 0) \pmod{8})$.
- (iv) (Q. Feng [40]) $(f, g) = (2^k, 1)$ where k is a positive integer.
- (v) (X.-H. Wang and S. Gou [194]) $(f, g) = (2^k, p)$, where k is a positive integer and p is an odd prime.

3 Terai-Jeśmanowicz conjecture

Let p, q, r be fixed positive integers such that

$$a^p + b^q = c^r, \min\{p, q, r\} > 1. \quad (3.1)$$

In 1994 N. Terai [176] proposed an important generalization on Jeśmanowicz' conjecture as follows:

Conjecture 3.1. *If a, b, c, p, q, r satisfy (3.1), then (1.1) has only one solution $(x, y, z) = (p, q, r)$.*

In 1999 Z.-F. Cao [15] showed that Conjecture 3.1 is clearly false. He suggested that the condition $\max\{a, b, c\} > 7$ should be added to the hypotheses of Conjecture 3.1, and used the term "Terai-Jeśmanowicz conjecture" for the resulting statement. In the same year, N. Terai [179] gave a similar modification for Conjecture 3.1. However, Z.-F. Cao and X.-L. Dong [17], and M.-H. Le [93], [94] independently found infinitely many counterexamples to the Terai-Jeśmanowicz conjecture. Therefore, they stated the following conjecture:

Conjecture 3.2. *If a, b, c, p, q, r satisfy (3.1), then the only solution to (1.1) with $\min\{x, y, z\} > 1$ is $(x, y, z) = (p, q, r)$.*

A history of the various versions of this conjecture is also given in [160] along with a history of the Jeśmanowicz' conjecture itself. We clarify a potentially confusing point in the history of the various versions of Conjecture 3.2 as given in [160]: there it is stated that [93] corrects [17], whereas [93] actually corrects [15]; [17] also corrects [15], and gives a version of the conjecture identical to that given in [93], which is Conjecture 3.2; in [160], Conjecture 3.2 is accidentally given twice (as Conjectures 4.3 and 4.4).

The existing results concerning (1.1) all apply to Conjecture 3.2. But, in general, Conjecture 3.2 is not yet solved. In 2015, by using the upper bound (1.2) with some elementary methods, Y.-Z. Hu and M.-H. Le [69] proved that if $\max\{a, b, c\}$ is large enough and a, b, c satisfy certain divisibility conditions, then (1.1) has at most one solution (x, y, z) with $\min\{x, y, z\} > 1$.

Recently, R. Scott and R. Styer [157] proposed a more precise conjecture as follows.

Conjecture 3.3. *Let $a < b$ with $\gcd(a, b) = 1$. Then we have $N(a, b, c) \leq 1$, except for the following cases:*

- (i) $(a, b, c) = (3, 5, 2)$, $(x, y, z) = (1, 1, 3)$, $(3, 1, 5)$, and $(1, 3, 7)$.
- (ii) $(a, b, c) = (3, 13, 2)$, $(x, y, z) = (1, 1, 4)$ and $(5, 1, 8)$.
- (iii) $(a, b, c) = (3, 13, 4)$, $(x, y, z) = (1, 1, 2)$ and $(5, 1, 4)$.
- (iv) $(a, b, c) = (3, 13, 16)$, $(x, y, z) = (1, 1, 1)$ and $(5, 1, 2)$.
- (v) $(a, b, c) = (3, 13, 2200)$, $(x, y, z) = (1, 3, 1)$ and $(7, 1, 1)$.
- (vi) $(a, b, c) = (2, 3, 11)$, $(x, y, z) = (1, 2, 1)$ and $(3, 1, 1)$.
- (vii) $(a, b, c) = (2, 3, 35)$, $(x, y, z) = (3, 3, 1)$ and $(5, 1, 1)$.
- (viii) $(a, b, c) = (2, 3, 259)$, $(x, y, z) = (4, 5, 1)$ and $(8, 1, 1)$.
- (ix) $(a, b, c) = (2, 5, 3)$, $(x, y, z) = (2, 1, 2)$ and $(1, 2, 3)$.
- (x) $(a, b, c) = (2, 5, 133)$, $(x, y, z) = (3, 3, 1)$ and $(7, 1, 1)$.
- (xi) $(a, b, c) = (2, 7, 3)$, $(x, y, z) = (1, 1, 2)$ and $(5, 2, 4)$.
- (xii) $(a, b, c) = (2, 89, 91)$, $(x, y, z) = (1, 1, 1)$ and $(13, 1, 2)$.
- (xiii) $(a, b, c) = (2, 91, 8283)$, $(x, y, z) = (1, 2, 1)$ and $(13, 1, 1)$.
- (xiv) $(a, b, c) = (3, 4, 259)$, $(x, y, z) = (1, 4, 1)$ and $(5, 2, 1)$.
- (xv) $(a, b, c) = (3, 10, 13)$, $(x, y, z) = (1, 1, 1)$ and $(7, 1, 3)$.
- (xvi) $(a, b, c) = (3, 16, 259)$, $(x, y, z) = (1, 2, 1)$ and $(5, 1, 1)$.
- (xvii) $(a, b, c) = (2, 2^k - 1, 2^k + 1)$, $(x, y, z) = (1, 1, 1)$ and $(k + 2, 2, 2)$ where k is a positive integer with $k \geq 2$.

Obviously, Conjecture 3.3 contains Conjecture 3.2 and Conjecture 2.1 for $n = 1$. Undoubtedly, we can look upon Conjecture 3.3 as the ultimate aim of solving (1.1). Although the known results concerning (1.1) show Conjecture 3.3 to be true in many cases, the complete resolution of the conjecture is a very difficult problem.

3.1 Earlier Results

Since the 1950s, methods that arise in elementary number theory and algebraic number theory have been used to find all solutions (x, y, z) of (1.1) for the following small values of a, b, c with $a < b$.

- (i) (T. Nagell [149]) $\max\{a, b, c\} \leq 7$.
- (ii) (A. Makowski [129]) $(a, b, c) = (2, 11, 5)$.
- (iii) (T. Hadano [59]) $11 \leq \max\{a, b, c\} \leq 17$ and $(a, b, c) \neq (3, 13, 2)$.
- (iv) (S. Uchiyama [188]) $(a, b, c) = (3, 13, 2)$.
- (v) (Z.-F. Cao and C.-S. Cao [16]) $\max\{a, b, c\} = 14$.
- (vi) (J.-H. Ren and J.-K. Zhang [154]) $14 \leq \max\{a, b, c\} \leq 16$.
- (vii) (Q. Sun and X.-M. Zhou [167]) $\max\{a, b, c\} = 19$.
- (viii) (X.-Z. Yang [202]) $\max\{a, b, c\} = 23$.
- (ix) (Z.-F. Cao [11], [13]) a, b, c distinct primes with $29 \leq \max\{a, b, c\} \leq 97$.
- (x) (X.-G. Guan [53], [54]) $a = 2$, b and c are distinct odd primes with $100 < \max\{b, c\} < 150$.
- (xi) (Z.-F. Cao, R.-Z. Tong, and Z.-J. Wang [19]) $a = 2$, b and c are distinct odd primes with $100 < \max\{b, c\} < 200$.
- (xii) (X.-E. Zhou [209]) $c = 2$, a and b are distinct odd primes with $200 < \max\{a, b\} < 300$.

Note that Theorem 2 of R. Scott [155] gives a general method for finding all solutions to (1.1) for a given (a, b, c) when c is odd or $c = 2$.

Using the above mentioned results (i)-(ix), Z.-F. Cao [14] conjectured that if a, b, c are distinct primes with $\max\{a, b, c\} > 7$, then $N(a, b, c) \leq 1$. But, owing to the fact that S. Uchiyama [188] missed the solution $(x, y, z) = (5, 1, 8)$ for $(a, b, c) = (3, 13, 2)$, Cao's conjecture is false. In 1985, under the parity classification of solutions (x, y, z) of (1.1), M.-H. Le [83] proved that if a, b, c are distinct primes with $a = 2$, then every class has at most one solution (x, y, z) . In the same paper, he proved that if $c = 2$ and a, b , are distinct odd primes with $\max\{a, b\} > 13$, then $N(a, b, c) \leq 1$. Using a different method, R. Scott [155] obtained similar results for a and b not necessarily prime.

Let $a = 2$ and let b, c be a pair of twin primes with $2 + b = c$. Q. Sun and X.-M. Zhou [167] proved that if $b > 3$ and $b^2 + b + 1$ is an odd prime with $c^{b+1} \not\equiv 1 \pmod{b^2 + b + 1}$, then (1.1) has only one solution $(x, y, z) = (1, 1, 1)$. Afterwards, Z.-F. Cao [12], H. Yang and R.-Q. Fu [197] independently solved this case. They proved that if $a = 2$ and b, c are twin primes with $c > b > 3$, then (1.1) has only one solution $(x, y, z) = (1, 1, 1)$.

Let k be a fixed positive integer. H. Edgar asked how many solutions (y, z) there are to the equation

$$c^z - b^y = 2^k, y, z \in \mathbb{N} \quad (3.2)$$

for fixed distinct odd primes b and c (see R. K. Guy [58, Problem D9]). In 1987, Z.-F. Cao and D.-Z. Wang [20], using an earlier result of Z.-F. Cao [10], proved that (3.1).

has at most two solutions (y, z) . In 2004, R. Scott and R. Styer [156] improved this to at most one solution (y, z) . In this last result, we can allow b composite if, instead of using the result from [10], we use a more general (but non-elementary) result in which b is not restricted to be prime. Such a result has been obtained independently by three different methods: the most direct approach is that of F. Luca [120], who uses results of Y. Bilu, G. Hanrot, and P. M. Voutier [7]; a second proof was given by M.-H. Le [92], using linear forms in logarithms and several auxiliary lemmas; the result can also be obtained by considering the equation $x^n + 2^\alpha y^n = Cz^2$, handled by M. A. Bennett and C. Skinner [4, Theorem 1.2] using (in the words of those authors) “combinations of every technique we have currently available.” This last result is often useful in treating other exponential diophantine equations.

M. Perisastri [151], M. Toyozumi [187], M.-H. Le [84], Z.-F. Cao [9] and [10], Y.-S. Cao [8], J.-M. Chen [25] gave certain solvability conditions to (1.1) for $a = 2$ or $2l$ where l is an odd prime.

3.2 Generalized primitive Pythagorean triples

For $p = q = 2$ and $r > 2$, by (3.1), we have

$$a^2 + b^2 = c^r. \tag{3.2}$$

Then (a, b, c) is called a generalized primitive Pythagorean triple. Since $\gcd(a, b) = 1$ and $r > 2$, we see from (3.2) that $c \equiv 1 \pmod 4$ and a, b have opposite parity. We may therefore assume without loss of generality that $a \equiv 0 \pmod 2$ and $b \equiv 1 \pmod 2$. By L. J. Mordell [148, Section 15.2], every generalized primitive Pythagorean triple (a, b, c) can be expressed as

$$a = f \left| \sum_{i=0}^{(r-1)/2} \binom{r}{2i} f^{r-2i-1} (-g^2)^i \right|, b = g \left| \sum_{i=0}^{(r-1)/2} \binom{r}{2i+1} f^{r-2i-1} (-g^2)^i \right|, c = f^2 + g^2, \\ f, g \in \mathbb{N}, \gcd(f, g) = 1, f \equiv 0 \pmod 2 \tag{3.4}$$

or

$$a = fg \left| \sum_{i=0}^{r/2-1} \binom{r}{2i+1} f^{r-2i-2} (-g^2)^i \right|, b = \left| \sum_{i=0}^{r/2-1} \binom{r}{2i} f^{r-2i} (-g^2)^i \right|, c = f^2 + g^2, \\ f, g \in \mathbb{N}, \gcd(f, g) = 1, fg \equiv 0 \pmod 2, \tag{3.5}$$

according as $r \equiv 1 \pmod 2$ or not.

Many recent works on (1.1) concern generalized primitive Pythagorean triples (a, b, c) . These results can be divided into the following:

Case I. If $r \in \{3, 5, 7, 9\}$ and f, g satisfy one of the following conditions, then (1.1) has only one solution $(x, y, z) = (2, 2, r)$.

(i) (N. Terai [176]; M.-H. Le [87]) $r = 3, g = 1$, and f satisfies certain congruence and divisibility conditions.

(ii) (M.-H. Le [91]) $r = 3, g = 1, f \equiv 2 \pmod 4$, and $f \geq 206$.

- (iii) (Z.-F. Cao and X.-L. Dong [18]) $r = 3$ and $g = 1$.
- (iv) (M.-H. Le [98]; Y.-Z. Hu and P.-Z. Yuan [73]) $r = 3$ and b is an odd prime power.
- (v) (N. Terai [177]; X.-L. Dong and Z.-F. Cao [39]) $r = 5$, $g = 1$, and b is an odd prime.
- (vi) (M.-H. Le [96]) $r = 5$, $g = 1$, and $f \geq 542$.
- (vii) (Z.-F. Cao and X.-L. Dong [18]; Y.-Z. Hu and P.-Z. Yuan [72]) $r = 5$ and $g = 1$.
- (viii) (J.-P. Chen [26]) $r = 7$, $g = 1$, and $f \geq 2.4 \times 10^9$.
- (ix) (X.-G. Guan [55]) $r = 7$, $g = 1$, and b is an odd prime. $r = 9$, $g = 1$, and $f > 2863$.

Case II. If $g = 1$, a , b , f , r satisfy one of the following conditions, then (1.1) has only one solution $(x, y, z) = (2, 2, r)$.

- (i) (N. Terai [178]; N. Terai and K. Takakuwa [186]) r is an odd prime, $a \equiv 2 \pmod{4}$, and b is an odd prime with $b \equiv 3 \pmod{4}$.
- (ii) (Z.-F. Cao and X.-L. Dong [17]) $r \equiv 1 \pmod{2}$, $a \equiv 2 \pmod{4}$, and b is an odd prime with $b \equiv 3 \pmod{4}$.
- (iii) (Z.-F. Cao [15]) $r \equiv 1 \pmod{2}$, $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{4}$, and c is an odd prime.
- (iv) (M.-H. Le [97]) $r \equiv 1 \pmod{2}$, $a \equiv 2 \pmod{4}$, $f > 41r^{3/2}$, and c is an odd prime.
- (v) (Z.-F. Cao and X.-L. Dong [18]) $r \equiv 1 \pmod{2}$, $a \equiv 2 \pmod{4}$, $f \geq 200$, and $f/\sqrt{825 \log(f^2 + 1) - 1} > r$.
- (vi) (J. Y. Xia and P.-Z. Yuan [195]) $r \equiv 1 \pmod{2}$, $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{4}$, and $f/\sqrt{1.5 \log_3(f^2 + 1) - 1} > r$.
- (vii) (M. Cipu and M. Mignotte [29]) $r \equiv 1 \pmod{2}$, $a \equiv 2 \pmod{4}$, and $b \equiv 3 \pmod{4}$, except for finitely many tetrads (a, b, c, r) with $r < 770$.
- (viii) (M.-H. Le [99]) $r \equiv 5 \pmod{8}$ and

$$f > \begin{cases} r^2, & \text{if } r < 11500, \\ \frac{2r}{\pi}, & \text{otherwise.} \end{cases}$$

- (ix) (B. He, A. Togbé, and S.-C. Yang [63]) r is an odd prime with $r \equiv 5 \pmod{8}$ or $r \equiv 19 \pmod{24}$, $f > 2r/\pi$.
- (x) (M.-H. Le [101]) $r \equiv 1 \pmod{2}$ and $f > 10^6 r^6$.
- (xi) (T. Miyazaki [132], [136]) $r \equiv 4$ or $6 \pmod{8}$, $f^2 \log(2)/\log(f^2 + 1) \geq r^3$.
- (xii) (X.-G. Guan [56]) $r \equiv 0 \pmod{2}$ and $f > 48400r^2(\log(r))^2$.
- (xiii) (M.-H. Le, A. Togbé, and H.-L. Zhu [104]) $f > \max\{10^{15}, 2r^3\}$.

In 2012, F. Luca [121] basically solved this case. He proved that if (a, b, c) is a generalized primitive Pythagorean triple with $g = 1$, then (1.1) has only one solution $(x, y, z) = (2, 2, r)$, except for finitely many pairs (r, f) . Two years later, T. Miyazaki

[139] showed that the exceptional cases (r, f) of Luca's result satisfy $r < 10^{74}$ and $f < 10^{34}$.

Similarly, T. Miyazaki and F. Luca [142] proved that if $p \equiv 2 \pmod{4}$, $q = r = 2$, and $a = f^2 - 1$ in (3.1), where f is a fixed positive integer with $f \equiv 0 \pmod{2}$, then (1.1) has only one solution $(x, y, z) = (p, 2, 2)$, except for finitely many pairs (p, f) .

Case III. If $r \equiv 1 \pmod{2}$, and if a, b , and c satisfy one of the following conditions, then (1.1) has only one solution $(x, y, z) = (2, 2, r)$.

(i) (M.-H. Le [90]) $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{8}$, and c is an odd prime power.

(ii) (N. Terai [179]) $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{8}$, $b \geq 41a$, and $\left(\frac{a}{b}\right) = -1$ where $\left(\frac{a}{b}\right)$ is the Jacobi symbol.

(iii) (N. Terai [179]) $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{8}$, $b \geq 30a$, and b has an odd prime divisor l with $\left(\frac{a}{l}\right) = -1$.

(iv) (Q. Feng and D. Han [41]) $r \equiv \pm 3 \pmod{8}$, $f > 2r/\pi$, and b is an odd prime.

(v) (Z.-F. Cao and X.-L. Dong [17]) $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{4}$, and $b \geq 25.1a$.

(vi) (M.-H. Le [93]) $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{4}$, and $\frac{b}{a} > 1/\sqrt{e^{r/1856} - 1}$. This implies that $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{4}$, and either $r \geq 1287$ or $b \geq 25a$.

(vii) (M.-H. Le [96]) $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{4}$, $r > 7200$, and $c > 3 \times 10^{37}$.

(viii) (J.-Y. Xia and P.-Z. Yuan [195]) $a \equiv 2 \pmod{4}$, $b \equiv 3 \pmod{4}$, $b > a$, and $c - 1$ is not a square.

3.3 Miscellaneous Results

In this subsection we introduce the other works on (1.1). Let $(p, q, r) = (1, 1, 1)$ in (3.1). In 2012, T. Miyazaki and A. Togbé [145] proved that, if $a = 2$, then (1.1) has only one solution $(x, y, z) = (1, 1, 1)$, except for $b = 89$. Afterwards, M. Tang and Q.-H. Yang [174] discussed a general equation as follows:

$$(2n)^x + (bn)^y = ((b+2)n)^z, x, y, z \in \mathbb{N}, \quad (3.3)$$

where b is an odd integer with $b > 1$. They proved that if (x, y, z) is a solution of (3.3) with $(x, y, z) \neq (1, 1, 1)$, then either $x > z \geq y$ or $y > z > x$. Obviously, for any b , (3.3) has solutions $(x, y, z, n) = (1, 1, 1, t)$ and $(3, 2, 2, (b+1)/2)$, where t is an arbitrary positive integer. In 2014, Y.-H. Yu and Z.-P. Li [205] proved that, for any b , (3.3) has only finitely many solutions (x, y, z, n) with $x > z = y$. In the same year, W.-J. Guan and S. Che [52] found all solutions (x, y, z, n) of (3.3) with $x > z = y$ for $b \not\equiv 7 \pmod{8}$. They proved that if $b \not\equiv 7 \pmod{8}$, then (3.3) with $x > z = y$ has only the solutions

$$(x, y, z, n) = \begin{cases} (3, 2, 2, \frac{b+1}{2}) & \text{if } b \equiv 1 \pmod{4}, \\ (3, 2, 2, \frac{b+1}{2}) \text{ and } (5, 4, 4, \frac{b+1}{4}(b^2 + 2b + 2)) & \text{if } b \equiv 3 \pmod{4} \text{ and } \frac{b+1}{4} \text{ is not a square,} \\ (3, 2, 2, \frac{b+1}{2}), (4, 2, 2, \sqrt{\frac{b+1}{4}}), \text{ and } (5, 4, 4, \frac{b+1}{4}(b^2 + 2b + 2)) & \text{otherwise.} \end{cases}$$

Moreover, T. Miyazaki, A. Togbé, and P.-Z. Yuan [146] deal with the equation

$$a^x + b^y = (b + 2)^z, x, y, z \in \mathbb{N} \quad (3.4)$$

for $a \neq 2$ and $\gcd(a, b) = 1$. They found all solutions (x, y, z) of (3.4) with $b \equiv -1 \pmod{a}$. Similarly, M.-H. Zhu and X.-X. Li [210] proved that if $a = 4$ and $c = b + 4$, then (1.1) has only one solution $(x, y, z) = (1, 1, 1)$.

Since $(p, q, r) = (1, 1, 1)$ in (3.1), we have $a + b = c$. Hence, for any positive integer n , the equation

$$(an)^x + (bn)^y = (cn)^z, x, y, z \in \mathbb{N} \quad (3.5)$$

has the solution $(x, y, z) = (1, 1, 1)$. A solution (x, y, z) of (3.5) with $(x, y, z) \neq (1, 1, 1)$ is called exceptional. Very recently, C.-F. Sun and M. Tang [165] showed that if $\min\{a, b\} > 2$ and (x, y, z) is an exceptional solution of (3.5), then either $x > z > y$ or $y > z > x$. In the same paper, they proved that if $(a, b) = (3, 5), (5, 8), (8, 13),$ or $(13, 21)$, then (3.5) has no exceptional solutions. In this respect, P.-Z. Yuan and Q. Han [206] proposed the following conjecture:

Conjecture 3.4. *If $a + b = c$ and $\min\{a, b\} \geq 4$, then (3.5) has no exceptional solutions (x, y, z) .*

In [206], they proved that if a and b are squares with $b \equiv 4 \pmod{8}$, then (3.5) has no solutions (x, y, z) with $y > z > x$, in particular, if $b = 4$, then Conjecture 3.4 is true. Very recently, M.-H. Le [103] proved that if a and b are squares with $a > 64b^3$, then (3.5) has no solutions (x, y, z) with $x > z > y$, in particular, if $b \equiv 4 \pmod{8}$, then Conjecture 3.4 is true.

Let $(p, q, r) = (2, k, 1)$ in (3.1), where k is an odd positive integer. If $a \equiv -1 \pmod{b^{k+1}}$, $b \equiv c \equiv 1 \pmod{2}$, and b, c, k satisfy one of the following conditions, then (1.1) has only one solution $(x, y, z) = (2, k, 1)$.

- (i) (N. Terai [180]) $k \in \{1, 3\}$ and b is an odd prime with $b \equiv 3 \pmod{4}$.
- (ii) (M.-H. Le [95]) $b \equiv 3 \pmod{4}$.
- (iii) (B.-L. Liu [113]) $b \equiv 5 \pmod{24}$.

In 2006, M.-Y. Lin [110] completely solved this case. He proved that if $(p, q, r) = (2, k, 1)$, $a \equiv -1 \pmod{b^{k+1}}$, and $b \equiv c \equiv k \equiv 1 \pmod{2}$, then (1.1) has only one solution $(x, y, z) = (2, k, 1)$.

Similarly, Z.-W. Liu [116] proved that if $(p, q, r) = (1, k, 2)$ where k is an odd positive integer, $b \equiv 5 \pmod{12}$, and c is an odd prime with $c \equiv -1 \pmod{b^{k+1}}$, then (1.1) has only one solution $(x, y, z) = (1, k, 2)$.

Let $(a, b, c) = (f, f + 1, f + 2)$, where f is a fixed positive integer with $f > 1$. By the result of W. Sierpiński [159], if $f = 3$, then (1.1) has only one solution $(x, y, z) = (2, 2, 2)$. In this respect, B. Leszczyński [105] and A. Makowski [130] gave some conditions for (1.1) to have solutions. In 2009, B. He and A. Togbé [61]

completely solved this case. They proved that if $f > 3$, then (1.1) has no solutions (x, y, z) .

Let $(a, b, c) = (f, 3f^2 - 1, 4f^2 - 1)$, where f is a fixed positive integer with $f > 1$. In 2007, Y.-Z. Hu [68] proved that (1.1) has only the solution $(x, y, z) = (2, 1, 1)$ with $x \equiv 0 \pmod{2}$. Afterwards, L.-M. Chen [27], and B. He and A. Togbé [62] independently showed that $(x, y, z) = (2, 1, 1)$ is the unique solution of (1.1).

Similarly, Y.-Z. Hu [67] proved that if $(a, b, c) = (8f^3 - 3f, 3f^2 - 1, 4f^2 - 1)$, where f is a fixed positive integer with $f > 3$, then (1.1) has only the solution $(x, y, z) = (2, 1, 3)$ with $x \equiv 0 \pmod{2}$. B. He and S.-C. Yang [64] showed that the solution $(x, y, z) = (2, 1, 3)$ is unique. Moreover, S.-C. Yang and B. He [201] proved that if $(a, b, c) = (8f^3 + 3f, 3f^2 + 1, 4f^2 + 1)$, where f is a fixed positive integer, then (1.1) has only one solution $(x, y, z) = (2, 1, 3)$.

Let $(a, b, c) = (2f, 2fg - 1, 2fg + 1)$, where f, g are fixed positive integers with $\min\{f, g\} > 1$. In 2012, T. Miyazaki and A. Togbé [145] proved that (1.1) has solutions (x, y, z) if and only if either $f = 2g$ or $g = f^2$. Further, (1.1) has only one solution $(x, y, z) = (2, 2, 2)$ if $f = 2g$ or $(3, 2, 2)$ if $g = f^2$.

Let $(a, b, c) = (fn^2 + 1, (g^2 - f)n^2 - 1, gn)$, where f, g, n are fixed positive integers. If f, g , and n satisfy one of the following conditions, then (1.1) has only one solution $(x, y, z) = (1, 1, 2)$.

- (i) (N. Terai [182]) $(f, g) = (4, 3)$, $n \leq 20$ or $n \not\equiv 3 \pmod{6}$.
- (ii) (J.-P. Wang, T.-T. Wang, and W.-P. Zhang [191]) $(f, g) = (4, 3)$ and $n \not\equiv 0 \pmod{3}$.
- (iii) (L.-J. Su and X.-X. Li [161]) $(f, g) = (4, 3)$, $n > 90$, and $n \equiv 3 \pmod{6}$.
- (iv) (C. Bertók [5]) $(f, g) = (4, 5)$, $20 < n \leq 90$, and $n \equiv 3 \pmod{6}$.
- (v) (N. Terai and T. Hibino [184]) $(f, g) = (12, 5)$, $n \not\equiv 17$ or $33 \pmod{40}$.
- (vi) (N. Terai and T. Hibino [185]) $(f, g) = (3l, l)$, where l is an odd prime with $l < 3784$ and $l \equiv 1 \pmod{4}$, $n \not\equiv 0 \pmod{3}$, and $n \equiv 1 \pmod{4}$.
- (vii) (T. Miyazaki and N. Terai [143]) $f = 1$, $g \equiv \pm 3 \pmod{8}$, $n \equiv \pm 1 \pmod{g}$, and $(g, n) \neq (3, 1)$.
- (viii) (X.-W. Pan [150]; R.-Q. Fu and H. Yang [44]) $f \equiv 4$ or $5 \pmod{8}$, $g \equiv 1 \pmod{2}$, $n > 6g^2 \log g$, and $\left(\frac{f+1}{g}\right) = -1$ where $\left(\frac{*}{*}\right)$ is the Jacobi symbol.

Let $F = \{F_m\}_{m=0}^\infty$ be the Fibonacci sequence, so we have

$$F_0 = 0, F_1 = 1, F_{m+2} = F_{m+1} + F_m, m \geq 0.$$

For any positive integer t with $t \geq 2$, if the sequence $F^{(t)} = \{F_m^{(t)}\}$ satisfies

$$F_0^{(t)} = \dots = F_{t-1}^{(t)} = 0, F_t^{(t)} = 1, F_{m+t+1}^{(t)} = F_{m+t}^{(t)} + \dots + F_m^{(t)}, m \geq 0,$$

then $F^{(t)}$ is called the t -generalized Fibonacci sequence. Correspondingly, F_m and $F_m^{(t)}$ are called Fibonacci numbers and t -generalized Fibonacci numbers, respectively. In 2002, N. Terai [181] asked if (1.1) has only one solution $(x, y, z) = (2, 2, 1)$ for

$(a, b, c) = (F_k, F_{k+1}, F_{2k+1})$, where k is a positive integer with $k \geq 3$. Thirteen years later, this question was answered by T. Miyazaki [141]. In addition, he proved the further result that if $(a, b, c) = (F_k, F_{2k+2}, F_{k+2})$, where k is a positive integer with $k \geq 3$, then (1.1) has only one solution $(x, y, z) = (2, 1, 2)$. Moreover, F. Luca and R. Oyono [122], A. P. Chaves and D. Marques [21] and [22], C. A. Gómez Ruiz and F. Luca [48], N. Hirata-Kohno and F. Luca [66] deal with (1.1) for some special Fibonacci numbers and generalized Fibonacci numbers a , b , and c .

3.4 A shuffle variant of the Terai-Jeśmanowicz conjecture

Finally, we introduce a shuffle variant of the Terai-Jeśmanowicz conjecture proposed by T. Miyazaki [138] as follows:

Conjecture 3.5. *Let a, b, c, p, q, r be fixed positive integers satisfying (3.1). Further, let $a < b$, $(a, b, c) \neq (2, 7, 3)$ and $(a, b, c) \neq (2, 2^k - 1, 2^k + 1)$, where k is a positive integer. If $q = r = 2$ and $b + 1 = c$, then the equation*

$$c^X + b^Y = a^Z, X, Y, Z \in \mathbb{N} \quad (3.8)$$

has only one solution $(X, Y, Z) = (1, 1, p)$. Otherwise, (3.8) has no solutions (X, Y, Z) .

In the same paper, T. Miyazaki proved that Conjecture 3.5 is true if $q = r = 2$ and $b + 1 = c$. In general, this problem has not been solved yet.

Acknowledgement. The authors thank Prof. P.-Z. Yuan for his valuable suggestions and recent papers.

References

- [1] X.-F. An. *On Jeśmanowicz' conjecture concerning Pythagorean numbers*. Master's thesis, Chongqing: Chongqing Normal Univ., 2015. (in Chinese).
- [2] Y. An. *On Jeśmanowicz' conjecture concerning Pythagorean numbers*. Master's thesis, Chongqing: Southwest Univ., 2014. (in Chinese).
- [3] M. A. Bennett and N. Billerey. *Sums of two S -units via Frey-Hellegouarch curves*. *Math. Comp.*, **86(305)**, (2017), 1375–1401. [MR3614021](#). [Zbl 06693286](#).
- [4] M. A. Bennett and C. Skinner. *Ternary diophantine equations via Galois representations and modular forms*. *Canad. J. Math.*, **56(1)**, (2004), 23–54. [MR2031121](#). [Zbl 1053.11025](#).
- [5] C. Bertók. *The complete solution of the diophantine equation $(4m^2 + 1)^x + (5m^2 - 1)^y = (3m)^z$* . *Period. Math. Hung.*, **72(1)**, (2016), 37–42. [MR3470802](#). [Zbl 1389.11085](#).

- [6] F. Beukers and H. P. Schlickewei. *The equation $x + y = 1$ in finitely generated groups*. Acta Arith., **78**(2), (1996), 189–199. [MR3470802](#). [Zbl 0880.11034](#).
- [7] Y. Bilu, G. Hanrot, and P. M. Voutier. *Existence of primitive divisors of Lucas and Lehmer numbers, with an appendix by M. Mignotte*. J. Reine Angew. Math., **539**, (2001), 75–122. [MR1863855](#). [Zbl 0995.11010](#).
- [8] Y.-S. Cao. *On the equation $a^x - b^y = (2p)^z$* . Northeast Math. J., **5**(4), (1989), 477–484. (in Chinese). [MR1053528](#). [Zbl 0713.11024](#).
- [9] Z.-F. Cao. *The equation $a^x - b^y = (2p^s)^z$ and Hugh Edgar's problem*. Chinese Sci. Bull., **30**(14), (1985), 1116–1117. (in Chinese). [MR0871578](#).
- [10] Z.-F. Cao. *The equation $x^2 + 2^m = y^n$ and Hugh Edgar's problem*. Chinese Sci. Bull., **31**(7), (1986), 555–556. (in Chinese).
- [11] Z.-F. Cao. *On the diophantine equation $a^x + b^y = c^z$ I*. Chinese Sci. Bull., **31**(22), (1986), 1688–1690. (in Chinese). [MR0886632](#).
- [12] Z.-F. Cao. *Exponential diophantine equations of the form $a^x + b^y = c^z$* . J. Harbin Inst. Tech., **4**, (1987), 113–121. (in Chinese). [MR0951789](#). [Zbl 0971.11517](#).
- [13] Z.-F. Cao. *On the diophantine equation $a^x + b^y = c^z$ II*. Chinese Sci. Bull., **33**(3), (1988), 237. (in Chinese).
- [14] Z.-F. Cao. *Introduction to diophantine equations*. Harbin Inst. Tech. Press, Harbin, 1989. (in Chinese). [Zbl 0849.11029](#).
- [15] Z.-F. Cao. *A note on the diophantine equation $a^x + b^y = c^z$* . Acta Arith., **91**(1), (1999), 85–93. [MR1726477](#). [Zbl 0946.11009](#).
- [16] Z.-F. Cao and C.-S. Cao. *On the equation $a^x + b^y = c^z$* . Pure Appl. Math., **4**(4), (1988), 98–100. (in Chinese). [Zbl 0921.11017](#).
- [17] Z.-F. Cao and X.-L. Dong. *On the Terai-Jeśmanowicz conjecture*. Publ. Math. Debrecen, **61**(3–4), (2002), 253–265. [MR1943694](#). [Zbl 1012.11025](#).
- [18] Z.-F. Cao and X.-L. Dong. *An application of a lower bound for linear forms in two logarithms to the Terai-Jeśmanowicz conjecture*. Acta Arith., **110**(2), (2003), 153–164. [MR2008082](#). [Zbl 1031.11020](#).
- [19] Z.-F. Cao, R.-Z. Tong, and Z.-J. Wang. *A conjecture on the exponential diophantine equations*. Chinese J. Nature, **14**(11), (1991), 872–873. (in Chinese).
- [20] Z.-F. Cao and D.-Z. Wang. *On Hugh Edgar's problem*. Chinese Sci. Bull., **32**(14), (1987), 1043–1046. [MR0940140](#).

- [21] A. P. Chaves and D. Marques. *A diophantine equation related to the sum of consecutive k -generalized Fibonacci numbers*. *Fibonacci Quart.*, **52**(1), (2014), 70–74. [MR3181100](#). [Zbl 1290.11021](#).
- [22] A. P. Chaves and D. Marques. *A diophantine equation related to the sum of powers of two consecutive generalized Fibonacci numbers*. *J. Number Theory*, **156**(1), (2015), 1–14. [MR3360324](#). [Zbl 1395.11031](#).
- [23] H. Che. *On the diophantine equation $(21n)^x + (220n)^y = (221n)^z$* . Master's thesis, Chongqing: Southwest Univ., 2011. (in Chinese).
- [24] F.-J. Chen. *On the diophantine equation $(na)^x + (nb)^y = (nc)^z$* . *Adv. Math. China*, **47**(3), (2018), 388–392. [Zbl 07028020](#).
- [25] J.-M. Chen. *On the diophantine equation $a^x + b^y = c^z$* . *J. Zhejiang Normal Univ., Nat. Sci.*, **17**(4), (1994), 17–20. (in Chinese).
- [26] J.-P. Chen. *The positive integer solutions of the diophantine equation $a^x + b^y = c^z$* . *J. Hainan Univ., Nat. Sci.*, **30**(4), (2012), 309–315. (in Chinese).
- [27] L.-M. Chen. *The odd integer solutions of the equation $n^x + (3n^2 - 1)^y = (4n^2 - 1)^z$* . *Adv. Math. Beijing*, **39**(4), (2010), 507–511. [MR2760537](#).
- [28] Z. Cheng, C.-F. Sun, and X.-N. Du. *On the diophantine equation $(20n)^x + (21n)^y = (29n)^z$* . *Math. Appl.*, **26**(1), (2013), 129–133. [MR3077060](#). [Zbl 1274.11088](#).
- [29] M. Cipu and M. Mignotte. *Bounds for counterexamples to Terai's conjecture*. *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)*, **53**(101)(3), (2010), 231–237. [MR2732411](#). [Zbl 1212.11046](#).
- [30] V. A. Dem'janenko. *On Jeśmanowicz' problem for Pythagorean numbers*. *Izv. Vysš Učebn. Zayed. Mat.*, **48**(1), (1965), 52–56. (in Russian). [MR0191865](#). [Zbl 0166.05103](#).
- [31] M.-J. Deng. *On Jeśmanowicz' conjecture*. *J. Harbin Inst. Tech.*, **25**(2), (1993), 14–17. (in Chinese). [MR1236132](#). [Zbl 0971.11514](#).
- [32] M.-J. Deng. *A note on the diophantine equation $(a^2 - b^2)^x + (2ab)^y = (a^2 + b^2)^z$* . *J. Nat. Sci. Heilongjiang Univ.*, **19**(3), (2002), 8–10. [MR1935472](#). [Zbl 1076.11508](#).
- [33] M.-J. Deng. *On the diophantine equation $(15n)^x + (112n)^y = (113n)^z$* . *J. Nat. Sci. Heilongjiang Univ.*, **24**(5), (2007), 617–620. (in Chinese). [Zbl 1150.11015](#).
- [34] M.-J. Deng. *A note on the diophantine equation $(na)^x + (nb)^y = (nc)^z$* . *Bull. Aust. Math. Soc.*, **89**(2), (2014), 316–321. [MR3182668](#). [Zbl 1372.11046](#).

- [35] M.-J. Deng and G. L. Cohen. *On the conjecture of Jeśmanowicz concerning Pythagorean triples.* Bull. Aust. Math. Soc., **57**(4), (1998), 515–524. [MR1623283](#). [Zbl 0916.11020](#).
- [36] M.-J. Deng and G. L. Cohen. *A note on a conjecture of Jeśmanowicz.* Colloq. Math., **86**(1), (2000), 25–30. [MR1799885](#). [Zbl 0960.11026](#).
- [37] M.-J. Deng and J. Guo. *A note on Jeśmanowicz' conjecture concerning primitive Pythagorean triples II.* Acta Math. Hung., **153**(2), (2017), 436–448. [MR3720980](#). [Zbl 1399.11099](#).
- [38] M.-J. Deng and D.-M. Huang. *A note on Jeśmanowicz conjecture concerning primitive Pythagorean triples.* Bull. Aust. Math. Soc., **95**(1), (2017), 5–13. [MR3592539](#). [Zbl 06717752](#).
- [39] X.-L. Dong and Z.-F. Cao. *The Terai-Jeśmanowicz conjecture on the equation $a^x + b^y = c^z$.* Chinese Ann. Math., Ser. A, **21A**(6), (2000), 709–714. (in Chinese). [Zbl 0987.11021](#).
- [40] Q. Feng. *The shuffle variant of Jeśmanowicz' conjecture on primitive Pythagorean numbers.* Math. Pract. Theory, **45**(16), (2015), 312–315. (in Chinese). [Zbl 1349.11072](#).
- [41] Q. Feng and D. Han. *On the diophantine system $a^2 + b^2 = c^r$ and $a^x + b^y = c^z$ for b a prime.* Int. J. Appl. Math. Stat., **52**(7), (2014), 65–73. [MR3256469](#). [Zbl 1339.11047](#).
- [42] C.-Y. Fu. *On Jeśmanowicz' conjecture concerning Pythagorean numbers.* Master's thesis, Haikou: Hainan Univ., 2016. (in Chinese).
- [43] C.-Y. Fu and M.-J. Deng. *On the diophantine equation $(n(7^{2r} - 4))^x + (n(4 \cdot 7^r))^y = (n(7^{2r} + 4))^z$.* J. Nat. Sci. Heilongjiang Univ., **32**(5), (2015), 596–599. (in Chinese). [Zbl 1349.11073](#).
- [44] R.-Q. Fu and H. Yang. *On the exponential diophantine equation $((am^2 + 1))^x + (bm^2 - 1)^y = (cm)^z$ with $c \mid m$.* Period. Math. Hung., **75**(2), (2017), 143–149. [MR3718506](#). [Zbl 06850216](#).
- [45] Y. Fujita and T. Miyazaki. *Jeśmanowicz' conjecture with congruence relations.* Colloq. Math., **128**(2), (2012), 211–222. [MR3002349](#). [Zbl 1318.11049](#).
- [46] Y. Fujita and T. Miyazaki. *Jeśmanowicz' conjecture with congruence relations II.* Canad. Math. Bull., **57**(3), (2014), 495–505. [MR3239111](#). [Zbl 1291.11071](#).
- [47] A. O. Gel'fond. *Sur la divisibilité de la différence des puissance de deux nombres entiers par une puissance d'un idéal premier.* Mat. Sb., **7**(49), (1940), 7–25. [MR0001755](#). [Zbl 0023.10405](#).

- [48] C. A. Gómez Ruiz and F. Luca. *An exponential diophantine equation related to the sum of powers of two consecutive k -generalized Fibonacci numbers*. *Colloq. Math.*, **137**(2), (2014), 171–188. [MR3286304](#). [Zbl 1338.11016](#).
- [49] S.-S. Gou. *On Jeśmanowicz' conjecture concerning Pythagorean numbers*. Master's thesis, Chongqing: Southwest Univ., 2016. (in Chinese).
- [50] S. S. Gou and H. Zhang. *On the diophantine equation $(16n)^x + (63n)^y = (65n)^z$* . *J. Southwest China Normal Univ., Nat. Sci.*, **40**(4), (2015), 4–7. (in Chinese).
- [51] W.-J. Guan. *The Jeśmanowicz conjecture on Pythagorean numbers*. *Basic Sci. J. Text. Univ.*, **24**(4), (2011), 557–559.
- [52] W.-J. Guan and S. Che. *On the diophantine equation $2^y n^{y-x} = (b+2)^x - b^x$* . *J. Northwest Univ., Nat. Sci.*, **44**(4), (2014), 534–536. (in Chinese). [MR3309624](#). [Zbl 1313.11072](#).
- [53] X.-G. Guan. *On a class of pure exponential diophantine equations*. *J. Hebei North Univ., Nat. Sci.*, **28**(4), (2012), 5–8. (in Chinese).
- [54] X.-G. Guan. *On the exponential diophantine equation $a^x + b^y = 2^z$* . *J. Zhoukou Normal Univ.*, **31**(2), (2014), 26–30. (in Chinese).
- [55] X.-G. Guan. *On Terai's conjecture concerning the diophantine equation $a^x + b^y = c^z$* . *Adv. Math. China*, **44**(6), (2015), 837–844. (in Chinese). [MR3493559](#). [Zbl 1349.11074](#).
- [56] X.-G. Guan. *On the pure exponential diophantine equation $a^x + b^y = (m^2 + 1)^z$* . *Adv. Math. China*, **45**(5), (2016), 687–699. (in Chinese). [Zbl 1374.11055](#).
- [57] Y.-D. Guo and M.-H. Le. *A note on Jeśmanowicz' conjecture concerning Pythagorean numbers*. *Commen. Math. Univ. St. Pauli*, **44**(2), (1995), 225–228. [MR1366530](#). [Zbl 0849.11036](#).
- [58] R. K. Guy. *Unsolved Problems in Number Theory*. Science Press, Beijing, 3rd edition, 2007. [MR2076335](#). [Zbl 1058.11001](#).
- [59] T. Hadano. *On the diophantine equation $a^x + b^y = c^z$* . *Math. J. Okayama Univ.*, **19**(1), (1976/1977), 25–29. [MR0432538](#). [Zbl 0349.10012](#).
- [60] Q. Han and P.-Z. Yuan. *A note on Jeśmanowicz' conjecture*. *Acta Math. Hungar.*, **156**(1), (2018), 220–225. [MR3856913](#). [Zbl 07011148](#).
- [61] B. He and A. Togbé. *The exponential diophantine equation $n^x + (n + 1)^y = (n + 2)^z$ revisited*. *Glasgow Math. J.*, **51**(5), (2009), 659–667. [MR2534015](#). [Zbl 1194.11044](#).

- [62] B. He and A. Togbé. *On the positive integer solutions of the exponential diophantine equation $a^x + (3a^2 - 1)^y = (4a^2 - 1)^z$* . Adv. Math. China, **40**(2), (2011), 227–234. [MR2841128](#).
- [63] B. He, A. Togbé, and S.-C. Yang. *On the solutions of the exponential diophantine equation $a^x + b^y = (m^2 + 1)^z$* . Quaes. Math., **36**(1), (2013), 119–135. [MR3043675](#). [Zbl 1274.11086](#).
- [64] B. He and S.-C. Yang. *The positive integer solutions of the diophantine equation $(8a^3 - 3a)^x + (3a^2 - 1)^y = (4a^2 - 1)^z$* . J. Sichuan Univ., Nat. Sci., **47**(1), (2010), 13–16. (in Chinese). [MR2643276](#). [Zbl 1240.11058](#).
- [65] N. Hirata-Kohno. *S-unit equations and integer solutions to exponential diophantine equations*. In Analytic number theory and surrounding areas, volume 1511, pages 92–97. Kyoto RIMS Kokyuroku, 2006.
- [66] N. Hirata-Kohno and F. Luca. *On the diophantine equation $F_n^x + F_{n+1}^y = F_m^y$* . Rocky Mt. J. Math., **45**(2), (2015), 509–538. [MR3356626](#). [Zbl 1332.11042](#).
- [67] Y.-Z. Hu. *The diophantine equation $(8a^3 - 3a)^{2x} + (3a^2 - 1)^y = (4a^2 - 1)^z$* . J. Sichuan Univ, Nat. Sci., **44**(2), (2007), 225–228. (in Chinese). [MR2340286](#). [Zbl 1164.11318](#).
- [68] Y.-Z. Hu. *On the exponential diophantine equation $a^{2x} + (3a^2 - 1)^y = (4a^2 - 1)^z$* . Adv. Math. China, **36**(4), (2007), 429–434. [MR2381768](#). [Zbl 1131.11328](#).
- [69] Y.-Z. Hu and M.-H. Le. *A note on ternary purely exponential diophantine equations*. Acta Arith., **171**(2), (2015), 173–182. [MR3414305](#). [Zbl 1379.11030](#).
- [70] Y.-Z. Hu and M.-H. Le. *An upper bound for the number of solutions of ternary purely exponential diophantine equations*. J. Number Theory, **183**, (2018), 62–73. [MR3715228](#). [Zbl 06802525](#).
- [71] Y.-Z. Hu and M.-H. Le. *An upper bound for the number of solutions of ternary purely exponential diophantine equations II*. arXiv 1808:06557, 2018.
- [72] Y.-Z. Hu and P.-Z. Yuan. *The exponential diophantine equation $a^x + b^y = c^z$* . Acta Math. Sinica, Chinese Ser., **48**(6), (2005), 1175–1178. (in Chinese). [MR2205060](#). [Zbl 1124.11307](#).
- [73] Y.-Z. Hu and P.-Z. Yuan. *The simultaneous diophantine equations $a^2 + b^2 = c^3$ and $a^x + b^y = c^z$* . Acta Math. Sinica, Chinese Ser., **52**(5), (2009), 1027–1032. (in Chinese). [MR2583775](#). [Zbl 1125.11309](#).
- [74] Y.-Z. Hu and P.-Z. Yuan. *Jeśmanowicz’ conjecture concerning Pythagorean numbers*. Acta Math. Sinica, Chinese Ser., **53**(2), (2010), 297–300. (in Chinese). [MR2666062](#). [Zbl 1224.11049](#).

- [75] L. Jeśmanowicz. *Several remarks on Pythagorean numbers*. Wiadom. Math., **1**(2), (1955/1956), 196–202. (in Polish). [MR0110662](#). [Zbl 0074.27205](#).
- [76] Z. Ke. *On Jeśmanowicz' conjecture*. J. Sichuan Univ., Nat. Sci., **4**(2), (1958), 81–90. (in Chinese).
- [77] Z. Ke. *On Pythagorean numbers*. J. Sichuan Univ., Nat. Sci., **4**(1), (1958), 73–80. (in Chinese).
- [78] Z. Ke. *On the diophantine equation $(a^2 - b^2)^x + (2ab)^y = (a^2 + b^2)^z$* . J. Sichuan Univ., Nat. Sci., **5**(3), (1959), 25–34. (in Chinese).
- [79] Z. Ke. *On the Pythagorean numbers $(2n + 1), 2n(n + 1), 2n(n + 1) + 1$ I*. J. Sichuan Univ., Nat. Sci., **9**(2), (1963), 9–14. (in Chinese).
- [80] Z. Ke. *On the Pythagorean numbers $(2n + 1), 2n(n + 1), 2n(n + 1) + 1$ III*. J. Sichuan Univ., Nat. Sci., **10**(4), (1964), 11–26. (in Chinese).
- [81] Z. Ke and Q. Sun. *On the Pythagorean numbers $(2n + 1), 2n(n + 1), 2n(n + 1) + 1$ II*. J. Sichuan Univ., Nat. Sci., **10**(3), (1964), 1–12. (in Chinese).
- [82] Z. Ke and Q. Sun. *Diophantine equations*. Shanghai Edu. Publ. House, Shanghai, 1980. (in Chinese). [Zbl 0603.10016](#).
- [83] M.-H. Le. *On the diophantine equation $a^x + b^y = c^z$* . J. Changchun Teachers College, Nat. Sci., **2**(1), (1985), 50–62. (in Chinese).
- [84] M.-H. Le. *On two problems of Hall and Edgar*. Northeast Math. J., **4**(4), (1988), 432–434. (in Chinese). [MR0987066](#). [Zbl 0695.10113](#).
- [85] M.-H. Le. *A note on Jeśmanowicz' conjecture*. Colloq. Math., **69**(1), (1995), 47–51. [MR1341681](#). [Zbl 0835.11015](#).
- [86] M.-H. Le. *On Jeśmanowicz' conjecture concerning Pythagorean numbers*. Proc. Japan Acad., Ser. A, **72A**(5), (1996), 97–98. [MR1404479](#). [Zbl 0876.11013](#).
- [87] M.-H. Le. *A note on the diophantine equation $(m^3 - 3m)^x + (3m^2 - 1)^y = (m^2 + 1)^z$* . Proc. Japan Acad., Ser. A, **73A**(7), (1997), 148–149. [MR1487581](#). [Zbl 0910.11010](#).
- [88] M.-H. Le. *A note on Jeśmanowicz' conjecture concerning Pythagorean numbers*. Bull. Aust. Math. Soc., **59**, (1999), 477–480. [MR1697985](#). [Zbl 0940.11021](#).
- [89] M.-H. Le. *An upper bound for the number of solutions of the exponential diophantine equation $a^x + b^y = c^z$* . Proc. Japan Acad., Ser. A, **75A**(6), (1999), 90–91. [MR1712652](#). [Zbl 0939.11018](#).

- [90] M.-H. Le. *On Terai's conjecture concerning Pythagorean numbers.* Bull. Aust. Math. Soc., **61**, (2000), 329–334. [MR1748713](#). [Zbl 0979.11021](#).
- [91] M.-H. Le. *On the exponential diophantine equation $(m^3 - 3m)^x + (3m^2 - 1)^y = (m^2 + 1)^z$.* Publ. Math. Debrecen, **58**(3–4), (2001), 461–466. [MR1831054](#). [Zbl 1062.11020](#).
- [92] M.-H. Le. *On Cohn's conjecture concerning the diophantine equation $x^2 + 2^m = y^n$.* Archiv der Mathematik, **78**, (2002), 26–35. [MR1887313](#). [Zbl 1006.11013](#).
- [93] M.-H. Le. *A conjecture concerning the exponential diophantine equation $a^x + b^y = c^z$.* Acta Arith., **106**(4), (2003), 345–353. [MR1957910](#). [Zbl 1023.11013](#).
- [94] M.-H. Le. *On Terai's conjecture concerning the exponential diophantine equation $a^x + b^y = c^z$.* Acta Math. Sinica, Chinese Ser., **46**(2), (2003), 245–250. (in Chinese). [MR1988161](#). [Zbl 1136.11306](#).
- [95] M.-H. Le. *A note on the exponential diophantine equation $a^x + b^y = c^z$.* Proc. Japan Acad., Ser. A, **80A**(4), (2004), 21–23. [MR2055070](#). [Zbl 1050.11040](#).
- [96] M.-H. Le. *A conjecture concerning the pure exponential diophantine equation $a^x + b^y = c^z$.* Acta Math. Sinica, English Ser., **20**(4), (2005), 943–948. [MR2156975](#). [Zbl 1159.11308](#).
- [97] M.-H. Le. *An open problem concerning the diophantine equation $a^x + b^y = c^z$.* Publ. Math. Debrecen, **68**(3–4), (2006), 283–295. [MR2212322](#). [Zbl 1111.11019](#).
- [98] M.-H. Le. *On the diophantine system $a^2 + b^2 = c^3$ and $a^x + b^y = c^z$ for b an odd prime.* Acta Math. Sinica, English Ser., **24**(6), (2008), 917–924. [MR2212322](#). [Zbl 1218.11037](#).
- [99] M.-H. Le. *A note on the diophantine system $a^2 + b^2 = c^r$ and $a^x + b^y = c^z$.* Acta Math. Sinica, Chinese Ser., **51**(4), (2008), 677–684. (in Chinese). [MR2454004](#). [Zbl 1174.11041](#).
- [100] M.-H. Le. *A note on Jeśmanowicz' conjecture concerning primitive Pythagorean triplets.* Acta Arith., **138**(2), (2009), 137–144. [MR2520132](#). [Zbl 1297.11015](#).
- [101] M.-H. Le. *The pure exponential diophantine equation $a^x + b^y = c^z$ for generalized Pythagorean triplets.* Acta Math. Sinica, Chinese Ser., **53**(6), (2010), 1239–1248. (in Chinese). [MR2789638](#). [Zbl 1240.11060](#).
- [102] M.-H. Le. *An application of Baker's method to Jeśmanowicz' conjecture on primitive Pythagorean triples.* arXiv:1811.00654

- [103] M.-H. Le. *An application of the BHV theorem to a new conjecture on exponential diophantine equations*. arXiv:1811.00609.
- [104] M.-H. Le, A. Togbé, and H.-L. Zhu. *On a pure ternary exponential diophantine equation*. Publ. Math. Debrecen, **85**(3–4), (2014), 395–411. [MR3291838](#). [Zbl 1340.11041](#).
- [105] B. Leszczyński. *On the equation $n^x + (n + 1)^y = (n + 2)^z$* . Wiadom Mat., **3**(1), (1959), 37–39. (in Polish) [MR0115973](#). [Zbl 0095.26205](#).
- [106] S.-Z. Li. *On Jeśmanowicz’ conjecture concerning Pythagorean numbers*. Master’s thesis, Chongqing: Southwest Univ., 2011. (in Chinese).
- [107] Y.-M. Li. *On the diophantine equation $(48n)^x + (55n)^y = (73n)^z$* . J. Chongqing Tech. Busin. Univ., Nat. Ser., **2**, (2018), 27–30. (in Chinese).
- [108] Z. Li. *On Jeśmanowicz’ conjecture concerning Pythagorean numbers*. J. Nat. Sci. Hailongjiang Univ., **20**(3), (2003), 54–55. (in Chinese). [MR2025903](#). [Zbl 1058.11026](#).
- [109] C.-N. Lin. *On the diophantine equation $(51n)^x + (1300n)^y = (1301)^z$* . Master’s thesis, Chongqing: Southwest Univ., 2017. (in Chinese).
- [110] M.-Y. Lin. *The positive integer solutions of an exponential diophantine equation*. J. Math. Wuhan, **26**(4), (2006), 409–414. (in Chinese). [MR2241997](#). [Zbl 1118.11017](#).
- [111] D.-R. Ling and J.-X. Weng. *On the diophantine equation $(195n)^x + (28n)^y = (197n)^z$* . Pure Appl. Math., **29**(4), (2013), 342–349. (in Chinese). [MR3154606](#). [Zbl 1299.11034](#).
- [112] B.-L. Liu. *A new conjecture on primitive Pythagorean numbers*. Math. Pract. Theory, **43**(9), (2013), 253–255. (in Chinese). [Zbl 1235.11091](#).
- [113] B.-L. Liu. *A note on ternary exponential diophantine equations*. J. Inner Mongolia Univ, Nat. Sci., **43**(4), (2014), 401–402. (in Chinese). [MR3243647](#). [Zbl 1313.11073](#).
- [114] B.-L. Liu. *On the diophantine equation $(143n)^x + (24n)^y = (145n)^z$* . Math. Pract. Theory, **47**(20), (2017), 178–182. (in Chinese). [MR3727287](#). [Zbl 1399.11102](#).
- [115] H.-L. Liu. *On Jeśmanowicz’ conjecture concerning Pythagorean numbers*. Master’s thesis, Chongqing: Southwest Univ., 2017. (in Chinese).
- [116] Z.-W. Liu. *On the ternary pure exponential diophantine equation $a^x + b^y = c^z$* . Pure Appl. Math., **23**(1), (2007), 28–30. (in Chinese).

- [117] W.-D. Lu. *On the Pythagorean numbers $4n^2 - 1$, $4n$ and $4n^2 + 1$* . J. Sichuan Univ. Nat. Sci., **5**(2), (1959), 39–42. (in Chinese).
- [118] W.-Y. Lu, L. Gao, and H.-F. Hao. *On the integer solutions of the diophantine equation $(44n)^x + (117n)^y = (125n)^z$* . Pure Appl. Math., **30**(6), (2014), 627–633. (in Chinese). [Zbl 1324.11035](#).
- [119] W.-Y. Lu, L. Gao, X.-H. Wang, and H.-F. Hao. *On the diophantine equation $(91n)^x + (4140n)^y = (4141n)^z$* . J. Guizhou Normal Univ., Nat. Sci., **33**(2), (2015), 48–53. (in Chinese).
- [120] F. Luca. *On the equation $x^2 + 2^a - 3^b = y^n$* . Int. J. Math. Math. Sci., **29**(4), (2002), 239–244. [MR1897992](#). [Zbl 1085.11021](#).
- [121] F. Luca. *On the system of diophantine equations $a^2 + b^2 = (m^2 + 1)^r$ and $a^x + b^y = (m^2 + 1)^z$* . Acta Arith., **153**(4), (2012), 373–392. [MR2925378](#). [Zbl 1272.11047](#).
- [122] F. Luca and R. Oyono. *An exponential diophantine equation related to powers of two consecutive Fibonacci numbers*. Proc. Japan Acad., Ser. A, **87A**(4), (2011), 45–50. [MR2803898](#). [Zbl 1253.11046](#).
- [123] J. Ma. *On the diophantine equation $(57n)^x + (1624n)^y = (1625n)^z$* . Master's thesis, Chongqing: Chongqing Southwest Univ., 2013. (in Chinese).
- [124] M.-M. Ma. *On Jeśmanowicz' conjecture*. Master's thesis, Nanjing: Nanjing Normal Univ., 2015. (in Chinese).
- [125] M.-M. Ma and Y.-G. Chen. *Jeśmanowicz' conjecture on Pythagorean triples*. Bull. Aust. Math. Soc., **96**(1), (2017), 30–35. [MR3668397](#). [Zbl 06757135](#).
- [126] M.-M. Ma and J.-D. Wu. *On the diophantine equation $(65n)^x + (72n)^y = (97n)^z$* . J. Nanjing Normal Univ., Nat. Sci., **37**(4), (2014), 28–30. (in Chinese). [MR3307760](#). [Zbl 1324.11036](#).
- [127] M.-M. Ma and J.-D. Wu. *On the diophantine equation $(an)^x + (bn)^y = (cn)^z$* . Bull. Korean Math. Soc., **52**(4), (2015), 1133–1138. [MR3385756](#). [Zbl 1335.11027](#).
- [128] K. Mahler. *Zur Approximation algebraischer Zahlen I: Über den grössten Primteiler binärer Formen*. Math. Ann., **107**, (1933), 691–730. [MR1512822](#).
- [129] A. Makowski. *On the diophantine equation $2^x + 11^y = 5^z$* . Norsk. Mat. Tidsskr., **7**(1), (1959), 81–96. [MR0109803](#). [Zbl 0084.27104](#).
- [130] A. Makowski. *On the equation $n^x + (n + 1)^y = (n + 2)^z$* . Wiadom. Mat., **9**(3), (1967), 221–224. [MR0213292](#). [Zbl 0153.06603](#).

- [131] T. Miyazaki. *On the conjecture of Jeśmanowicz concerning Pythagorean triples.* Bull. Aust. Math. Soc., **80**(3), (2009), 413–422. [MR2569916](#). [Zbl 1225.11038](#).
- [132] T. Miyazaki. *Exceptional cases of Terai’s conjecture on diophantine equations.* Arch. Math. Basel, **95**(6), (2010), 519–527. [MR2745461](#). [Zbl 1210.11047](#).
- [133] T. Miyazaki. *Generalizations of classical results on Jeśmanowicz’ conjecture concerning Pythagorean triples.* In T. Komatsu et al., editor, Diophantine Analysis and Related Fields, AIP Conf. Proc., **1264**, (2010), 41–51. New York. [MR2731813](#).
- [134] T. Miyazaki. *Jeśmanowicz’ conjecture on exponential diophantine equations.* Funct. Approx. Comment. Math., **45**(2), (2011), 207–229. [MR2895155](#). [Zbl 1266.11064](#).
- [135] T. Miyazaki. *The shuffle variant of Jeśmanowicz’ conjecture concerning Pythagorean triples.* J. Aust. Math. Soc., **90**(3), (2011), 355–370. [MR2833306](#). [Zbl 1225.11039](#).
- [136] T. Miyazaki. *Terai’s conjecture on exponential diophantine equations.* Int. J. Number Theory, **7**(4), (2011), 981–999. [MR2812648](#). [Zbl 1221.11092](#).
- [137] T. Miyazaki. *Generalizations of classical results on Jeśmanowicz’ conjecture concerning Pythagorean triples.* J. Number Theory, **133**(2), (2013), 583–589. [MR2994375](#). [Zbl 1309.11029](#).
- [138] T. Miyazaki. *The shuffle variant of Terai’s conjecture on exponential diophantine equations.* Publ. Math. Debrecen, **83**(1–2), (2013), 43–62. [MR3081225](#). [Zbl 1274.11087](#).
- [139] T. Miyazaki. *A note on the article by F. Luca “On the system of diophantine equations $a^2 + b^2 = (m^2 + 1)^r$ and $a^x + b^y = (m^2 + 1)^z$ ”.* Acta Arith., **164**(1), (2014), 31–42. [MR3223317](#). [Zbl 1300.11027](#).
- [140] T. Miyazaki. *A remark on Jeśmanowicz’ conjecture for the non-coprimality case.* Acta Math. Sinica, English Ser., **31**(8), (2015), 1255–1260. [MR3367686](#). [Zbl 1330.11021](#).
- [141] T. Miyazaki. *Upper bounds for solutions of an exponential diophantine equation.* Rocky Mt. J. Math., **45**(1), (2015), 303–344. [MR3334214](#). [Zbl 1378.11046](#).
- [142] T. Miyazaki and F. Luca. *On the system of diophantine equations $(m^2 - 1)^r + b^2 = c^2$ and $(m^2 - 1)^x + b^y = c^z$.* J. Number Theory, **153**, (2015), 321–345. [MR3327578](#). [Zbl 1365.11033](#).

- [143] T. Miyazaki and N. Terai. *On the exponential diophantine equation $(m^2 + 1)^x + (cm^2 - 1)^y = (am)^z$* . Bull. Aust. Math. Soc., **90**(1), (2014), 9–19. [MR3227125](#). [Zbl 1334.11019](#).
- [144] T. Miyazaki and N. Terai. *On Jeśmanowicz' conjecture concerning Pythagorean triples II*. Acta Math. Hung., **147**(2), (2015), 286–293. [MR3420578](#). [Zbl 1374.11056](#).
- [145] T. Miyazaki and A. Togbé. *The diophantine equation $(2am - 1)^x + (2m)^y = (2am + 1)^z$* . Int. J. Number Theory, **8**(8), (2012), 2035–2044. [MR2978854](#). [Zbl 1290.11068](#).
- [146] T. Miyazaki, A. Togbé, and P.-Z. Yuan. *On the diophantine equation $a^x + b^y = (a + 2)^z$* . Acta Math. Hung., **149**(1), (2016), 1–9. [MR3498942](#). [Zbl 1389.11088](#).
- [147] T. Miyazaki, P.-Z. Yuan, and D.-Y. Wu. *Generalizations of classical results on Jeśmanowicz' conjecture concerning Pythagorean triples II*. J. Number Theory, **141**(1), (2014), 184–201. [MR3195395](#). [Zbl 1310.11040](#).
- [148] L. J. Mordell. *Diophantine Equations*. Academic Press, London, 1969. [MR0249355](#). [Zbl 0188.34503](#).
- [149] T. Nagell. *Sur une classe d'équations exponentielles*. Ark. Mat., **3**(4), (1958), 569–582. [MR0103858](#). [Zbl 0083.03902](#).
- [150] X.-W. Pan. *A note on the exponential diophantine equation $(am^2 + 1)^x + (bm^2 - 1)^y = (cm)^z$* . Colloq. Math., **149**(2), (2017), 265–273. [MR3697141](#). [Zbl 06789358](#).
- [151] M. Perisastri. *A note on the equation $a^x - b^y = 10^z$* . Math. Stud., **37**(2), (1969), 211–212. [MR0263738](#). [Zbl 0207.35302](#).
- [152] Z. Rábai. *A note on the shuffle variant of Jeśmanowicz's conjecture*. Tokyo J. Math., **40**(1), (2017), 153–163. [MR3689983](#). [Zbl 1390.11071](#).
- [153] D.-M. Rao. *A note on the diophantine equation $(2n + 1)^x + (2n(n + 1))^y = (2n(n + 1) + 1)^z$* . J Sichuan Univ., Nat. Sci., **6**(1), (1960), 79–80. (in Chinese).
- [154] J.-H. Ren and J.-K. Zhang. *On the diophantine equation $a^x + b^y = c^z$* . J. Northwest Univ., Nat. Sci., **19**(1), (1989), 12–22. (in Chinese). [MR1018048](#). [Zbl 0695.10016](#).
- [155] R. Scott. *On the equation $p^x - q^y = c$ and $a^x + b^y = c^z$* . J. Number Theory, **44**(1), (1993), 153–165. [MR1225949](#). [Zbl 0786.11020](#).

- [156] R. Scott and R. Styer. *On $p^x - q^y = c$ and related three term exponential diophantine equations with prime bases.* J. Number Theory, **105**(2), (2004), 212–234. [MR2040155](#). [Zbl 1080.11032](#).
- [157] R. Scott and R. Styer. *Number of solutions to $a^x + b^y = c^z$.* Publ. Math. Debrecen, **88**(1–2), (2016), 131–138. [MR3452168](#). [Zbl 1374.11057](#).
- [158] T. N. Shorey and R. Tijdeman. *Exponential Diophantine Equations.* Cambridge Univ. Press, Cambridge, 1986. [MR0891406](#). [Zbl 1156.11015](#).
- [159] W. Sierpiński. *On the equation $3^x + 4^y = 5^z$.* Wiadom. Math., **1**(2), 1955/1956), 194–195. (in Polish). [MR0105384](#). [Zbl 0074.27204](#).
- [160] G. Soydan, M. Demirci, I. N. Cangul, and A. Togbé. *On the conjecture of Jeśmanowicz.* Int. J. Appl. Math. Stat., **56**(6), (2017), 46–72. [MR3685484](#).
- [161] L.-J. Su and X.-X. Li. *The exponential diophantine equation $(4m^2 + 1)^x + (5m^2 - 1)^y = (3m)^z$.* Abst. App. Anal., pages 1–5, April 2014. [MR3198228](#). [Zbl 07022844](#).
- [162] C.-F. Sun and Z. Cheng. *A note on Jeśmanowicz’ conjecture.* J. Math. Wuhan, **33**(5), (2013), 788–794. [MR3154659](#).
- [163] C.-F. Sun and Z. Cheng. *A conjecture of Jeśmanowicz concerning Pythagorean triples.* Adv. Math. China, 43(2):267–275, 2014. [MR3210692](#). [Zbl 1324.11039](#).
- [164] C.-F. Sun and Z. Cheng. *On Jeśmanowicz’ conjecture concerning Pythagorean triples.* J. Math. Res. Appl., **35**(2), (2015), 143–148. [MR3242773](#). [Zbl 1324.11039](#).
- [165] C.-F. Sun and M. Tang. *On the diophantine equation $(an)^x + (bn)^y = (cn)^z$.* Chinese Math. Ann., Ser. A, **39**(1), (2018), 87–94. (in Chinese). [MR3821058](#). [Zbl 06960844](#).
- [166] H.-N. Sun. *On the diophantine equation $(35n)^x + (612n)^y = (613n)^z$.* Master’s thesis, Chongqing: Southwest Univ., 2015. (in Chinese).
- [167] Q. Sun and X.-M. Zhou. *On the diophantine equation $a^x + b^y = c^z$.* Chinese Sci. Bull., **29**(1), (1984), 61. (in Chinese). [MR0838063](#). [Zbl 0565.10017](#).
- [168] K. Takakuwa. *A remark on Jeśmanowicz’ conjecture.* Proc. Japan Acad., Ser. A, **72A**(6), (1996), 109–110. [MR1404483](#). [Zbl 0863.11025](#).
- [169] K. Takakuwa and Y. Asaeda. *On a conjecture on Pythagorean numbers.* Proc. Japan Acad., Ser. A, **69A**(7), (1993), 252–255. [MR1249222](#). [Zbl 0796.11009](#).

- [170] K. Takakuwa and Y. Asaeda. *On a conjecture on Pythagorean numbers II*. Proc. Japan Acad., Ser. A, **69A**(8), (1993), 287–290. [MR1249439](#). [Zbl 0796.11010](#).
- [171] K. Takakuwa and Y. Asaeda. *On a conjecture on Pythagorean numbers III*. Proc. Japan Acad., Ser. A, **69A**(9), (1993), 345–349. [MR1261610](#). [Zbl 0822.11025](#).
- [172] G. Tang. *On the diophantine equation $(45n)^x + (28n)^y = (53n)^z$* . J. Southwest Nation. Univ., Nat. Sci., **40**(1). (2014), 101–104. (in Chinese).
- [173] M. Tang and J.-X. Weng. *Jeśmanowicz' conjecture with Fermat numbers*. Taiwan J. Math., **18**(3), (2014), 925–930. [MR3213395](#). [Zbl 1357.11041](#).
- [174] M. Tang and Q.-H. Yang. *The diophantine equation $(bn)^x + (2n)^y = ((b + 2)n)^z$* . Colloq. Math., **132**(1), (2013), 95–100. [MR3106090](#). [Zbl 1355.11026](#).
- [175] M. Tang and Z.-J. Yang. *Jeśmanowicz' conjecture revisited*. Bull. Aust. Math. Soc., **88**(3), (2013), 486–491. [MR3189299](#). [Zbl 1310.11041](#).
- [176] N. Terai. *The diophantine equation $a^x + b^y = c^z$* . Proc. Japan Acad, Ser. A, **70A**(1), (1994), 22–26. [MR1272664](#). [Zbl 0812.11024](#).
- [177] N. Terai. *The diophantine equation $a^x + b^y = c^z$ II*. Proc. Japan Acad, Ser. A, **71A**(6), (1995), 109–111. [MR1344658](#). [Zbl 0842.11009](#).
- [178] N. Terai. *The diophantine equation $a^x + b^y = c^z$ III*. Proc. Japan Acad, Ser. A, **72A**(1), (1996), 20–22. [MR1382780](#). [Zbl 0858.11017](#).
- [179] N. Terai. *Applications of a lower bound for linear forms in two logarithms to exponential diophantine equations*. Acta Arith., **90**(1), (1999), 17–35. [MR1708700](#). [Zbl 0933.11013](#).
- [180] N. Terai. *On the exponential diophantine equation $a^x + l^y = c^z$* . Proc. Japan Acad, Ser. A, **77A**(9), (2001), 151–154. [MR1869111](#). [Zbl 1009.11026](#).
- [181] N. Terai. *On an exponential diophantine equation concerning Fibonacci numbers*. In Abstracts and short communications and poster sessions, page 55, Beijing, 2002. Int. Conf. Math., Higher Edu. Press.
- [182] N. Terai. *On the exponential diophantine equation $(4m^2 + 1)^x + (5m^2 - 1)^y = (3m)^z$* . Int. J. Algebra, **6**(21–24), (2012), 1135–1146. [MR2974671](#). [Zbl 1271.11039](#).
- [183] N. Terai. *On Jeśmanowicz' conjecture concerning primitive Pythagorean triples*. J. Number Theory, **141**(2), (2014), 316–323. [MR3195402](#). [Zbl 1309.11030](#).

- [184] N. Terai and T. Hibino. *On the exponential diophantine equation $(12m^2 + 1)^x + (13m^2 - 1)^y = (5m)^z$* . Int. J. Algebra, **9**, (2015), 261–272.
- [185] N. Terai and T. Hibino. *The exponential diophantine equation $(3pm^2 - 1)^x + (p(p - 3)m)^y = (pm)^z$* . Period. Math. Hung., **74**(2), (2017), 227–234. [MR3645689](#). [Zbl 1399.11104](#).
- [186] N. Terai and K. Takakuwa. *A note on the diophantine equation $a^x + b^y = c^z$* . Proc. Japan Acad., Ser. A, **73A**(9), (1997), 161–164. [MR1606004](#). [Zbl 0910.11011](#).
- [187] M. Toyozumi. *On the equation $a^x - b^y = (2p)^z$* . Math. Stud., **46**(2–4), (1978), 113–115. [Zbl 0526.10014](#).
- [188] S. Uchiyama. *On the diophantine equation $2^x = 3^y + 13^z$* . Math. J. Okayama Univ., **19**(1), (1976/1977), 31–38. [MR0432539](#). [Zbl 0349.10013](#).
- [189] M. Waldschmidt. *Perfect powers: Pillai’s works and their developments*. arXiv:0908:4031v1, 27 August 2009.
- [190] J.-H. Wang and M.-J. Deng. *On the diophantine equation $(a^2 - b^2)^x + (2ab)^y = (a^2 + b^2)^z$* . J. Nat. Sci. Heilongjiang Univ., **13**(4), (1996), 23–25. (in Chinese). [MR1445399](#). [Zbl 1076.11511](#).
- [191] J.-P. Wang, T.-T. Wang, and W.-P. Zhang. *A note on the exponential diophantine equation $(4m^2 + 1)^x + (5m^2 - 1)^y = (3m)^z$* . Colloq. Math., **139**(1), (2015), 121–126. [MR3332737](#). [Zbl 1364.11087](#).
- [192] L.-L. Wang. *On the diophantine equation $(39n)^x + (760n)^y = (761n)^z$* . Master’s thesis, Chongqing: Southwest Univ., 2011. (in Chinese).
- [193] T.-T. Wang, X.-H. Wang, and Y.-Z. Jiang. *An application of the Baker method to Jeśmanowicz’ conjecture on Pythagorean triples*. Rev. R. Acad. Cienc. Exactas Fis. Nat., Ser. A Mat., RASCOM, **112**(2), (2018), 385–390. [MR3775276](#). [Zbl 06859078](#).
- [194] X.-H. Wang and S. Gou. *On Miyazaki’s conjecture on primitive Pythagorean numbers*. Math. Pract. Theory, **44**(8), (2014), 287–290. (in Chinese). [MR3237472](#). [Zbl 1340.11043](#).
- [195] J. Y. Xia and P.-Z. Yuan. *On the Terai-Jeśmanowicz conjecture*. Acta Math. Sinica, English Ser., **24**(12), (2008), 2061–2064. [MR2453085](#). [Zbl 1234.11035](#).
- [196] J.-J. Xing. *On Jeśmanowicz’ conjecture concerning Pythagorean numbers*. Master’s thesis, Chongqing: Southwest Univ., 2015. (in Chinese).

- [197] H. Yang and R.-Q. Fu. *A kind of an exponential diophantine system and its integer solutions.* J. Northwest Univ., Nat. Sci., **43**(4), (2013), 524–526. (in Chinese). [MR3183801](#). [Zbl 1299.11036](#).
- [198] H. Yang and R.-Q. Fu. *A note on Jeśmanowicz' conjecture concerning primitive Pythagorean triples.* J. Number Theory, **156**(1), (2015), 183–194. [MR3360336](#). [Zbl 1395.11066](#).
- [199] H. Yang and R.-Q. Fu. *Fermat primes and Jeśmanowicz' conjecture.* Adv. Math. China, **46**(6), (2017), 857–866. (in Chinese). [MR3778513](#). [Zbl 1399.11107](#).
- [200] H. Yang, R.-Z. Ren, and R.-Q. Fu. *On Jeśmanowicz' conjecture concerning Pythagorean numbers.* Math. J. Wuhan, **37**(3), (2017), 506–512. (in Chinese). [Zbl 1399.11108](#).
- [201] S.-C. Yang and B. He. *The solutions of a class of exponential diophantine equations.* Adv. Math. China, **41**(5), (2012), 565–573. (in Chinese). [MR3058640](#). [Zbl 1274.11097](#).
- [202] X.-Z. Yang. *On the diophantine equation $a^x + b^y = c^z$.* J. Sichuan Univ., Nat. Sci., **1985**(4), (1985), 151–158. (in Chinese). [MR0843518](#).
- [203] Z.-J. Yang and M. Tang. *On the diophantine equation $((8n)^x + (15n)^y = (17n)^z$.* Bull. Aust. Math. Soc., **86**(2), (2012), 348–352. [MR2979995](#). [Zbl 1272.11048](#).
- [204] Z.-J. Yang and J.-X. Weng. *On the diophantine equation $(12n)^x + (35n)^y = (37n)^z$.* Pure Appl. Math., **28**(5), (2012), 698–704. (in Chinese). [MR3053137](#). [Zbl 1274.11098](#).
- [205] Y.-H. Yu and Z.-P. Li. *The exceptional solutions of the exponential diophantine equation $(bn)^x + (2n)^y = ((b+2)n)^z$.* Math. Pract. Theory, **44**(18), (2014), 290–293. [MR3328405](#). [Zbl 1324.11034](#).
- [206] P.-Z. Yuan and Q. Han. *Jeśmanowicz' conjecture and related equations.* Acta Arith., **184**(1), (2018), 37–49. [MR3826639](#). [Zbl 06921720](#).
- [207] X.-W. Zhang and W.-P. Zhang. *The exponential diophantine equation $((2^{2m}-1)n)^x + (2^{m+1}n)^y = ((2^{2m}+1)n)^z$.* Bull. Math. Soc. Math. Roum., Nouv. Sér., **57**(3), (2014), 337–344. [MR3241772](#). [Zbl 1340.11044](#).
- [208] C.-Y. Zheng. *A note on coprime cases of Jeśmanowicz' conjecture.* J. Huaihai Engin. College, Nat. Ser., **3**, (2017), 1–3. (in Chinese).
- [209] X.-E. Zhou. *On the diophantine equation $p^x + q^y = 2^z$ ($200 < \max(p, q) < 300$).* J. Nat. Sci. Hainan Univ., **32**(3), (2014), 197–199. (in Chinese).

- [210] M.-H. Zhu and X.-X. Li. *The exponential diophantine equation $4^x + b^y = (b + 4)^z$* . *J. Math. Wuhan*, **36**(4), (2016), 782–786. (in Chinese). [Zbl 1363.11062](#).

Maohua Le
Institute of Mathematics, Lingnan Normal College
Zhangjiang, Guangdong 524048,
China.
e-mail: lemaohua2008@163.com

Reese Scott
Somerville, MA, USA.

Robert Styer
Department of Mathematics and Statistics,
Villanova University,
800 Lancaster Avenue
Villanova, PA 19085 USA.
e-mail: robert.styer@villanova.edu
<http://www41.homepage.villanova.edu/robert.styer/>

License

This work is licensed under a [Creative Commons Attribution 4.0 International License](#). 

Surveys in Mathematics and its Applications **14** (2019), 109 – 140
<http://www.utgjiu.ro/math/sma>