



Corrigendum: Inter- and Intra-specific variation in egg size among reef fishes across the Isthmus of Panama

OPEN ACCESS

Edited and reviewed by:

Michael Arthur St. John,
 Technical University of Denmark,
 Denmark

*Correspondence:

D. Ross Robertson
 dr@stri.org

†These authors have contributed
 equally to this work

Specialty section:

This article was submitted to
 Marine Ecosystem Ecology,
 a section of the journal
 Frontiers in Ecology and Evolution

Received: 20 February 2019

Accepted: 05 March 2019

Published: 27 March 2019

Citation:

Robertson DR and Collin R (2019)
 Corrigendum: Inter- and Intra-specific
 variation in egg size among reef fishes
 across the Isthmus of Panama.
 Front. Ecol. Evol. 7:83.
 doi: 10.3389/fevo.2019.00083

D. Ross Robertson*† and Rachel Collin†

Smithsonian Tropical Research Institute, Balboa, Panama

Keywords: primary productivity, upwelling, tropical reef-fish, egg-size variation, temperature-size rule, life history evolution, Isthmus of Panama

A Corrigendum on

Inter- and Intra-specific variation in egg size among reef fishes across the Isthmus of Panama
 by Robertson, D. R., and Collin, R. (2015). *Front. Ecol. Evol.* 2:84. doi: 10.3389/fevo.2014.00084

In the original article, there was a mistake in **Supplementary Table 1** as published. The unit for Egg Dry Weight in **Supplementary Table 1** should be “milligrams (mg)” not “micrograms (μg).” The corrected **Supplementary Table 1** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2019 Robertson and Collin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Supplementary Table 1 | Summary of egg volume, dry weight, and energy content of eggs of 24 neotropical reef fishes averaged for the cool and the warm seasons.

Species	Ocean	Season	Egg volume (μ l)			Egg dry weight (mg)			Egg energy (Joules)		
			Mean	St Dev	N	Mean	St Dev	N	Mean	St Dev	N
POMACENTRIDAE											
<i>Abudefduf saxatilis</i>	Caribbean	Warm	0.190	0.012	259	0.045	0.0024	45	1.034	0.071	31
		Cool	0.199	0.013	115	0.046	0.0028	21	1.075	0.047	17
<i>Abudefduf troschelii</i>	Pacific	Warm	0.146	0.007	45	0.035	0.0035	32	0.780	0.092	22
		Cool	0.153	0.010	38	0.038	0.0044	22	0.861	0.105	22
<i>Chromis multilineata</i>	Caribbean	Warm	–	–	–	0.018	0.002	8	0.438	0.020	7
		Cool	–	–	–	0.019	0.001	7	0.443	0.051	7
<i>Chromis atrilobata</i>	Pacific	Warm	0.076	0.002	14	0.016	0.001	8	–	–	–
		Cool	0.078	0.006	7	0.016	0.001	7	0.346	0.043	6
<i>Microspathodon bairdii</i>	Pacific	Warm	0.137	0.015	27	0.027	0.0037	14	0.665	0.075	10
		Cool	0.157	0.004	4	0.027	0.0075	2	–	–	–
<i>Microspathodon chrysurus</i>	Caribbean	Warm	0.123	0.007	246	0.027	0.004	35	0.634	0.086	21
		Cool	0.132	0.006	104	0.028	0.003	17	0.674	0.091	17
<i>Microspathodon dorsalis</i>	Pacific	Warm	0.127	0.012	37	0.025	0.0035	20	0.594	0.046	16
		Cool	0.152	0.018	47	0.031	0.0033	12	0.751	0.083	8
<i>Stegastes adustus</i>	Caribbean	Warm	0.095	0.008	55	0.020	0.0046	34	0.465	0.116	25
		Cool	0.105	0.008	22	0.021	0.0012	15	0.518	0.030	15
<i>Stegastes diencaeus</i>	Caribbean	Warm	0.142	0.010	261	0.029	0.0039	28	0.722	0.054	16
		Cool	0.150	0.010	77	0.026	0.0009	2	0.640	0.028	2
<i>Stegastes leucostictus</i>	Caribbean	Warm	0.226	0.014	57	0.045	0.006	37	1.175	0.118	22
		Cool	0.238	0.016	34	0.051	0.0028	19	1.227	0.096	19
<i>Stegastes partitus</i>	Caribbean	Warm	0.073	0.008	252	0.015	0.0027	28	0.349	0.083	18
		Cool	0.079	0.005	109	0.017	0.0012	23	0.401	0.032	22
<i>Stegastes planifrons</i>	Caribbean	Warm	0.133	0.007	23	0.026	0.0032	22	0.617	0.071	10
		Cool	0.147	0.006	14	0.029	0.0048	14	0.752	0.127	9
<i>Stegastes variabilis</i>	Caribbean	Warm	0.134	0.012	33	0.025	0.005	20	0.626	0.136	11
		Cool	0.145	0.012	32	0.031	0.003	20	0.763	0.056	13
<i>Stegastes acapulcoensis</i>	Pacific	Warm	0.099	0.004	64	0.019	0.0029	31	0.448	0.065	22
		Cool	0.122	0.014	35	0.020	0.0014	6	–	–	–
<i>Stegastes flavilatus</i>	Pacific	Warm	0.080	0.004	65	0.016	0.0040	25	0.384	0.079	17
		Cool	0.101	0.011	75	0.020	0.0038	13	0.461	0.085	10
BLENNIIDAE											
<i>Ophioblennius macclurei</i>	Caribbean	Warm	0.137	0.007	27	0.030	0.0027	27	0.728	0.062	12
		Cool	0.142	0.006	27	0.035	0.0013	16	0.859	0.041	16
<i>Ophioblennius steindachneri</i>	Pacific	Warm	0.110	0.008	37	0.025	0.0031	12	0.572	0.073	12
		Cool	0.134	0.013	46	0.031	0.0034	12	0.699	0.062	12
CHAENOPSIDAE											
<i>Acanthemblamaria rivasi</i>	Caribbean	Warm	0.582	0.051	31	0.116	0.013	31	2.80	0.266	28
		Cool	0.584	0.036	9	0.138	0.012	8	3.20	0.289	8
<i>Acanthemblamaria handcocki</i>	Pacific	Warm	0.357	0.026	37	0.077	0.008	34	1.802	0.199	30
		Cool	0.344	0.018	13	0.088	0.007	12	2.034	0.160	12
GOBIIDAE											
<i>Coryphopterus dicrus</i>	Caribbean	Warm	0.095	0.009	16	0.013	0.0014	14	0.297	0.020	11
		Cool	–	–	–	–	–	–	–	–	–
<i>Coryphopterus glaucofrenum</i>	Caribbean	Warm	0.077	0.011	42	0.009	0.0020	26	0.201	0.049	20
		Cool	0.097	0.033	23	0.011	0.0015	22	–	–	–
<i>Coryphopterus personatus</i>	Caribbean	Warm	0.072	0.007	28	0.010	0.0012	28	0.211	0.030	28
		Cool	–	–	–	–	–	–	–	–	–
<i>Coryphopterus tortugae</i>	Caribbean	Warm	0.106	0.009	29	0.011	0.0013	27	0.25	0.035	27
		Cool	–	–	–	–	–	–	–	–	–
<i>Coryphopterus urospilus</i>	Pacific	Warm	0.051	0.009	20	0.006	0.0004	14	0.178	0.031	30
		Cool	0.078	0.027	33	0.008	0.0012	31	–	–	–