



Ukrainian Journal of Nephrology and Dialysis

Scientific and Practical, Medical Journal

Founders:

- State Institution «Institute of Nephrology NAMS of Ukraine»
- National Kidney Foundation of Ukraine

ISSN 2304-0238;
eISSN 2616-7352

Journal homepage: <https://ukrjnd.com.ua>

Research Article

N. M. Voroshylova, M. D. Timchenko, S. V. Verevka

doi: 10.31450/ukrjnd.1(61).2019.05

Bence-Jones protein as the form of nano-scaled β -stacked supramolecular aggregates

State Institution «O.C. Kolomyichenko Institute of Otolaryngology of the National Academy of Medical Sciences of Ukraine»

Citation:

N. Voroshylova, M. Timchenko, S. Verevka. Bence-Jones protein as the form of nano-scaled β -stacked supramolecular aggregates. Ukr J Nephrol Dial. 2019;1(61):39-44. doi: 10.31450/ukrjnd.1(61).2019.05

Abstract. *The formation in β -structured protein aggregates in tissues and fluids of the body is one of the most dangerous complications of various diseases. The most famous of them are amyloidoses, but they such deposits are observed at other, much more widespread, diseases. The generally accepted approach to amyloids' detection is based on high-specific coloring by Congo Red dye. However, the Abbe's diffraction limit excludes the seeing of the objects smaller than 0.61 wavelengths (about 240 nm). Such nanoscale formations are capable to disrupt the functioning of surrounding tissues, to cause the complications and recurrences of the disease, and to pass through biological barriers with the following accumulation in body's fluids. It's likely that these conditions are the cause of the urinary congophilia, that is associated with preeclampsia at pregnancy and chronic kidney disease. Nor the less suspicious object is the Bence-Jones protein that appears in the urine at multiple myeloma and some other diseases, which are in more or less extent, are related to the disturbance of protein metabolism.*

The purpose of this study was to clarify the aggregate state of the Bence-Jones protein as a possible β -structured supramolecular associate.

Methods. The subject of the study was the freshly received urine from a patient with multiple myeloma. The presence of the Bence-Jones protein was checked by thermoprecipitation of the acidified sample. For control, the urine was used by a healthy person with the addition of certain amounts of human serum albumin ("Reanal", Hungary) with a concentration of 0, 0.01, 0.1 and 1%.

Result. The obtained data testify to the appropriateness of such a point of view and create prerequisites for the expanding of diagnostic possibilities.

Conclusions. The results obtained during the study testify to the peculiarity of the structure of the Bence-Jones protein, which is nano-sized beta-structured supramolecular

Key words: protein aggregation, amyloidoses, β -aggregates, congophilia, Bence-Jones protein.

Conflict of interest statement: the authors declared no competing interests.

© N. Voroshylova, M. Timchenko, S. Verevka, 2019. All rights reserved.

Correspondence should be addressed to Sergiy Verevka: verevka.biochem@gmail.com

Article history:

Received January 16, 2019

Received in revised form

February 03, 2019

Accepted February 18, 2019



© Н.М. Ворошилова, М.Д. Тімченко, С.В. Верьовка

УДК: 577.112:616-003.821

Н.М. Ворошилова, М.Д. Тімченко, С.В. Верьовка

Білок Бенс-Джонса як різновид β -структурованого супрамолекулярного агрегату

ДУ «Інститут отоларингології ім. проф. О.С. Коломійченка
Національної академії медичних наук України»

Резюме. Формування в тканинах та рідинах організму β -структурованих білкових агрегатів належать до найтяжчих ускладнень різноманітних захворювань. Провідне місце серед них посідають амілоїдози, однак подібні депозити спостерігаються й за інших, значно поширеніших, захворювань. Класичний підхід до виявлення амілоїдів ґрунтується на їх високовибірковому забарвленні барвником Конго червоним. Однак дифракційне обмеження Аббе робить неможливим виявлення агрегатів, менших за 0.61 довжини світлової хвилі (порядку 240 нм). Подібні нанорозмірні утворення повній мірі здатні порушувати функціонування оточуючих тканин, викликати ускладнення та рецидив захворювання, проходити через біологічні бар'єри та накопичуватись в рідинах організму. Ймовірно, що саме цими обставинами обумовлено уриниру конгофілію, пов'язану з преекламсією вагітних та хронічним захворюванням нирок. Не менш загадковим об'єктом є білок Бенс-Джонса, що з'являється в сечі за мієломної хвороби та групи інших захворювань, в тій чи іншій мірі пов'язаних з порушенням білкового обміну.

Метою даного дослідження є дослідження агрегатного стану білка Бенс-Джонса як можливого β -структурованого супрамолекулярного асоціата. Отримані дані свідчать про слушність подібної точки зору та обумовлюють нові діагностичні можливості.

Методи. Предметом дослідження була щойно отримана сеча пацієнта з множинною мієломою. Наявність білка Бенс-Джонса перевіряли термопацікацією підкисленого зразка. Для контролю використовували сечу здорової людини з додаванням певних кількостей людського сироваткового альбуміну («Reanal», Угорщина) з концентрацією 0, 0,01, 0,1 і 1%.

Результат. Отримані дані свідчать про доцільність такої точки зору та створення попередніх умов для розширення діагностичних можливостей.

Висновки. Отримані в ході дослідження результати свідчать про особливість структури білка Bence-Jones, який є нанорозмірним бета-структурованим супрамолекулярним

Ключові слова: агрегація білків, амілоїдоз, β -агрегати, конгофілія, білок Бенс-Джонса.

Introduction. Determination of protein content in urine belongs to the most mass clinical trials. The norm for urinary protein excretion does not exceed 30 mg, which is not registered by normal laboratory methods and is defined as «traces or lack of protein in the urine.» The protein content of more than 150 mg / l is evaluated as proteinuria, which may indicate significant impaired functioning of the kidneys, organs and tissues of the body [1]. Not only quantitative, but also qualitative definition of protein in the urine is of great diagnostic value, since it, together with clinical signs, allows both to detect the presence of trouble in the body, and to specify the nature of the disease [2].

It is known that the formation and maintenance of proteins in the native conformation is a complex energy-intensive process, which leads to the formation

of the only true structure for each protein. An integral circumstance of many diseases is the development of endogenous intoxication caused by the formation and accumulation of various structurally damaging derivatives of native proteins. Such structures are prone to aggregation. We have emphasized that β -structured protein aggregates are a kind of «energy bottom» for an immense number of conformational states of misfolded proteins [3]. It is the energy utility that gives the formation of such associate self-sufficient autochthonous nature. Similar processes are observed both for pathological changes, and for the normal state of many living organisms [4,5]. The most severe manifestations of the formation of β -structured protein deposits in the body include amyloidosis – a rather large group of diseases that are not curable [6,7]. Formation of amyloid fibrils and aggregates formed by them is a regular multistage process due to the formation of several levels of supramolecular structures [8]. In the association thus formed, the distance between adjacent peptide chains in the β -sheet is 4.7 Å, and the distance between individual sheets is 10 Å [9]. Aggregated β -fold structures are capable of sorption and rearrangement on their own kind of soluble proteins, which ensures the growth of

Sergiy Verevka
verevka.biochem@gmail.com

the aggregate [10]. In the case of most amyloidosis, the starting matrix of this process is the damaged structures of insoluble proteins – collagen and reticulin, which is responsible for the periclogenic and periereticular nature of most amyloidosis [11]. The starting matrix can be formed as a result of membrane holding of unstructured or structurally damaged proteins [12]. Structurally unbalanced proteins are also capable of spontaneous β -aggregation from the solution [13]. In the latter case, the transition from a pool of individual molecules to an insoluble aggregate must pass through the formation of nanosized groups [8]. Due to Abbe's diffraction limit (0.61λ), similar supramolecular associates are not visible in the optical wavelength range. It means that the protein aggregates become visible on exceeding their linear dimensions more than 240 nm. The total molecular weight of such an aggregate is at least 7,000,000 kDa and it becomes visible in the optical wavelength range on exceeding of this value only [14]. It is known that nanoscale particles easily overcome biological barriers and affect significantly the course of various physiological processes [15].

Therefore, the identification of such deposits is of considerable scientific, educational and practical interest. The most generally accepted approaches to identifying β -stacked structures are based on their interaction with specific dyes. Congo Red and Thioflavin T are the most used among them (Figure 1).




Fig. 1. Structural model of conjugation of the Congo red (top right) and Thioflavin T with the β -folded sheet of the protein aggregate [16].

The most widespread method for amyloid testing is based on Congo red staining [17]. Absorption level of this dye on intravenous injection is considered as a faultless diagnostic criterion of amyloidosis [18]. This dye stains amyloids in different shades of red. However, there is a risk to obtain false-positive results during direct visualization at histological research, because Congo red is able to stain elastin. In addition, intact and hemolyzed erythrocytes and various cell constituents stained with Congo red give similar color. Hence, along with examination of stained tissues with using of light microscope, utilization of polarizing microscopy

is required [19]. Due to regular structure, which is a feature of amyloid aggregates, they exhibit an optical activity. Consequently, amyloid formations stained with Congo red are observed in polarizing microscope as apple-green crystal-like structures that drastically differ from shaded tissues, which have no stereospecificity and thus are not able to deflect polarized light [20]. That is the cause why the reliable identification of amyloid entities with using of Congo red staining should combine both light and polarizing microscopy techniques. In addition, Congo Red is capable of effectively depositing microparticles of artificially-generated β -folded protein aggregates [21]. The latter circumstance prompts a closer look at the case of the urinary Congo red, which constitutes the essential circumstance of preeclampsia, chronic kidney disease and lupus nephritis [22]. The degree of binding of Congo red to urine proteins not only makes it possible to diagnose the corresponding complications, but also suggests the presence of nanoscale protein aggregates in the urine. No less suspicious in this regard is the Bence-Jones protein (B-JP). This protein is absent in healthy people, and its appearance in the urine is the result of prerenal proteinuria caused by myeloma disease, as well as Waldenström macroglobulinemia, plasmacytoma, primary amyloidosis, idiomatic monoclonal gammopathy, chronic lymphocytic leukemia, endotheliomas, paraproteinemic hemoblastomas, and lymphogranulomatosis. B-JP is able to provoke urolithiasis due to the formation of protein cylinders in the renal tubules, that makes it similar with damage-associated molecular patterns.

Quantitative B-JP in urine and blood plasma is an important indicator for assessing the patient's condition and the effectiveness of treatment. In clinical and laboratory practice B-JP determination is expedient at the positive probe for total protein by precipitation of sulphosalicylic acid only. It destroys the hydrated cover of the protein molecules, that cause the aggregation and precipitation. The classic method for detecting B-JP is based on the thermal precipitation in a slightly acidic medium. Filtered urine in a quantity of 4 ml is mixed with 1 ml of 2 M sodium acetate buffer pH 4.9 and heated for 15 minutes in a water bath at 56°C. In the presence of the Bence-Jones protein it rapidly precipitates, but dissolves again when cooled. B-JP consists of the light chains of immunoglobulins, that are monoclonal mainly [23]. But the monoclonal light chains are the basis of protein deposits for the most common type of amyloidosis – AL [6,7]. The reverse nature of the thermoprecipitation, the molecular composition, and the damage of the nephron tubules by the protein deposits suggests for the possibility of staying B-JP in a water-soluble high-stabilized state of β -structured protein aggregates. All these considerations led us to test the aggregate state of B-JP by simple and accessible laboratory methods.

The purpose of our study was to verify the possible aggregate state of the Bence-Jones protein as a set of nanosized β -folded groups.

Materials and methods. To accomplish this task, it was considered appropriate to use Congo Red and Thioflavin T dyes that are specific for the β -folded structures. Unlike the small-specific dyes, which are used more or less successfully for colorimetric or turbidimetric determination of protein content in urine [1], as Congo red, and Thiophlavin T are characterized by exceptionally high selectivity of affinity precisely to β -structured protein groups [19, 24]. Therefore, the definition of amyloid-like inclusions in the tissues with the help of these dyes by light, polarization and luminescent microscopy refers to classical methods of histology.

The subject of the study was the freshly received urine from a patient with multiple myeloma. The presence of the Bens-Jones protein was checked by thermopacification of the acidified sample as discussed above. For control, the urine was used by a healthy person with the addition of certain amounts of human serum albumin («Reanal», Hungary) with a concentration of 0, 0.01, 0.1 and 1%. The total protein precipitation was performed by adding of an equal volume of 10% trichloroacetic acid to the test samples, followed by centrifugation and dissolving of the precipitate in 0.1% solution of sodium hydroxide for the determining of the protein amount of by Lowry method [25]. Specific dyes of Congo Red and Thioflavin T (Acros Organics, USA) were used in the form of a 0.1% solution in 0.9% sodium chloride.

Results. When added to the filtered sample of the urine of a patient with multiple myeloma of equal volume of 5% sulphosalicylic acid, the protein aggregate was precipitated. For quantitative evaluation of the protein content in urine, an equal volume of 10% solution of trichloroacetic acid was added to the urine specimen and the precipitate was separated by centrifugation using an OPN-8 centrifuge for 15 minutes at 3000 rpm. The precipitate was dissolved in 0.1 N solution NaOH and the protein content was determined by the Lowry method. It was 0.740 mg / ml of patient's urine. None protein deposition was observed in the urine specimen of a healthy person. On addition of 1 ml of 3 M sodium acetate buffer pH 4.9 to 4 ml of the patient's filtered urine and incubation of the mixture at 56°C in the water thermostat, the formation of a loose precipitate was achieved. This precipitate dissolved quickly at room temperature that is typical for Bens-Jones protein. The heating of the same sample for 15 minutes in a boiling water bath deposition of the protein was slow, but irreversible. When added to the sample of urine equal to 0.1% of the Congo solution, there were no immediate changes, but at heating to 56°C an irreversible deposition of a dark red sediment was observed. The same deposits were formed at the storage of the mixture at 2°C for two days. Both precipitates when viewed in a light microscope were turned to be an array of red-colored particles that had a bright green color in the polarization microscope (Figure 2).




Fig. 2. Aggregates of the Bens-Jones protein precipitated by Congo Red dye. Polarization microscopy, $\times 200$.

In other words, the formed precipitate gave a color reaction, typical for amyloid proteins. The attempt to conduct a fluorimetric study of the interaction of the Bens-Jones protein with Thioflavin T was unsuccessful, since the optical properties of the urine themselves blocked the wavelength of the emission of fluorophore formed by the interaction of this dye with amyloid structures. At the same time, it has been noted that adding the patient's urine to an equal volume of 0.1% Thioflavin T at room temperature or at store for 2 days at 20°C were resulted in the formation of a precipitate. In the case of a warming of such a sample at 45°C the formation of the precipitate was significantly accelerated, and at 55°C it was immediate and irreversible. The heating of the same samples in a boiling water bath, both with the Congo Red, and with Thiophlavin T, led to a rapid and irreversible deposition of proteins.

For control, 1%, 0.1% and 0.01% of human serum albumin solutions in the urine of the healthy person were used. Adding to the test samples equal volumes of 0.1% solutions of both Congo Red and Thiophlavin T at room temperature, neither when heated to 56°C, nor with prolonged storage at 2°C did not lead to precipitation. The heating of the same samples in a boiling water bath for 15 minutes both with dyes and in their absence led to precipitation of denatured protein. The precipitate sorbed Congo red, but no optical activity was detected in the polarization microscope. On these reasons we may conclude about the absence of any induction effects of the used dyes on the confirmation status as a Bens-Jones protein in the urine of the patient with multiple myeloma and human serum albumin in the urine of a healthy person.

Discussion. The formation of structured fibrils is a complex, multi-step and long process (Fig. 3). Therefore, instant precipitation of the protein by the dye when heated to 56°C is a consequence of the violation of the hydration shell of the ready nanoaggregates.




Fig. 3. The key stages of self-assembly of β -structured protein aggregate [8].

Due to the peculiarities of the nano-sized objects, such aggregates are not visible in the optical range and are not precipitated from the aquatic environment by themselves. The addition of the dye leads to the formation of macro-aggregates and sedimentation, similar to that described in the literature for artificial β -structured microparticles [26]. The revealed features of the interaction of Congo Red and Thioflavin T with the Bens-Jones protein can be recognized as evidence of the existence of the latter in the form of nano-sized β -structured aggregates. In favor of this point of view, the abolition of the nature of the depositional action of both dyes indicates that they are in good agreement with the given data regarding the specificity of their interaction with β -folded structures [16].

It is probable that the continuation of the aggregation processes B-JP is the molecular basis for the formation of protein cylinders in the renal tubules and due to their urolithiasis. The differences in the precipitation of B-JP by Congo Red and Thioflavin T may be explained by the differences in the mode of interaction of these dyes with β -stacked structures (Fig. 1). These peculiarities may be used both for the following studies of B-JP and for its quantifying in urine. The resulting precipitate can be investigated for the total protein content by any method that does not coincide with the optical range of the precipitating dye. It should be emphasized that the formation of nanosized β -aggregated supramolecular structures is unlikely to be limited to the Bens-Jones protein. The possible presence of such kind of nanosized protein aggregates in medicated drugs and food products [27] deserves close attention. No less significant consequences may be caused by the presence of such particles in body's tissues were as the possibility of a violation by them of the normal course of physiological processes is obvious. In particular, the forma-

tion and incorporation of highly stabilized protein aggregates into the structure of cell membranes may be the main cause of fibrosis, that is the routine characteristic of processes associated with a violation of the synthesis and exchange of proteins. No less important is their possible role in the relapse of various abnormal tissues. It is clear that the study of these tissues requires for the use of a set of physico-chemical methods that aren't limited by Abbe diffraction limit. Thus, the great attention has been paid to the creation of new fluorescence probes for the determination of β -structured protein aggregates, first of all – with Alzheimer's disease [28]. The use of this kind of probes for the determination of nanosized protein aggregates in blood circulation can also be used to quantify the excessive amounts of such formations, which appears to be a promising direction of molecular diagnostics.

Conclusions. The results obtained during the study testify to the peculiarity of the structure of the Bens-Jones protein, which is nano-sized β -structured supramolecular aggregates. The possible participation of such kind of formations in the disruption of the normal course of physiological processes is under consideration.

Conflict of Interest: The authors state that there is no conflict of interest.

Information about the contribution of each participant.

N.M. Voroshylava: precipitation of proteins, determination of their concentration, and participation in the writing of manuscript,

M.D. Timchenko: microscopic examination of samples and participation in the writing of manuscript,

S.V. Verevka: general research guides and participation in the writing of manuscript.

References:

1. Pupkova VI, Prasolova L. Opredelenie belka v moche i spinnomozgovej zhidkosti. – Kol'covo, 2007. – 43 s. [In Russian].
2. Vel'kov VV, Reznikova OI. Sovremennaja laboratornaja diagnostika renal'nyh patologij: ot rannih stadij do ostroj pochechnoj nedostatochnosti. Laboratorna diagnostika. 2010; 4 (54): 59-65. [In Russian].
3. Jahn T, Radford S. The Yin and Yang of protein folding. FEBS Journal. 2005; 272 (23): 5962-70. doi: 10.1111/j.1742-4658.2005.05021.

4. *Zabolotnyi DI, Bielousova AO, Zarytska IS, et al.* Autokhtonna β -ahrehatsiia bilkiv: prychny, molekuliarni mekhanizmy ta patolohichni naslidky. Zhurnal NAMN Ukrainy. 2014; 24 (4): 385-92. (In Ukrainian).
5. *Hamley IW.* Peptide fibrillization. *Angew Chem Int Ed.* 2007; 46 (43):8128-47. doi: 10.1002/anie.200700861.
6. *Gertz MA.* Immunoglobulin light chain amyloidosis: 2018 update on diagnosis, prognosis, and treatment. *Am J Hematol.* 2018; 93 (9): 1169-80. doi: 10.1002/ajh.25149.
7. *Sideras K. and Gertz M.* Amyloidosis. *Adv Clin Chem.* 2009; 47: 1-44. PMID: 19634775. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19634775>
8. *Semerdzhiyev SA, Dekker DR, Subramaniam V, Cklaessens MMAE.* Self-assembly of protein fibrils into suprafibrillar aggregates: bridging the nano- and mesoscale. *ACS Nano.* 2014. 8(6): 5543-51. doi: 10.1021/nn406309c.
9. *Sunde M, Serpell LC, Bartlam M, Fraser PE, Pepys MB, Balke CC.* Common core structure of amyloids fibrils by synchrotron X-ray diffraction. *J Mol Biol.* 1997; 273(3): 729-39. doi: 10.1006/jmbi.1997.1348.
10. *Koga T, Taguchi K, Kogiso M, Kobuke Y, Kinoshita T, Higuchi M.* Amyloid formation of native folded protein induced by peptide-based graft copolymer. *FEBS Lett.* 2002; 531 (2): 137-40. PMID: 12417301. doi: 10.1016/S0014-5793(02)03438-5
11. *Smirnov VP, Fadeev MJu.* Bolezninakoplenija (tezaurosomozy). N. Novnogod, Izd-vo NGMA, 2007. 104 s. [In Russian].
12. *Verevka SV.* Parametabolic -Aggregation of proteins: familiar mechanisms with diverse sequels. In: Berhardt LV, editor. *Advances in Medicine and Biology.* New York: Nova Science Publishers; 2013. Vol. 72. p. 29-48. Available from: http://www.novapublishers.org/catalog/product_info.php?products_id=45363
13. *Krebs MRH, Bromley EHC, Rogers SS, Donald AM.* The mechanism of amyloid spherulite formation by bovine insulin. *Biophys J.* 2005; 88 (3): 2013-21. doi: 10.1529/biophysj.104.051896.
14. *Tsvirinko IR, Voroshylova NM, Timchenko MD, Kruhlyk OV, Verevka SV.* β -ahrehovani bilki v patolohichno zminenykh tkanynakh. VI. Osoblyvosti struktury bilka Bens-Dzhonsa ta obumovleni nymy diahnostychni mozhlyvosti. *Laboratorna diahnostyka.* 2018; 2 (81): 3-8. (In Ukrainian).
15. *Zabolotnyi DI, Gogunskaya IV, Zabolotnaya DD, et al.* Suicide antigens: induced denaturation of proteins in the development of allergic reactions. In: Berhardt LV, editor. *Advances in Medicine and Biology.* New York: Nova Science Publishers, 2012, Vol. 53, p.217-32. Available from: http://www.novapublishers.org/catalog/product_info.php?products_id=36951
16. *Sapozhnikov SP, Karyshev PB, Sheptuhina AI, Nikolaeva OV, Avrujskaja AA, Mitrasov JuN, Kozlov VA.* Novye fljuorescentnye zondy dlja vyjavlenija amiloida. *Sovremennye tehnologii v medicine.* 2017; 9 (2): 91-8. [In Russian].
17. *Puchtler H, Sweat F, Levine M.* On the binding of Congo red by amyloid. *Journ Histochem Cytochem.* 1962; 10: 355-64. doi: 10.1177/10.3.355
18. *Kyle RA.* Amyloidosis: a convoluted story. Historical review. *Brit Journ Haematol.* 2001; 114 (3): 529-38. PMID: 11552976. doi: 10.1046/j.1365-2141.2001.02999.x
19. *Brigger D, Muckle T.* Comparison of Sirius red and Congo red as stains for amyloid in animal tissues. *Journ Histochem Cytochem.* 1975; 23 (1): 84-8. doi: 10.1177/23.1.46874.
20. *Westermarck GT, Westermarck P.* Serum amyloid A and protein AA: molecular mechanisms of transmissible amyloidosis. *FEBS Lett.* 2009; 583 (16): 2685-90. doi: 10.1016/j.febslet.2009.04.026.
21. *Buxbaum J, Linke R.* A molecular history of the amyloidosis. *J. Mol. Biol.* 2012; 421 (2-3): 142-59. doi: 10.1016/j.jmb.2012.01.024.
22. *McCarthy F, Adetoba A, Gill C, Bramham K, Bertollacini M, Burton G, Girardi G, Seed P, Poston L, Chappell L.* Urinary comgophilia in women with hypertensive disorders of pregnancy and preexisting proteinuria or hypertension. *Am J Obst Gynecol.* 2016; 215 (4): 1-7. doi: 10.1016/j.ajog.2016.04.041.
23. *Beetham R.* Detection of Bence-Jones protein in practice. *Ann Clin Biochem.* 2000; 37(5): 563-70. doi: 10.1258/0004563001899690.
24. *Saeed SM, Fine G.* Thioflavin-T for amyloid detection. *Am J Chem Pathol.* 1967; 47(5): 588-593. PMID: 4164576. doi:10.1093/ajcp/47.5.588.
25. *Lowry OH, Rosebrough NI, Farr AL, Randall RI.* Protein measurement with the Folin phenol reagent. *J Biol Chem.* 1951; 193 (1): 265-275. PMID: 14907713. Available from: <http://www.jbc.org/content/193/1/265.long>
26. *Anders H-J, Schaefer L.* Beyond tissue injury – damage associated molecular patterns, toll-like receptors, and inflammasomes also drive regeneration and fibrosis. *JAmSocNephrol.* 2014; 25 (7): 1387-1400. doi: 10.1681/ASN.2014010117.
27. *Yildiz G, Andrade J, Engeseth NE, Feng H.* Functionalizing soy protein nanoaggregates with pH-shifting and mano-thermo-sonication. *J. Colloid Interface Sci.* 2017; 505: 836-846. doi: 10.1016/j.jcis.2017.06.088
28. *Kovalska V, Chernii S, Cherepanov S, Losytskyy M, Chernii V, Varzatskii O, Naumovets A, Yarmoluk S.* The impact of binding of macrocyclic metal complexes on amyloid fibrillization of insulin and lysozyme. *Journ Mol Recognit.* 2017; 30 (8): e2622. doi: 10.1002/jmr.2622.