



Are Fungal Endophytes Merely Mycorrhizal Copycats? The Role of Fungal Endophytes in the Adaptation of Plants to Metal Toxicity

Agnieszka Małgorzata Domka¹, Piotr Rozpadek^{2*} and Katarzyna Turnau¹

¹Institute of Environmental Sciences, Jagiellonian University, Kraków, Poland, ²Malopolska Centre of Biotechnology, Jagiellonian University, Kraków, Poland

OPEN ACCESS

Edited by:

Raul Antonio Sperotto,
University of Taquari Valley, Brazil

Reviewed by:

Josélli Schwambach,
University of Caxias do Sul, Brazil
Nuria Ferrol,
Spanish National Research Council
(CSIC), Spain

*Correspondence:

Piotr Rozpadek
piotr.rozpadek@uj.edu.pl

Specialty section:

This article was submitted to
Plant Microbe Interactions,
a section of the journal
Frontiers in Microbiology

Received: 19 November 2018

Accepted: 12 February 2019

Published: 15 March 2019

Citation:


Domka AM, Rozpadek P and
Turnau K (2019) Are Fungal
Endophytes Merely Mycorrhizal
Copycats? The Role of Fungal
Endophytes in the Adaptation of
Plants to Metal Toxicity.
Front. Microbiol. 10:371.
doi: 10.3389/fmicb.2019.00371

The contamination of soil with toxic metals is a worldwide problem, resulting in the disruption of plant vegetation and subsequent crop production. Thus, remediation techniques for contaminated soil and water remain a constant interest of researchers. Phytoremediation, which utilizes plants to remove or stabilize contaminants, is perceived to be a promising strategy. However, phytoremediation's use to date is limited because of constraints associated with such factors as slow plant growth rates or metal toxicity. Microbial-assisted phytoremediation serves as an alternative solution, since the impact of the microbial symbionts on plant growth and stress tolerance has frequently been described. Endophytic fungi occur in almost every plant in the natural environment and contribute to plant growth and tolerance to environmental stress conditions. Although this group of symbiotic fungi was found to form association with a wide range of hosts, including the non-mycorrhizal Brassicaceae metallophytes, their role in the response of plants to metal toxicity has not been thoroughly elucidated to date. This review summarizes the current knowledge regarding the role of endophytic fungi in the tolerance of plants to toxic metals and highlights the similarities and differences between this group of symbiotic fungi and mycorrhizal associations in terms of the survival of the plant during heavy metal stress.

Keywords: fungal endophytes, toxic metals, mycorrhiza, phytoremediation, phytomining

INTRODUCTION

The deposition of toxic metals (TMs) in the topsoils of a significant acreage of land has become a major problem over a wide range of countries from both highly developed and developing regions of the world. Increasing quantities of Zn, Cd, Pb, and Fe that are produced by sewage discharges, mining operations, and runoff from metal-refining industries have severely limited vegetation what negatively affects numerous branches of human activity including food production/agriculture, urbanization, tourism, and other human practices. Difficulties in restoration of metal polluted environments arise from: (1) metal toxicity, (2) low nutrient content, (3) poor physical structure of the substrate, and (4) degraded microbial communities (Rajkumar et al., 2009).



In the natural environment, plants interact with a multitude of symbiotic microorganisms. This interaction includes mycorrhizal fungi (ecto- and endomycorrhizae), other rhizosphere-borne plant growth promoting microorganisms (PGPMs), such as rhizobial and endophytic bacteria, including plant growth promoting rhizobacteria (PGPR) and endophytic fungi (Vandenkoornhuysen et al., 2015). The role of beneficial fungal symbionts in plant metal accumulation and tolerance has been underestimated for many years; however, it has attracted more interest recently (Rajkumar et al., 2009, 2012; Weyens et al., 2009b; Burges et al., 2017). Significant progress has been made in understanding the role of mycorrhizae in metal stress tolerance, and several comprehensive reviews have recently been published (Meier et al., 2012; Cabral et al., 2015; Ferrol et al., 2016; Coninx et al., 2017; Das et al., 2017; Miransari, 2017; Rozpądek et al., 2017). Besides mycorrhiza, another group of symbiotic fungi – fungal endophytes – has been attracting the interest of the scientific community due to their potential beneficial impact on vegetation, including in the facilitation of plant growth metal polluted environments. Endophytic fungi are a taxonomically diverse group of ubiquitous cryptic microorganisms that reside inside their host without causing any visible symptoms of infection for at least part of their life cycle (Rodríguez et al., 2009; Purahong and Hyde, 2011; Sridhar, 2012). Based on the host range, *in planta* colonization, mode of the transmission and biodiversity, two primary groups (divided into 4 classes) of endophytic fungi have been distinguished. The group of clavicipitaceae (C) endophytes (class 1) is closely related species, colonizing systemically cool- and warm-season grasses (Bischoff and White, 2005). While, among non-clavicipitaceae (NC) endophytes, three broad-host range classes can be distinguished, including fungi colonizing systemically host

tissues (class 2), fungi growing exclusively in plant above-ground tissues (class 3), and those restricted to plant roots (class 4) that include dark septate endophytes (DSEs) belonging to Ascomycota and non-mycorrhizal members of Sebaciniales, Basidiomycota (Rodríguez et al., 2009; Andrade-Linares and Franken, 2013).

Fungal endophytes positively affect vegetation as follows: (1) indirectly by soil formation, which is particularly important in degraded environments and (2) directly by fine tuning the adaptation of the plant to metal toxicity and improving the plant biomass yield (Figure 1) (Li et al., 2012b; Coninx et al., 2017; Rozpądek et al., 2018). These traits of fungal endophytes make them attractive for plant-based environment restoration technologies (phytoremediation).

In this review, we summarized the studies available on the role of endophytic fungi in conferring plant toxic metal tolerance, with a particular emphasis on NC endophytes. We also attempted to address the issue of the differences between this group of symbionts and other groups of fungal mutualists, particularly mycorrhizae, in the context of fungal-dependent mechanisms that enable the plant-microbe consortia to survive under the challenge of toxic metals.

ENDOPHYTIC FUNGI: LIFESTYLE AND TAXONOMY

Endophytic fungi exhibit a high degree of lifestyle versatility, and depending on the genetic traits of the partners, developmental stage, nutritional status, and other environmental factors, they can interact with their host in mutualistic, commensalic, or as latent pathogenic as summarized by Schulz et al. (2002).

Such versatility and the loss of virulence can be the result of a single mutation or an epigenetic change due to changes in the environment. However, plants colonized by the endophytic fungi show improved growth, better performance, induced resistance, and the biocontrol of nematodes, fungi, and pathogenic bacteria and fungi. The asymptotic colonization of the plant relies on the balance of the antagonism between the host and the endophyte. This balance can be destroyed when the environmental factors change or the plant reaches the stage of senescence (Deckert et al., 2001).

NC endophytes include Ascomycota, Basidiomycota, and Mucoromycotina (Huang et al., 2001; Rosa et al., 2010; Andrade-Linares and Franken, 2013) that were shown to be able to inhabit bryophytes, ferns, gymnosperms, and angiosperms (trees, shrubs, and herbs), including the non-mycorrhizal Brassicaceae, particularly abundant in metallophytes and hyper-accumulators. The association of Mucoromycotina with ancient bryophytes and liverworts is thought to represent an ancestral plant-fungal interaction and probably played a crucial role in the terrestrialization of plants (Bidartondo et al., 2011). These fungi could have developed from saprophytes that first adapted to become endophytes, and later, those inhabiting the below-ground parts of the plants could have developed into mycorrhizal fungi (discussed in Strullu-Derrien et al., 2018). The interaction certainly took place several times during evolution; thus, the endophytes evolved within distant phylogenetic groups of fungi in a manner similar to that of the mycorrhizal fungi, which facilitated the interaction with almost all plants. The beneficial interaction of the host and its microbiome, including bacteria, archaea, and fungi, is responsible for maintaining the health of the plant (Syamala and Sivaji, 2017).

Endophytic fungi are facultative plant symbionts (facultative biotrophs) and according to Brundrett (2002), in contrast to mycorrhizal fungi, their development is not synchronized with the development of their hosts. Thus, fungal endophytes may complete their lifecycle outside the host organism and thus are able to grow on artificial media (Petrini, 1996), which facilitate the manufacture of pure inoculum under sterile conditions and eliminate the difficulty of their propagation. This fact is particularly important in relation to phytoremediation and the large-scale production of inoculum.

Another distinct feature of the fungal endophytes (species belonging to classes 1, 2, and 3) that distinguishes them from mycorrhizae is their ability to colonize the above-ground organs of the plant. In contrast to mycorrhizal fungi, NC endophytic fungi were found in plant leaves, stems, flowers, and seeds (Yuan et al., 2010; Hardoim et al., 2015), while mycorrhizae are restricted to the roots of plants. In addition, a large number of species can exist in the form of mycelia or form yeast-like structures or mycosomes (Atsatt and Whiteside, 2014). This aspect of endophytic fungal biology is particularly interesting because the morphological changes described may occur during the colonization of the plant host. This may facilitate the colonization process by evading the detection system of the hosts. The invader would lose virulence factors present on the surface of the cell wall that allows detection by the plant. This step would resemble the entrance of the arbuscule

into the mycorrhizal plant cells that result in the formation of the specific interface between the partners that facilitate the transfer of substances between the partners (Brundrett, 2002). Interestingly, endophytic fungi lack this specialized interface and communicate with their hosts using relatively unspecialized hyphae (Brundrett, 2006).

In contrast to mycorrhizal fungi, endophytic fungi possess the ability to colonize plants belonging to the Brassicaceae, which are particularly abundant in metal hyper-accumulators and metallophytes; hyper-accumulators have been found in 45 families of plants with a large number belonging to this family (van der Ent et al., 2013). Similar to the mycorrhizal fungi, endophytic fungi were shown to promote plant growth under nutrient-limiting conditions (Hiruma et al., 2018). Such conditions are frequently encountered in metalliferous soils. In general, the Brassicaceae do not associate with mycorrhizae, probably due to the loss of symbiotic genes and the lowered expression of the nucleotide-binding site leucine-rich repeat resistance proteins (NLRs) required for the arbuscular mycorrhizal fungi (AMF) (reviewed in Hiruma et al., 2018). However, there were a few reports of AM fungi present in some Brassicaceae, such as the Cd hyper-accumulator *Biscutella laevigata* (Orłowska et al., 2002) or the Zn hyper-accumulator *Thlaspi caerulescens* (Regvar et al., 2003) collected from sites rich in TM. These cases are presently interpreted as the endophytic growth of mycorrhizal fungi, since the presence of arbuscules is mostly ephemeral, often only observed at the flowering stage, and the cases are rare, being observed only under field conditions and not confirmed under greenhouse conditions (Brundrett and Tedersoo, 2018). While endophytic fungi dominate non-mycorrhizal plants, mycorrhizal plants are also inhabited by numerous endophytes, including species of fungi that inhabit the Brassicaceae (Chen et al., 2012; Wężowicz et al., 2014; Sim et al., 2018). This finding is particularly interesting in the context of symbiotic fungi function. If the roles of these different symbionts overlap, how do they affect each other's function? Another pending question is what are the specific functions of the endophyte that distinguish them from mycorrhiza? It appears that the response of the plants to inoculation with a single fungus (mycorrhizal or endophyte) differs from the response to inoculation with species of both groups of fungi (Wężowicz et al., 2017; Berthelot et al., 2018; Ważny et al., 2018; Zhou et al., 2018). The costs of harboring multiple fungal symbionts are higher but increased gains compensate for this (Field et al., 2016). The interaction between these two groups of fungi *in planta* or the effect of the presence of mycorrhizal fungi on the diversity and function of fungal endophytes has not been thoroughly investigated. Such studies would broaden our understanding of the role of different fungi in plant biology and the plasticity of the plant-fungus interaction. Available studies indicate that fungal endophytes may play a role similar to that of mycorrhizae (Rodriguez et al., 2009; Rudgers and Swafford, 2009; Andrade-Linares and Franken, 2013; Hiruma et al., 2018). How this role is affected by the presence of mycorrhizal fungi and how the fungal lifestyles change during the presence of a potential competitor remain to be investigated.

TOXIC METAL TOLERANCE OF ENDOPHYTIC FUNGI

Certain strains of endophytic fungi exhibit extraordinarily high resistance to toxic metals. In culture, endophytic fungi can withstand TM concentrations in the mM ranges (Deng et al., 2011; Khan et al., 2017a; Domka et al., 2019). This adaptation provides a competitive edge over non-adapted fungi in colonizing plants in metalliferous habitats (Gadd, 2016). The role of symbiosis in the evolution of the fungal TM tolerance is unknown. It remains to be elucidated whether the TM tolerant plant served as shelter for the fungus, allowing it to evolve metal tolerance under the TM stress, or whether the metal-tolerant fungus facilitates the growth of the host plant in metalliferous soils. In the case of the former, the ability to colonize plant tissues in addition to its metal tolerance would play a role in determining the adaptation of the endophytic fungi to metal toxicity.

Many TM-tolerant fungal species, such as representatives of the genera *Phomopsis* and *Bipolaris* (DSE, class 4), have been isolated from many plant families (Ban et al., 2012; Wężowicz et al., 2014; Zhang et al., 2014; Sim et al., 2018; Domka et al., 2019). Studies indicate that endophytic fungi isolated from plants growing in areas polluted with TM are more tolerant to pollution and thus can be potentially more appealing for use in microbial-assisted phytoremediation. For example, DSEs isolated from *Alnus nepalensis* were shown to possess a higher tolerance to Cd than isolates from less polluted stands (Xu et al., 2015). In another study, 50% of the endophytic fungi isolated from six dominant plant species growing on Pb-Zn wastelands were shown to be tolerant to Pb²⁺ or Zn²⁺ (Li et al., 2012a). In addition, the growth of some of these fungi was even stimulated by the TM. The authors suggested that certain fungal species could adapt to metal toxicity due to their long-time exposure to the TM. Two strains of the endophytic fungi isolated from *Solanum nigrum* L. growing in a Cd-contaminated environment possessed the highest tolerance to Cd concentrations of up to 2 mM (Khan et al., 2017b). Similarly, Sim et al. (2018) isolated highly tolerant fungal endophytes species from *Phragmites* sp. utilized in wastewater cleanup. The majority of the fungi were able to withstand Pb, Zn, and Cu concentrations over 200 mg l⁻¹. Interestingly, Gadd (2016) reported that certain fungi that are not necessarily fungal endophytes not adapted to TM can exhibit significant TM tolerance and become dominant in environments polluted with heavy metals; thus, resistance is not necessarily related to their environmental origin. The TM tolerance of some non-adapted fungi was shown to be similar to that of the fungal strains from the sites polluted with TM (Gadd, 2016). This finding suggests that inherent tolerance (probably not TM-related) mechanisms may have been the determining factor in the selection of the fungal endophytes that inhabit metalliferous environments.

High quantities of toxic metals deposited in soils cause qualitative and quantitative changes in microbial populations (Chodak et al., 2013; Corneo et al., 2013). Li et al. (2016) investigated the diversity of endophytic fungi isolated from

Dysphania ambrosioides occurring at two TM-polluted stands. These researchers identified significant differences in the fungal population structure and dominant genera that were present between the sites analyzed. The colonization rate of the plants selected from stands highly polluted with TM was significantly lower than those from less polluted locations. Among all the strains isolated, regardless of origin, 50% were shown to be TM tolerant. Dominant genera exhibited higher tolerance. However, more metal tolerant isolates were found in plants from the less polluted stand. There were no differences in the Shannon index (H') between the sites, which indicate similarity in fungal diversity. Interestingly, according to Wężowicz et al. (2014) fungal populations isolated from *Verbascum lychnitis* growing on post-mining tailings were more diverse than those from the non-polluted sites.

Mechanisms of Fungal Metal Tolerance – The Role of Glutathione

Metal tolerance mechanisms in fungi that are described most frequently involve glutathione (GSH) and GSH-related tolerance that are induced in response to a variety of environmental factors (Na and Salt, 2011; Saraswat and Rai, 2011). Thus, the tolerant fungal strains, not necessarily TM adapted, that possessed the ability to colonize plants would be the driving force in the plant-fungal symbiosis in habitats polluted with heavy metals. Although endophytic fungi were shown to be able to accumulate extremely high quantities of TM in their mycelia, there is a lack of a comprehensive description of the mechanisms of metal transport, sequestration, and detoxification for this group of fungi (Deng et al., 2014; Zahoor et al., 2017). Zhao et al. (2015a) performed a transcriptome analysis of *Exophiala pisciphila* (DSE) growing in the presence of Cd using RNA-Seq. These researchers found several upregulated genes associated with TM binding, transport and detoxification, reactive oxygen species scavenging, redox homeostasis maintenance, and sulfur assimilation. The authors suggested that sulfate-containing molecules could increase during Cd stress, including glutathione (GSH), phytochelatin, and metallothioneins (Na and Salt, 2011). After uptake, the toxic metals are neutralized by complexation in the cytosol and translocated for storage in the vacuole. The vacuolar compartmentalization of metal chelates enables the fungi to isolate potentially dangerous pollutants from sensitive cellular compartments. Out of the numerous metal chelating molecules present in living organisms, the thiol-containing substances GSH (glutathione) and metallothioneins were shown to be present in, e.g., mycorrhizal fungi (Saraswat and Rai, 2011). The tripeptide GSH (γ -glu-cys-gly) is a non-protein thiol found abundantly in all cell compartments. In addition to its ability to sequester metals, it acts as a potent antioxidant and removes the damage caused by oxidative stress (Freeman, 2004). In addition, Ban et al. (2012, 2017) reported that Pb treatment leads to a transient GSH increase in non-pathogenic *Gaeumannomyces cylindrosporus* (DSE) that improves tolerance to Pb. Apart from GSH-dependent tolerance, Wei et al. (2016) analyzed the role of an *Nramp* (*natural resistance-associated macrophage protein*) gene from the DSE *E. pisciphila*. This gene encodes the plasma membrane bivalent cation transporter.

The expression of *EpNramp* is downregulated after Cd exposure. Its heterologous expression in yeast Fe-uptake mutants leads to growth complementation. The authors suggested that this protein is partially involved in the tolerance of *E. pisciphila* to cadmium.

Fungal Endophytes Limit Metal Toxicity by Exuding Metal-Binding Molecules Into the Rhizosphere


In addition to immobilizing the TM within its hyphae, fungi can limit the toxicity of TM by secreting various types of agents that chelate metals (Meharg, 2003; Bellion et al., 2006; Cabral et al., 2015), such as citrates or oxalic acid (Gadd, 1999). Available reports indicate that AMF are not able to secrete these types of organic acids but were shown to synthesize and release the glycoprotein glomalin (reviewed in Meharg, 2003; González-Chávez et al., 2004). Chelator exudation is an important mechanism of TM stabilization in the soil and prevents the entry of TM into the plant roots. The synthesis and secretion of metal-chelating molecules, including organic acids, siderophores, exopolysaccharides (EPSs), and phenolic compounds by fungal endophytes similarly to other mycorrhizal fungi with the exception of AMF, were reported (Bartholdy et al., 2001; Koulman et al., 2012; Zhao et al., 2015b; Garza et al., 2016; Gupta and Joia, 2016; Yamaji et al., 2016; Tong et al., 2017). In addition, a small number of reports indicate that fungal endophytes may bind TM ions to their cell wall components, similar to mycorrhizae (Tong et al., 2017). The production of siderophores by endophytic fungi was reported on a few occasions. Koulman et al. (2012) showed that the clavicipitaceous endophyte *Epichloë festucae* synthesizes siderophores with an unusual structure. In another study, Johnson et al. (2013) showed that the *E. festucae sid* gene, encoding a siderophore synthase, is necessary to establish symbiosis between the fungus and *Lolium perenne*. The plants infected with the $\Delta sidN$ mutants were stunted, and detailed microscopic studies revealed abnormalities in the distribution and localization of the mutant hyphae (Johnson et al., 2013). Bartholdy et al. (2001) reported the production of siderophores by the DSE fungus *Phialocephala fortinii*. HPLC analysis identified ferricrocin, ferrirubin, and ferrichrome C in the culture filtrate. Experiments performed by Yamaji et al. (2016) showed that the two DSE strains *P. fortinii* and *Rhizoderma veluwensis* were able to produce siderophores that are probably involved in the TM immobilization in the rhizosphere. Ban et al. (2012) reported that the melanin content in *G. cylindrosporus* increased following supplementation with Pb. Melanin is considered to be the most important component of the fungal cell wall involved in the alleviation of TM stress. There are numerous studies reporting its ability to bind metal ions; however, the role of melanin in the TM tolerance of the DSE remains unclear (Bruenger et al., 1967; Gadd, 1993; Fogarty and Tobin, 1996; Ban et al., 2012).

In addition, most endophytic fungi studied to date were shown to possess the ability to form nanoparticles (NPs), what may subsequently reduce the metal toxicity exerted on the host plants, as well as on the fungi (Durán et al., 2011).

The efficiency of NPs mycosynthesis differed in between species of fungi (Shankar et al., 2003; Qian et al., 2013; Devi and Joshi, 2015). Recent studies indicate that NPs may be used in agriculture as nanofertilizers (Liu and Lal, 2015) that facilitate uptake of micronutrients (Dimkpa et al., 2015). Fungi utilize cellular enzymes, proteins, and membrane-bound molecules as electron donors in the reduction reaction. Reduced metal ions can be precipitated as NPs intracellularly or extracellularly (reviewed in El Enshasy et al., 2018). It was proposed that during intracellular synthesis metal ions are at first electrostatically bound to the fungal cell wall or diffuse through it and are subsequently reduced and precipitated by enzymes present in the cytoplasmic membrane (e.g., ATPases, hydrogenases) (Vahabi and Dorcheh, 2014). This leads to nanoparticle formation. The extracellular synthesis of nanoparticles occurs through the release of reductase enzymes (e.g., nitrate reductase, NADPH-dependent reductases, and FAD-dependent glutathione reductase) by metal exposed fungi and subsequent reduction of metal ions to form NPs (Kashyap et al., 2013; Singh et al., 2016; El Enshasy et al., 2018). Devi and Joshi (2015) reported the ability of three endophytic fungi to biosynthesize silver NPs. Similar properties were shown for the endophytic *Epicoccum nigrum* (DSE) and *Colletotrichum* sp. (Shankar et al., 2003; Qian et al., 2013). Also, Netala et al. (2016) have shown that the synthesis of silver NPs by the endophytic fungus *Pestalotiopsis microspora* VJ1/VS Yeasts from the *Cryptococcus* and *Rhodotorula* genus was also shown to produce silver NPs on several occasions (Salvadori et al., 2014; Fernández et al., 2016). Data concerning the ability of mycorrhizal fungi to synthesize NPs are scarce. Manceau et al. (2008) investigated copper speciation at the soil-root interface and found that copper was biotically reduced to NPs by mycorrhizal plants. Also, González-Chávez et al. (2002) reported the formation of electron-dense Cu granules within hyphae of AMF isolated from Cu- and As-contaminated soil, what suggests that mycorrhizal fungi also can produce nanoparticulate copper. However, more studies are required to understand the synthesis of the NPs in the plant rhizosphere, their effect on plants and microbial tolerance to TM, and their potential use in phytoremediation strategies.

PLANT METAL TOLERANCE INDUCED BY THE FUNGI

Plants inhabiting sites rich in toxic metals (TMs) developed a variety of mechanisms that allow them to survive in extremely adverse environments. Two primary strategies have evolved as an adaptation to metal toxicity. Plants may either accumulate high quantities of TM and internally neutralize toxic elements by a highly effective and complex detoxifying system or employ avoidance/exclusion strategies that limit metal toxicity (reviewed in DalCorso et al., 2013). According to available studies, fungal endophytes may impact both strategies depending on the strategy employed by its host (Jiang et al., 2008; Deng et al., 2013, 2014; Zhu et al., 2015; Zahoor et al., 2017; Rozpądek et al., 2018) (Figure 2) (Table 1). For example, Rozpądek et al. (2018) showed that endophytic *Mucor* sp. affects TM avoidance strategy of



Arabidopsis arenosa, leading to the decrease of Zn and Fe uptake by the plant and simultaneously to upregulation of root to shoot translocation of Zn, Fe, and Cd. During interaction with the endophyte, plant genes implicated in metal homeostasis were affected. In turn, Jiang et al. (2008) revealed that *Brassica juncea* inoculated with Acacia-derived endophytic fungi was able to accumulate more Cd and Ni, depending on TM present in the soil, than wild type. These findings imply that the metal tolerance mechanisms of the host affected by the symbiotic fungal endophytes are not specific or novel but rather are differentially expressed and regulated. An important aspect of plant metal stress tolerance is the optimal metal distribution between the plant organs and the cell compartments that enable effective metal detoxification and protect the sensitive cellular compartments from metal toxicity. The accumulation of high quantities of TM has received a substantial amount of attention due to the importance of hyper-accumulating plants in phytoremediation and phytomining (reviewed in Verbruggen et al., 2009; van der Ent et al., 2013). Plant metallophytes have evolved highly effective metal detoxification mechanisms. In the case of hyper-accumulators, these mechanisms allow them to translocate and accumulate high quantities of metals in their shoots, which are associated with the upregulation of the genes encoding metal transporters enabling them to optimize metal distribution within the plant (Verbruggen et al., 2009). The number of studies describing the role of fungi, primarily AMF (Chen et al., 2003, 2004; Leung et al., 2006; Vogel-Mikuš et al., 2006; Punamiya et al., 2010), in the adaptation of hyper-accumulators is limited, and to our knowledge, the endophytic mycobiota of this group of plants has not been thoroughly investigated. In the work of Vogel-Mikuš et al. (2006), the influence of AM fungi on uptake of Zn, Cd, and Pb by *Thlaspi praecox* was investigated. The authors

revealed that AM colonization resulted in higher nutrient uptake and lower TM accumulation by plants. In contrast, Punamiya et al. (2010) showed that *Chrysopogon zizanioides* accumulated more Pb in association with the AMF *Funneliformis mosseae*.

An Extension/Filter of the Root?

Fungi can efficiently immobilize TM and limit metal uptake by the plant due to the large surface of the mycelia and the development of various mechanisms to tolerate and detoxify metals. This finding is particularly important because the hyphae of symbiotic fungi spread out from the root into the surrounding soil and significantly extend the root surface area. It was suggested that the extraradical hyphae of the AMF could represent up to 90% of the total AMF biomass and account for approximately 25% of the total microbial biomass in agricultural soils (Leake et al., 2004). This allows the efficient supply of the plant with the simultaneous filtering of the uptake of nutrients and water (Joner, 2000). Arbuscular mycorrhizal symbiosis was shown to significantly alter the root morphology, usually leading to a reduction in the root/shoot ratio. These changes result in a high degree of dependence of the plant on the symbiosis; AMF are responsible for the majority of nutrient absorption and delivery to the host plant (Sanders et al., 1977; Hetrick, 1991). In ectomycorrhizal roots (ECM), as an adaptation to symbiosis, the number and length of the root hairs are significantly limited. This finding implies that as in AMF, the majority of water and nutrient acquisition is transmitted through the fungal hyphae (Hetrick, 1991). Turnau et al. (1996) revealed the filtering role of the ectomycorrhizal fungus *Rhizopogon roseolus* hyphal mantle formed on *Pinus sylvestris* roots. The authors investigated the distribution of potentially toxic metals within mycorrhizae and showed the enhanced accumulation of Cd and Al in the

TABLE 1 | Comparison of plant toxic metal tolerance mechanisms activated by symbiotic fungi.

Mechanism		Endomycorrhizal fungi (ECM)	Arbuscular mycorrhizal fungi (AMF)	Endophytic fungi
TM avoidance	Reduced TM transfer to the plant	<ul style="list-style-type: none"> Biosorption (Chalot et al., 2000; Colpaert et al., 2011) Bioaccumulation (Turnau et al., 1996) 	<ul style="list-style-type: none"> Biosorption (Cabral et al., 2015) Bioaccumulation (Janoušková et al., 2006; Meier et al., 2012; Corneo et al., 2013) 	<ul style="list-style-type: none"> Biosorption (Tong et al., 2017) Bioaccumulation (Zahoor et al., 2017)
	Reduced TM bioavailability	<ul style="list-style-type: none"> Metal transformation (Gadd, 1993) Secretion of chelating agents (e.g., organic acids) (Bellion et al., 2006; Ray and Adholeya, 2009) 	<ul style="list-style-type: none"> Metal transformation (Gadd, 1993; Wu et al., 2015) Secretion of chelating agents (e.g., organic acids,glomalin) (González-Chávez et al., 2004; Cabral et al., 2015) 	<ul style="list-style-type: none"> Metal transformation (Gadd, 1993) Secretion of chelating agents (e.g., organic acids, siderophores, phenolic compounds) (Bartholdy et al., 2001; Koulman et al., 2012; Zhao et al., 2015b; Yamaji et al., 2016; Tong et al., 2017)
Increased TM concentration in the plant	Increased availability of TM	<ul style="list-style-type: none"> Secretion of chelating agents (Fomina et al., 2005; Machuca et al., 2007; Vamerali et al., 2010) Metal transformation (Gadd, 1993) 	<ul style="list-style-type: none"> Secretion of chelating agents (Göhre and Paszkowski, 2006) Metal transformation (Gadd, 1993; Li et al., 2016) 	<ul style="list-style-type: none"> Metal transformation (Gadd, 1993)
	Increased TM uptake and translocation	<ul style="list-style-type: none"> Indirectly through plant growth promotion (Sell et al., 2005) Regulation of the expression of host genes encoding metal transporters (Ma et al., 2014) 	<ul style="list-style-type: none"> Metal transport from fungal hyphae to plant cells (Göhre and Paszkowski, 2006) Indirectly through plant growth promotion (Usman and Mohamed, 2009) Regulation of the expression of host genes encoding metal transporters (Burleigh et al., 2003; Bona et al., 2010) 	<ul style="list-style-type: none"> Indirectly through plant growth promotion (Zhu et al., 2015)

fungal mantle and their gradual decrease along the Hartig net toward the inside of the root. A similar filtering effect was observed in the *Suillus luteus/P. sylvestris* mycorrhizae in which Ca, Fe, Zn, and Pb were concentrated in the mantle and rhizomorph (Turnau et al., 2001). A limited number of studies indicate that some species of root-bound endophytic fungi (e.g., *Colletotrichum tofieldiae*, *Serendipita indica*, Helotiales F229) may behave similarly to mycorrhizae and extend out of the root to increase the root surface area and transmit nutrients to the plant (Yadav et al., 2010; Hiruma et al., 2016; Almario et al., 2017). However, this phenomenon has not been confirmed in the presence of TM. The fungi's role as a physical barrier preventing contact with TM cannot be ruled out, but to the best of our knowledge, there are no studies that would provide clear evidence that endophytic fungi entwine plant roots and serve as a physical protection against TM. Additionally, several distantly related groups of endophytic fungi were shown to activate root hair elongation and induce other alterations in the root architecture leading to an increase in the root surface (Harman et al., 2004; Stewart and Hill, 2014; Hiruma et al., 2016). One type of fungus, *Mucor* sp., was shown to activate the root hair elongation of its host *A. arenosa* in non-polluted soil, as well as in substrates polluted with TM (Rozpądek et al.,

2018). In mycorrhizal plants, the fungal hyphae partly take over the root functions, enabling the plant to grow relatively smaller roots that are less specialized for water and nutrient uptake. Endophytic fungi activate root growth and thus allow the plant to penetrate the surrounding soil more thoroughly but with its own tissues. It cannot be ruled out that fungal endophytes increase the surface of the plant root and simultaneously extend into the rhizosphere and serve as a filter that binds and prevents TM uptake by the plant. This situation may be true in the case of non-mycorrhizal plants, but in nature, most plants are inhabited by endophytic fungi and other fungal symbionts, including mycorrhizae (Wężowicz et al., 2017). Fungal endophytes were shown to occur within the roots and form complex webs with mycorrhizal hyphae (Horton et al., 1998; Agler et al., 2016). The majority of the studies available describe plant-fungal interactions in single inoculation experiments. This approach has several advantages but unfortunately neglects the full spectrum of the plant-fungus and symbiotic fungus-fungus interactions that occur in natural environments. The plant response to symbiotic microorganisms appears to be far more complex than the responses described in single inoculation experiments, and its description needs to include the complexity of microorganisms inhabiting its host.

Endophytic Fungi Affect Toxic Metal Uptake and Distribution Within the Plant

After overcoming the physical barriers of the root, the TM efflux into the cell, vacuolar sequestration, and xylem loading is coordinated by a number of heavy metal transporters, a broad group of different proteins, such as CPx-ATPases for Cu or Cd, ABC transporters for Cd transport into the vacuole, ZIP transporters (ZRT- and IRT-related proteins for Fe or Zn, respectively), and Nramp transporters. Although this network is thought to balance the concentration and partitioning of essential metals, such as Zn, it also unselectively transports toxic elements, such as Cd (reviewed in DalCorso et al., 2013). Metal-specific and non-specific transporter proteins enable metal compartmentalization and subsequent stress alleviation by toxic metal complexation with organic molecules, such as metallothioneins, organic acids, and phytochelatins (PCs) that are synthesized by the plant. The abundance of these molecules based on studies with overexpression lines and hyper-accumulators correlates with the ability to accumulate TM (Freeman, 2004; Janoušková et al., 2005).

There are several studies that confirm the influence of endophytic fungi on the TM uptake and distribution in the host plants. Inoculation with the fungal endophyte *Mucor* sp. (class 2) isolated from plants growing on soils heavily polluted with TM resulted in a decrease in Cr uptake of up to 90% by *Brassica campestris*. The fungus was able to bio-transform the metals deposited in the soil and accumulate them in its hyphae, thereby making them less available to the plant (Zahoor et al., 2017). Recently, similar results were obtained for a strain of *Mucor* sp. isolated from *A. arenosa* inhibiting the “Boleslaw” Zn, Cd, and Pb mine dump in southern Poland. Inoculation with the fungal strain UNIJAG.PL50 improved the production of plant biomass, decreased the accumulation of metals, and significantly affected the metal distribution within the plant, leading to more effective TM translocation to the above-ground parts and subsequently to more uniform TM distribution (Rozpądek et al., 2018). The same *Mucor* sp. strain was simultaneously used with *Rhizoglosum intraradices* and induced increased Zn uptake in *Lactuca serriola* growing on TM-rich industrial wastes. The concentration of Zn was higher in both the roots and shoots of the symbiotic plants (Ważny et al., 2018). In turn, the inoculation of maize with the TM-tolerant DSE *E. pisciphila* strain H93 alleviated the deleterious effects of the presence of TM by reducing the translocation of the HM ions from the roots to the shoots (Li et al., 2011). Similarly, He et al. (2017) reported the positive effect of *E. pisciphila* on maize growth under high Cd and the reduced translocation of the Cd to the shoots. Yamaji et al. (2016) showed that *Clethra barbinervis* inoculated with its fungal endophytes accumulated less TM than the control plants, while Zhu et al. (2015) reported that two species belonging to the *Mucor* genus, *M. circinelloides* (Z4) and *M. racemosus* (Z8), improved the growth and the accumulation of Pb and Cd of Guizhou oilseed rape. Both strains increased the accumulation of the TM in the plant by 117.6% for Cd and 63.48% for Pb, simultaneously reducing their concentrations in the substratum by 60.57% and 27.12%, respectively. Another endophyte that was shown to improve the growth of *Brassica napus* and *B. campestris* in TM-polluted soil was *Rhodotorula* sp. This group of yeast endophytes (class 2) is particularly

interesting due to their ability to produce exopolysaccharides (EPSs) with a high potential to absorb pollutants, including TM (Garza et al., 2016; Fernández et al., 2018). Inoculation with this *Rhodotorula* sp. increased the efficiency of the plant to extract Cd, Cu, and Pb (Wang et al., 2013). In addition, Deng et al. (2014) demonstrated that *Lasiodiplodia* sp. was an endophytic fungus of rape that showed promise for bioremediation. The fungal strain increases the translocation rate of the Cd from the roots to the shoots. *Solanum nigrum* was shown to accumulate more Cd in the roots and shoots after inoculation with the DSE *Phomopsis fukushii* strain PDL-10. He et al. (2017) showed similar results. Studies reported the influence of *E. pisciphila* on maize root-colonizing DSE and Cd tolerance. Plants inoculated with the fungus accumulated more Cd in the roots, and the metal translocation to the above-ground parts was limited (He et al., 2017). Ni uptake and most importantly extraction efficiency were positively affected in *Brassica juncea* inoculated with *Trichoderma atroviride* strain F6 (also belonging to class 4, root inhabiting Hypocreales). In addition, the inoculated *B. juncea* exhibited a higher tolerance to metal toxicity (Cao et al., 2008).

Despite the increasing number of reports examining the influence of endophytic fungi on TM uptake and translocation in the host plant, only a small number of studies attempted to investigate the role of endophytic fungi with respect to the mechanism of this aspect of plant TM metabolism. Rozpądek et al. (2018) showed that the reduction of metal accumulation and changes in their distribution within *A. arenosa* were accompanied by the upregulation of several metal homeostasis-related genes involved in metal exclusion, sequestration in the vacuole, and ROS (reactive oxygen species) scavenging. The upregulated genes include *hma3* (heavy metal associated 3), *mtp1* (metal transport protein 1), *zif1* (zinc induced facilitator 1), and *cax2* (cation exchanger 2) that encode tonoplast-bound vacuole carrier proteins (Rozpądek et al., 2018). The overexpression or complementation with these genes conferred tolerance to metal stress to plants and yeast, respectively (Haydon and Cobbett, 2007; Koren'kov et al., 2007; Morel et al., 2009). In hyper-accumulating *T. caerulea* and *Arabidopsis halleri*, phloem loading is accomplished by the highly expressed *hma2* (heavy metal associated 2) and *hma4* (heavy metal associated 4). To date, there have been no reports indicating that fungal endophytes may affect metal translocation from the root to the shoot by interfering with the expression of these metal transporters. However, the expression of *pcr2* (plant cadmium resistance 2) involved in Zn phloem loading and metal efflux out of the root epidermal cells (Song et al., 2010) was upregulated, and metal accumulation was higher in the shoots of inoculated plants (Rozpądek et al., 2018). In another study, Wang et al. (2016) analyzed the response of maize inoculated with DSE in the presence of Cd. These researchers showed that inoculation with the endophytes led to the inhibition of an unspecified *zip* gene, which coincided with the decreased uptake of Cd following DSE infection. In addition, *mtp1* expression was also activated in the plants inoculated with DSE during Cd toxicity.

Mycorrhizal fungi were also shown to affect the uptake of TM and their subsequent translocation and accumulation by the host plants (Table 1). The expression of metal transporter-encoding

genes, including proteins belonging to Nramp, Zip families, HMA4 transporter, or arsenic putative transporter POR 29, was activated in the AMF plants (Burleigh et al., 2003; Ouziad et al., 2005; Bona et al., 2010; Ma et al., 2014). Based on the current knowledge, the AM fungi, depending on the fungal and plant species, may inhibit the TM uptake by the host plant or increase the metal translocation to the above-ground parts of the plant (Joner and Leyval, 1997; Chen et al., 2007). Ectomycorrhizal fungi were proven to serve as a barrier against TM uptake, since they were reported to inhibit the uptake, translocation, and accumulation of the TM in the upper parts of the plant (Adriaensen et al., 2004, 2005, 2006; Walker et al., 2004). Depending on the fungal species, the TMs were shown to accumulate mostly in the hyphal mantle or in the extramatrical mycelia (Turnau et al., 1996, 2002).

The Influence of Endophytic Fungi on TM Detoxification by the Plant

Endophytic fungi were also shown to affect the expression of the stress-related genes of the host plant, thus triggering resistance against TM toxicity. In maize, inoculation with the DSE *E. pisciphila* (Wang et al., 2016) led to the upregulation of *pcs* expression. The authors suggested that this could facilitate the transpeptidation of glutathione, as well as other thiols. Khan and Lee (2013) reported that soybean plants inoculated with the class 3 endophytic *Penicillium funiculosum* produced more reduced glutathione during Cu stress than control non-inoculated plants. In another study by the same authors, the interaction of the plant with endophytic strain RSF-6L led to increased catalase activity in *S. nigrum* during Cd toxicity (Khan et al., 2017a). The effect of inoculation on the activity and abundance of antioxidants was confirmed in several other studies (Wang et al., 2016). Endophytic *Trichoderma atroviride* and *T. virens* (class 4) were shown to influence the levels of expression of the *adc1* and *adc2* genes implicated in spermidine biosynthesis in *A. thaliana* in the control (Salazar-Badillo et al., 2015). The synthesis of the stress protective polyamines spermidine and spermine is often activated following exposure to TM, and their abundance often correlates with stress tolerance. However, there is a lack of studies indicating the role of polyamines in the resistance to TM triggered by the fungal endophytes.

Gene expression studies confirm the role of mycorrhizae in conferring plant tolerance against TM by regulating the expression of the plant stress response genes. The genes that encode plant defense compounds that are described the most frequently are those involved in the upregulation of sulfur metabolism (sulfur containing compounds), such as metallothionein, GSH-synthase, ascorbate peroxidase, or glutathione S-transferase (Bona et al., 2010; Cicatelli et al., 2012; Lingua et al., 2012; Shahabivand et al., 2017).

Plant Growth Promotion Imposed by the Fungi

There is no unequivocal “mode” of fungal action in terms of the effect on metal uptake and distribution. Some species of fungi protect the plant by decreasing metal uptake, while other species increase uptake and translocation from the root to the

shoot. In addition, there is no consensus regarding the mechanism of the alleviation of metal toxicity by fungi. It remains to be elucidated whether symbiotic fungi improve plant growth during metal toxicity by facilitating water and nutrient uptake by the plant and thus indirectly by fine tuning the metal tolerance of the host or directly by upregulating specific mechanisms of metal tolerance. In the majority of studies available, the plants inoculated with endophytic fungi produce more biomass during metal toxicity than the non-inoculated plants (Zhu et al., 2015; Wang et al., 2016; Rozpądek et al., 2018). However, it is not clear whether this is a result of fungal dependent stress alleviation or a strategy that allows the plants to “dilute” the accumulated metals due to improved growth as a result of enhanced nutrition. Interestingly, the ability to facilitate nutrient uptake was a trait that distinguished the fungal endophytes from the mycorrhizae. Endophytes were thought to be unable to improve nutrient uptake by the plant, but in recent years, fungal endophytes were shown to participate in plant nutrition (Brundrett et al., 2002). Fungal endophytes improve plant water and nutrient uptake. It has been reported that toxic metals often interfere with the root uptake of nutrients, such as Fe, P, Mg, K, Ca, and Zn, and with the metabolic functions of the essential nutrients, leading to plant growth retardation (Ouzounidou et al., 2006). In addition, soils augmented with toxic metals are often highly deficient in nutrients, which impose additional restraints to plant growth. Under such conditions, the fungal endophytes improve the acquisition of plant nutrients by mobilizing nutrients and making them available to the plant roots (Cui and Nobel, 2006; Weyens et al., 2009a; Rajkumar et al., 2012). The Glomeromycota are obligatory biotrophs and lack saprotrophic capabilities. Thus, their ability to decompose organic matter and to liberate compounds that are absorbable for the plant into the soil is limited. In contrast, endophytic fungi are saprotrophic microorganisms that can improve plant nutrition by decomposing organic matter (Promputtha et al., 2007; Purahong and Hyde, 2011). This trait can be particularly useful in environments polluted with TMs that are devoid of available nutrients.

AMF develop specialized structures – arbuscules – that allow an exchange between the fungus and the plant. Arbuscules are branched intracellular hyphae that are present in the root cortex. The absence of these structures in fungal endophytes suggests that the mechanism of nutrient transfer at the plant-endophyte interface differs from that employed by the AMF. Interestingly, the Mucoromycotina of the bryophytes formed characteristic intracellular coils of an unknown function (Field et al., 2015). We can only hypothesize that these structures may play a similar role in Mucoromycotina to that of the arbuscules in the AMF. Another possible way to improve plant nutrition by fungal endophytes is the rhizophagy cycle hypothesized to act in the interaction between plants and endophytic bacteria and yeast (Paungfoo-Lonhienne et al., 2010; White et al., 2018). The rhizophagy theory suggests that fungi and endophytic bacteria can be the prey of the plant root cells and serve as nourishing factors that enable the plants to survive even without available nutrients in the surroundings.

Endophytic fungi are associated with a wide range of plant growth promoting activities, including P solubilization and the

production of siderophores and phytohormones (Chhabra and Dowling, 2017). The role of fungal endophytes in the delivery of P to the plant was described in numerous studies. Hiruma et al. (2016) showed that *C. tofieldiae* transfers P to the host plant and that the phosphate starvation response (PSR) mechanism that is active in plant tissues is responsible for the control of root colonization. In another study, Rozpadek et al. (2018) showed that *Mucor* sp. improved host plant nutrition by inducing the PSR and probably stimulating the plant to produce enzymes implicated in phosphorus acquisition. Yadav et al. (2010) reported that maize inoculated with *S. indica* accumulates higher levels of P than control plants. In addition, plants inoculated with *S. indica* with knocked down *PiPT* accumulated less P than plants inoculated with a wild-type fungus. This indicates that *PiPT* is involved in phosphorus transport and that *S. indica* improves the nutrition of the host plant. Improved host phosphorus acquisition by *S. indica* was conferred by Wu et al. (2018). The class 2 endophytic *Fusarium tricinctum* strain isolated from *S. nigrum* was shown to produce IAA and stimulate the growth of the shoots and roots during Cd toxicity (Khan et al., 2017a). Yamaji et al. (2016) reported that *Clethra barbinervis*, naturally occurring at mine sites, could tolerate high TM concentrations due to the support of DSE, including *P. fortinii*, *Rhizoderma veluwensis*, and *Rhizoscyphus* sp. Inoculation tests revealed that these fungal endophytes enhance plant growth by promoting the uptake of K.

HYPER-ACCUMULATORS AND SYMBIOTIC FUNGI

The role of microorganisms in the phenomenon of hyper-accumulation remains unknown, even though much progress in this matter has recently been accomplished (reviewed in Furini et al., 2015). Only 10% of the metal hyper-accumulating species have been examined for their ability to interact with rhizospheric microorganisms, mostly for their associations with bacteria (Li et al., 2007; Pal et al., 2007; Kidd et al., 2018).

Turnau and Mesjasz-Przybyłowicz (2003) reported several Ni hyper-accumulating plants from ultramafic soils in South Africa that were abundantly colonized by AMF. Orłowska et al. (2013) reported that the Ni hyper-accumulating perennial *Berkheya coddii* inoculated with mycorrhizal fungi accumulated P, K, Mn, and Zn in the cortical layer of the lateral roots and in the vascular stele more strongly, while Ni was detected particularly in the vascular tissue in non-inoculated plants. The inoculated plants translocated Ni to the above-ground parts of the plant more efficiently. The effect of the endophytic fungi on these plants that are strongly dependent on mycorrhizae still awaits investigation, which could be useful for phytoextraction.

In recent years, there has been an increase in studies on the influence of endophytic fungi on hyper-accumulating plant species. Khan et al. (2017b) characterized the endophytic fungal community associated with Cd-hyper-accumulating *S. nigrum*. They isolated 42 culturable fungal strains that belonged to the Ascomycota. In another study, Khan et al. (2017a) tested endophytic fungal strains isolated from *S. nigrum* for Cd

tolerance and accumulation potential. They showed that one of the strains isolated RSF-6L exhibited the ability to reduce Cd uptake in the plant. Both the translocation factor (TF) and the bio-concentration factor were lower in the plants inoculated with endophytes than the control (Khan et al., 2017a). Interestingly, earlier studies showed that inoculation with the mycorrhizal fungi *Claroideoglossum claroideum* and *R. intraradices* enhanced the accumulation of Zn in the plant tissues with the primary reservoir of Zn in the shoots (Marques et al., 2006). It would be interesting to compare these results with those of double inoculation and various metals. In addition, inoculation with the fungi conferred protection to the host plant, leading to an improvement in tolerance.

CONCLUSIONS

The pollution of soils with toxic metals not only affects the composition and diversity of microbial communities, leading to a reduction in the overall abundance of microbial species, but also results in the enrichment of metal-tolerant microbial strains (Yao et al., 2003; Stefanowicz et al., 2008; Azarbad et al., 2013; Chen et al., 2014). These strains are extremely tolerant to metal toxicity and based on the studies available may facilitate plant growth in environments polluted with metals. The role of mycorrhizae in the adaptation of plants to metal pollution has been extensively studied over the last three decades. However, the role of other microbial components of the mycorrhizosphere cannot be overlooked. This is particularly important due to the fact that plants in their natural environments simultaneously interact with a wide array of microorganisms. In case of the Brassicaceae metallophytes and hyper-accumulators that lost the ability to interact with mycorrhizae fungal endophytes may play a role resembling mycorrhizae in the adaptation of non-mycorrhizal plants to metalliferous soils; however, the mechanisms of the action of endophytes may distinguish them from other symbiotic fungi (Table 1). In nature plants, both mycorrhizal and non-mycorrhizal and microorganisms form complex assemblages termed the holobiont. The functioning of this super organism is determined by relationships between the interacting organisms, thus understanding the complexity and routes of mutual communication will allow us to utilize the potential of plant-based clean-up technologies more effectively. In general, the influence of toxic metals on fungi and plants encompasses induced changes on the level of the epigenome, transcriptome, proteome, metabolome, and secretome. In addition, these changes further influence the interactions between the plant and symbiotic fungi and other soil microorganisms (Figure 1). Endophytic fungi may alleviate TM stress in the host plants in various manners. They can serve as a barrier and prevent the entry of metal ions into plant tissues and optimize metal distribution within the plant. These roles of fungi have been described on several occasions; however, very little is known about how the symbiont induces the metal phenotype in plants. One of the most pending questions is whether the metal adapted fungus directly affects metal specific tolerance mechanisms or is improved adaptation to metal toxicity

a result of indirect action such as improving water and nutrient availability in the soil and their subsequent uptake by plants that confer plant fitness? Another question is how much the endophyte influence on the plant TM tolerance differs from the analogous impact of mycorrhizal fungi? To answer these questions more studies concerning the role of fungal symbionts in plant response to TM are needed. Bioremediation is a dynamically growing area of green biotechnology. Endophytes may have considerable potential applications not only in microbial-assisted phytoremediation practice but also in mycoremediation of the areas that cannot be used for plant growth.

REFERENCES

- Adriaensen, K., van der Lelie, D., Van Laere, A., Vangronsveld, J., and Colpaert, J. V. (2004). A zinc-adapted fungus protects pines from zinc stress. *New Phytol.* 161, 549–555. doi: 10.1046/j.1469-8137.2003.00941.x
- Adriaensen, K., Vralstad, T., Noben, J.-P., Vangronsveld, J., and Colpaert, J. V. (2005). Copper-adapted *Suillus luteus*, a symbiotic solution for pines colonizing Cu mine spoils. *Appl. Environ. Microbiol.* 71, 7279–7284. doi: 10.1128/AEM.71.11.7279-7284.2005
- Adriaensen, K., Vangronsveld, J., and Colpaert, J. V. (2006). Zinc-tolerant *Suillus bovinus* improves growth of Zn-exposed *Pinus sylvestris* seedlings. *Mycorrhiza* 16, 553–558. doi: 10.1007/s00572-006-0072-7
- Agler, M. T., Ruhe, J., Kroll, S., Morhenn, C., Kim, S.-T., Weigel, D., et al. (2016). Microbial hub taxa link host and abiotic factors to plant microbiome variation. *PLoS Biol.* 14:e1002352. doi: 10.1371/journal.pbio.1002352
- Almario, J., Jeena, G., Wunder, J., Langen, G., Zuccaro, A., Coupland, G., et al. (2017). Root-associated fungal microbiota of nonmycorrhizal *Arabidopsis thaliana* and its contribution to plant phosphorus nutrition. *Proc. Natl. Acad. Sci. USA* 114, E9403–E9412. doi: 10.1073/pnas.1710455114
- Andrade-Linares, D. R., and Franken, P. (2013). Fungal endophytes in plant roots: taxonomy, colonization patterns, and functions. *Symbiotic Endophytes. Soil biology* 37, 311–334. doi: 10.1007/978-3-642-39317-4_16
- Atsatt, P. R., and Whiteside, M. D. (2014). Novel symbiotic protoplasts formed by endophytic fungi explain their hidden existence, lifestyle switching, and diversity within the plant kingdom. *PLoS One* 9:e95266. doi: 10.1371/journal.pone.0095266
- Azarbad, H., Niklińska, M., van Gestel, C. A. M., van Straalen, N. M., Röling, W. F. M., and Laskowski, R. (2013). Microbial community structure and functioning along metal pollution gradients. *Environ. Toxicol. Chem.* 32, 1992–2002. doi: 10.1002/etc.2269
- Ban, Y., Tang, M., Chen, H., Xu, Z., Zhang, H., and Yang, Y. (2012). The response of dark septate endophytes to heavy metals in pure culture. *PLoS One* 7:e47968. doi: 10.1371/journal.pone.0047968
- Ban, Y., Xu, Z., Yang, Y., Zhang, H., Chen, H., and Tang, M. (2017). Effect of dark septate endophytic fungus *Gaeumannomyces cylindrosporus* on plant growth, photosynthesis and Pb tolerance of maize (*Zea mays* L.). *Pedosphere* 27, 283–292. doi: 10.1016/S1002-0160(17)60316-3
- Bartholdy, B. A., Berreck, M., and Haselwandter, K. (2001). Hydroxamate siderophore synthesis by *Phialocephala fortinii*, a typical dark septate fungal root endophyte. *Biomaterials* 14, 33–42. doi: 10.1023/A:1016687021803
- Bellion, M., Courbot, M., Jacob, C., Blaudez, D., and Chalot, M. (2006). Extracellular and cellular mechanisms sustaining metal tolerance in ectomycorrhizal fungi. *FEMS Microbiol. Lett.* 254, 173–181. doi: 10.1111/j.1574-6968.2005.00044.x
- Berthelot, C., Blaudez, D., Beguiristain, T., Chalot, M., and Leyval, C. (2018). Co-inoculation of *Lolium perenne* with *Funnelformis mosseae* and the dark septate endophyte *Cadophora* sp. in a trace element-polluted soil. *Mycorrhiza* 28, 301–314. doi: 10.1007/s00572-018-0826-z
- Bidartondo, M. I., Read, D. J., Trappe, J. M., Merckx, V., Ligrone, R., Duckett, G., et al. (2011). The dawn of symbiosis between plants and fungi. *Biol. Lett.* 7, 574–577. doi: 10.1098/rsbl.2010.1203
- Bischoff, J. F., and White, J. F. (2005). “Evolutionary development of the Clavicipitaceae” in *The fungal community its organization and role in the ecosystem*. 3rd edn. eds. J. Dighton, J. White, and P. Oudemans (Boca Raton, FL, USA: Taylor & Francis), 508–518.
- Bona, E., Cattaneo, C., Cesaro, P., Marsano, F., Lingua, G., Cavaletto, M., et al. (2010). Proteomic analysis of *Pteris vittata* fronds: two arbuscular mycorrhizal fungi differentially modulate protein expression under arsenic contamination. *Proteomics* 10, 3811–3834. doi: 10.1002/pmic.200900436
- Bruenger, F. W., Stover, B. J., and Atherton, D. R. (1967). The incorporation of various metal ions into in vivo- and in vitro-produced melanin. *Radiat. Res.* 32, 1, 12.
- Brundrett, M. C. (2002). Coevolution of roots and mycorrhizas of land plants. *New Phytol.* 154, 275–304. doi: 10.1046/j.1469-8137.2002.00397.x
- Brundrett, M. C. (2006). “Understanding the roles of multifunctional mycorrhizal and endophytic fungi” in *Microbial root endophytes*. eds. B. J. E. Schulz, C. J. C. Boyle, and T. N. Sieber (Berlin, Heidelberg: Springer), 281–298.
- Brundrett, M. C., and Tedersoo, L. (2018). Evolutionary history of mycorrhizal symbioses and global host plant diversity. *New Phytol.* 220, 1108–1115. doi: 10.1111/nph.14976
- Burges, A., Epelde, L., Blanco, F., Becerril, J. M., and Garbisu, C. (2017). Ecosystem services and plant physiological status during endophyte-assisted phytoremediation of metal contaminated soil. *Sci. Total Environ.* 584–585, 329–338. doi: 10.1016/j.scitotenv.2016.12.146
- Burleigh, S. H., Kristensen, B. K., and Bechmann, I. E. (2003). A plasma membrane zinc transporter from *Medicago truncatula* is up-regulated in roots by Zn fertilization, yet down-regulated by arbuscular mycorrhizal colonization. *Plant Mol. Biol.* 52, 1077–1088. doi: 10.1023/A:1025479701246
- Cabral, L., Soares, C. R. F. S., Giachini, A. J., and Siqueira, J. O. (2015). Arbuscular mycorrhizal fungi in phytoremediation of contaminated areas by trace elements: mechanisms and major benefits of their applications. *World J. Microbiol. Biotechnol.* 31, 1655–1664. doi: 10.1007/s11274-015-1918-y
- Cao, L., Jiang, M., Zeng, Z., Du, A., Tan, H., and Liu, Y. (2008). *Trichoderma atroviride* F6 improves phytoextraction efficiency of mustard (*Brassica juncea* (L.) Coss. var. *foliosa* Bailey) in Cd, Ni contaminated soils. *Chemosphere* 71, 1769–1773. doi: 10.1016/j.chemosphere.2008.01.066
- Chalot, M., Botton, B., and Blaudez, D. (2000). Cadmium uptake and subcellular compartmentation in the ectomycorrhizal fungus *Paxillus involutus*. *Microbiology* 146, 1109–1117. doi: 10.1099/00221287-146-5-1109
- Chen, B., Li, X., Tao, H., Christie, P., and Wong, M. (2003). The role of arbuscular mycorrhiza in zinc uptake by red clover growing in a calcareous soil spiked with various quantities of zinc. *Chemosphere* 50, 839–846. doi: 10.1016/S0045-6535(02)00228-X
- Chen, B., Shen, H., Li, X., Feng, G., and Christie, P. (2004). Effects of EDTA application and arbuscular mycorrhizal colonization on growth and zinc uptake by maize (*Zea mays* L.) in soil experimentally contaminated with zinc. *Plant Soil* 261, 219–229. doi: 10.1023/B:PLSO.0000035538.09222.f
- Chen, B. D., Zhu, Y.-G., Duan, J., Xiao, X. Y., and Smith, S. E. (2007). Effects of the arbuscular mycorrhizal fungus *Glomus mosseae* on growth and metal uptake by four plant species in copper mine tailings. *Environ. Pollut.* 147, 374–380. doi: 10.1016/j.envpol.2006.04.027
- Chen, J., Wang, H., and Guo, S. X. (2012). Isolation and identification of endophytic and mycorrhizal fungi from seeds and roots of *Dendrobium* (Orchidaceae). *Mycorrhiza* 22, 297–307. doi: 10.1007/s00572-011-0404-0

AUTHOR CONTRIBUTIONS

AD, PR, and KT wrote the manuscript. AD and KT prepared the figures. AD and PR contributed equally to the manuscript.

FUNDING

This work was supported by Polish National Centre for Research and Development (FACCE SURPLUS/I/AGRONICKEL/02/2016) and the National Science Center (2017/27/B/NZ8/01199).

- Chen, J., He, F., Zhang, X., Sun, X., Zheng, J., and Zheng, J. (2014). Heavy metal pollution decreases microbial abundance, diversity and activity within particle-size fractions of a paddy soil. *FEMS Microbiol. Ecol.* 87, 164–181. doi: 10.1111/1574-6941.12212
- Chhabra, S., and Dowling, D. N. (2017). “Endophyte-promoted nutrient acquisition: phosphorus and iron” in *Functional importance of the plant microbiome*. ed. S. Doty (Cham: Springer International Publishing), 21–42.
- Chiappello, M., Martino, E., and Perotto, S. (2015). Common and metal-specific proteomic responses to cadmium and zinc in the metal tolerant ericoid mycorrhizal fungus *Oidiodendron maius* Zn. *Metallomics* 7, 805–815. doi: 10.1039/C5MT00024F
- Chodak, M., Gołbiewski, M., Morawska-Płoskonka, J., Kuduk, K., and Niklińska, M. (2013). Diversity of microorganisms from forest soils differently polluted with heavy metals. *Appl. Soil Ecol.* 64, 7–14. doi: 10.1016/j.apsoil.2012.11.004
- Cicatelli, A., Lingua, G., Todeschini, V., Biondi, S., Torrigiani, P., and Castiglione, S. (2012). Arbuscular mycorrhizal fungi modulate the leaf transcriptome of a *Populus alba* L. clone grown on a zinc and copper-contaminated soil. *Environ. Exp. Bot.* 75, 25–35. doi: 10.1016/j.envexpbot.2011.08.012
- Colpaert, J. V., Wevers, J. H. L., Krzmaric, E., and Adriaensen, K. (2011). How metal-tolerant ecotypes of ectomycorrhizal fungi protect plants from heavy metal pollution. *Ann. For. Sci.* 68, 17–24. doi: 10.1007/s13595-010-0003-9
- Coninx, L., Martinova, V., and Rineau, F. (2017). Mycorrhiza-assisted phytoremediation. *Adv. Bot. Res.* 83, 127–188. doi: 10.1016/bs.abr.2016.12.005
- Corneo, P. E., Pellegrini, A., Cappellin, L., Roncador, M., Chierici, M., Gessler, C., et al. (2013). Microbial community structure in vineyard soils across altitudinal gradients and in different seasons. *FEMS Microbiol. Ecol.* 84, 588–602. doi: 10.1111/1574-6941.12087
- Cui, M., and Nobel, P. S. (2006). Nutrient status, water uptake and gas exchange for three desert succulents infected with mycorrhizal fungi. *New Phytol.* 122, 643–649. doi: 10.1111/j.1469-8137.1992.tb00092.x
- DalCorso, G., Manara, A., and Furini, A. (2013). An overview of heavy metal challenge in plants: from roots to shoots. *Metallomics* 5, 1117–1132. doi: 10.1039/c3mt00038a
- Das, M., Jakkula, V. S., and Adholeya, A. (2017). “Role of mycorrhiza in phytoremediation processes: a review” in *Mycorrhiza - Nutrient uptake, bioremediation, ecorestoration*. eds. A. Varma, R. Prasad, and N. Tuteja (Cham: Springer International Publishing), 271–286.
- Deckert, R. J., Melville, L. H., and Peterson, R. L. (2001). Structural features of a *Lophodermium* endophyte during the cryptic life-cycle phase in the foliage of *Pinus strobus*. *Mycol. Res.* 105, 991–997. doi: 10.1016/S0953-7562(08)61957-7
- Deng, Z., Cao, L., Huang, H., Jiang, X., Wang, W., Shi, Y., et al. (2011). Characterization of Cd- and Pb-resistant fungal endophyte *Mucor* sp. CBRF59 isolated from rapes (*Brassica chinensis*) in a metal-contaminated soil. *J. Hazard. Mater.* 185, 717–724. doi: 10.1016/j.jhazmat.2010.09.078
- Deng, Z., Zhang, R., Shi, Y., Hu, L., Tan, H., and Cao, L. (2013). Enhancement of phytoremediation of Cd- and Pb-contaminated soils by self-fusion of protoplasts from endophytic fungus *Mucor* sp. CBRF59. *Chemosphere* 91, 41–47. doi: 10.1016/j.chemosphere.2012.11.065
- Deng, Z., Zhang, R., Shi, Y., Hu, L., Tan, H., and Cao, L. (2014). Characterization of Cd-, Pb-, Zn-resistant endophytic *Lasiodiplodia* sp. MXSF31 from metal accumulating *Portulaca oleracea* and its potential in promoting the growth of rape in metal-contaminated soils. *Environ. Sci. Pollut. Res.* 21, 2346–2357. doi: 10.1007/s11356-013-2163-2
- Devi, L., and Joshi, S. (2015). Ultrastructures of silver nanoparticles biosynthesized using endophytic fungi. *J. Microsc. Ultrastruct.* 3, 29–37. doi: 10.1016/j.jmau.2014.10.004
- Dimkpa, C. O., McLean, J. E., Britt, D. W., and Anderson, A. J. (2015). Nano-CuO and interaction with nano-ZnO or soil bacterium provide evidence for the interference of nanoparticles in metal nutrition of plants. *Ecotoxicology* 24, 119–129. doi: 10.1007/s10646-014-1364-x
- Domka, A., Rozpądek, P., Ważny, R., and Turnau, K. (2019). *Mucor* sp.—an endophyte of Brassicaceae capable of surviving in toxic metal-rich sites. *J. Basic Microbiol.* 59, 24–37. doi: 10.1002/jobm.201800406
- Durán, N., Marcato, P. D., Durán, M., Yadav, A., Gade, A., and Rai, M. (2011). Mechanistic aspects in the biogenic synthesis of extracellular metal nanoparticles by peptides, bacteria, fungi, and plants. *Appl. Microbiol. Biotechnol.* 90, 1609–1624. doi: 10.1007/s00253-011-3249-8
- El Enshasy, H. A., El Marzugi, N. A., Elsayed, E. A., Ling, O. M., Malek, R. A., Kepki, A. N., et al. (2018). “Medical and cosmetic applications of fungal nanotechnology: production, characterization, and bioactivity” in *Fungal nanobionics: Principles and applications*. eds. R. Prasad, V. Kumar, M. Kumar, and S. Wang (Singapore: Springer), 21–59.
- Fernández, J. G., Fernández-Baldo, M. A., Berni, E., Camí, G., Durán, N., Raba, J., et al. (2016). Production of silver nanoparticles using yeasts and evaluation of their antifungal activity against phytopathogenic fungi. *Process Biochemistry* 51, 1306–1313. doi: 10.1016/j.procbio.2016.05.021
- Fernández, P. M., Viñarta, S. C., Bernal, A. R., Cruz, E. L., and Figueroa, L. I. C. (2018). Bioremediation strategies for chromium removal: current research, scale-up approach and future perspectives. *Chemosphere* 208, 139–148. doi: 10.1016/j.chemosphere.2018.05.166
- Ferrol, N., Tamayo, E., and Vargas, P. (2016). The heavy metal paradox in arbuscular mycorrhizas: from mechanisms to biotechnological applications. *J. Exp. Bot.* 67, 6253–6265. doi: 10.1093/jxb/erv403
- Field, K. J., Rimington, W. R., Bidartondo, M. I., Allinson, K. E., Beerling, D. J., Cameron, D. D., et al. (2015). First evidence of mutualism between ancient plant lineages (Haplomitriopsida liverworts) and Mucoromycotina fungi and its response to simulated Palaeozoic changes in atmospheric CO₂. *New Phytol.* 205, 743–756. doi: 10.1111/nph.13024
- Field, K. J., Rimington, W. R., Bidartondo, M. I., Allinson, K. E., Beerling, D. J., Cameron, D. D., et al. (2016). Functional analysis of liverworts in dual symbiosis with Glomeromycota and Mucoromycotina fungi under a simulated Palaeozoic CO₂ decline. *ISME J.* 10, 1514–1526. doi: 10.1038/ismej.2015.204
- Fogarty, R. V., and Tobin, J. M. (1996). Fungal melanins and their interactions with metals. *Enzyme Microb. Technol.* 19, 311–317. doi: 10.1016/0141-0229(96)00002-6
- Fomina, M. A., Alexander, I. J., Colpaert, J. V., and Gadd, G. M. (2005). Solubilization of toxic metal minerals and metal tolerance of mycorrhizal fungi. *Soil Biol. Biochem.* 37, 851–866. doi: 10.1016/j.soilbio.2004.10.013
- Freeman, J. L. (2004). Increased glutathione biosynthesis plays a role in nickel tolerance in *Thlaspi* nickel hyperaccumulators. *Plant Cell* 16, 2176–2191. doi: 10.1105/tpc.104.023036
- Furini, A., Manara, A., and DalCorso, G. (2015). Editorial: environmental phytoremediation: plants and microorganisms at work. *Front. Plant Sci.* 6:520. doi: 10.3389/fpls.2015.00520
- Gadd, G. M. (1993). Interactions of fungus with toxic metals. *New Phytol.* 124, 25–60. doi: 10.1111/j.1469-8137.1993.tb03796.x
- Gadd, G. M. (1999). Fungal production of citric and oxalic acid: importance in metal speciation, physiology and biogeochemical processes. *Adv. Microb. Physiol.* 41, 47–92. doi: 10.1016/S0065-2911(08)60165-4
- Gadd, G. M. (2016). “5 Fungi and industrial pollutants” in *Environmental and microbial relationships*. eds. I. Druzhinina, and C. Kubicek (Cham: Springer International Publishing), 99–125.
- Gadd, G. M., Rhee, Y. J., Stephenson, K., and Wei, Z. (2012). Geomycology: metals, actinides and biominerals. *Environ. Microbiol. Rep.* 4, 270–296. doi: 10.1111/j.1758-2229.2011.00283.x
- Garza, M. T. G., Perez, D. B., Rodriguez, A. V., Garcia-Gutierrez, D. I., Zarate, X., Cardenas, M. E. C., et al. (2016). Metal-induced production of a novel bioadsorbent exopolysaccharide in a native *Rhodotorula mucilaginosa* from the Mexican northeastern region. *PLoS One* 11, 1–18. doi: 10.1371/journal.pone.0148430
- Göhre, V., and Paszkowski, U. (2006). Contribution of the arbuscular mycorrhizal symbiosis to heavy metal phytoremediation. *Planta* 223, 1115–1122. doi: 10.1007/s00425-006-0225-0
- González-Chávez, M. C., D’Haen, J., Vangronsveld, J., and Dodd, J. C. (2002). Copper sorption and accumulation by the extraradical mycelium of different *Glomus* spp. (arbuscular mycorrhizal fungi) isolated from the same polluted soil. *Plant Soil* 240, 287–297. doi: 10.1023/A:1015794622592
- González-Chávez, M. C., Carrillo-González, R., Wright, S. F., and Nichols, K. A. (2004). The role of glomalin, a protein produced by arbuscular mycorrhizal fungi, in sequestering potentially toxic elements. *Environ. Pollut.* 130, 317–323. doi: 10.1016/j.envpol.2004.01.004
- Gupta, A., and Joia, J. (2016). Microbes as potential tool for remediation of heavy metals: a review. *J. Microb. Biochem. Technol.* 8, 364–372. doi: 10.4172/1948-5948.1000310
- Hardoim, P. R., van Overbeek, L. S., Berg, G., Pirttilä, A. M., Compant, S., Campisano, A., et al. (2015). The hidden world within plants: ecological and evolutionary considerations for defining functioning of microbial endophytes. *Microbiol. Mol. Biol. Rev.* 79, 293–320. doi: 10.1128/MMBR.00050-14

- Harman, G. E., Petzoldt, R., Comis, A., and Chen, J. (2004). Interactions between *Trichoderma harzianum* strain T22 and maize inbred line Mo17 and effects of these interactions on diseases caused by *Pythium ultimum* and *Colletotrichum graminicola*. *Phytopathology* 94, 147–153. doi: 10.1094/PHYTO.2004.94.2.147
- Hasan, M. K., Cheng, Y., Kanwar, M. K., Chu, X.-Y., Ahammed, G. J., and Qi, Z.-Y. (2017). Responses of plant proteins to heavy metal stress—a review. *Front. Plant Sci.* 8, 1492. doi: 10.3389/fpls.2017.01492
- Haydon, M. J., and Cobbett, C. S. (2007). A novel major facilitator superfamily protein at the tonoplast influences zinc tolerance and accumulation in *Arabidopsis*. *Plant Physiol.* 143, 1705–1719. doi: 10.1104/pp.106.092015
- He, Y., Yang, Z., Li, M., Jiang, M., Zhan, F., Zu, Y., et al. (2017). Effects of a dark septate endophyte (DSE) on growth, cadmium content, and physiology in maize under cadmium stress. *Environ. Sci. Pollut. Res.* 24, 18494–18504. doi: 10.1007/s11356-017-9459-6
- Hetrick, B. A. D. (1991). Mycorrhizas and root architecture. *Experientia* 47, 355–362. doi: 10.1007/BF01972077
- Hiruma, K., Gerlach, N., Sacristán, S., Nakano, R. T., Hacquard, S., Kracher, B., et al. (2016). Root endophyte *Colletotrichum tofieldiae* confers plant fitness benefits that are phosphate status dependent. *Cell* 165, 464–474. doi: 10.1016/j.cell.2016.02.028
- Hiruma, K., Kobae, Y., and Toju, H. (2018). Beneficial associations between Brassicaceae plants and fungal endophytes under nutrient-limiting conditions: evolutionary origins and host–symbiont molecular mechanisms. *Curr. Opin. Plant Biol.* 44, 145–154. doi: 10.1016/j.pbi.2018.04.009
- Horton, T. R., Cázares, E., and Bruns, T. D. (1998). Ectomycorrhizal, vesicular-arbuscular and dark septate fungal colonization of bishop pine (*Pinus muricata*) seedlings in the first 5 months of growth after wildfire. *Mycorrhiza* 8, 11–18. doi: 10.1007/s005720050205
- Huang, Y., Wang, J., Li, G., Zheng, Z., and Su, W. (2001). Antitumor and antifungal activities in endophytic fungi isolated from pharmaceutical plants *Taxus mairei*, *Cephalataxus fortunei* and *Torreya grandis*. *FEMS Immunol. Med. Microbiol.* 31, 163–167. doi: 10.1111/j.1574-695X.2001.tb00513.x
- Illmer, P., and Buttinger, R. (2006). Interactions between iron availability, aluminium toxicity and fungal siderophores. *BioMetals* 19, 367–377. doi: 10.1007/s10534-005-3496-1
- Jaeckel, P., Krauss, G., Menge, S., Schierhorn, A., Rücknagel, P., and Krauss, G.-J. (2005). Cadmium induces a novel metallothionein and phytochelatin 2 in an aquatic fungus. *Biochem. Biophys. Res. Commun.* 333, 150–155. doi: 10.1016/j.bbrc.2005.05.083
- Janoušková, M., Pavlíková, D., Macek, T., and Vosátka, M. (2005). Influence of arbuscular mycorrhiza on the growth and cadmium uptake of tobacco with inserted metallothionein gene. *Appl. Soil Ecol.* 29, 209–214. doi: 10.1016/j.apsoil.2004.12.006
- Janoušková, M., Pavlíková, D., and Vosátka, M. (2006). Potential contribution of arbuscular mycorrhiza to cadmium immobilisation in soil. *Chemosphere* 65, 1959–1965. doi: 10.1016/j.chemosphere.2006.07.007
- Jiang, M., Cao, L., and Zhang, R. (2008). Effects of *Acacia auriculaeformis* A. Cunn.-associated fungi on mustard (*Brassica juncea* (L.) Coss. var. *foliosa* Bailey) growth in Cd- and Ni-contaminated soils. *Lett. Appl. Microbiol.* 47, 561–565. doi: 10.1111/j.1472-765X.2008.02454.x
- Johnson, L. J., Koulman, A., Christensen, M., Lane, G. A., Fraser, K., Forester, N., et al. (2013). An extracellular siderophore is required to maintain the mutualistic interaction of *Epichloë festucae* with *Lolium perenne*. *PLoS Pathog.* 9:e1003332. doi: 10.1371/journal.ppat.1003332
- Joner, E. J. (2000). Metal-binding capacity of arbuscular mycorrhizal mycelium. *Plant Soil* 226, 227–234. doi: 10.1023/A:1026565701391
- Joner, E. J., and Leyval, C. (1997). Uptake of Cd by roots and hyphae of a *Glomus mosseae*/*Trifolium subterraneum* mycorrhiza from soil amended with high and low concentrations of cadmium. *New Phytol.* 135, 353–360. doi: 10.1046/j.1469-8137.1997.00633.x
- Kashyap, P. L., Kumar, S., Srivastava, A. K., and Sharma, A. K. (2013). Myconanotechnology in agriculture: a perspective. *World J. Microbiol. Biotechnol.* 29, 191–207. doi: 10.1007/s11274-012-1171-6
- Khan, A. L., and Lee, I.-J. (2013). Endophytic *Penicillium funiculosum* LHL06 secretes gibberellin that reprograms *Glycine max* L. growth during copper stress. *BMC Plant Biol.* 13:86. doi: 10.1186/1471-2229-13-86
- Khan, A. R., Ullah, I., Waqas, M., Park, G.-S., Khan, A. L., Hong, S.-J., et al. (2017a). Host plant growth promotion and cadmium detoxification in *Solanum nigrum*, mediated by endophytic fungi. *Ecotoxicol. Environ. Saf.* 136, 180–188. doi: 10.1016/j.ecoenv.2016.03.014
- Khan, A. R., Waqas, M., Ullah, I., Khan, A. L., Khan, M. A., Lee, I. J., et al. (2017b). Culturable endophytic fungal diversity in the cadmium hyperaccumulator *Solanum nigrum* L. and their role in enhancing phytoremediation. *Environ. Exp. Bot.* 135, 126–135. doi: 10.1016/j.envexpbot.2016.03.005
- Khouja, H. R., Abbà, S., Lacerat-Didier, L., Daghino, S., Doillon, D., Richaud, P., et al. (2013). OmZnT1 and OmFET, two metal transporters from the metal-tolerant strain Zn of the ericoid mycorrhizal fungus *Oidiodendron maius*, confer zinc tolerance in yeast. *Fungal Genet. Biol.* 52, 53–64. doi: 10.1016/j.fgb.2012.11.004
- Kidd, P. S., Bani, A., Benizri, E., Gonnelli, C., Hazotte, C., Kisser, J., et al. (2018). Developing sustainable agromining systems in agricultural ultramafic soils for nickel recovery. *Front. Environ. Sci.* 6:44. doi: 10.3389/fenvs.2018.00044
- Koren'kov, V., Park, S., Cheng, N. H., Sreevidya, C., Lachmansingh, J., Morris, J., et al. (2007). Enhanced Cd²⁺-selective root-tonoplast-transport in tobaccos expressing *Arabidopsis* cation exchangers. *Planta* 225, 403–411. doi: 10.1007/s00425-006-0352-7
- Koulman, A., Lee, T. V., Fraser, K., Johnson, L., Arcus, V., Lott, J. S., et al. (2012). Identification of extracellular siderophores and a related peptide from the endophytic fungus *Epichloë festucae* in culture and endophyte-infected *Lolium perenne*. *Phytochemistry* 75, 128–139. doi: 10.1016/j.phytochem.2011.11.020
- Leake, J., Johnson, D., Donnelly, D., Muckle, G., Boddy, L., and Read, D. (2004). Networks of power and influence: the role of mycorrhizal mycelium in controlling plant communities and agroecosystem functioning. *Can. J. Bot.* 82, 1016–1045. doi: 10.1139/b04-060
- Leung, H. M., Ye, Z. H., and Wong, M. H. (2006). Interactions of mycorrhizal fungi with *Pteris vittata* (As hyperaccumulator) in As-contaminated soils. *Environ. Pollut.* 139, 1–8. doi: 10.1016/j.envpol.2005.05.009
- Li, W. C., Ye, Z. H., and Wong, M. H. (2007). Effects of bacteria on enhanced metal uptake of the Cd/Zn-hyperaccumulating plant, *Sedum alfredii*. *J. Exp. Bot.* 58, 4173–4182. doi: 10.1093/jxb/erm274
- Li, T., Liu, M. J., Zhang, X. T., Zhang, H. B., Sha, T., and Zhao, Z. W. (2011). Improved tolerance of maize (*Zea mays* L.) to heavy metals by colonization of a dark septate endophyte (DSE) *Exophiala pisciphila*. *Sci. Total Environ.* 409, 1069–1074. doi: 10.1016/j.scitotenv.2010.12.012
- Li, D., Xu, H., Mei, T., and Li, H. (2012a). Diversity of endophytic fungi from six dominant plant species in a Pb-Zn mine wasteland in China. *Shengtai Xuebao/Acta Ecol. Sin.* 32, 2288–2293. doi: 10.5846/stxb201105260705
- Li, H. Y., Wei, D. Q., Shen, M., and Zhou, Z. P. (2012b). Endophytes and their role in phytoremediation. *Fungal Divers.* 54, 11–18. doi: 10.1007/s13225-012-0165-x
- Li, X., Li, W., Chu, L., White, J. F., Xiong, Z., and Li, H. (2016). Diversity and heavy metal tolerance of endophytic fungi from *Dysphania ambrosioides*, a hyperaccumulator from Pb-Zn contaminated soils. *J. Plant Interact.* 11, 186–192. doi: 10.1080/17429145.2016.1266043
- Lingua, G., Bona, E., Todeschini, V., Cattaneo, C., Marsano, F., Berta, G., et al. (2012). Effects of heavy metals and arbuscular mycorrhiza on the leaf proteome of a selected poplar clone: a time course analysis. *PLoS One* 7:e38662. doi: 10.1371/journal.pone.0038662
- Liu, R., and Lal, R. (2015). Potentials of engineered nanoparticles as fertilizers for increasing agronomic productions. *Sci. Total Environ.* 514, 131–139. doi: 10.1016/j.scitotenv.2015.01.104
- Ma, Y., He, J., Ma, C., Luo, J., Li, H., Liu, T., et al. (2014). Ectomycorrhizas with *Paxillus involutus* enhance cadmium uptake and tolerance in *Populus × canescens*. *Plant. Cell Environ.* 37, 627–642. doi: 10.1111/pce.12183
- Machuca, A., Pereira, G., Aguiar, A., and Milagres, A. M. F. (2007). Metal-chelating compounds produced by ectomycorrhizal fungi collected from pine plantations. *Lett. Appl. Microbiol.* 44, 7–12. doi: 10.1111/j.1472-765X.2006.02046.x
- Manceau, A., Nagy, K. L., Marcus, M. A., Lanson, M., Geoffroy, N., Jacquet, T., et al. (2008). Formation of metallic copper nanoparticles at the soil–root interface. *Environ. Sci. Technol.* 42, 1766–1772. doi: 10.1021/es072017o
- Marques, A. P. G. C., Oliveira, R. S., Rangel, A. O. S. S., and Castro, P. M. L. (2006). Zinc accumulation in *Solanum nigrum* is enhanced by different arbuscular mycorrhizal fungi. *Chemosphere* 65, 1256–1263. doi: 10.1016/j.chemosphere.2006.03.022
- Masood, A., Iqbal, N., and Khan, N. A. (2012). Role of ethylene in alleviation of cadmium-induced photosynthetic capacity inhibition by sulphur in mustard. *Plant. Cell Environ.* 35, 524–533. doi: 10.1111/j.1365-3040.2011.02432.x

- Mathys, W. (1977). The role of malate, oxalate, and mustard oil glucosides in the evolution of zinc-resistance in herbage plants. *Physiol. Plant.* 40, 130–136. doi: 10.1111/j.1399-3054.1977.tb01509.x
- Meharg, A. A. (2003). The mechanistic basis of interactions between mycorrhizal associations and toxic metal cations. *Mycol. Res.* 107, 1253–1265. doi: 10.1017/S0953756203008608
- Meier, S., Borie, F., Bolan, N., and Cornejo, P. (2012). Phytoremediation of metal-polluted soils by arbuscular mycorrhizal fungi. *Crit. Rev. Env. Sci. Technol.* 42, 741–775. doi: 10.1080/10643389.2010.528518
- Min Yang, Z., and Chen, J. (2013). A potential role of microRNAs in plant response to metal toxicity. *Metallomics* 5, 1184–1190. doi: 10.1039/c3mt00022b
- Miransari, M. (2017). “Arbuscular mycorrhizal fungi and heavy metal tolerance in plants” in *Arbuscular mycorrhizas and stress tolerance of plants*. ed. Q.S. Wu (Singapore: Springer), 147–161.
- Morel, M., Crouzet, J., Gravot, A., Auroy, P., Leonhardt, N., Vavasseur, A., et al. (2009). AtHMA3, a P1B-ATPase allowing Cd/Zn/Co/Pb vacuolar storage in *Arabidopsis*. *Plant Physiol.* 149, 894–904. doi: 10.1104/pp.108.130294
- Na, G., and Salt, D. E. (2011). The role of sulfur assimilation and sulfur-containing compounds in trace element homeostasis in plants. *Environ. Exp. Bot.* 72, 18–25. doi: 10.1016/j.envexpbot.2010.04.004
- Netala, V., Bethu, M., Pushpalatha, B., Baki, V., Aishwarya, S., Rao, J., et al. (2016). Biogenesis of silver nanoparticles using endophytic fungus *Pestalotiopsis microspora* and evaluation of their antioxidant and anticancer activities. *Int. J. Nanomed.* 11, 5683–5696. doi: 10.2147/IJN.S112857
- Orłowska, E., Zubek, S., Jurkiewicz, A., Szarek-Lukaszewska, G., and Turnau, K. (2002). Influence of restoration on arbuscular mycorrhiza of *Biscutella laevigata* L. (Brassicaceae) and *Plantago lanceolata* L. (Plantaginaceae) from calamine spoil mounds. *Mycorrhiza* 12, 153–159. doi: 10.1007/s00572-001-0155-4
- Orłowska, E., Przybyłowicz, W., Orłowski, D., Mongwaketsi, N. P., Turnau, K., and Mesjasz-Przybyłowicz, J. (2013). Mycorrhizal colonization affects the elemental distribution in roots of Ni-hyperaccumulator *Berkheya coddii* Roessler. *Environ. Pollut.* 175, 100–109. doi: 10.1016/j.envpol.2012.12.028
- Ouziad, F., Hildebrandt, U., Schmelzer, E., and Bothe, H. (2005). Differential gene expressions in arbuscular mycorrhizal-colonized tomato grown under heavy metal stress. *J. Plant Physiol.* 162, 634–649. doi: 10.1016/j.jplph.2004.09.014
- Ouzounidou, G., Moustakas, M., Symeonidis, L., and Karataglis, S. (2006). Response of wheat seedlings to Ni stress: effects of supplemental calcium. *Arch. Environ. Contam. Toxicol.* 50, 346–352. doi: 10.1007/s00244-005-5076-3
- Pal, A., Wauters, G., and Paul, A. K. (2007). Nickel tolerance and accumulation by bacteria from rhizosphere of nickel hyperaccumulators in serpentine soil ecosystem of Andaman, India. *Plant Soil* 293, 37–48. doi: 10.1007/s11104-007-9195-7
- Pandey, G., Sharma, N., Pankaj Sahu, P., and Prasad, M. (2016). Chromatin-based epigenetic regulation of plant abiotic stress response. *Curr. Genomics* 17, 490–498. doi: 10.2174/1389202917666160520103914
- Paungfoo-Lonhienne, C., Rentsch, D., Robatzek, S., Webb, R. I., Sagulenko, E., Näsholm, T., et al. (2010). Turning the table: plants consume microbes as a source of nutrients. *PLoS One* 5, 1–11. doi: 10.1371/journal.pone.0011915
- Petrini, O. (1996). “Ecological and physiological aspects of host-specificity in endophytic fungi” in *Endophytic fungi in grasses and woody plants: Systematics, ecology, and evolution*. eds. S. Redlin and L. Carris (St. Paul: APS Press), 87–100.
- Prasad, M. N. V. (1999). “Metallothioneins and metal binding complexes in plants” in *Heavy metal stress in plants*. ed. M. N. V. Prasad (Berlin, Heidelberg: Springer), 51–72.
- Promptutha, I., Lumyong, S., Dhanasekaran, V., McKenzie, E. H. C., Hyde, K. D., and Jeewon, R. (2007). A phylogenetic evaluation of whether endophytes become saprotrophs at host senescence. *Microb. Ecol.* 53, 579–590. doi: 10.1007/s00248-006-9117-x
- Punamiya, P., Datta, R., Sarkar, D., Barber, S., Patel, M., and Das, P. (2010). Symbiotic role of *Glomus mosseae* in phytoextraction of lead in vetiver grass [*Chrysopogon zizanioides* (L.)]. *J. Hazard. Mater.* 177, 465–474. doi: 10.1016/j.jhazmat.2009.12.056
- Purahong, W., and Hyde, K. D. (2011). Effects of fungal endophytes on grass and non-grass litter decomposition rates. *Fungal Divers.* 47, 1–7. doi: 10.1007/s13225-010-0083-8
- Qian, Y., Yu, H., He, D., Yang, H., Wang, W., Wan, X., et al. (2013). Biosynthesis of silver nanoparticles by the endophytic fungus *Epicoccum nigrum* and their activity against pathogenic fungi. *Bioprocess. Biosyst. Eng.* 36, 1613–1619. doi: 10.1007/s00449-013-0937-z
- Rajkumar, M., Ae, N., and Freitas, H. (2009). Endophytic bacteria and their potential to enhance heavy metal phytoextraction. *Chemosphere* 77, 153–160. doi: 10.1016/j.chemosphere.2009.06.047
- Rajkumar, M., Sandhya, S., Prasad, M. N. V., and Freitas, H. (2012). Perspectives of plant-associated microbes in heavy metal phytoremediation. *Biotechnol. Adv.* 30, 1562–1574. doi: 10.1016/j.biotechadv.2012.04.011
- Rausser, W. E. (1999). Structure and function of metal chelators produced by plants. *Cell Biochem. Biophys.* 31, 19–48. doi: 10.1007/BF02738153
- Ray, P., and Adholeya, A. (2009). Correlation between organic acid exudation and metal uptake by ectomycorrhizal fungi grown on pond ash in vitro. *BioMetals* 22, 275–281. doi: 10.1007/s10534-008-9163-6
- Regvar, M., Vogel, K., Irgel, N., Wraber, T., Hildebrandt, U., Wilde, P., et al. (2003). Colonization of pennycresses (*Thlaspi* spp.) of the Brassicaceae by arbuscular mycorrhizal fungi. *J. Plant Physiol.* 160, 615–626. doi: 10.1078/0176-1617-00988
- Rodriguez, R. J., White, J. F. Jr., Arnold, A. E., and Redman, R. S. (2009). Fungal endophytes: diversity and functional roles. *New Phytol.* 182, 314–330. doi: 10.1111/j.1469-8137.2009.02773.x
- Rosa, L. H., Almeida Vieira, M. d. L., Santiago, I. F., and Rosa, C. A. (2010). Endophytic fungi community associated with the dicotyledonous plant *Colobanthus quitensis* (Kunth) Bartl. (Caryophyllaceae) in Antarctica. *FEMS Microbiol. Ecol.* 73, 178–189. doi: 10.1111/j.1574-6941.2010.00872.x
- Rozpądek, P., Domka, A., and Turnau, K. (2017). “Mycorrhizal fungi and accompanying microorganisms in improving phytoremediation techniques” in *The fungal community*. ed. J. F. W. J. Dighton (Boca Raton, London, New York: CRC Press Taylor & Francis Group 6000), 419–432.
- Rozpądek, P., Domka, A., Ważny, R., Nosek, M., Jędrzejczyk, R., Tokarz, K., et al. (2018). How does the endophytic fungus *Mucor* sp. improve *Arabidopsis arenosa* vegetation in the degraded environment of a mine dump? *Environ. Exp. Bot.* 147, 31–42. doi: 10.1016/j.envexpbot.2017.11.009
- Rudgers, J. A., and Swafford, A. L. (2009). Benefits of a fungal endophyte in *Elymus virginicus* decline under drought stress. *Basic Appl. Ecol.* 10, 43–51. doi: 10.1016/j.baec.2007.12.004
- Salazar-Badillo, F. B., Sánchez-Rangel, D., Becerra-Flora, A., López-Gómez, M., Nieto-Jacobo, F., Mendoza-Mendoza, A., et al. (2015). *Arabidopsis thaliana* polyamine content is modified by the interaction with different *Trichoderma* species. *Plant Physiol. Biochem.* 95, 49–56. doi: 10.1016/j.plaphy.2015.07.003
- Salvadori, M. R., Ando, R. A., Oller do Nascimento, C. A., and Corrêa, B. (2014). Intracellular biosynthesis and removal of copper nanoparticles by dead biomass of yeast isolated from the wastewater of a mine in the Brazilian Amazonia. *PLoS One* 9:e87968. doi: 10.1371/journal.pone.0087968
- Sanders, F. E., Tinker, P. B., Black, R. L. B., and Palmerley, S. M. (1977). The development of endomycorrhizal root systems: I. Spread of infection and growth-promoting effects with four species of vesicular-arbuscular endophyte. *New Phytol.* 78, 257–268. doi: 10.1111/j.1469-8137.1977.tb04829.x
- Saraswat, S., and Rai, J. P. N. (2011). “Mechanism of metal tolerance and detoxification in mycorrhizal fungi” in *Bio-management of metal-contaminated soils. Environmental pollution*. eds. M. Khan, A. Zaidi, R. Goel, and J. Musarrat (Dordrecht: Springer), 225–240.
- Sarwar, N., Imran, M., Shaheen, M. R., Ishaque, W., Kamran, M. A., Matloob, A., et al. (2017). Phytoremediation strategies for soils contaminated with heavy metals: modifications and future perspectives. *Chemosphere* 171, 710–721. doi: 10.1016/j.chemosphere.2016.12.116
- Schulz, B., Boyle, C., Draeger, S., Rimmert, A.-K., and Krohn, K. (2002). Endophytic fungi: a source of novel biologically active secondary metabolites. *Mycol. Res.* 106, 996–1004. doi: 10.1017/S0953756202006342
- Sell, J., Kayser, A., Schulin, R., and Brunner, I. (2005). Contribution of ectomycorrhizal fungi to cadmium uptake of poplars and willows from a heavily polluted soil. *Plant Soil* 277, 245–253. doi: 10.1007/s11104-005-7084-5
- Shahavivand, S., Parvaneh, A., and Aliloo, A. A. (2017). Root endophytic fungus *Piriformospora indica* affected growth, cadmium partitioning and chlorophyll fluorescence of sunflower under cadmium toxicity. *Ecotoxicol. Environ. Saf.* 145, 496–502. doi: 10.1016/j.ecoenv.2017.07.064
- Shankar, S. S., Ahmad, A., Pasricha, R., and Sastry, M. (2003). Bioreduction of chloroaurate ions by geranium leaves and its endophytic fungus yields gold nanoparticles of different shapes. *J. Mater. Chem.* 13, 1822. doi: 10.1039/b303808b

- Sharma, S. S., and Dietz, K.-J. (2006). The significance of amino acids and amino acid-derived molecules in plant responses and adaptation to heavy metal stress. *J. Exp. Bot.* 57, 711–726. doi: 10.1093/jxb/erj073
- Sim, C. S. F., Cheow, Y. L., Ng, S. L., and Ting, A. S. Y. (2018). Discovering metal-tolerant endophytic fungi from the phytoremediator plant *Phragmites*. *Water Air Soil Pollut.* 229, 1–11. doi: 10.1007/s11270-018-3733-1
- Singh, V. P., Singh, S., Kumar, J., and Prasad, S. M. (2015). Investigating the roles of ascorbate-glutathione cycle and thiol metabolism in arsenate tolerance in ridged *Luffa* seedlings. *Protoplasma* 252, 1217–1229. doi: 10.1007/s00709-014-0753-6
- Singh, P., Kim, Y.-J., Zhang, D., and Yang, D.-C. (2016). Biological synthesis of nanoparticles from plants and microorganisms. *Trends Biotechnol.* 34, 588–599. doi: 10.1016/j.tibtech.2016.02.006
- Song, W.-Y., Choi, K. S., Kim, D. Y., Geisler, M., Park, J., Vincenzetti, V., et al. (2010). *Arabidopsis* PCR2 is a zinc exporter involved in both zinc extrusion and long-distance zinc transport. *Plant Cell* 22, 2237–2252. doi: 10.1105/tpc.109.070185
- Sridhar, K. R. (2012). “Aspect and prospect of endophytic fungi” in *Microbes: Diversity and biotechnology*. eds. S. C. Sati and M. Belwal (New Delhi: Daya Publishing House), 43–62.
- Stefanowicz, A. M., Niklińska, M., and Laskowski, R. (2008). Metals affect soil bacterial and fungal functional diversity differently. *Environ. Toxicol. Chem.* 27, 591. doi: 10.1897/07-288.1
- Stewart, A., and Hill, R. (2014). “Applications of *Trichoderma* in plant growth promotion” in *Biotechnology and biology of Trichoderma*. eds. V. K. Gupta, M. S. A. Herrera-Estrella, R. S. Upadhyay, I. Druzhinina, and M. G. Tuohy (Oxford, United Kingdom: Elsevier), 415–428.
- Strullu-Derrien, C., Spencer, A. R. T., Goral, T., Dee, J., Honegger, R., Kenrick, P., et al. (2018). New insights into the evolutionary history of fungi from a 407 Ma Blastocladiomycota fossil showing a complex hyphal thallus. *Philos. Trans. R. Soc. B Biol. Sci.* 373, 20160502. doi: 10.1098/rstb.2016.0502
- Syamala, M., and Sivaji, M. (2017). Functional characterization of various plant growth promoting activity of *Pseudomonas fluorescens* and *Bacillus subtilis* from *Aloe vera* rhizosphere. *J. Pharmacogn. Phytochem.* 6, 120–122.
- Tong, J., Miaowen, C., Juhui, J., Jinxian, L., and Baofeng, C. (2017). Endophytic fungi and soil microbial community characteristics over different years of phytoremediation in a copper tailings dam of Shanxi, China. *Sci. Total Environ.* 574, 881–888. doi: 10.1016/j.scitotenv.2016.09.161
- Turnau, K., and Mesjasz-Przybyłowicz, J. (2003). Arbuscular mycorrhiza of *Berkheya coddii* and other Ni-hyperaccumulating members of Asteraceae from ultramafic soils in South Africa. *Mycorrhiza* 13, 185–190. doi: 10.1007/s00572-002-0213-6
- Turnau, K., Kotkka, L., and Dexheimer, J. (1996). Toxic element filtering in *Rhizopogon roseolus*/*Pinus sylvestris* mycorrhizas collected from calamine dumps. *Mycol. Res.* 100, 16–22. doi: 10.1016/S0953-7562(96)80094-3
- Turnau, K., Przybyłowicz, W., and Mesjasz-Przybyłowicz, J. (2001). Heavy metal distribution in *Suillus luteus* mycorrhizas – as revealed by micro-PIXE analysis. *Nucl. Instr. Meth. Phys. Res. Sect. B Beam Interact. Mater. Atoms* 181, 649–658. doi: 10.1016/S0168-583X(01)00631-0
- Turnau, K., Mleczko, P., Blaudez, D., Chalot, M., and Botton, B. (2002). Heavy metal binding properties of *Pinus sylvestris* mycorrhizas from industrial wastes. *Acta Soc. Bot. Pol.* 71, 253–261. doi: 10.5586/asbp.2002.030
- Usman, A. R. A., and Mohamed, H. M. (2009). Effect of microbial inoculation and EDTA on the uptake and translocation of heavy metal by corn and sunflower. *Chemosphere* 76, 893–899. doi: 10.1016/j.chemosphere.2009.05.025
- Vahabi, K., and Dorcheh, S. K. (2014). “Biosynthesis of silver nano-particles by *Trichoderma* and its medical applications” in *Biotechnology and biology of Trichoderma*. eds. V. K. Gupta, M. S. A. Herrera-Estrella, R. S. Upadhyay, I. Druzhinina, and M. G. Tuohy (Oxford, United Kingdom: Elsevier), 393–404.
- Vamerali, T., Bandiera, M., and Mosca, G. (2010). Field crops for phytoremediation of metal-contaminated land. A review. *Environ. Chem. Lett.* 8, 1–17. doi: 10.1007/s10311-009-0268-0
- van der Ent, A., Baker, A. J. M., Reeves, R. D., Pollard, A. J., and Schat, H. (2013). Hyperaccumulators of metal and metalloid trace elements: facts and fiction. *Plant Soil* 362, 319–334. doi: 10.1007/s11104-012-1287-3
- Vandenkoornhuyse, P., Quaiser, A., Duhamel, M., Le Van, A., and Dufresne, A. (2015). The importance of the microbiome of the plant holobiont. *New Phytol.* 206, 1196–1206. doi: 10.1111/nph.13312
- Verbruggen, N., Hermans, C., and Schat, H. (2009). Molecular mechanisms of metal hyperaccumulation in plants. *New Phytol.* 181, 759–776. doi: 10.1111/j.1469-8137.2008.02748.x
- Viti, C., Marchi, E., Decorosi, F., and Giovannetti, L. (2014). Molecular mechanisms of Cr(VI) resistance in bacteria and fungi. *FEMS Microbiol. Rev.* 38, 633–659. doi: 10.1111/1574-6976.12051
- Vogel-Mikuš, K., Pongrac, P., Kump, P., Nečemer, M., and Regvar, M. (2006). Colonisation of a Zn, Cd and Pb hyperaccumulator *Thlaspi praecox* Wulfen with indigenous arbuscular mycorrhizal fungal mixture induces changes in heavy metal and nutrient uptake. *Environ. Pollut.* 139, 362–371. doi: 10.1016/j.envpol.2005.05.005
- Walker, R. F., McLaughlin, S. B., and West, D. C. (2004). Establishment of sweet birch on surface mine spoil as influenced by mycorrhizal inoculation and fertility. *Restor. Ecol.* 12, 8–19. doi: 10.1111/j.1061-2971.2004.00255.x
- Wang, Y., Hu, H., Zhu, L.-Y., and Li, X.-X. (2012). Response to nickel in the proteome of the metal accumulator plant *Brassica juncea*. *J. Plant Interact.* 7, 230–237. doi: 10.1080/17429145.2011.603060
- Wang, W., Deng, Z., Tan, H., and Cao, L. (2013). Effects of Cd, Pb, Zn, Cu-resistant endophytic *Enterobacter* sp. CBSB1 and *Rhodotorula* sp. CBSB79 on the growth and phytoextraction of Brassica plants in multimetal contaminated soils. *Int. J. Phytorem.* 15, 488–497. doi: 10.1080/15226514.2012.716101
- Wang, J., Li, T., Liu, G., Smith, J. M., and Zhao, Z. (2016). Unraveling the role of dark septate endophyte (DSE) colonizing maize (*Zea mays*) under cadmium stress: physiological, cytological and genic aspects. *Sci. Rep.* 6, 22028. doi: 10.1038/srep22028
- Ważny, R., Rozpądek, P., Jędrzejczyk, R. J., Śliwa, M., Stojakowska, A., Anielska, T., et al. (2018). Does co-inoculation of *Lactuca serriola* with endophytic and arbuscular mycorrhizal fungi improve plant growth in a polluted environment? *Mycorrhiza* 28:235. doi: 10.1007/s00572-018-0819-y
- Wei, Y. F., Li, T., Li, L. F., Wang, J. L., Cao, G. H., and Zhao, Z. W. (2016). Functional and transcript analysis of a novel metal transporter gene EpNramp from a dark septate endophyte (*Exophiala pisciphila*). *Ecotoxicol. Environ. Saf.* 124, 363–368. doi: 10.1016/j.ecoenv.2015.11.008
- Weyens, N., van der Lelie, D., Taghavi, S., Newman, L., and Vangronsveld, J. (2009a). Exploiting plant-microbe partnerships to improve biomass production and remediation. *Trends Biotechnol.* 27, 591–598. doi: 10.1016/j.tibtech.2009.07.006
- Weyens, N., van der Lelie, D., Taghavi, S., and Vangronsveld, J. (2009b). Phytoremediation: plant-endophyte partnerships take the challenge. *Curr. Opin. Biotechnol.* 20, 248–254. doi: 10.1016/j.copbio.2009.02.012
- Wężowicz, K., Rozpądek, P., and Turnau, K. (2014). The diversity of endophytic fungi in *Verbascum lychnitis* from industrial areas. *Symbiosis* 64, 139–147. doi: 10.1007/s13199-015-0312-8
- Wężowicz, K., Rozpądek, P., and Turnau, K. (2017). Interactions of arbuscular mycorrhizal and endophytic fungi improve seedling survival and growth in post-mining waste. *Mycorrhiza* 27, 499–511. doi: 10.1007/s00572-017-0768-x
- White, J. F., Kingsley, K. L., Verma, S. K., and Kowalski, K. P. (2018). Rhizophagy cycle: an oxidative process in plants for nutrient extraction from symbiotic microbes. *Microorganisms*. doi: 10.3389/fphys.2018.01865
- Wu, F., and Zhang, G. (2002). Genotypic variation in kernel heavy metal concentrations in barley and ALS as affected by soil factors. *J. Plant Nutr.* 25, 1163–1173. doi: 10.1081/PLN-120004380
- Wu, S., Zhang, X., Sun, Y., Wu, Z., Li, T., Hu, Y., et al. (2015). Transformation and Immobilization of Chromium by Arbuscular Mycorrhizal Fungi as Revealed by SEM-EDS, TEM-EDS, and XAFS. *Environ. Sci. Technol.* 49, 14036–14047. doi: 10.1021/acs.est.5b03659
- Wu, C., Li, B., Wei, Q., Pan, R., and Zhang, W. (2018). Endophytic fungus *Serendipita indica* increased nutrition absorption and biomass accumulation in *Cunninghamia lanceolata* seedlings under low phosphate. *Acta Ecol. Sin.* 39, 21–29. doi: 10.1016/j.chnaes.2018.06.005
- Xu, R., Li, T., Cui, H., Wang, J., Yu, X., Ding, Y., et al. (2015). Diversity and characterization of Cd-tolerant dark septate endophytes (DSEs) associated with the roots of Nepal alder (*Alnus nepalensis*) in a metal mine tailing of southwest China. *Appl. Soil Ecol.* 93, 11–18. doi: 10.1016/j.apsoil.2015.03.013
- Yadav, V., Kumar, M., Deep, D. K., Kumar, H., Sharma, R., Tripathi, T., et al. (2010). A phosphate transporter from the root endophytic fungus *Piriformospora indica* plays a role in phosphate transport to the host plant. *J. Biolumin. Chemilumin.* 285, 26532–26544. doi: 10.1074/jbc.M110.111021

- Yamaji, K., Watanabe, Y., Masuya, H., Shigeto, A., Yui, H., and Haruma, T. (2016). Root fungal endophytes enhance heavy-metal stress tolerance of *Clethra barbinervis* growing naturally at mining sites via growth enhancement, promotion of nutrient uptake and decrease of heavy-metal concentration. *PLoS One* 11, 1–15. doi: 10.1371/journal.pone.0169089
- Yao, H., Xu, J., and Huang, C. (2003). Substrate utilization pattern, biomass and activity of microbial communities in a sequence of heavy metal-polluted paddy soils. *Geoderma* 115, 139–148. doi: 10.1016/S0016-7061(03)00083-1
- Yin, Y., Hu, Y., and Xiong, F. (2011). Sorption of Cu(II) and Cd(II) by extracellular polymeric substances (EPS) from *Aspergillus fumigatus*. *Int. Biodeterior. Biodegrad.* 65, 1012–1018. doi: 10.1016/j.ibiod.2011.08.001
- Yuan, Z. L., Zhang, C. L., and Lin, F. C. (2010). Role of diverse non-systemic fungal endophytes in plant performance and response to stress: progress and approaches. *J. Plant Growth Regul.* 29, 116–126. doi: 10.1007/s00344-009-9112-9
- Zahoor, M., Irshad, M., Rahman, H., Qasim, M., Afridi, S. G., Qadir, M., et al. (2017). Alleviation of heavy metal toxicity and phytostimulation of *Brassica campestris* L. by endophytic *Mucor* sp. MHR-7. *Ecotoxicol. Environ. Saf.* 142, 139–149. doi: 10.1016/j.ecoenv.2017.04.005
- Zhang, Z.-C., Chen, B.-X., and Qiu, B.-S. (2010). Phytochelatin synthesis plays a similar role in shoots of the cadmium hyperaccumulator *Sedum alfredii* as in non-resistant plants. *Plant Cell Environ.* 33, 1248–1255. doi: 10.1111/j.1365-3040.2010.02144.x
- Zhang, Q., Zhang, J., Yang, L., Zhang, L., Jiang, D., Chen, W., et al. (2014). Diversity and biocontrol potential of endophytic fungi in *Brassica napus*. *Biol. Control* 72, 98–108. doi: 10.1016/j.biocontrol.2014.02.018
- Zhao, D., Li, T., Shen, M., Wang, J., and Zhao, Z. (2015a). Diverse strategies conferring extreme cadmium (Cd) tolerance in the dark septate endophyte (DSE), *Exophiala pisciphila*: evidence from RNA-seq data. *Microbiol. Res.* 170, 27–35. doi: 10.1016/j.micres.2014.09.005
- Zhao, W., Han, J., and Long, D. (2015b). Effect of copper-induced oxidative stress on sclerotial differentiation, endogenous antioxidant contents, and antioxidative enzyme activities of *Penicillium thomii* PT95. *Ann. Microbiol.* 65, 1505–1514. doi: 10.1007/s13213-014-0989-6
- Zhou, Y., Li, X., Gao, Y., Liu, H., Gao, Y. B., van der Heijden, M. G. A., et al. (2018). Plant endophytes and arbuscular mycorrhizal fungi alter plant competition. *Funct. Ecol.* 32, 1168–1179. doi: 10.1111/1365-2435.13084
- Zhu, S., Tang, J., Zeng, X., Wei, B., Yang, S., and Huang, B. (2015). Isolation of *Mucor circinelloides* Z4 and *Mucor racemosus* Z8 from heavy metal-contaminated soil and their potential in promoting phytoextraction with Guizhou oilseed rap. *J. Cent. South Univ.* 22, 88–94. doi: 10.1007/s11771-015-2498-6

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Domka, Rozpałek and Turnau. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.