



Stoichiometric Shifts in Soil C:N:P Promote Bacterial Taxa Dominance, Maintain Biodiversity, and Deconstruct Community Assemblages

Zachary T. Aanderud^{1*}, Sabrina Saurey¹, Becky A. Ball², Diana H. Wall³, John E. Barrett⁴, Mario E. Muscarella⁵, Natasha A. Griffin¹, Ross A. Virginia⁶, Albert Barberán⁷ and Byron J. Adams⁸

¹ Department of Plant and Wildlife Sciences, Brigham Young University, Provo, UT, United States, ² School of Mathematical and Natural Sciences, Arizona State University, Phoenix, AZ, United States, ³ Department of Biology, School of Global Environmental Sustainability, Colorado State University, Fort Collins, CO, United States, ⁴ Department of Biological Sciences, Virginia Polytechnic Institute, Blacksburg, VA, United States, ⁵ Department of Plant Biology, University of Illinois Urbana-Champaign, Champaign, IL, United States, ⁶ Environmental Studies Program, Dartmouth College, Hanover, NH, United States, ⁷ Department of Soil, Water and Environmental Science, University of Arizona, Tucson, AZ, United States, ⁸ Evolutionary Ecology Laboratories, and Monte L. Bean Museum, Department of Biology, Brigham Young University, Provo, UT, United States

OPEN ACCESS

Edited by:

Thulani Peter Makhalanyane,
University of Pretoria, South Africa

Reviewed by:

David Anthony Pearce,
Northumbria University,
United Kingdom
Richard Allen White III,
RAW Molecular Systems (RMS) LLC,
United States

*Correspondence:

Zachary T. Aanderud
zachary_aanderud@byu.edu

Specialty section:

This article was submitted to
Extreme Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 21 February 2018

Accepted: 07 June 2018

Published: 03 July 2018

Citation:

Aanderud ZT, Saurey S, Ball BA, Wall DH, Barrett JE, Muscarella ME, Griffin NA, Virginia RA, Barberán A and Adams BJ (2018) Stoichiometric Shifts in Soil C:N:P Promote Bacterial Taxa Dominance, Maintain Biodiversity, and Deconstruct Community Assemblages. *Front. Microbiol.* 9:1401. doi: 10.3389/fmicb.2018.01401

Imbalances in C:N:P supply ratios may cause bacterial resource limitations and constrain biogeochemical processes, but the importance of shifts in soil stoichiometry are complicated by the nearly limitless interactions between an immensely rich species pool and a multiple chemical resource forms. To more clearly identify the impact of soil C:N:P on bacteria, we evaluated the cumulative effects of single and coupled long-term nutrient additions (i.e., C as mannitol, N as equal concentrations NH_4^+ and NO_3^- , and P as Na_3PO_4) and water on communities in an Antarctic polar desert, Taylor Valley. Untreated soils possessed relatively low bacterial diversity, simplified organic C sources due to the absence of plants, limited inorganic N, and excess soil P potentially attenuating links between C:N:P. After 6 years of adding resources, an alleviation of C and N colimitation allowed one rare Micrococcaceae, an *Arthrobacter* species, to dominate, comprising 47% of the total community abundance and elevating soil respiration by 136% relative to untreated soils. The addition of N alone reduced C:N ratios, elevated bacterial richness and diversity, and allowed rare taxa relying on ammonium and nitrite for metabolism to become more abundant [e.g., nitrite oxidizing *Nitrospira* species (Nitrosomonadaceae), denitrifiers utilizing nitrite (Gemmatimonadaceae) and members of Rhodobacteraceae with a high affinity for ammonium]. Based on community co-occurrence networks, lower C:P ratios in soils following P and CP additions created more diffuse and less connected communities by disrupting 73% of species interactions and selecting for taxa potentially exploiting abundant P. Unlike amended nutrients, water additions alone elicited no lasting impact on communities. Our results suggest that as soils become nutrient rich a wide array of outcomes are possible from species dominance and the deconstruction of species interconnectedness to the maintenance of biodiversity.

Keywords: ecological stoichiometry, Lake Fryxell Basin, McMurdo Dry Valleys, network community modeling, nutrient colimitation, Solirubrobacteriaceae

INTRODUCTION

Environmental conditions dramatically structure soil bacterial communities; however, only a few environmental variables such as pH, salinity, and C substrate quality and quantity are known to drive community assemblages (Lozupone and Knight, 2007; Fierer et al., 2009; Baldrian et al., 2012). Borrowing from foundations in plant community ecology, following light and water, nutrient additions dramatically alter species abundance within communities often allowing certain species to dominate and biodiversity to decline (Bedford et al., 1999; Bobbink et al., 2010). For example, in a grassland, even low continual additions of N reduced species richness in just 2 years and depressed the number of species for as long as 20 years (Clark and Tilman, 2008). Classically, soil bacterial metabolism and growth is limited by water availability, then the quality and quantity of C substrates, and finally nutrient concentrations. But the role of major nutrients, such as N and P, remains incomplete even though changes in nutrient availability shape the responses of specific bacterial species or species interactions within soil communities (Marschner et al., 2003; Ramirez et al., 2010). Unlike plants, bacterial responses to resource constraints are complicated by interactions within consortia or communities where functionally disparate taxa (e.g., decomposers, nitrifiers, and methanogens) potentially dictate the form and availability of specific C substrates and nutrients necessary for other bacteria to become metabolically active and grow. In most soil communities such interactions are further complicated because soil organic C substrates are extremely numerous and diverse, containing both labile and more recalcitrant sources structuring the availability of N and P (Orwin et al., 2006; Hernandez and Hobbie, 2010). Furthermore, the C, N, and P requirements of bacterial biomass differ among species and ecosystems, and are not homeostatic through time (Cleveland and Liptzin, 2007; Hartman and Richardson, 2013).

Ecological stoichiometry is a unifying body of theory in ecology predicting relationships between the organismal biochemistry of plants, invertebrates, and microorganisms and the availability and recycling of nutrient elements in the environment (Elser and Hamilton, 2007). Ecological stoichiometry may also help identify the resource requirements of bacterial taxa and the conditions allowing certain bacteria to become metabolically active. Ecological stoichiometric theory was developed in aquatic ecosystems, but is universally valid, and over the last decades was also successfully applied to terrestrial systems (Redfield, 1958; Reiners, 1986; Cleveland and Liptzin, 2007; Austin and Vitousek, 2012). Soil (186:13:1) and soil microbial (60:7:1) C:N:P stoichiometry, like Redfield ratios for planktonic biomass (C:N:P = 106:16:1) (Redfield, 1958), are well-constrained across multiple biomes (Cleveland and Liptzin, 2007) offering incredible utility in understanding bacterial resource limitations and constraints on biogeochemical processes (Tian et al., 2010; Xu et al., 2013). The C:N:P stoichiometry of plant residues, soil organic matter, and bacterial biomass influence litter decomposition rates (Aneja et al., 2006; Zechmeister-Boltenstern et al., 2015), N and P mineralization

rates (Mooshammer et al., 2012), and C-use-efficiency of bacteria, determining metabolic activity and trace gas flux (Keiblinger et al., 2010). Community structure is intimately connected to C:N:P ratios (Elser et al., 2000). Specifically, soil C:N:P stoichiometry sheds light on the potential for N and P availability to influence bacterial community structure. For example, higher N:P ratios in afforested soils of the Loess Plateau in China reflected P deficiencies among bacteria, leading to lower diversity but a higher abundance of Proteobacteria, Acidobacteria, and Nitrospirae (Ren et al., 2016). Further, a decrease in soil C:P ratios caused Gram-positive bacterial biomass to increase by 22% and the abundance of arbuscular mycorrhizal fungi to increase by 46% in a pasture following slash-and-burn agriculture in the South Ecuadorian Andes (Tischer et al., 2015). Taken together, investigating soil C:N, C:P, and N:P ratios are instrumental in identifying patterns of ecological coherence among responding bacteria under varying resource conditions.

Soil ecosystems of the McMurdo Dry Valleys, Eastern Antarctica are a model system for investigating stoichiometric controls over soil communities and ecosystem processes (Barrett et al., 2006). The extreme environment limits biota to cryptogrammic vegetation, a few taxa of metazoan invertebrates, and microbial dominated food webs. Phylum-level bacterial diversity in Antarctic soils is surprisingly high considering the environmental extremes and dearth of resources, i.e., organic matter and available nutrients (Cary et al., 2010; Lee et al., 2011). However, these soils host comparatively low diversity at the family or genus level relative to other biomes (Fierer et al., 2012), attenuating the nearly limitless possibilities of links between C:N:P stoichiometry and communities potentially present in high diversity ecosystems. Further, soil C:N:P ratios in most systems are necessarily complicated by plant residues with multiple different stoichiometric ratios that may mask links between C sources and release rate of N and P (Elser et al., 2000). Alternatively, due to the absence of vascular plants, Antarctic Dry Valley soils have some of the lowest soil organic matter concentrations on Earth (Burkins et al., 2000; Jobbagy and Jackson, 2000) with much of the soil organic matter being “legacy” C accumulated over thousands of years by cryptoendolithic bacteria, paleolake deposition, and minor inputs of contemporary algae and cyanobacteria from lakes, intermittent streams and saturated zones (Burkins et al., 2000). The concentrations of inorganic N and P are relatively low with N entering the system via atmospheric deposition (Michalski et al., 2005), endolithic and hypolithic cyanobacterial N₂ fixation over millions of years (Cowan et al., 2014), and from dust, while P enters soils through mineral weathering (Barrett et al., 2007; Heindel et al., 2017). Both N and P concentrations vary among soils occurring on glacial tills with distinct exposure age and mineralogy (Barrett et al., 2007). Accompanying nutrient limitations, water is a universal resource essential for polar bacteria. In the McMurdo Dry Valleys, low precipitation inputs and sublimation and ablation processes (Fountain et al., 1999) cause dehydration stress and limit substrate diffusion to bacterial cells (Stark

and Firestone, 1995). Water additions to Antarctic soils do create higher bacterial growth rates, elevate soil respiration, and decrease soil community diversity (Schwartz et al., 2014; Buelow et al., 2016). Therefore, water, in addition to nutrients, may have direct and indirect effects on community composition.

In this study, we explored the effects of long-term, coupled resource additions and water on bacterial species responses and ecosystem processes in a cold desert of Antarctica. After treating soils with six different resource additions including combinations of water, C as mannitol, N as equal NH_4^+ NO_3^- , and P as Na_3PO_4 annually over 6 years in the field, we evaluated shifts in bacterial community composition metrics such as richness, alpha diversity and evenness, taxa co-occurrence patterns, and soil respiration. Based on polar desert resource conditions, our initial soil C:N:P ratio (mean = 167:8:1, $n = 8$; more initial soil chemistry data is provided in the first section of the “Materials and Methods”), and the modal C:N:P ratio of soils (186:13:1, Cleveland and Liptzin, 2007), we hypothesized that C, N, CN, and CP additions will alleviate resource limitations and provide organic C and inorganic N for a subset of the community to exploit, while P is not limiting and adding more inorganic P will not alter community composition.

MATERIALS AND METHODS

Study Site and Initial Soil Chemistry

Our study was conducted in a polar desert of the McMurdo Dry Valleys (76°30′ – 78°00′S, 160°00′ – 165°00′E) at the McMurdo Long Term Ecological Research (LTER) site in Antarctica. The site was located in the Lake Fryxell basin (77°36.5′S, 163°14.9′E) of Taylor Valley, on Ross Sea drift soils (late-Quaternary) (Bockheim et al., 2008).

Dominant soil microflora include multiple species of algae from the division Chlorophyta and Heterokontophyta; microfauna comprised of nematodes, tardigrades and rotifers; and cyanobacteria, such species as, *Leptolyngbya frigida* and *Nostoc commune* in aquatic and terrestrial habitats (Adams et al., 2006). The basin experiences fewer than 50 days where average temperatures exceed 0°C within the summer months of December, January, February when mean annual temperature is $-4.21^\circ\text{C} \pm 0.80$ SD ($n = 24$). Soils receive less than 10 cm per yr^{-1} of effective mean annual precipitation falling as snow (Doran et al., 2002). Soils are Typic Haploturbels with shallow surface layer (≈ 0 –10 cm depth), which experience continual cryoturbation, and a perennial permafrost layer (≈ 30 –300 cm depth) (Bockheim and McLeod, 2008; Bockheim et al., 2008). All soils are poorly developed silty-loams with an average pH of 9.69 ± 0.12 SEM ($n = 8$) and an electrical conductivity of 258 ± 115 $\mu\text{S cm}^{-1}$ ($n = 8$). Initial soil chemistry demonstrated that soils were generally extremely low in organic C (organic C = $0.03\% \pm 0.003$, total soil C = $0.13\% \pm 0.01$, $n = 8$) and possessed relatively high soil P (2.35 ± 0.18 $\mu\text{g g}^{-1}$ soil, $n = 8$) but low levels of soil N of ($0.003\% \pm 0.0004$, $n = 8$).

Stoichiometry and Water Long-Term Manipulations

To gain insights into the resource controls on microbial community assembly, we conducted a 6-year field stoichiometry experiment (austral summer field season 2006 – 2007 to 2011 – 2012) altering the stoichiometry of major nutrients (i.e., C, N, and P) and water availability. The experiment was a randomized block design with plots (1 m × 1 m) consisting of six treatments and an un-amended control: water only (W); C as mannitol and water (C); N as equal concentrations NH_4^+ and NO_3^- with water (N); P as Na_3PO_4 and water (P); C, N, and water (CN); C, P, with water (CP); and the untreated control (U). The C additions, as mannitol, mirrored nutrient inputs from more contemporary algae and cyanobacteria. The N and P additions closely followed the Redfield ratio (106:16:1) to mimic new biomass from photosynthetic organisms entering organic matter-impooverished soils (Hecky et al., 1993). Annually, all nutrients were delivered as aqueous solutions to bring the soils to field capacity with concentrations of 15.3 g C m^{-2} , 2.69 g N m^{-2} as NH_4NO_3 , and 0.37 g P m^{-2} as $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ and water of 12.7 L $\text{H}_2\text{O m}^{-2}$. For more information on the treatments and treatment application in the field see Ball et al. (2018). The present study comprises an analysis of the Fryxell basin site only.

Soil C:N:P and Chemistry

To measure post-amendment changes in C:N:P stoichiometry we calculated C:N:P ratios from total C and N, and extractable P, and measured soil organic C and inorganic N. Soils were collected from all treatments (5 nutrient additions with water, 1 water addition, and a control × 8 replicates = 56) with a plastic scoop to a soil depth of 10 cm (approximately 500 g), sieved (2 mm sieve), and frozen until processing. Total C and N were measured on a Elantech Flash EA 1200 (CE Elantech, NJ, United States). Extractable soil P, as phosphate, was measured in 10 g soil with 0.5 M NaHCO_3 (1:5 w/v) at pH 8.5, acidified with 3 mL of 6 N HCl, and analyzed on a Lachat Autoanalyzer (Barrett et al., 2007). We measured dissolved organic C on a Shimadzu TOC-5000A (Shimadzu Corporation, Columbia, MD, United States). Inorganic N ($\mu\text{g N-NH}_4^+$ g soil $^{-1}$, $\mu\text{g N-NO}_3^-$ g soil $^{-1}$) was evaluated from 20 g of soil extracted with 2 M KCl extraction (1:2.5 w/v), passed through a Whatman #1 filter, and measured on an a Lachat Quikchem 8500 (Lachat Instruments, Loveland, CO, United States). We tested for the effect of the additions on our response variables and soil C:N:P ratios using one-way ANOVA and Tukey’s HSD test to identify significant differences among the treatments in R (R Development Core Team, 2013). For stoichiometric analyses data were converted into molar ratios.

Bacterial Community Responses to Changes in Soil C:N:P and Water

After maintaining the treatments for more than half a decade, we characterized soil communities in treatment soils using barcoded sequencing of the 16S rRNA gene. Soils were collected from three randomly selected replicates in all treatments (5 nutrient additions with water, 1 water addition, and a control × 3 replicates = 21) to a depth of 10 cm using sterile plastic scoops.

All soils were transported from the field in an insulated chest, sieved to 2 mm, and stepped-down to -20°C over 24 h. Nucleic acids were extracted from 1.5 g of soil using a PowerSoil DNA Isolation Kit (Mo Bio Corporation, Carlsbad, CA, United States). We PCR amplified the V4 region of the 16S rRNA gene using bacterial specific primer set 515F and 806R with unique 12-nt error correcting Golay barcodes (Caporaso et al., 2010; Aanderud et al., 2013). The thermal cycle conditions consisted of an initial denaturing step at 94°C for 3 min followed by 35 cycles of denaturing at 94°C for 45 s, annealing at 50°C for 30 s, and amplifying at 72°C for 90 s. After purifying (Agencourt AMPure XP PCR Purification Beckman Coulter Inc., Brea, CA, United States) and pooling PCR amplicons at approximately equimolar concentrations, samples were sequenced at the Brigham Young University DNA Sequencing Center¹ using a 454 Life Sciences genome sequencer FLX (Roche, Branford, CT, United States). All sequences were trimmed and cleaned using mothur [v. 1.31.2; All sequences were trimmed and cleaned using mothur (v. 1.31.2²; Schloss et al., 2009)]. After removing barcodes and primers, we eliminated sequences that were <250 bp in length or possessed homopolymers longer than 8 bp. We then denoised the sequences with AmpliconNoise (Quince et al., 2011), removed chimeras using UCHIME (Edgar et al., 2011), and eliminated chloroplast, mitochondrial, archaeal, and eukaryotic 16S rRNA gene sequences based on reference sequences from the Ribosomal Database Project (Cole et al., 2009). We aligned the sequences against the SILVA database (Pruesse et al., 2007) with the SEED aligner, created operational taxonomic units (OTUs) based on uncorrected pairwise distances at 97% sequence similarity, and determined the phylogenetic identity of OTUs using the SILVA database.

To assess the effects of resources on soil bacterial communities, we first visualized differences in community composition using Principal Coordinates Analysis (PCoA) based on a Bray–Curtis distance matrix with the ‘vegan’ package in R (R Development Core Team, 2013). We then quantitated the effects of the different resource treatments [i.e., W, C, single nutrients N and P, and combined C and nutrients (CN and CP)] on community compositions using permutational multivariate analyses of variance (PERMANOVA) (Anderson, 2001) performed with the *adonis* function in the vegan package in R (Oksanen et al., 2013). Second, we quantified and constructed 95% confidence intervals for estimated richness as the total number of OTUs, alpha diversity as the inverse Shannon index, and taxa evenness using Pielou’s evenness based on 1000 iterations of 900 random resampled sequences from each replicate (Muscarella et al., 2014). Last, we calculated the relative recovery of nine phyla and three subclasses to identify differences in the distribution of major taxonomical groups (recovery $\geq 1.0\%$) due to the resource additions. Taxonomic trends of 24 families (recovery $\geq 1.0\%$ in at least one replicate) were visualized in a heat map with hierarchical clustering using the *heatmap* function in the ‘gplot’ package in R (Warnes et al., 2014).

¹<http://dnac.byu.edu/>

²https://www.mothur.org/wiki/454_SOP

Bacterial Community Network Models

To assess changes within communities at the OTU taxonomical level, we created network co-occurrence models for combinations of resources based on maximal information coefficient (MIC) analysis. We calculated all possible linear and non-linear associations between OTUs using the *mic* function and the ‘minerva’ package in R (Filosi et al., 2017), which belongs to a class of maximal information-based non-parametric exploration statistics for identifying and classifying relationships (Reshef et al., 2011). The nodes in the networks represented individual OTUs at 97% identity, while edges corresponded to valid or significant co-occurrence connections that occurred in at least 75% of all samples and had a MIC that was both >0.7 and statistically significant (P -value = 0.01) (Barberan et al., 2012). This filtering facilitated the determination of the OTUs interacting within the treatments and removed poorly represented OTUs reducing network complexity (Barberan et al., 2012). We described the network through a series of topological parameters: mean path length, mean degree, mean clustering coefficient, density, and modularity (Freedman et al., 2016). Network graphs in the *graphml* format were generated using ‘igraph’ package in R (Csardi and Nepusz, 2006) and visualized with the interactive platform Gephi (v. 0.8.2-beta) (Bastian et al., 2009). To identify the taxonomy of bacteria within the networks, we elevated nodes at the order taxonomical rank. We calculated the node number as the total number of nodes within each of the nineteen orders comprising the networks, and the relative recovery of nodes as the summation of the mean relative recoveries of the nodes within an a given order from P and CP or U and W communities.

Soil Respiration

To investigate the links between microbial communities and ecosystem processes, we measured soil respiration ($\mu\text{moles C-CO}_2 \text{ m}^{-2} \text{ soil sec}^{-1}$) in all resource treatments (5 nutrient additions with water, 1 water addition, and a control $\times 8$ replicates = 56). Within 1 week of resource additions, soil CO_2 flux in the field was evaluated using a Li-COR 8100 (LI-COR Biosciences, Lincoln, NE, United States) with a 10-cm diameter PVC ring inserted 2 cm into the soil at least 1 h prior to measurement (Ball et al., 2018).

RESULTS

N and P Additions Altered Soil C:N:P

Resource additions clearly altered soil C:N:P leading to increases in N and P availability. Following multiple years of N additions, C:N were lower and N:P were higher in N and CN than P and CP soils (Table 1). The shifts in ratios were highlighted by inorganic NH_4^+ (one-way ANOVA, $df = 6$, $F = 39.1$, $P < 0.001$) and NO_3^- (one-way ANOVA, $df = 6$, $F = 14.9$, $P < 0.001$) concentrations being at least forty-four- and nine-times higher, respectively, in N and CN than all other soil treatment (Table 2). Accompanying soil P additions, C:P were lower in P and CP soils than all other treatments besides CN (Table 1). Similarly, extractable P increased more than 53.0% in P and CP relative to

TABLE 1 | Molar C:N:P ratios of soils following 6 years of six resource additions in Fryxell lake basin of the Taylor Valleys in Antarctica.

	C:N:P	C:N	C:P	N:P
U	182:7:1	28.2 ± 2.69 ab	182 ± 29.7 ab	6.85 ± 1.01 bcd
W	247:8:1	31.6 ± 5.76 ab	247 ± 45.4 a	8.28 ± 0.768 bcd
C	218:8:1	30.0 ± 4.82 ab	218 ± 21.0 ab	8.43 ± 1.31 bc
N	223:16:1	14.4 ± 1.17 b	223 ± 30.8 a	15.5 ± 1.89 a
P	90:3:1	33.0 ± 49 a	90.3 ± 4.65 c	3.02 ± 0.400 d
CN	165:12:1	14.4 ± 1.68 b	165 ± 26.5 abc	11.9 ± 1.85 ab
CP	108:3:1	37.8 ± 6.00 a	108 ± 5.93 bc	3.47 ± 0.675 cd

Treatments abbreviations include: un-amended control (U); water only (W); C as mannitol and water (C); N as equal concentrations NH_4^+ and NO_3^- with water (N); P as Na_3PO_4 and water (P); C, N, and water (CN); C, P, with water (CP). Data are mean (\pm SEM, $n = 8$) with different letters indicating differences among treatments ($P < 0.05$) from ANOVA and Tukey's HSD.

W, C, and N (one-way ANOVA, $df = 6$, $F = 6.26$, $P < 0.001$, **Table 2**). Conversely, the additions of C had no apparent effect on total soil C only slightly increasing SOC in the C compared to P and W treatments (one-way ANOVA, $df = 6$, $F = 3.80$, $P = 0.004$, **Table 2**). The C:N:P ratios for each treatment are listed in **Table 1**.

CN Reduced Evenness and Diversity but N Alone Enhanced Richness and Diversity

Bacterial evenness and diversity was reduced following CN additions, while diversity and richness was enhanced by N additions (**Figure 1**). Specifically, the addition of CN dramatically depressed taxa evenness by at least 20.1% relative to all other treatments, and alpha diversity by at least 24.5% relative to the C, N, and CP treatments. In contrast, N additions stimulated OTU richness by 48.4 and 38.9% relative to bacterial communities in C and P soils, respectively. Diversity also increased by more than 13% in N compared to C, P, and CN treatments. In general, all resource additions reduced the variability surrounding richness and diversity metrics. All community inferences were based on the recovery of 138,458 quality sequences and 1,450 unique OTUs with samples possessing an average sequencing coverage of $98.4\% \pm 0.21$ (mean and SEM). All sequence data were submitted to NCBI and are available as BioProject PRJNA476992.


All Nutrient Additions Created Distinct Communities, Especially CN

The CN treatment dramatically influenced bacterial communities, most notably by the separation of communities along axis one, which explained 31.6% of the variation among communities (**Figure 2**). Further, the addition of any nutrient (i.e., C, N, P, and CP) reduced the variability among communities compared to the untreated control and water only addition along axis 2, which explained 19.6% of the variation. PERMANOVA results supported these interpretations, as communities were distinct among treatments ($F = 3.5$, $R^2 = 0.60$, $P < 0.001$, $df = 6$).

TABLE 2 | Soil chemistry following 6 years of resource additions.

	U	W	C	N	P	CN	CP
pH	9.7 ± 0.15	9.9 ± 0.05	9.9 ± 0.12	9.5 ± 0.20	9.8 ± 0.10	9.6 ± 0.17	9.8 ± 0.12
EC (ds m^{-1})	220 ± 42.1	183 ± 45.7	224 ± 40.9	331 ± 146	220 ± 42.0	301 ± 97.4	266 ± 73.8
Total C (%)	0.14 ± 0.01	0.12 ± 0.01	0.14 ± 0.01	0.14 ± 0.02	0.13 ± 0.01	0.15 ± 0.01	0.16 ± 0.02
Total N (%)	0.004 ± 0.001 bc	0.004 ± 0.001 c	0.005 ± 0.001 bc	0.008 ± 0.001 ab	0.004 ± 0.001 c	0.01 ± 0.001 a	0.004 ± 0.001 bc
P ($\mu\text{g N g soil}^{-1}$)	3.6 ± 0.86 ab	2.2 ± 0.37 b	2.7 ± 0.39 b	2.5 ± 0.23 b	5.6 ± 0.67 a	4.2 ± 0.74 ab	5.8 ± 0.54 a
SOC ($\mu\text{g N g soil}^{-1}$)	0.03 ± 0.004 ab	0.03 ± 0.004 b	0.06 ± 0.008 a	0.03 ± 0.004 ab	0.03 ± 0.004 b	0.04 ± 0.007 ab	0.05 ± 0.005 ab
N-NH ₄ ⁺ ($\mu\text{g N g soil}^{-1}$)	0.08 ± 0.09 b	0.27 ± 0.05 b	0.04 ± 0.01 b	17 ± 1.6 a	0.29 ± 0.10 b	13 ± 2.6 a	0.05 ± 0.01 b
N-NO ₃ ⁻ ($\mu\text{g N g soil}^{-1}$)	0.64 ± 0.30 c	1.1 ± 0.57 c	0.09 ± 0.03 c	22 ± 4.2 a	0.77 ± 0.26 c	10 ± 2.5 b	1.1 ± 1.1 c

For treatment abbreviations see **Table 1**. Data are mean (\pm SEM, $n = 8$) with different letters indicating significant differences among treatments ($P < 0.05$) based on ANOVA and Tukey's HSD.



Family-Specific Responses to Nutrients

All resource additions, except water, promoted taxonomic shifts in bacterial OTU abundance in 11 families across five phyla (Figure 3). The most pronounced increase in relative recovery occurred in the Micrococcaceae (Actinobacteria) in CN soils where a single bacterium, an *Arthrobacter species*, was relatively rare ($0.06\% \pm 0.05$) in untreated soils but constituted $47\% \pm 5.6$ of the community in CN-amended soils. The only other bloom occurred in the Trueperaceae (*Deinococcus*), where one OTU was classified as intermediate ($0.42\% \pm 0.09$) in untreated but became abundant ($9.6\% \pm 3.8$) in CP soils. In general, Actinobacteria were abundant in all soils and accounted for at least 36% of the community composition in all treatments (Supplementary





Figure S1), but not all Actinobacteria responded positively to CN. For example, three Actinobacteria families, Solirubrobacteraceae, Solirubrobacterales unclassified, and Rubrobacteriaceae, decreased from 2.1- to 4.9-fold in CN compared to all other treatments (Figure 3). CN additions also stimulated Xanthomonadaceae ($2.7\% \pm 1.2$, Gammaproteobacteria) and Sphingobacteriaceae ($1.8\% \pm 0.48$, Bacteroidetes). With the addition of N, the Nitrosomonadaceae (Betaproteobacteria) increased in recovery 5.2-times allowing the N treatment to have the highest recovery of Betaproteobacteria ($3.1\% \pm 0.17$). Annual N additions also enhanced the recovery of Rhodobacteraceae ($0.51\% \pm 0.16$, Alphaproteobacteria) and Gemmatimonadaceae ($2.7\% \pm 0.27$, Gemmatimonadetes) by at least 1.8-times relative to all other treatments. Both CP and P additions enhanced the recovery of Chitinophagaceae (Bacteroidetes) and Spartobacteria unclassified (Verrucomicrobia) in particular. The recovery of these families was at least 1.5-times higher in P and CP than all other soils and caused the recovery of Bacteroidetes to increase upward of 100% ($P = 6.3\% \pm 2.2$, $CP = 6.8\% \pm 1.8$) in both treatments.

Deinococcus-Thermus were present in all soils (Supplementary Figure S1), but the addition of CP, C, and P stimulated the recovery of these taxa causing the Deinococcaceae to range from 5.3 to 11% (Figure 3). The addition of C alone promoted different families with Sphingomonadaceae (Alphaproteobacteria) and the Geodermatophilaceae (Actinobacteria) increasing upward of 3.0- and 39-times, respectively, but combined these two taxa only accounted for less than 4.1% of the community in C-amended soils.

Network Community Modeling

Nutrient additions disrupted interactions among community assemblages. Due to the requirement of more than three samples



to create reliable community network models with MIC, we only created community network models for two combined treatments (U and W, and P and CP) that were relatively similar (PERMANOVA: U and W, $F = 5.8$, $R^2 = 0.59$, $P = 0.1$, $df = 1$; PW and CPW, $F = 1.2$, $R^2 = 0.22$, $P = 0.5$, $df = 1$). After years of P and CP additions, multiple aspects of the community broke down relative to the untreated and water only soils (**Figure 4** and **Table 3**). For example, the number of significant nodes or taxa, and edges or connections between taxa was 51 and 73% lower, respectively, in P and CP networks. The mean degree (number of connections per node to other nodes) declined twofold from U and W to P and CP networks, and mean path length (number of nodes needed to link any one node to any other in the network) decreased from 3.5 in the U

and W to 2.8 in P and CP models. Within the two networks, specific orders were favored and the nodes were often major contributors to the recovery of the community. For example, the Trueperaceae represented 3 nodes in the combined P and CP network and 7.7% relative recovery in the P and the CP communities, but only 1 taxon in U and W network and 0.61% of the recovery in U and W communities. Alternatively, in the U and W network, Phycisphaerae (unclassified; 10 nodes, 2.1% relative recovery), Intraspangiaceae (2 nodes, 0.13% relative recovery), and Xanthomonadaceae (2 nodes, 0.55% relative recovery) were present but completely absent from the P and CP network. The Spartobacteria (unclassified), Micrococcaceae, and Chitinophagaceae had similar numbers of nodes in all models, but contributed substantially more in abundance in the P and CP communities, 14, 3.2, and 3.1%, respectively. The Solirubrobacteriaceae and Solirubrobacterales (unclassified) consistently contributed to both models with 14 and 10 nodes in the P and CP, and the U and W networks, respectively, and comprised no less than 20% of the relative recovery from either community.

CN Elevated Soil Respiration

Only the CN resource addition elevated soil respiration 1 week following the nutrient additions. As a result, we observed a 114–234% increase in soil respiration in CN soils compared to all other treatments (one-way ANOVA, $df = 6$, $F = 11.6$, $P < 0.001$, **Figure 5**). The soil treatments exhibited no differences in temperature at the time of sampling (one-way ANOVA, $df = 6$, $F = 1.46$, $P = 0.24$). The mean temperature of all treatments was $7.2^\circ\text{C} \pm 0.32$ SD ($n = 56$).

DISCUSSION

Stoichiometric shifts of C:N:P in the coldest and driest soils on Earth (Virginia and Wall, 1999) alleviated resource limitations, created species-specific bacterial responses, and altered ecosystem processes. Our long-term coupled resource




TABLE 3 | Community network model characteristics for soil bacteria in P-amended and control soils.

Network topographical parameters	U and W		P and CP	
Nodes, edges (#)	113, 1321		75, 433	
Mean path length	2.8		3.5	
Mean degree	23		11.5	
Mean clustering coefficient	0.84		0.74	
Density	0.21		0.16	
Modularity	0.54		0.51	

Network taxonomy (order)	U and W		P and CP	
	Node #	Relative recovery of nodes (%)	Node #	Relative recovery of nodes (%)
Acidobacteriaceae	10	14	7	13
Intrasporangiaceae	2	0.13	0	0
Micrococcaceae	1	0.06	1	3.2
Nocardioideaceae	1	0.10	0	0
Rubrobacteriaceae	4	1.0	3	0.83
Conexibacteraceae	1	0.40	1	0.16
Solirubrobacteriaceae	2	1.3	4	2.0
Solirubrobacteriales Unclassified	8	22	10	18
Chitinophagaceae	5	0.32	4	3.1
Sphingobacteriaceae	0	0	1	0.90
Caldilineaceae	1	0.28	1	1.5
Trueperaceae	1	0.51	3	7.7
Gemmatimonadaceae	1	0.65	1	0.16
Phycisphaerae unclassified	10	2.1	0	0
Sphingomonadaceae	1	0.98	2	1.8
Alcaligenaceae	1	0.12	1	0.14
Oxalobacteraceae	1	0.18	1	0.50
Xanthomonadaceae	2	0.55	0	0
Spartobacteria unclassified	4	7.3	4	14


For treatment abbreviations see **Table 1** and for an explanation of network topographical parameters see network community modeling in the results section. For network taxonomy, node # is the total number of nodes or OTUs in a given order for a network. The relative recovery of nodes is the summation of mean relative recoveries of the nodes within the order from P and CP or U and W communities.

additions dramatically altered soil C:N:P leading to increases in inorganic N and P availability, but only a slight increase in soil organic C content, which was presumably consumed by bacteria. As hypothesized, C, N, CN, and CP additions created unique communities, relative to untreated soils, with CN and N having the most pronounced effect on bacterial species responses. We found that the alleviation of a C and N co-limitation facilitated the dominance of an *Arthrobacter species* (family, Micrococcaceae) that ultimately elevated soil respiration, and that shifts in C:N ratios may remove nutrient constraints on bacteria enhancing species richness and diversity. Contrary to our hypothesis, the addition of P, even to our relatively P-rich soils (Aislabie et al., 2006; Barrett et al., 2007), helped create unique communities for all single and coupled resource additions.

Colimitation of CN Facilitates Species Dominance and Enhanced Respiration

The colimitation of organic C, N, and/or P are common in marine and freshwater systems where the abundance of

photoautotrophs to organoheterotrophs is often influenced by two or more nutrients (Bouvy et al., 2004; Howarth and Marino, 2006; Cherif and Loreau, 2007; Saito et al., 2008). In soils, co-limitation exists but is harder to identify due to the high levels of bacterial diversity and the wide variety of resource substrates for species to exploit, from extremely labile C substrates to recalcitrant soil organic matter. Over the 6 years of the present study, as CN limitations were eased, (e.g., C:N decreased and N:P increased in CN amended soils), an *Arthrobacter species* (family = Micrococcaceae, Actinobacteria) went from being rare ($0.06\% \pm 0.05$) to dominant ($47\% \pm 5.6$). *Arthrobacter species* are common psychrotrophs found in Adelie penguin guano (Zdanowski et al., 2004) and Antarctic epilithic lichens (Selbmann et al., 2010) seeming to capitalize on the localized nutrient-rich penguin feces in the otherwise nutrient poor landscape. *Arthrobacter* strains can respond quickly to changes in nutrient conditions by breaking dormancy and growing within an hour of the removal of starvation stress (Solyanikova et al., 2017). Compared to temperate *Arthrobacter species*, Antarctic



Arthrobacter possess lower metabolic versatility (Dsouza et al., 2015) but similar genes to many psychrophilic/psychrotolerant species (e.g., cold active hydrolytic enzymes; sigma factors; signal transduction pathways; carotenoid biosynthesis pathway; and genes induced by cold-shock, oxidative, and osmotic stresses (Dsouza et al., 2015; Singh et al., 2016). Further, our unclassified *Arthrobacter* OTU falls within a genus whose members readily decompose almost any algal and cyanobacterial bioproducts, from cyanotoxins (Lawton et al., 2011) to cellobiose, the final derivative of cellulose utilization (Schellenberger et al., 2011). Thus, the *Arthrobacter* we recovered is likely a well-adapted psychrophile poised to exploit common bacterial and algal derived C sources when N is available.

The functional consequences of *Arthrobacter* dominance were easily distinguishable. Often the functional consequences of soil bacterial community change is exceptionally difficult due to discern due to levels of functional redundancy (Yin et al., 2000; Miki et al., 2014) and large fractions of bacterial diversity being dormant or metabolically inactive at any given time (Lennon and Jones, 2011). In contrast, only in soils where *Arthrobacter* bacteria achieved dominance did soil respiration dramatically increase (114–234%). Our results are consistent with the findings of Hopkins et al. (2006) who showed that the addition of glucose, glycine, and ammonium stimulated the mineralization of lacustrine detritus and soil organic matter across different geomorphically defined landscapes in Garwood Valley, Antarctica. Even though the link between our dominant species and respiration is implied rather than explicit, dominant or abundant bacteria often contribute proportionally to universal soil processes such as respiration (Pedros-Alio, 2012). Thus, *Arthrobacter* most likely exploited mannitol and ammonium or nitrate to a greater extent than other species to become more metabolically active. Bacterial competition for essential resources may loosely be classified into two competition categories, scramblers and contesters

(Hibbing et al., 2010). Scramble competition or exploitation competition involves rapid utilization of resources without directly interacting with other bacteria, while contest competition or interference competition involves direct antagonistic interactions between competitors. While both scramblers and contesters occur in most soils, the effects of competition for resources on bacterial taxa are often only implied (Zhou et al., 2002; Freilich et al., 2011) and potential interactions among limiting nutrients are often neglected (Fanin et al., 2015). In our CN-enriched soils, *Arthrobacter* is most likely a scrambler, better suited to capitalize on emerging resources. Its rise to dominance resulted in a decline in bacterial evenness and diversity while allowing for the persistence of rare taxa, as evidenced by similar species richness levels exhibited among the different resource treatments.

Inorganic N Opened New Bacterial Niches for Rare Taxa

The removal of ammonium limitations opened new niches for once rare taxa to exploit. With the immense elevation of ammonium levels following N additions, bacterial richness increased upward of 48% in comparison to soils that receive only C or P additions. Higher levels of ammonium increased bacterial richness directly and indirectly by potentially stimulating nitrifying bacteria relying on ammonium and nitrite. We found that ammonium additions enhanced the abundance of two nitrite oxidizing *Nitrospira* species from the family Nitrosomonadaceae. Antarctic *Nitrospira* species may also contain amoA sequences (Magalhaes et al., 2014) and participate in complete nitrification (Daims et al., 2015). Our findings are consistent with the suggestion that the extreme abiotic severity of the McMurdo Dry Valley soil habitat drives the presence of ammonia oxidizing bacteria (AOB) like *Nitrospira* (Geyer et al., 2014; Magalhaes et al., 2014). Only after the soils became considerably less harsh with N additions were AOB found in soils. Presumably the increase in nitrite triggered certain species to increase in abundance. For example, six Gemmatimonadaceae species with the ability to reduce nitrite to nitric oxide via the NirK gene (clade II) (Decleyre et al., 2016) increased in abundance at least 1.8-fold under N additions relative to all other treatments. Also, four Rhodobacteraceae taxa with a high affinity for ammonium, characterized by high transcript abundance for ammonium transporters (Pfreundt et al., 2016), increased in abundance as N limitations were lifted. Many of these taxa responding to ammonium and possibly nitrite were rare (Sogin et al., 2006) with an abundance in untreated soils <0.1% of the total recovery (Aanderud et al., 2015). Thus, the enhanced availability of reduced forms of N following N-amendments stimulates microbial biodiversity in Antarctic soils.

CP and P Deconstructed Species Assemblages

Lower soil C:P following P additions disrupted community co-occurrence patterns and facilitated new nutrient-related

interactions among taxa. Even an increase in soil P, a nutrient that was not predicted to be limiting based on the initial soil C:N:P ratios, influenced species interactions. Excess soil P disrupted potential interactions among community assemblages, as evidenced by more than 50% of the network (i.e., significant species and interactions) disappearing and 14 taxa from three families (i.e., Phycisphaerae unclassified, Intrasporangiaceae, and Xanthomonadaceae) vanishing from the co-occurrence network. However, as P limitations were alleviated a more diffuse and less connected network potentially centered on P availability emerged. For example, two new Trueperaceae species (*Deinococcus*) were incorporated into the model and the collective abundance of the three Trueperaceae species was 13-times higher in P amended soils. Members of the Trueperaceae family are remarkably resistant to ionizing radiation and able to grow under multiple extreme conditions, including alkaline, moderately saline, and high temperature environments (Ivanova et al., 2011). *Deinococcus-Thermus* taxa in general possess a remarkable number of genes encoding for catabolic enzymes including phosphatases (Daly et al., 2010), suggesting that the access to P is potentially linked to radiation resistance and/or helps boost the survival of Trueperaceae in extreme soils. Additionally, multiple shared taxa between our two networks [i.e., Spartobacteria unclassified (Verrucomicrobia), Micrococcaceae (Actinobacteria), and Chitinophagaceae (Bacteroidetes)] increased in abundance under P additions. The abundance of taxa within these families may track the availability of P in soils and water (Johnson et al., 2017; Wang et al., 2017). Even with P additions communities seemed to remain partially reliant on Solirubrobacteriaceae taxa. Members of the Solirubrobacteriaceae family may enhance the weathering of volcanic rocks, which are common in soils occurring on Ross Sea Till (Cockell et al., 2013). Despite P additions, Solirubrobacteriaceae taxa were integral to all models constituting upward of 10% of species and 20% of the recovery suggesting that mineral weathering is essential to enhance micronutrient availability under both high and low nutrient conditions. As soils transition from a nutrient-poor to a nutrient-rich state, the excess soil P potentially disrupted interactions among bacterial taxa. For example, the addition of water and organic matter in McMurdo Dry Valley soils caused certain bacteria (members of the and Actinobacteria, Proteobacteria, and Firmicutes) to become active demonstrating a potential taxonomical shift from species adapted to dry oligotrophic to moist copiotrophic conditions (Buelow et al., 2016). Thus, future climate-driven changes that ameliorate the current stoichiometric imbalances of the dry valley soils (Nielsen and Ball, 2014; Gooseff et al., 2017), may deconstruct current bacterial communities and reorganizing them into communities dominated by more copiotrophic taxa. Co-occurrence networks do provide insights into potential interactions among taxa within a community (Freilich et al., 2011; Freedman and Zak, 2015), but to fully understand interactions among bacteria a more direct approach is needed.

Annual Water Additions Failed to Elicit Bacterial Response

Water is necessary for imbalances in stoichiometric nutrient ratios to influence communities; however, our one-time water addition alone was not enough to create lasting effects on bacteria community structure. Frequent water additions do influence bacterial activity across Dry Valleys (Schwartz et al., 2014; Buelow et al., 2016) where soil moisture is ephemeral and extremely patchy. Surface hydrogeological features such as water tracks in soils (Levy et al., 2014), lateral margins of stream and lake margins (Zeglin et al., 2011), and discontinuous patches of soils often form in the same location due to wind sheltering and microtopography (Gooseff et al., 2003) enhance bacterial metabolic activity and alter species distributions.

CONCLUSION

Stoichiometric additions of C, N, and P reduced resource limitations, created species-specific bacterial responses, and in one case altered a fundamental ecosystem process. The most dramatic effects of changes in ecosystem stoichiometry occurring around C and N additions in our initially N limited soils. C as mannitol and N as equal molar concentrations ammonium and nitrate induced an almost twofold reduction in C:N ratios; caused bacterial evenness and diversity to decline; allowed one rare Micrococcaceae, an *Arthrobacter* species, to dominate community abundance; and elevated soil respiration by 136% compared to untreated soils. N additions alone also reduced C:N ratios, and in contrast to CN-additions, increased species richness and diversity by at least 48 and 13%, respectively, compared to soils receiving a single resource as C or P, and enhanced the abundance of rare taxa dependent on N for metabolism and growth. The addition of P to levels well below the C:P ratio necessary for balanced microbial growth also influenced soil microbial communities. Based on community co-occurrence networks, lower C:P ratios in soils following P and CP additions reduced the number of taxa interacting with one another by 51% and the number of interactions among taxa by 73% relative to untreated and watered soils. Our results suggest that the alleviation of C and N co-limitation facilitated the dominance of single species ultimately altering ecosystem processes; the reduced forms of inorganic N open multiple niches for bacteria to exploit, and excess soil P disrupted interactions within communities.

AUTHOR CONTRIBUTIONS

ZA, SS, BB, DW, JB, RV, and BA conducted the experiments. ZA, SS, BB, DW, JB, MM, NG, RV, AB, and BA analyzed and interpreted the data. ZA, SS, DW, JB, NG, MM, RV, AB, and BA helped to write and review the manuscript. ZA agrees to be accountable for

all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

FUNDING

Our research was supported by the National Science Foundation Office of Polar Programs grants to the McMurdo Long-Term Ecological Research Program ANT-0423595 and OPP-1115245.

REFERENCES

- Aanderud, Z. T., Jones, S. E., Fierer, N., and Lennon, J. T. (2015). Resuscitation of the rare biosphere contributes to pulses of ecosystem activity. *Front. Microbiol.* 6:24. doi: 10.3389/fmicb.2015.00024
- Aanderud, Z. T., Jones, S. E., Schoolmaster, D. R., Fierer, N., and Lennon, J. T. (2013). Sensitivity of soil respiration and microbial communities to altered snowfall. *Soil Biol. Biochem.* 57, 217–227. doi: 10.1016/j.soilbio.2012.07.022
- Adams, B. J., Bardgett, R. D., Ayres, E., Wall, D. H., Aislabie, J., Bamforth, S., et al. (2006). Diversity and distribution of Victoria Land biota. *Soil Biol. Biochem.* 38, 3003–3018. doi: 10.1016/j.soilbio.2006.04.030
- Aislabie, J. M., Chhour, K. L., Saul, D. J., Miyauchi, S., Ayton, J., Paetzold, R. F., et al. (2006). Dominant bacteria in soils of Marble Point and Wright Valley, Victoria Land, Antarctica. *Soil Biol. Biochem.* 38, 3041–3056. doi: 10.1016/j.soilbio.2006.02.018
- Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* 26, 32–46.
- Aneja, M. K., Sharma, S., Fleischmann, F., Stich, S., Heller, W., Bahnweg, G., et al. (2006). Microbial colonization of beech and spruce litter - Influence of decomposition site and plant litter species on the diversity of microbial community. *Microb. Ecol.* 52, 127–135. doi: 10.1007/s00248-006-9006-3
- Austin, A. T., and Vitousek, P. M. (2012). Introduction to a virtual special issue on ecological stoichiometry and global change. *New Phytol.* 196, 649–651. doi: 10.1111/j.1469-8137.2012.04376
- Baldrian, P., Kolarik, M., Stursova, M., Kopecky, J., Valaskova, V., Vetrovsky, T., et al. (2012). Active and total microbial communities in forest soil are largely different and highly stratified during decomposition. *ISME J.* 6, 248–258. doi: 10.1038/ismej.2011.95
- Ball, B. A., Adams, B. J., Barrett, J. E., Wall, D. H., and Virginia, R. A. (2018). Soil biological responses to C, N and P fertilization in a polar desert of Antarctica. *Soil Biol. Biochem.* 122, 7–18. doi: 10.1016/j.soilbio.2018.03.025
- Barberan, A., Bates, S. T., Casamayor, E. O., and Fierer, N. (2012). Using network analysis to explore co-occurrence patterns in soil microbial communities. *ISME J.* 6, 343–351. doi: 10.1038/ismej.2011.119
- Barrett, J. E., Virginia, R. A., Hopkins, D. W., Aislabie, J., Bargagli, R., Bockheim, J. G., et al. (2006). Terrestrial ecosystem processes of Victoria Land, Antarctica. *Soil Biol. Biochem.* 38, 3019–3034. doi: 10.1016/j.soilbio.2006.04.041
- Barrett, J. E., Virginia, R. A., Lyons, W. B., McKnight, D. M., Priscu, J. C., Doran, P. T., et al. (2007). Biogeochemical stoichiometry of Antarctic Dry Valley ecosystems. *J. Geophys. Res.* 112:G01010. doi: 10.1029/2005JG000141
- Bastian, M., Heymann, S., and Jacomy, M. (2009). “Gephi: an open source software for exploring and manipulating networks,” in *Proceedings of the 3rd International AAAI Conference on Weblogs and Social Media*. Available at: <http://www.aaai.org/ocs/index.php/ICWSM/09/paper/view/154>
- Bedford, B. L., Walbridge, M. R., and Aldous, A. (1999). Patterns in nutrient availability and plant diversity of temperate North American wetlands. *Ecology* 80, 2151–2169. doi: 10.1890/0012-9658(1999)080[2151:PINAAP]2.0.CO;2
- Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., et al. (2010). Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecol. Appl.* 20, 30–59. doi: 10.1890/08-1140.1
- Bockheim, J. G., and McLeod, M. (2008). Soil distribution in the McMurdo Dry Valleys, Antarctica. *Geoderma* 144, 43–49. doi: 10.1016/j.geoderma.2007.10.015
- Bockheim, J. G., Prentice, M. L., and McLeod, M. (2008). Distribution of glacial deposits, soils, and permafrost in Taylor Valley, Antarctica. *Arctic Antarct. Alp. Res.* 40, 279–286. doi: 10.1657/1523-0430(06-057)[BOCKHEIM]2.0.CO;2
- Bouvy, M., Troussellier, M., Got, P., and Arfi, R. (2004). Bacterioplankton responses to bottom-up and top-down controls in a West African reservoir (Selingue, Mali). *Aquat. Microb. Ecol.* 34, 301–307. doi: 10.3354/ame035301
- Buelow, H. N., Winter, A. S., Van Horn, D. J., Barrett, J. E., Gooseff, M. N., Schwartz, E., et al. (2016). Microbial community responses to increased water and organic matter in the arid soils of the McMurdo Dry Valleys, Antarctica. *Front. Microbiol.* 7:1040. doi: 10.3389/fmicb.2016.01040
- Burkins, M. B., Virginia, R. A., Chamberlain, C. P., and Wall, D. H. (2000). Origin and distribution of soil organic matter in Taylor Valley, Antarctica. *Ecology* 81, 2377–2391. doi: 10.1890/0012-9658
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P. J., et al. (2010). Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc. Natl. Acad. Sci. U.S.A.* 108, 4516–4522. doi: 10.1073/pnas.1000080107
- Cary, S. C., McDonald, I. R., Barrett, J. E., and Cowan, D. A. (2010). On the rocks: the microbiology of Antarctic Dry Valley soils. *Nat. Rev. Microbiol.* 8, 129–138. doi: 10.1038/nrmicro2281
- Cherif, M., and Loreau, M. (2007). Stoichiometric constraints on resource use, competitive interactions, and elemental cycling in microbial decomposers. *Am. Nat.* 169, 709–724. doi: 10.1086/516844
- Clark, C. M., and Tilman, D. (2008). Loss of plant species after chronic low-level nitrogen deposition to prairie grasslands. *Nature* 451, 712–715. doi: 10.1038/nature06503
- Cleveland, C. C., and Liptzin, D. (2007). C : N : P stoichiometry in soil: is there a “Redfield ratio” for the microbial biomass? *Biogeochemistry* 85, 235–252. doi: 10.1007/s10533-007-9132-0
- Cockell, C. S., Kelly, L. C., and Marteinsson, V. (2013). Actinobacteria—an ancient phylum active in volcanic rock weathering. *Geomicrobiol. J.* 30, 706–720. doi: 10.1080/01490451.2012.758196
- Cole, J. R., Wang, Q., Cardenas, E., Fish, J., Chai, B., Farris, R. J., et al. (2009). The ribosomal database project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res.* 37, D141–D145. doi: 10.1093/nar/gkn879
- Cowan, D. A., Makhalyane, T. P., Dennis, P. G., and Hopkins, D. W. (2014). Microbial ecology and biogeochemistry of continental Antarctic soils. *Front. Microbiol.* 5:154. doi: 10.3389/fmicb.2014.00154
- Csardi, G., and Nepusz, T. (2006). The igraph software package for complex network research. *InterJournal Complex Syst.* 1695, 1–9.
- Daims, H., Lebedeva, E. V., Pjevac, P., Han, P., Herbold, C., Albertsen, M., et al. (2015). Complete nitrification by Nitrospira bacteria. *Nature* 528, 504–509. doi: 10.1038/nature16461
- Daly, M. J., Gaidamakova, E. K., Matrosova, V. Y., Kiang, J. G., Fukumoto, R., Lee, D. Y., et al. (2010). Small-molecule antioxidant proteome-shields in *Deinococcus radiodurans*. *PLoS One* 5:e12570. doi: 10.1371/journal.pone.0012570
- Decleyre, H., Heylen, K., Tytgat, B., and Willems, A. (2016). Highly diverse *nirK* genes comprise two major clades that harbour ammonium-producing denitrifiers. *BMC Genomics* 17:155. doi: 10.1186/s12864-016-2465-0
- Doran, P. T., McKay, C. P., Clow, G. D., Dana, G. L., Fountain, A. G. A., Nylan, T., et al. (2002). Valley floor climate observations from the

ACKNOWLEDGMENTS

We thank Ashley Shaw providing valuable comments that improved the paper.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2018.01401/full#supplementary-material>

- McMurdo Dry Valleys, Antarctica, 1986-2000. *J. Geophys. Res.* 107, 4772–4784. doi: 10.1029/2001JD002045
- Dsouza, M., Taylor, M. W., Turner, S. J., and Aislabie, J. (2015). Genomic and phenotypic insights into the ecology of *Arthrobacter* from Antarctic soils. *BMC Genomics* 16:18. doi: 10.1186/s12864-015-1220-2
- Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., and Knight, R. (2011). UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27, 2194–2200. doi: 10.1093/bioinformatics/btr381
- Elser, J. J., and Hamilton, A. (2007). Stoichiometry and the new biology - The future is now. *PLoS Biol.* 5:1403–1405. doi: 10.1371/journal.pbio.0050181
- Elser, J. J., Sterner, R. W., Gorokhova, E., Fagan, W. F., Markow, T. A., and Cotner, J. B. (2000). Biological stoichiometry from genes to ecosystems. *Ecol. Lett.* 3, 540–550. doi: 10.1046/j.1461-0248.2000.00185.x
- Fanin, N., Hattenschwiler, S., Schimann, H., and Fromin, N. (2015). Interactive effects of C, N and P fertilization on soil microbial community structure and function in an Amazonian rain forest. *Funct. Ecol.* 29, 140–150. doi: 10.1111/1365-2435.12329
- Fierer, N., Leff, J. W., Adams, B. J., Nielsen, U. N., Bates, S. T., Lauber, C. L., et al. (2012). Cross-biome metagenomic analyses of soil microbial communities and their functional attributes. *Proc. Natl. Acad. Sci. U.S.A.* 109, 21390–21395. doi: 10.1073/pnas.1215210110
- Fierer, N., Strickland, M. S., Liptzin, D., Bradford, M. A., and Cleveland, C. C. (2009). Global patterns in belowground communities. *Ecol. Lett.* 12, 1238–1249. doi: 10.1111/j.1461-0248.2009.01360
- Filosi, M., Visintainer, R., and Albanese, D. (2017). *Minerva: Maximal Information-Based Nonparametric Exploration for Variable Analysis. R Package Version 1.4.7*. Available at: <https://CRAN.R-project.org/package=minerva>
- Fountain, A. G., Lyons, W. B., Burkins, M. B., Dana, G. L., Doran, P. T., Lewis, K. J., et al. (1999). Physical controls on the Taylor Valley Ecosystem, Antarctica. *BioScience* 49, 961–971. doi: 10.1525/bisi.1999.49.12.961
- Freedman, Z. B., Upchurch, R. A., and Zak, D. R. (2016). Microbial potential for ecosystem N loss is increased by experimental N deposition. *PLoS One* 11:e0164531. doi: 10.1371/journal.pone.0164531
- Freedman, Z. B., and Zak, D. R. (2015). Atmospheric N deposition alters connectance, but not functional potential among saprotrophic bacterial communities. *Mol. Ecol.* 24, 3170–3180. doi: 10.1111/mec.13224
- Freilich, S., Zarecki, R., Eilam, O., Segal, E. S., Henry, C. S., Kupiec, M., et al. (2011). Competitive and cooperative metabolic interactions in bacterial communities. *Nat. Commun.* 2:589. doi: 10.1038/ncomms1597
- Geyer, K. M., Altrichter, A. E., Takacs-Vesbach, C. D., Van Horn, D. J., Gooseff, M. N., and Barrett, J. E. (2014). Bacterial community composition of divergent soil habitats in a polar desert. *FEMS Microbiol. Ecol.* 89, 490–494. doi: 10.1111/1574-6941.12306
- Gooseff, M. N., Barrett, J. E., Adams, B. J., Doran, P. T., Fountain, A. G., and Lyons, W. B. (2017). Decadal ecosystem response to an anomalous melt season in a polar desert in Antarctica. *Nat. Ecol. Evol.* 1, 1334–1338. doi: 10.1038/s41559-017-0253-0
- Gooseff, M. N., Barrett, J. E., Doran, P. T., Fountain, A. G., Lyons, W. B., Parsons, A. N., et al. (2003). Snow-patch influence on soil biogeochemical processes and invertebrate distribution in the McMurdo Dry Valleys, Antarctica. *Arctic Antarct. Alp. Res.* 35, 91–99. doi: 10.1657/1523-0430(2003)035[0091:SPIOSB]2.0.CO;2
- Hartman, W. H., and Richardson, C. J. (2013). Differential nutrient limitation of soil microbial biomass and metabolic quotients (qCO₂): Is there a biological stoichiometry of soil microbes? *PLoS One* 8:e57127. doi: 10.1371/journal.pone.0057127
- Hecky, R. E., Campbell, P., and Hendzel, L. L. (1993). The stoichiometry of carbon, nitrogen, and phosphorus in particulate matter of lakes and oceans. *Limnol. Oceanogr.* 38, 709–724. doi: 10.4319/lo.1993.38.4.0709
- Heindel, R. C., Spickard, A. M., and Virginia, R. A. (2017). Landscape-scale soil phosphorus variability in the McMurdo Dry Valleys, Antarctica. *Antarct. Sci.* 29, 252–263. doi: 10.1017/S0954102016000742
- Hernandez, D. L., and Hobbie, S. E. (2010). The effects of substrate composition, quantity, and diversity on microbial activity. *Plant Soil* 335, 397–411. doi: 10.1007/s11104-010-0428-9
- Hibbing, M. E., Fuqua, C., Parsek, M. R., and Peterson, S. B. (2010). Bacterial competition: surviving and thriving in the microbial jungle. *Nat. Rev. Microbiol.* 8, 15–25. doi: 10.1038/nrmicro2259
- Hopkins, D. W., Sparrow, A. D., Elberling, B., Gregorich, E. G., Novis, P. M., Greenfield, L. G., et al. (2006). Carbon, nitrogen and temperature controls on microbial activity in soils from an Antarctic dry valley. *Soil Biol. Biochem.* 38, 3130–3140. doi: 10.1016/j.soilbio.2006.01.012
- Howarth, R. W., and Marino, R. (2006). Nitrogen as the limiting nutrient for eutrophication in coastal marine ecosystems: Evolving views over three decades. *Limnol. Oceanogr.* 51, 364–376. doi: 10.4319/lo.2006.51.1_part_2.0364
- Ivanova, N., Rohde, C., Munk, C., Nolan, M., Lucas, S., Del Rio, T. G., et al. (2011). Complete genome sequence of *Truepera radiovictrix* type strain (RQ-24(T)). *Stand. Genomic Sci.* 4, 91–99. doi: 10.4056/signs.1563919
- Jobbagy, E. G., and Jackson, R. B. (2000). The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecol. Appl.* 10, 423–436. doi: 10.1890/1051-0761(2000)010[0423:TVDOSO]2.0.CO;2
- Johnson, W. H., Douglas, M. R., Lewis, J. A., Stuecker, T. N., Carbonero, F. G., Austin, B. J., et al. (2017). Do biofilm communities respond to the chemical signatures of fracking? A test involving streams in North-central Arkansas. *BMC Microbiol.* 17:12. doi: 10.1186/s12866-017-0926-5
- Keiblinger, K. M., Hall, E. K., Wanek, W., Szukics, U., Hammerle, I., Ellersdorfer, G., et al. (2010). The effect of resource quantity and resource stoichiometry on microbial carbon-use-efficiency. *FEMS Microbiol. Ecol.* 73, 430–440. doi: 10.1111/j.1574-6941.2010.00912
- Lawton, L. A., Welgamage, A., Manage, P. M., and Edwards, C. (2011). Novel bacterial strains for the removal of microcystins from drinking water. *Water Sci. Technol.* 63, 1137–1142. doi: 10.2166/wst.2011.352
- Lee, C. K., Lozier, J. D., Barbier, B. A., Mills, N. J., Bottos, E. M., and Cary, S. C. (2011). The Inter-Valley Soil Comparative Survey: the ecology of Dry Valley edaphic microbial communities. *ISME J.* 6, 1046–1057. doi: 10.1038/ismej.2011.170
- Lennon, J. T., and Jones, S. E. (2011). Microbial seed banks: the ecological and evolutionary implications of dormancy. *Nat. Rev. Microbiol.* 9, 119–130. doi: 10.1038/nrmicro2504
- Levy, J. S., Fountain, A. G., Gooseff, M. N., Barrett, J. E., Vantreesse, R., Welch, K. A., et al. (2014). Water track modification of soil ecosystems in the Lake Hoare basin, Taylor Valley, Antarctica. *Antarct. Sci.* 26, 153–162. doi: 10.1017/S0954102013000045
- Lozupone, C. A., and Knight, R. (2007). Global patterns in bacterial diversity. *Proc. Natl. Acad. Sci. U.S.A.* 104, 11436–11440. doi: 10.1073/pnas.0611525104
- Magalhaes, C. M., Machado, A., Frank-Fahle, B., Lee, C. K., and Car, S. C. (2014). The ecological dichotomy of ammonia-oxidizing archaea and bacteria in the hyper-arid soils of the Antarctic Dry Valleys. *Front. Microbiol.* 5:515. doi: 10.3389/fmicb.2014.00515
- Marschner, P., Kandeler, E., and Marschner, B. (2003). Structure and function of the soil microbial community in a long-term fertilizer experiment. *Soil Biol. Biochem.* 35, 453–461. doi: 10.1016/S0038-0717(02)00297-3
- Michalski, G., Bockheim, J. G., Kendall, C., and Thiemens, M. (2005). Isotopic composition of Antarctic Dry Valley nitrate: implications for NO_y sources and cycling in Antarctica. *Geophys. Res. Lett.* 32:L13817. doi: 10.1029/2004GL022121
- Miki, T., Yokokawa, T., and Matsui, K. (2014). Biodiversity and multifunctionality in a microbial community: a novel theoretical approach to quantify functional redundancy. *Proc. Biol. Sci.* 281:20132498. doi: 10.1098/rspb.2013.2498
- Mooshammer, M., Wanek, W., Schnecker, J., Wild, B., Leitner, S., Hofhansl, F., et al. (2012). Stoichiometric controls of nitrogen and phosphorus cycling in decomposing beech leaf litter. *Ecology* 93, 770–782. doi: 10.1890/11-0721.1
- Muscarella, M. E., Bird, K. C., Larsen, M. L., Placella, S. A., and Lennon, J. T. (2014). Phosphorus resource heterogeneity in microbial food webs. *Aquat. Microb. Ecol.* 73, 259–272. doi: 10.3354/ame01722
- Nielsen, U. N., and Ball, B. A. (2014). Impacts of altered precipitation regimes on soil communities and biogeochemistry in arid and semi-arid ecosystems. *Glob. Change Biol.* 21, 1407–1421. doi: 10.1111/gcb.12789
- Oksanen, J., Guillaume, F., Kindt, B., Kindt, R., Legendre, P., Minchin, P. R., et al. (2013). *Vegan: Community Ecology Package*. Available at: <https://cran.r-project.org/web/packages/vegan/index.html>
- Orwin, K. H., Wardle, D. A., and Greenfield, L. G. (2006). Ecological consequences of carbon substrate identity and diversity in a laboratory study. *Ecology* 87, 580–593. doi: 10.1890/05-0383
- Pedros-Alio, C. (2012). The rare bacterial biosphere. *Ann. Rev. Mar. Sci.* 4, 449–466. doi: 10.1146/annurev-marine-120710-100948

- Pfreundt, U., Spungin, D., Bonnet, S., Berman-Frank, I., and Hess, W. R. (2016). Global analysis of gene expression dynamics within the marine microbial community during the VAHINE mesocosm experiment in the southwest Pacific. *Biogeosciences* 13, 4135–4149. doi: 10.5194/bg-13-4135-2016
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W. G., Peplies, J., et al. (2007). SILVA: A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res.* 35, 7188–7196. doi: 10.1093/nar/gkm864
- Quince, C., Lanzen, A., Davenport, R. J., and Turnbaugh, P. J. (2011). Removing noise from pyrosequenced amplicons. *BMC Bioinformatics* 12, doi: 10.1186/1471-2105-12-38
- R Development Core Team (2013). *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Ramirez, K. S., Lauber, C. L., Knight, R., Bradford, M. A., and Fierer, N. (2010). Consistent effects of nitrogen fertilization on soil bacterial communities in contrasting systems. *Ecology* 91, 3463–3470. doi: 10.1890/10-0426.1
- Redfield, A. C. (1958). The biological control of chemical factors in the environment. *Am. Sci.* 46, 205–221.
- Reiners, W. A. (1986). Complementary models for ecosystems. *Am. Nat.* 46, 205–221. doi: 10.1086/284467
- Ren, C. J., Zhao, F. Z., Kang, D., Yang, G. H., Han, X., Tong, H. G., et al. (2016). Linkages of C:N:P stoichiometry and bacterial community in soil following afforestation of former farmland. *Forest Ecol. Manag.* 376, 59–66. doi: 10.1016/j.foreco.2016.06.004
- Reshef, D. N., Reshef, H. K., Finucane, S. R., Grossman, G., McVean, P. J., Turnbaugh, P. J., et al. (2011). Detecting novel associations in large data sets. *Science* 334, 1518–1524. doi: 10.1126/science.1205438
- Saito, M. A., Goepfert, T. J., and Ritt, J. T. (2008). Some thoughts on the concept of colimitation: three definitions and the importance of bioavailability. *Limnol. Oceanogr.* 53, 276–290. doi: 10.4319/lo.2008.53.1.0276
- Schellenberger, S., Drake, H. L., and Kolb, S. (2011). Functionally redundant cellulose-degrading soil bacteria respond differentially to oxygen. *Appl. Environ. Microbiol.* 77, 6043–6048. doi: 10.1128/AEM.00564-11
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al. (2009). Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537–7541. doi: 10.1128/AEM.01541-09
- Schwartz, E., Van Horn, D. J., Buelow, H. N., Okie, J. G., Gooseff, M. N., Barrett, J. E., et al. (2014). Characterization of growing bacterial populations in McMurdo Dry Valley soils through stable isotope probing with ^{18}O -water. *FEMS Microb. Ecol.* 89, 415–425. doi: 10.1111/1574-6941.12349
- Selbmann, L., Zucconi, L., Ruisi, S., Grube, M., Cardinale, M., and Onofri, S. (2010). Culturable bacteria associated with Antarctic lichens: affiliation and psychrotolerance. *Polar Biol.* 33, 71–83. doi: 10.1007/s00300-009-0686-2
- Singh, R. N., Gaba, S., Yadav, A. N., Gaur, P., Gulati, S., Kaushik, R., et al. (2016). First high quality draft genome sequence of a plant growth promoting and cold active enzyme producing psychrotrophic *Arthrobacter agilis* strain L77. *Stand. Genomic Sci.* 11:54. doi: 10.1186/s40793-016-0176-4
- Sogin, M. L., Morrison, H. G., Huber, J. A., Welch, M. D., Huse, S. M., Neal, P. R., et al. (2006). Microbial diversity in the deep sea and the unexplored "rare biosphere." *Proc. Natl. Acad. Sci. U.S.A.* 32, 12115–12120. doi: 10.1073/pnas.060512710
- Solyanikova, I. P., Suzina, N. E., Egozarian, N. S., Polivtseva, V. N., Prisyazhnaya, N. V., El-Registan, G. I., et al. (2017). The response of soil *Arthrobacter agilis* lush13 to changing conditions: transition between vegetative and dormant state. *J. Environ. Sci. Health B* 52, 745–751. doi: 10.1080/03601234.2017.1356665
- Stark, J. M., and Firestone, M. K. (1995). Mechanisms for soil moisture effects on the activity of nitrifying bacteria. *Appl. Environ. Microbiol.* 61, 218–221.
- Tian, H. Q., Chen, G. S., Zhang, C., Melillo, J. M., and Hall, C. A. S. (2010). Pattern and variation of C:N:P ratios in China's soils: a synthesis of observational data. *Biogeochemistry* 98, 139–151. doi: 10.1007/s10533-009-9382-0
- Tischer, A., Werisch, M., Dobbelin, F., Camenzind, T., Rillig, M. C., Potthast, K., et al. (2015). Above- and belowground linkages of a nitrogen and phosphorus co-limited tropicalmountain pasture system-responses to nutrient enrichment. *Plant Soil* 391, 333–352. doi: 10.1007/s11104-015-2431-7
- Virginia, R. A., and Wall, D. H. (1999). How soils structure communities in the Antarctic dry valleys. *Bioscience* 49, 973–983. doi: 10.1525/bisi.1999.49.12.973
- Wang, R., Sun, Q. Q., Wang, Y., Liu, Q. F., Du, L. L., Zhao, M., et al. (2017). Temperature sensitivity of soil respiration: synthetic effects of nitrogen and phosphorus fertilization on Chinese Loess Plateau. *Sci. Total Environ.* 574, 1665–1673. doi: 10.1016/j.scitotenv.2016.09.001
- Warnes, G. R., Bolker, B., Lodewijk, B., Gentleman, R., Liaw, W. H. A., Lumley, T., et al. (2014). *gplots: Various R Programming Tools for Plotting Data Package*. Available at: <https://cran.r-project.org/package=gplots>
- Xu, X. F., Thornton, P. E., and Post, W. M. (2013). A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in terrestrial ecosystems. *Glob. Ecol. Biogeogr.* 22, 737–749. doi: 10.1111/geb.12029
- Yin, B., Crowley, D., Sparovek, G., De Melo, W. J., and Borneman, J. (2000). Bacterial functional redundancy along a soil reclamation gradient. *Appl. Environ. Microbiol.* 66, 4361–4365. doi: 10.1128/AEM.66.10.4361-4365.2000
- Zdanowski, M. K., Weglenski, P., Golik, P., Sasin, J. M., Borsuk, P., Zmuda, M. J., et al. (2004). Bacterial diversity in Adelie penguin, *Pygoscelis adeliae*, guano: molecular and morpho-physiological approaches. *FEMS Microbiol. Ecol.* 50, 163–173. doi: 10.1016/j.femsec.2004.06.012
- Zechmeister-Boltenstern, S., Keiblinger, K. M., Mooshammer, M., Penuelas, J., Richter, A., Sardans, J., et al. (2015). The application of ecological stoichiometry to plant-microbial-soil organic matter transformations. *Ecol. Monograph.* 85, 133–155. doi: 10.1890/14-0777.1
- Zeglin, L. H., Dahm, C. N., Barrett, J. E., Gooseff, M. N., Fitzpatrick, S. K., and Takacs-Vesbach, C. D. (2011). Bacterial community structure along moisture gradients in the Parafluvial sediments of two ephemeral desert streams. *Microb. Ecol.* 61, 543–556. doi: 10.1007/s00248-010-9782-7
- Zhou, J. Z., Xia, B. C., Treves, D. S., Wu, L. Y., Marsh, T. L., O'Neill, R. V., et al. (2002). Spatial and resource factors influencing high microbial diversity in soil. *Appl. Environ. Microbiol.* 68, 326–334. doi: 10.1128/AEM.68.1.326-334.2002

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Aanderud, Saurey, Ball, Wall, Barrett, Muscarella, Griffin, Virginia, Barberán and Adams. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.