



RESEARCH NOTE

A simple formula for enumerating comparisons in trials and network meta-analysis [version 1; peer review: 2 approved]

Farhad Shokraneh , Clive E. Adams

Cochrane Schizophrenia Group, Division of Psychiatry and Clinical Psychology, Institute of Mental Health, School of Medicine, University of Nottingham, Nottingham, NG7 2TU, UK

V1 First published: 09 Jan 2019, 8:38 (<https://doi.org/10.12688/f1000research.17352.1>)Latest published: 03 Apr 2019, 8:38 (<https://doi.org/10.12688/f1000research.17352.2>)**Abstract**

We present use of a simple formula to calculate the number of pairwise comparisons of interventions within a single trial or network meta-analyses. We used the data from our previous network meta-analysis to build a study-based register and enumerated the direct pairwise comparisons from the trials therein. We then compared this with the number of comparisons predicted by use of the formula and finally with the reported number of comparisons (indirect or direct) within the network meta-analysis. A total of 133 trials included in the network generated 163 comparisons (16 unique direct comparisons for 8 interventions). The formula predicted an expected 28 indirect or direct comparisons and this is the number that were indeed reported. The formula produces an accurate enumeration of the potential comparisons within a single trial or network meta-analysis. Its use could help transparency of reporting should a shortfall occur between comparisons actually used and the potential total.


Keywords

Pairwise Comparisons, Study-Based Registers, Clinical Trials, Randomised Controlled Trials, Network Meta-Analysis, Systematic Reviews

Open Peer ReviewReviewer Status  

Invited Reviewers

1 2

REVISED**version 2**published
03 Apr 2019**version 1**published
09 Jan 2019 report report1 **Julie Broderick** , Trinity College Dublin, Dublin, Ireland2 **G. Mustafa Soomro**, St. James Hospital, Portsmouth, UK

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Farhad Shokraneh (Farhad.Shokraneh@nottingham.ac.uk)**Author roles:** **Shokraneh F:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Adams CE:** Investigation, Methodology, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing**Competing interests:** No competing interests were disclosed.**Grant information:** The author(s) declared that no grants were involved in supporting this work.**Copyright:** © 2019 Shokraneh F and Adams CE. This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**How to cite this article:** Shokraneh F and Adams CE. **A simple formula for enumerating comparisons in trials and network meta-analysis [version 1; peer review: 2 approved]** F1000Research 2019, 8:38 (<https://doi.org/10.12688/f1000research.17352.1>)**First published:** 09 Jan 2019, 8:38 (<https://doi.org/10.12688/f1000research.17352.1>)

Introduction

The pairwise comparisons reported within each randomized controlled trial are being documented in study-based registers¹. This lends itself to accurate indexing and enumeration of these comparisons within the studies and then subsequent supply of immediate, highly sensitive and highly specific search results to those wishing to investigate one or more particular comparisons within systematic reviews and meta-analyses or overviews and network meta-analysis^{1,2}.

To gain a perspective on the absolute effectiveness of a treatment it is ideal to compare all the existing medications with placebo and for relative effects with each other in pairwise comparison trials. However, some of pairwise comparisons of the medications have not been tested within trials at all. Finally, even if some of the possible pairwise comparisons have been directly tested within trials not all may be eligible for inclusion in a network meta-analysis³. This leaves a gap between the research has been done and the research that should or could have been undertaken and finding this highlights gaps in the fair testing of treatments⁴.

A two-arm trial will generate one pairwise comparison. A three-arm trial, however, will generate three, and a six-arm study, 15 pairwise comparisons. It is easy to lose track of how many comparisons one study can generate. This is more likely when it comes to the many direct, indirect or mixed comparisons within a network. This paper describes a simple formula for enumerating the possible number of comparisons within a single trial or planned network meta-analysis in advance.

Methods

The formula

The formula below solves this where *n* is the number of arms in a single study or network and *N* is the number of pairwise comparisons:

$$N = (n*(n-1))/2$$

Where *n* > 0;

n is a natural number;

Then every intervention is compared to every other intervention except itself so: *n*(n-1)*;

Because *N* is a bidirectional comparison (*X* vs. *Y* = *Y* vs. *X*) so: *(n*(n-1))/2*;

N is a triangular number.

A visual proof of a network of five interventions and $(5*(5-1))/2=10$ pairwise comparisons is presented in **Figure 1**.

Adding any new intervention to the trial or network will create *n-1* new pairwise comparisons. For example, where there are 6 arms in a trial—or 6 nodes in network meta-analysis—there will be $(6*(6-1))/2=15$ comparisons; adding a new intervention ($6+1=7$) will create $7-1=6$ new pairwise direct comparisons in




Figure 1. Network of five interventions and $(5*(5-1))/2=10$ pairwise comparisons.

an individual trial and 6 direct or indirect comparisons in a network meta-analysis. Although this formula has been used for other purposes such as Metcalfe’s law in telecommunication, its use in the current context is novel.

Testing the formula: working back from existing network meta-analyses

We used the open data⁵ from our previously published network meta-analysis⁶ to re-create and enumerate the comparisons within the network. Using the direct comparisons reported in the trials within the network, we applied the formula and then compared the number of potential or expected comparisons (formula-derived) and the actual or observed number reported within the network analysis.

Results

Number of direct and indirect comparisons

We built a small study-based register based—thus avoiding the pitfall of multiple counting—containing all 133 included studies in our previous network meta-analysis^{6,7}. These trials reported comparisons from 8 interventions. Using our formula, 8 interventions should create 28 unique comparisons: $(8*(8-1))/2=28$ (Figure 2).

Reported comparisons within the trials

We extracted the separate intervention arms from the open data to re-create the direct comparisons from within trials. There trials had either two or three arms so each study could create either two or three comparisons. As a result the 133 studies had 163 comparisons, the majority of which were duplicated. After removing these duplicates, this created 16 unique direct

comparisons with between 1 and 47 studies per comparison for 8 interventions (Table 1). These 16 observed comparisons are 57% of the 28 expected by use of the formula above.

Direct comparisons eligible for network meta-analysis

Among five networks reported in the final paper, the number of comparisons in these five network meta-analyses, however, varies from 6 (for 3 networks) to 11 (for 1 network) and 13 (for 1 network) (Figure 3). As visualized in Figure 3, only 21.42% to 46.42% of comparisons were eligible for pairwise meta-analysis (Table 2).

Comparisons in network meta-analysis plots

From Figure 3 we can calculate that about 42% of comparisons expected though use of the formula have not been tested directly in trials. This is a direct evidence-gap. The number of missing comparisons varies between nine out of 15 in three networks with six interventions, 17 out of 28 in one network with eight interventions, and 15 out of 28 in another network with eight interventions (Figure 3). However, all 28 comparisons expected by use of the formula were utilized and reported within the network meta-analysis. It is possible that some of the comparisons predicted by the formula would have been deemed ineligible—either by adherence to a network review protocol or though *post hoc* exclusions—but this was not the case in this particular review (Figure 4).

Discussion

This formula can be employed when estimating the total number of comparisons (direct and indirect combined) theoretically possible within a proposed network meta-analysis. It would be possible that there would sometimes be a discrepancy between




Figure 2. All the possible unique bidirectional comparisons of 8 ADHD medications. Only 16 out of 28 comparisons have been directly compared in trials (green lines).

Table 1. Direct comparisons extracted from trials and their associated studies.

Comparison	Number of studies	Study tag
Amphetamines vs. Atomoxetine	1	Wigal 2005 (SLI381-404, NCT00506727)
Amphetamines vs. Guanfacine	1	Taylor 2001
Amphetamines* vs. Methylphenidate	6	Coghill 2013 (SPD489-325); Efron 1997; Pilzka 2000; SPD489-405 (NCT01552915); SPD489-406 (NCT01552902); Stein 2011 (NCT00393042)
Amphetamines vs. Modafinil	1	Taylor 2000
Amphetamines vs. Placebo	21	Adler 2008b (NRP104-303, NCT00334880); Adler 2013 (SPD489-403, NCT01101022); Biederman 2002 (SLI 381-301); Biederman 2007 (NRP104-301, NCT00248092); Biederman 2012 (2008P000971, NCT00801229); Coghill 2013 (SPD489-325); Findling 2011 (SPD 489-305, NCT00735371); Frick 2017 (SPD465-303, NCT00152022); Kay 2009a; Paterson 1999; Pilzka 2000; Spencer 2001; SPD489-405 (NCT01552915); SPD489-406 (NCT01552902); Spencer 2006 (SLI381-314, NCT00507065); Spencer 2008 (SPD465-301, NCT00150579); Stein 2011 (NCT00393042); Taylor 2000; Taylor 2001; Weisler 2006 (SLI381-303); Winhusen 2010 (NCT00253747)
Atomoxetine vs. Guanfacine	1	Hervas 2014 (SPD503-316, NCT01244490, EudraCT: 2010- 018579-12)
Atomoxetine vs. Methylphenidate	8	Bedard 2015 (NCT00183391); Newcorn 2008 (B4Z-MC-LYBI); Sangal 2006 (B4Z-US-LYAV); Schulz 2012; Spencer 2002a (B4Z-MC-HFBD); Spencer 2002b (B4Z-MC-HFBK); Wang 2007 (NCT00486083, B4Z-MC-LYBR (6934)); Weisler 2012 (NCT00880217)
Atomoxetine vs. Placebo	41	Adler 2008a (B4Z-MC-LYBV, NCT00190931); Adler 2009a (B4Z-US-LYDQ, NCT00190879); Adler 2009b (B4Z-US-LYCU, NCT00190736); NCT00190736); Allen 2005 (B4Z-MC-LYAS); Arnold 2006; Bain 2013 (NCT00429091); Bangs 2007 (B4Z-MC-LYAX); Bangs 2008 (B4Z-MC-LYBX, NCT00191698); Block 2009 (B4Z-US-LYCC, NCT00486122); Dell'Agnello 2009; Dittman 2011; Durell 2013 (B4Z-US-LYDZ, NCT00510276); Gau 2007 (B4Z-TW-S010, NCT00485459); Geller 2007 (B4Z-US-LYBP); Goto 2017 (B4ZJE-LYEE, NCT00962104); Harfterkamp 2012 (NCT00380692); Hervas 2014 (SPD503-316, NCT01244490, EudraCT: 2010- 018579-12); Kay 2009b; Kelsey 2004 (B4Z-US-LYBG); Lin 2016 (NCT00917371); Martenji 2010 (B4Z-MW-LYCZ, NCT00386581); McRae-Clark 2010 (R21DAO18221, NCT00360269); Michelson 2001 (B4Z-MC-LYAC); Michelson 2002 (B4Z-MC-LYAT); Michelson 2003a; Michelson 2003b; Montoya 2009 (B4Z-XM-LYDM, NCT00191945); Newcorn 2008 (B4Z-MC-LYBI); Spencer 1998; Spencer 2002a (B4Z-MC-HFBD); Spencer 2002b (B4Z-MC-HFBK); Sutherland 2012 (NCT00174226); Svanborg 2009 (B4Z-SO-LY15, EUCR2004-003941-42-SE, NCT00191542); Takahashi 2009 (B4Z-JE-LYBC, NCT00191295); Wehmeier 2012 (B4Z-SB-LYDV, NCT00546910); Weisler 2012 (NCT00880217); Weiss 2005 (B4Z-MC-LYAW); Wietecha 2013 (NCT00607919); Wilens 2008 (B4Z-MC-LYBY, NCT00190957); Wilens 2011 (NCT00528697); Young 2011 (B4Z-US-LYCW, NCT00190775)
Bupropion vs. Methylphenidate	2	Jafarinia 2012; Moharari 2012 (IRCT201012295500N1)
Bupropion vs. Placebo	4	Casat 1989; Reimherr 2005; Wilens 2001; Wilens 2005 (NCT00048360)
Clonidine vs. Methylphenidate	4	Connor 2000; Kurlan 2002; Palumbo 2008 (NCT00031395); Singer 1995; van der Meere 1999
Clonidine vs. Placebo	5	Jain 2011 (NCT00556959); Kurlan 2002; Palumbo 2008 (NCT00031395); Singer 1995; van der Meere 1999
Guanfacine vs. Placebo	12	Biederman 2008 (SPD503-301, NCT00152009); Connor 2010 (SPD503-307, NCT00367835); Hervas 2014 (SPD503-316, NCT01244490, EudraCT: 2010- 018579-12); Kollins 2011 (SPD503-206, NCT00150592); McCracken 2016; NCT01069523; Newcorn 2013 (SPD503-314, NCT00997984); Ruginio 2014 (NCT01156051); Sallee 2009 (SPD503-304, NCT00150618); Schahill 2001 (NCT00004376); Taylor 2001; Wilens 2015 (SPD503-312, EUCR2011-002221-21, NCT01081132)
Methylphenidate vs. Modafinil	1	Amiri 2008
Methylphenidate vs. Placebo	47	Abikoff 2009; Adler 2009c (CR011560, NCT00326391); Biederman 2006a (subsample of NCT00181571); Biehl 2016; Bron 2014; Buitelaar 1996; Casas 2013 (EudraCT: 2007-002111-82); Childress 2009 (CRIT124E2305, NCT00301236); Coghill 2013 (SPD489-325); Cook 1993; CRIT124DUS02; Dopfner 2003; Findling 2008 (NCT00444574); Ginsberg 2012 (EUCR2006-002553-80-SE); Goodman 2016 (NCT00937040); Greenhill 2002; Greenhill 2006b (CRIT124E2301); Grizenko 2012; Herring 2012 (NCT00475735); Huss 2014 (CRIT124D2302, EUCR2010-021533-31-DE, NCT01259492); Kooij 2004; Kurlan 2002; Lin 2014 (NCT00922636); Medori 2008 (LAIMDA-I EUCR2004-000730-37, NCT00246220); Newcorn 2008 (B4Z-MC-LYBI); Palumbo 2008 (NCT00031395); Philippen 2015 (EUCR2006-000222-31-DE, ISRCTN54096201); Pilzka 2000; Reimherr 2007; Rosler 2009; Schranter 2016 (NTR3103, EUCR2010-023654-37-NL); Simonoff 2013 (ISRCTN683849); SPD489-405 (NCT01552915); SPD489-406 (NCT01552902); Spencer 1995; Spencer 2002a (B4Z-MC-HFBD); Spencer 2002b (B4Z-MC-HFBK); Spencer 2005; Spencer 2007 (CRIT124E2302); Stein 2011 (NCT00393042); Takahashi 2014 (NCT01323192); Taylor 1987; van der Meere 1999; Weisler 2012 (NCT00880217); Wender 2011; Wigal 2004; Wigal 2015 (NCT01239030)
Modafinil vs. Placebo	8	Arnold 2014 (C1538/2027/AD/US, NCT00315276); Biederman 2005 (Study 311 Cephalon); Biederman 2006a (Study 309 Cephalon); Kanbazi 2009; Ruginio 2003; Swanson 2006; Taylor 2000

* Amphetamines include Lisdexamfetamine.




Figure 3. Direct and indirect comparisons in the network meta-analysis of 8 interventions for primary outcome. (Dark lines are eligible comparisons for pairwise meta-analysis, added dotted blue lines show indirect comparisons). This image has been modified from Cotese *et al.* 2018⁶ under Creative Commons Attribution License (CC BY).




Figure 4. Venn diagram showing the coverage of comparisons by the network meta-analysis (from formula), and pairwise meta-analysis (from network plots), and trials (from study-based register).

Table 2. Comparisons from the body of evidence.

Source of comparisons	Type of comparisons		Eligibility for analyses		# of comparisons	% of comparisons
	Direct	Indirect	Eligible	Ineligible		
Formula	√	√	√	√	28=(8*(8-1))/2	100.00
Randomised trials	√	×	√	√	16 (Table 1)	57.14
Pairwise meta-analysis	√	×	√	×	6-13 (Figure 3)*	21.42 to 46.42
Network meta-analysis	√	√	√	√	28 (Figure 2)	100.00

* There are five networks in Figure 3 and each has 6, 11, or 13 eligible comparisons. Three out of 16 comparisons from trials have not been included in any of five network plots.

the number of comparisons *theoretically* possible and those actually employed within any given network meta-analysis. The formula would highlight this for researchers and readers and, before and after analyses, facilitate descriptions of why particular comparisons have not been included.

Conclusion

The formula produces an accurate enumeration of the potential comparisons within a single trial or network meta-analysis.

Any shortfall between the full potential of the data and the actual number of comparisons within a network meta-analysis

should be possible to explain through reference to pre-stipulated eligibility criteria or *post hoc* exclusions.

Data availability

The data analyzed in the present study have been published previously^{6,7}.

Grant information

The authors declared that no grants were involved in funding this work.

References

- Shokraneh F, Adams CE: **Study-based registers of randomized controlled trials: Starting a systematic review with data extraction or meta-analysis.** *BiolImpacts.* 2017; 7(4): 209–217. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Shokraneh F, Adams CE: **Gallstone, snake venom and witchcraft for schizophrenia: the challenges of classifying [schizophrenia] trials.** *Evidence-Based Medicine.* 2018; 23(Suppl. 1): A18. [Publisher Full Text](#)
- Li T, Puhan MA, Vedula SS, *et al.*: **Network meta-analysis-highly attractive but more methodological research is needed.** *BMC Med.* 2011; 9: 79. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Evans I, Thornton H, Chalmers I, *et al.*: **Testing Treatments: Better Research for**

Better Healthcare. Pinter and Martin Publishers. 2011.

[PubMed Abstract](#)

5. Cipriani A: **Cortese et al_LancetPsyc 2018_OPEN DATA.** *Mendeley Data.* 2018; v1.
[Publisher Full Text](#)
6. Cortese S, Adamo N, Del Giovane C, *et al.*: **Comparative efficacy and tolerability of medications for attention-deficit hyperactivity disorder in children,**

adolescents, and adults: a systematic review and network meta-analysis.

Lancet Psychiatry. 2018; 5(9): 727–38.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

7. Cortese S, Adamo N, Mohr-Jensen C, *et al.*: **Comparative efficacy and tolerability of pharmacological interventions for attention-deficit/hyperactivity disorder in children, adolescents and adults: protocol for a systematic review and network meta-analysis.** *BMJ Open.* 2017; 7(1): e013967.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 04 March 2019

<https://doi.org/10.5256/f1000research.18976.r44117>

© 2019 Soomro G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



G. Mustafa Soomro

St. James Hospital, Portsmouth, UK

This is a useful paper for demonstrating and discussing how an established formula could be used for calculating all possible pairs of comparisons for interventions in a network meta-analysis. Thus, reviewers would be able to find out how many potential comparisons have not been carried out.

I have some minor suggestions numbered from 1) to 4):

1. In the abstract the following sentence should be amended for clarity perhaps as follows: "A total of 133 trials included in the network generated 163 comparisons (16 unique direct comparisons for 8 interventions)". Amendment: "A total of 133 trials of 8 interventions were selected which included 163 comparisons. The network of these showed 16 unique direct comparisons."
2. On page 2, it says that N is a triangular number. Either this point is not relevant, or why being a triangular number is important should be described.
3. I think they should say that the formula they have used is an established formula from combinatorics for calculating number of pairs for a number of items in a set.
4. On page 3 under "Comparisons in network meta-analysis plots", line 2 and line 11 "though" should be changed to "through".

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Systematic reviews and meta-analysis and secondary data analysis.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 27 Mar 2019

Farhad Shokraneh, University of Nottingham, Nottingham, UK

Dear Dr G. Mustafa Soomro,

Thanks for spending your time for reviewing our work and commenting on it. In the following lines, we replied to your comments and made changes to the paper to cover your suggestions.

COMMENT: This is a useful paper for demonstrating and discussing how an established formula could be used for calculating all possible pairs of comparisons for interventions in a network meta-analysis. Thus, reviewers would be able to find out how many potential comparisons have not been carried out. I have some minor suggestions: In the abstract the following sentence should be amended for clarity perhaps as follows: "A total of 133 trials included in the network generated 163 comparisons (16 unique direct comparisons for 8 interventions)". Amendment: "A total of 133 trials of 8 interventions were selected which included 163 comparisons. The network of these showed 16 unique direct comparisons."

REPLY: Thank you. We revised the sentence as suggested.

CHANGE: "A total of 133 trials of 8 interventions were selected which included 163 comparisons. The network of these showed 16 unique direct comparisons".

COMMENT: On page 2, it says that N is a triangular number. Either this point is not relevant, or why being a triangular number is important should be described.

REPLY: We agree. We deleted this part.

CHANGE: "N is a triangular number" was deleted.

COMMENT: I think they should say that the formula they have used is an established formula from combinatorics for calculating number of pairs for a number of items in a set.

REPLY: Thank you for adding this. We agree and we added your suggestion in the text right after formula.

CHANGE: "This is an established formula from combinatorics for calculating number of pairs for a number of items in a set".

COMMENT: On page 3 under "Comparisons in network meta-analysis plots", line 2 and line 11 "though" should be changed to "through".

REPLY: Thank you for detecting these errors. We collected both.

CHANGE: we replaced 'though' with 'through'.

Thanks again for your valuable comments.

Best Regards,
Farhad Shokraneh

Competing Interests: No competing interests were disclosed.

Reviewer Report 13 February 2019

<https://doi.org/10.5256/f1000research.18976.r43427>

© 2019 Broderick J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Julie Broderick 

Department of Medicine, Division of Physiotherapy, Trinity Centre for Health Sciences, Trinity College Dublin, Dublin, Ireland

In my opinion, this is a novel approach which allows the quantification of comparators in trials and network meta-analyses. Following on from this, the comparators which are possible versus how many were reported can be discussed in further depth. This formula has great applicability in terms of network meta-analysis methodology which is still very much a developing area.

Some very minor points;

1. Please add a table which demonstrates the use of the formula to quantify how many comparators are possible for trials from 2 interventions up to 10.
2. Pg 3, paragraph with the heading 'Reported comparisons within the trials' - line 2 change 'there to 'the'.
3. Explanation of Figure 4 in text is not clear.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Systematic reviews, overview methodology, physical activity in chronic disease

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 27 Mar 2019

Farhad Shokraneh, University of Nottingham, Nottingham, UK

Dear Dr Julie Broderick,

Thanks for spending your time for reviewing our work and commenting on it. In the following lines, we replied to your comments and made changes to the paper to cover your suggestions.

COMMENT: In my opinion, this is a novel approach which allows the quantification of comparators in trials and network meta-analyses. Following on from this, the comparators which are possible versus how many were reported can be discussed in further depth. This formula has great applicability in terms of network meta-analysis methodology which is still very much a developing area. Some very minor points; please add a table which demonstrates the use of the formula to quantify how many comparators are possible for trials from 2 interventions up to 10.

REPLY: Thanks for your positive comment. We added a single visual proof for this formula to show how it works. The networks of 2 to 10 interventions will create networks in shapes of line, triangle, rectangle, pentagon, hexagon, heptagon, octagon, and decagon.

CHANGE: We added: "The networks of 2 to 10 interventions will create networks in shapes of line, triangle, rectangle, pentagon, hexagon, heptagon, octagon, and decagon".

COMMENT: Pg 3, paragraph with the heading 'Reported comparisons within the trials' - line 2 change 'there' to 'the'.

REPLY: Thank you and sorry for this typo.

CHANGE: We change 'there' to 'the'.

COMMENT: Explanation of Figure 4 in text is not clear.

REPLY: We added a few sentences in the text to clarify it.

CHANGE: "This diagram shows that only some of the comparisons from trials in study-based register could be included in pairwise meta-analysis. In addition, the number of comparisons in network meta-analysis (calculated by formula) is larger and inclusive of all the comparisons in the network of interventions and includes all the possible unique comparisons even if the comparisons are not in trials or in pairwise meta-analysis".

Thanks again for your valuable comments.

Best Regards,
Farhad Shokraneh

Competing Interests: None.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research