



RESEARCH ARTICLE

An extension of the Alma-Ata vision for primary health care in light of twenty-first century evidence and realities [version 1; peer review: 1 approved, 2 approved with reservations]

Henry B. Perry

Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, 21205, USA

v1 First published: 14 Dec 2018, 2:70 (<https://doi.org/10.12688/gatesopenres.12848.1>)Latest published: 14 Dec 2018, 2:70 (<https://doi.org/10.12688/gatesopenres.12848.1>)

Abstract

This paper builds upon and extends the definition of primary health care in the 1978 Declaration of Alma-Ata. The definition proposes a stronger role for community-based delivery of services and community mobilization, participation and empowerment. It calls for a stronger integration with vertical, disease-specific programs. And, finally, it calls for a strong role for certain curative services (including basic and essential surgery) that many today would not consider as part of primary health care. There is growing evidence that communities can and should play a stronger role than has traditionally been the case, that community-level workers who are properly trained and supported can provide effective services outside of health facilities, and that primary health centers staffed with non-specialist physicians and even non-physician clinicians can perform many of the lower-level inpatient services now performed at first-level referral hospitals. An approach to primary health care that is appropriate to the local context and that merges local epidemiological priorities with the communities' perceived priorities will make it possible to engage communities as partners. Currently, essential and basic health care services are available to only one-half of the world's population. The full development of primary health care as envisioned here will accelerate progress in achieving Health for All as envisioned at the International Conference on Primary Health Care in 1978.

Keywords

primary health care, community health, community-based primary health care

Open Peer Review

Reviewer Status

	Invited Reviewers		
	1	2	3
version 1			
published 14 Dec 2018	report	report	report

- Peter S. Hill** , University of Queensland, Brisbane, Australia
- Wim Van Damme** , Institute of Tropical Medicine Antwerp, Antwerp, Belgium
Juan Antonio Ricarte , University of the Philippines, Manila, Philippines
- Peter Byass** , Umeå University, Umeå, Sweden

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Henry B. Perry (hperry2@jhu.edu)

Author roles: Perry HB: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by The Bill and Melinda Gates Foundation (grant numbers OPP1152330 and 24123). *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2018 Perry HB. This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Perry HB. **An extension of the Alma-Ata vision for primary health care in light of twenty-first century evidence and realities [version 1; peer review: 1 approved, 2 approved with reservations]** Gates Open Research 2018, 2:70 (<https://doi.org/10.12688/gatesopenres.12848.1>)

First published: 14 Dec 2018, 2:70 (<https://doi.org/10.12688/gatesopenres.12848.1>)

Introduction

Primary health care in the context of global health and the reality of underserved populations in low-income countries, as opposed to primary medical care in the context of well-developed health systems of high-income countries¹, addresses practical approaches to the provision of basic ambulatory health services. Thus, instead of focusing on curative care at health facilities by clinicians, the global context of primary health care builds on the definition of primary health care as defined in the Declaration of Alma-Ata at the International Conference on Primary Health Care in Alma-Ata (now Almaty), Kazakhstan. This paper proposed an expanded definition of primary health care that builds on this original Alma-Ata definition in light of current evidence and current realities.

The World Health Organization reports that one-half of the world's population does not have access to essential and basic health services². Progress in improving the health and nutritional status of the poorest segments of many low-income countries continues to lag behind that of the rest of the world; only 4 of the 74 countries with 97% of the world's maternal and child deaths actually met both Millennium Development Goals for reduction in child and maternal mortality³. The best single indicator we have of the quality and coverage of primary health care services in a low-income country is the mortality rate of children younger than 5 years of age. Second, the population coverage of the most basic and essential services that fall within the realm of primary health care among mothers and children in low-income populations remains surprisingly low: 60% or less for 13 out of 21 interventions and 40% or less for 6 of 21 interventions³. Basic health-promoting behaviors such as exclusive breastfeeding and hand washing, which we know can be effectively promoted through community-based primary health care programs, are still not the norm. Third, the limits of the decades-long emphasis on specific disease-control programs and selective top-down programs are being increasingly recognized. The lack of funding for, and emphasis on, health systems strengthening and, in particular, the lack of emphasis on strengthening of primary health care programs and the community health service delivery component of primary health care, are becoming recognized. Fourthly, health promotion, community participation, community mobilization, and community empowerment are now part of the domain of primary health care and are recognized as essential for improving the health of impoverished populations, for increasing the coverage of basic and essential services, and for adopting healthy behaviors⁴. Fifthly, with the emergence of chronic diseases as the major global disease burden of the future and the need for persons with chronic diseases to have access to health services, the need for a functioning primary health care system is becoming ever more obvious. Sixthly, and perhaps most importantly, 1 billion people have never seen a health provider⁵.

The desire—and right—to live a long, healthy and productive life is one of the most fundamental of human hopes and aspirations, and one of the most important functions that societies strive to fulfil. Without the full development of primary health care, Health for All cannot be achieved.

An extension of the Alma-Ata definition of primary health care for underserved populations in light of 21st evidence and realities

I propose here a definition of primary health care that builds on the historical tradition of global primary health care, but at the same time pushes the borders of the definition in three directions: (1) a stronger role for community-based delivery of services and community mobilization/participation/empowerment, (2) a stronger role for disease prevention with links to vertical programs, and (3) a stronger role for certain curative services that many today would not consider as part of primary health care.

Primary health care consists of those services that people seek and that providers (individual and organizational) deliver to protect health and to treat basic and uncomplicated illness, disease and injuries—especially those that are public health priorities in terms of disease burden that can be alleviated through cost-effective and affordable interventions and programs. These services include those that can be provided in communities outside of facilities by community-based workers, as well as services provided at facilities by frontline health workers (including auxiliary nurses, graduate nurses, and physicians) without advanced specialized training and without expensive diagnostic and laboratory support. The presence of a basic laboratory, other diagnostic equipment, including basic x-ray and ultrasound capability, and an operating room is appropriate for a primary health care center along with the staff and support required to utilize them.

This definition implies that low-cost diagnostic and laboratory support would be available at a health center. Given the fact that technology is evolving rapidly, some of this support available today would have been considered advanced 20 years ago, and what is considered advanced today may be low-cost and usable in primary health care centers in 20 years into the future. Furthermore, what is considered to be “low-cost” at any given point in time will vary from one setting to another. X-ray capabilities and an operating room to perform certain uncomplicated surgical procedures were in fact included in 1920 as elements of primary health centers proposed by the Health Survey and Development Committee led by Lord Dawson of Penn in 1920 in what is known today as the Dawson Report⁶, which contains the first known use of the term “primary health center.”

Primary health care is composed of three types of activities: disease-oriented primary health care, services-oriented primary health care, and community-oriented primary health care. Disease-oriented primary health care consists of local efforts to control diseases that constitute a significant health burden in the population and for which disease-control interventions exist. While these activities normally have strong technical and funding support from higher levels of government, they are in fact mostly carried out in the community and should be integrated with the primary health care system. Services-oriented primary health care consists of efforts to extend basic personal health care services to the entire population. Community-oriented

primary health care consists of efforts to work in partnership with communities to improve their health. These domains are not mutually exclusive and overlap in many instances.

Each of these three types of primary health care is equally important, and together they function like the three legs of a stool on which the “seat” of primary health care rests. If one leg is poorly developed, the primary health care system is sub-optimally constructed to reach its full potential. For the most underserved populations in low-income countries, disease-oriented primary health care has been the strongest leg of this stool by far, with services-oriented primary health care receiving less attention and community-oriented primary health care receiving far less attention than the other two legs.

Box 1 lists the components that make up an ideal primary health care system for underserved populations in low-income countries.

Box 1. Categories of primary health care for underserved populations in low-income countries*

- Disease prevention
- Screening for priority infectious diseases and other priority health-related conditions
- Treatment of priority infectious diseases (HIV/AIDS, tuberculosis, and malaria)
- Health promotion
- Basic and essential services for women, mothers and children.
- Appropriate treatment of common diseases and injuries
- Basic and essential surgical care
- Community engagement
- Attention to the social determinants of ill health (illiteracy and lack of education, lack of an adequate food supply and adequate housing).

*Some, but not all of these were mentioned specifically in the Declaration of Alma-Ata

Disease prevention: Education about prevailing preventable health problems and methods of preventing and controlling them; provision of preventive services such as immunizations against major infectious diseases, provision of clean water and sanitation, micronutrient supplementation, distribution of insecticide-treated bed nets and intermittent preventive treatment of malaria in pregnant women and children in malaria-endemic areas.

Screening for priority infectious diseases and other health-related conditions: HIV infection, cervical and breast cancer, under- and over-nutrition.

Treatment of priority infectious diseases: HIV/AIDS, tuberculosis and malaria.

Health promotion: Education about behaviors and health care utilization that promote good health, including hand washing,

clean water, sanitation and household cleanliness as well as dental hygiene, menstrual hygiene, safe sex, and birth spacing; assurance of good nutrition, especially in mothers and infants; promotion of smoking cessation, weight reduction for those who are obese, physical activity for those who are sedentary, and moderation of alcohol intake; and awareness of warning signs of serious illness for which medical care should be sought.

Basic and essential services for women, mothers and children:

Provision of first-line family planning services (birth control pills, condoms, injectable contraceptives, subcutaneous contraceptive implants; detection and treatment of reproductive tract infections; screening and providing initial treatment for breast of, cervical cancer; antenatal care; education about recognition and prevention of, and need to obtain care for complications of pregnancy and childbirth, proper care of newborns and young children, and warning signs of newborn and childhood illness for which attention from properly trained health care providers should be sought; provision of safe and hygienic delivery and treatment of common obstetrical complications, including infection, retained products of conception or retained placenta, management of obstructed labor, prevention and treatment of post-partum hemorrhage, treatment of pre-eclampsia and eclampsia, and management of abnormal fetal presentation/prolonged labor; prevention, detection and treatment of pre-term birth, of birth asphyxia, and of neonatal infection; prevention, detection and treatment of childhood undernutrition, pneumonia, and diarrhea; and recognition of serious neonatal and childhood illness. Management of obstetrical complications requires the capacity for giving parenteral medications and blood transfusions and performing evacuation of the uterus and cesarean section, all of which can be performed in an adequately equipped primary health care center staffed with health personnel who do not have highly specialized training.

Appropriate treatment of common diseases and injuries:

Provision of first aid, acute illness management, chronic disease management, pain management, and mental illness care with the full complement of essential medicines; treatment of eye and skin infections, acute respiratory infection/uncomplicated pneumonia; and recognition and referral of life-threatening conditions.

Basic and essential surgical care:

“First-line” surgical care (repair of uncomplicated hernias, initial surgical management of the acute surgical abdomen (appendectomy, management of intestinal perforation or uncomplicated intestinal obstruction, and so forth), circumcision, and cataract surgery, all of which can be performed by non-specialist clinicians in primary health centers that can also serve as a first-line basic hospital. Other surgical procedures can be performed outside of hospitals under local anesthesia by appropriately trained and supervised non-physician surgeons during “campaigns”, in which large numbers of patients come to specially formed teams to receive these services. Examples of such procedures include tubal ligation, vasectomy, circumcision, cataract extraction, and herniorrhaphy) in addition to burn, wound, and fracture management; and cesarean section.

Assistance in the rehabilitation of people with long-term disabilities: Helping people with physical and mental disabilities to become as functional as possible in the community.

Community engagement: Engaging the community in understanding and addressing their health priorities and the determinants of these problems through community-based workers, participatory women's groups, and participation in community oversight and advocacy activities; and monitoring of local population health status through registration of vital events and periodic population-based health surveys.

Attention to the social determinants of ill health: Literacy and basic education, promotion of adequate food supply and appropriate dietary intake, and provision of adequate housing.

Further elaboration on this extended definition of primary health care

This updated definition of primary health care seeks to give emphasis to the importance of extending services into the community and households and helping communities to engage in activities that will improve health. Including the term “care” in the phrase primary health care conveys, unfortunately, the notion that a service is being provided to treat an illness or that a service is being provided to prevent an illness. However, as stated previously, the concept of health promotion to adopt healthy household behaviors and appropriate utilization of health services is now fully embedded in the emerging new vision of primary health care.

Disease prevention at the local level, promotion of healthy behaviors, and timely utilization of priority health services fall within the domain of primary health care. Primary health care includes the provision of health-related services provided to persons in their homes, communities, and at ambulatory health facilities by primary health care practitioners. Primary health care is increasingly provided by health teams. I am also proposing that the concept of primary health care should include first-line, lower-level inpatient care that can be provided at a health center, including basic and essential surgical services such as caesarean sections.

In its conceptual and functional sense, primary health care is not tied to any specific type of health facility or to any type of health care provider. Mothers are primary health care providers in their homes for their children – probably the most important of all primary health care providers in impoverished populations. They prevent and treat more illnesses than the formal health system does. In addition to mothers, primary health care is provided by community health workers (CHWs), auxiliaries, nurses, physicians, traditional healers, drug sellers, and informal or non-formally trained practitioners.

In its narrowest traditional sense, primary health care is simply ambulatory health services that are provided by frontline health workers, both formal and informal. Thus, in the narrow traditional sense, primary health care is what I am calling services-oriented primary health care. But this concept of primary

health care fails to include disease-oriented primary health care and community-oriented primary health care.

The 1978 International Conference on Primary Health Care and its Declaration of Alma-Ata served as the much-needed catalyst to shift the paradigm of primary health care away from the Western notion of primary medical care (with its focus on the provision of ambulatory health services provided by non-specialist physicians or mid-level clinicians such as nurse practitioners and physician assistants at health facilities such as medical offices, health centers, or outpatient clinics of hospitals) to the notion of “essential care” that could be provided in settings where such infrastructure is non-existent. The Declaration states that primary health care is:

[E]ssential health care based on practical, scientifically sound and socially acceptable methods and technology made universally accessible to individuals and families in the community through their full participation and at a cost that the community and country can afford to maintain at every stage of their development in the spirit of self-reliance and self-determination⁷.

The 1978 International Conference on Primary Health Care was the largest and most representative global health conference that had been held up to that time, with representatives of 134 governments and 67 international organizations⁸. It is, then, a definition that had broad legitimacy when issued and has consistently been affirmed since as the “gold standard” for primary health care. It includes the three types of primary health care I have specified. The Conference also called for the achievement of Health for All through primary health care by the year 2000—a goal obviously not yet met, but which will remain with us for at least most of the 21st century before it is achieved.

Primary health care responds to patient-defined needs for illness care and also proactively addresses local epidemiological priorities in the community. Primary health care is therefore important for improving the health status of the community. Local epidemiological priorities are the most frequent, serious, readily preventable or treatable conditions in the community. Since epidemiological priorities vary from one locality to another, they are best determined from locally acquired surveillance data. Admittedly, this is not readily accomplished, but epidemiographic surveillance has long been proposed⁹ and is feasible to incorporate into primary health care systems that utilize community-based workers to visit all households on a regular basis. And, in fact, it provides an opportunity for linking civil registration of vital events with primary health care services and thus improving registration of births and deaths in low-income countries, a recognized global priority¹⁰. Addressing epidemiological priorities also involves addressing the basic underlying physical and social determinants of ill health—access to safe water, sanitation, good nutrition, adequate housing, and basic education.

There is no general global standard of what constitutes a primary health care center in low-income settings. At present,

many of these health centers have inpatient beds and provide inpatient care but do not commonly have the capability to provide—at least for now—the basic laboratory and diagnostic procedures or the basic surgical procedures that should become standard for primary health care programs throughout the world, as will be discussed later.

Households and communities are resources, not just targets, for implementing primary health care. Because of the scarcity and expense of physical health facilities in low-income countries and the higher-level professional personnel who staff them, much of primary health care needs to be provided in homes and communities by community-based staff who are properly trained and supervised in implementing evidence-based interventions and carrying out procedures that have been scientifically validated for provision by lower-level health workers, including illiterate CHWs in populations with high levels of illiteracy.

Health teams led by higher-level health personnel are necessary to train and support CHWs, and a strong system of logistical support for supplies, medicines and commodities is also required. Primary health care is that element of the health systems that can make the greatest contribution to improving population health. As such, it should be the foundation of health systems, particularly in resource-constrained settings. Strengthening the provision of primary health care services should be the first health priority of governments.

How can a core package of PHC services that are appropriate for the local context be defined?

The updated definition of primary health care provided at the outset contains a full expression of what services need to be provided to disadvantaged populations of low-income countries (and of course to all people everywhere). Out of this full set of services, what portion might be identified as a core set of priority services to be provided?

Efforts to do this have usually used cost-effectiveness criteria for population health benefits, but this approach ignores the obvious fact that people have always needed, and will always need, curative medical services for symptomatic conditions, whether or not they have significance for mortality or serious disability. Urinary infections, arthritis, wrist fractures, toothaches and eye infections, to mention only a few examples, even if they have a low likelihood of causing death or long-term serious disability, will produce a desire for treatment from some kind of primary health care provider because of the symptoms they produce. So, a core set of PHC services needs to include curative care along with services that have demonstrable benefit for reduce the risk of death.

How might one go about defining such a core package? To my mind, the most logical way to go about this is through the application of the principles of what are now called the census-based, impact-oriented approach¹¹. This involves a health program developing a partnership with a population, using local surveillance carried out by routine home visits to define epidemiological priorities (the most frequent, serious, readily

preventable or treatable conditions in the population) and to help the communities in the population identify what they consider their health priorities to be (and in low-income settings the community's health priorities often revolve around improving curative care services). Then, with the available resources (financial, infrastructure, and human), develop a plan that addresses program priorities, which are a combination of epidemiological and community-defined priorities. Thus, the actual content of the services would vary from place to place and over time, depending on the local situation. This approach represents a paradigm shift from what has been the norm in global health for the past half-century or more, namely top-down vertical approaches with no meaningful engagement of local populations or local data in establishing program priorities¹².

The neglected demand-side approach to strengthening primary health care

The concept of community participation in primary health care is strongly enunciated in the Declaration of Alma-Ata, with its call for the following:

The people have the right and duty to participate individually and collectively in the planning and implementation of their health care....

Primary health care ... requires and promotes maximum community and individual self-reliance and participation in the planning, organization, operation and control of primary health care, ... and to this end develops through appropriate education the ability of communities to participate.

Primary health care ... relies ... on health workers ... suitably trained socially and technically to work as a health team and to respond to the expressed health needs of the community¹³.

In the extended definition of primary health care presented earlier in this paper, community-oriented primary health care is one of the three “pillars” (or legs of the stool) of primary health care, and this involves working in partnership with communities to help them improve their health.

Roles of higher- and lower-level personnel

The updated definition of primary health care proposed here calls for the inclusion of the full spectrum of primary health care providers, from mothers in the home to community health workers, auxiliary workers, nurses, and physicians. In resource-constrained settings and even in resource-rich settings such as the United States, the team approach to provision of health services leads to the best quality of services and to the most cost-efficient use of available resources. Even in the United States, the use of CHWs as members of health teams is growing rapidly^{14,15}. There is now a vast global evidence base regarding the suitability and advisability of delegating to lesser-trained staff many tasks and responsibilities that were once the sole purview of physicians and graduate nurses¹⁶. Although not without controversy still, the weight of the evidence continues to strongly support the full engagement of

carefully selected lower-level workers to carry out many well-defined tasks and responsibilities as long as these persons are suitably trained and well-supervised¹⁷.

Remuneration and other incentives for frontline workers

The remuneration of frontline health workers is a current source of controversy, particularly with respect to whether or not it is appropriate for health programs to engage CHWs on a voluntary, non-salaried basis. Although the amounts of money paid to frontline workers are modest, the vast number of these workers means that the implications of payment policies for CHWs are substantial, especially when the funding comes from a central government source. On the one hand, there is the frequently made charge that engaging CHWs without remuneration is unjust exploitation. Nonetheless, field experiences have demonstrated that communities commonly have people who are eager to serve their neighbors on a voluntary basis. If funding scarcity were not a problem, then payment would of course be desirable. However, there are examples of CHW programs that made initial commitments to pay their workers but then could not maintain that commitment, leading to a crumbling of the program^a. If the creation of an expectation for a salary is established and then the program cannot sustain that expectation over time, the program is worse off than if it had originally started with volunteer CHWs who had no expectation of a salary.

In NGO child survival programs there are many examples of volunteer CHWs who receive no formal salary but who receive some kind of “incentives,” in the form of special recognition from the community, release from certain community responsibilities, special privileges in accessing health services, and so forth. Many top-down, highly selective vertical programs provide a short-term daily incentive for a short period of time. The most balanced approach to this issue is to not expect from volunteers more than a modest amount of ongoing weekly work (e.g., no more than 4–5 hours per week) and, if on-going financial remuneration is to be provided, to be confident that this support can be maintained on a sustainable basis.

Impact of global trends on the evolution of primary health care for the disadvantaged in low-income countries

Anticipating the specific influences of global forces and trends on primary health care is, of course, impossible. We can be sure that more and more of the disadvantaged in low-income countries will be living in slums in urban area. We can be sure that technological advances will bring new diagnostic and laboratory tests within the reach of low-cost primary health care programs, and mobile health (mHealth) will make it easier for people to communicate with their health care provider and easier for different members of the health team to communicate among themselves, providing opportunities for improving the quality of care. As the population ages, chronic diseases will increasingly

dominate the disease burden. AIDS will have become a chronic disease, and primary health care will be heavily concerned with care of the elderly. Socioeconomic development will lead to higher living standards and higher educational levels, and this will affect the demand for and consumption of primary health care services. No doubt the trend for women’s empowerment will continue, and this will also affect the demand for and consumption of primary health care services.

Towards a rebirth and revision of primary health care

In recent articles and writings related to primary health care, there has been a persistent theme: the concept of primary health care as articulated at Alma-Ata, Kazakhstan is valid and needs to be maintained. But at the same time there is a broadly held view that we are now at a time for a renewal and revitalization of primary health care and perhaps even for a re-definition of primary health care for the 21st century, leading to an even stronger commitment to achieve the vision of Alma-Ata.

In recognition of the 25th anniversary of the Alma-Ata conference in 2003, the Pan American Health Organization convened a series of events and dialogues, culminating in a 2007 position paper on renewing primary health care in the Americas¹⁸. Their report states that there is “a growing recognition that PHC is an approach to strengthen society’s ability to reduce inequities in health; and a growing consensus that PHC represents a powerful approach to addressing the causes of poor health and inequality”¹⁸ [p. 2]. In celebration of the 30th anniversary of the Declaration of Alma-Ata, *The Lancet* published a series of papers entitled “Alma-Ata: Rebirth and Revision.” The lead editorial by *The Lancet* team stated that “The Alma-Ata Declaration revolutionized the world’s interpretation of health. Its message was that inadequate and unequal health care was unacceptable: economically, socially, and politically.”¹⁹ Margaret Chan, Director General of the World Health Organization, in her lead editorial to the series, remarked: “With an emphasis on local ownership, primary health care honoured the resilience and ingenuity of the human spirit and made space for solutions created by communities, owned by them, and sustained by them.”²⁰ One of the articles in the series expressed the same idea this way:

The very idea of health for all energised workers and fueled new efforts in many countries to improve service coverage, especially for previously underserved communities. The inherent focus on equity, the necessity of reaching the unreached and involving them not only in the benefits of health care, but more importantly, in the decisions and actions that collectively make health, was at once novel and revolutionary. Thus, the precepts of social justice became an integral part of health planning²¹ [p. 919].

The Second International Conference on Primary Health Care²², held in October 2018 in Astana, Kazakhstan, to commemorate the 40th anniversary of the Declaration of Alma-Ata, and its Declaration of Astana²³, reaffirmed the central importance of primary health care for achieving universal health coverage, the health-related Sustainable Development Goals, and Health for

^aThis happened in Nepal in the early 1980s, and later these workers, who had become inactive, were recruited back later as Female Community Health Volunteers.

All. It is time to ignite a global renaissance in primary health care as the centrepiece of efforts to end preventable child and maternal deaths by the year 2030. Implementing the updated definition of primary health care provided here will contribute to the achievement of the highest attainable standard of health for all people and especially for the “bottom billion” of our global family.

Now is the time for a renewed focus on the principles of primary health care outlined at Alma-Ata in 1978. Unfortunately, Health for All was not achieved by the year 2000 and may not be achieved by the year 2100. But the full implementation of primary health care as proposed here for the disadvantaged in

low-income countries will help to achieve this global human aspiration sooner rather than later.

Data availability

No data are associated with this article.

Grant information

This work was supported by The Bill and Melinda Gates Foundation (grant numbers OPP1152330 and 24123).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Starfield B: **Primary Care: Concept, Evaluation, and Policy**. New York: Oxford University Press; 1992.
[Reference Source](#)
- WHO: **Universal Health Coverage (UHC)**. 2018; (accessed 31 October 2018).
[Reference Source](#)
- Victora CG, Requejo JH, Barros AJ, *et al.*: **Countdown to 2015: a decade of tracking progress for maternal, newborn, and child survival**. *Lancet*. 2015; **387**(10032): 2049–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rosato M, Laverack G, Grabman LH, *et al.*: **Community participation: lessons for maternal, newborn, and child health**. *Lancet*. 2008; **372**(9642): 962–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- GHWA: **Imagine**. 2011; (accessed 31 October 2018).
[Reference Source](#)
- Dawson BE: **Interim Report on the Future Provision of Medical and Allied Services 1920 (Lord Dawson of Penn)**. London: Ministry of Health, Consultative Council of Medical and Allied Services, 1920.
[Reference Source](#)
- World Health Organization, UNICEF: **International Conference on Primary Health Care, 6-12 September 1978**. 1978; Alma-Ata, USSR: World Health Organization; 1978.
[Reference Source](#)
- Cueto M: **The origins of primary health care and selective primary health care**. *Am J Public Health*. 2004; **94**(11): 1864–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fredericksen HS: **Epidemiographic Surveillance**. In: Fredericksen HS, Dunn FL, eds. *Epidemiographic Surveillance: A Symposium*. Chapel Hill, NC: Carolina Population Center; 1971.
- Setel PW, Macfarlane SB, Szreter S, *et al.*: **A scandal of invisibility: making everyone count by counting everyone**. *Lancet*. 2007; **370**(9598): 1569–77.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Perry H, Robison N, Chavez D, *et al.*: **Attaining health for all through community partnerships: principles of the census-based, impact-oriented (CBIO) approach to primary health care developed in Bolivia, South America**. *Soc Sci Med*. 1999; **48**(8): 1053–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Packard R: **A History of Global Health: Interventions into the Lives of Other People**. Baltimore, MD: Johns Hopkins University Press; 2016.
[Reference Source](#)
- WHO, UNICEF: **Declaration of Alma-Ata International Conference on Primary Health Care, Alma-Ata, USSR, 6-12 September 1978**. 1978; (accessed 31 October 2018).
[Reference Source](#)
- Singh P, Chokshi DA: **Community health workers—a local solution to a global problem**. *N Engl J Med*. 2013; **369**(10): 894–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Perry HB, Zulliger R, Rogers MM: **Community health workers in low-, middle-, and high-income countries: an overview of their history, recent evolution, and current effectiveness**. *Annu Rev Public Health*. 2014; **35**: 399–421.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dawson AJ, Buchan J, Duffield C, *et al.*: **Task shifting and sharing in maternal and reproductive health in low-income countries: a narrative synthesis of current evidence**. *Health Policy Plan*. 2014; **29**(3): 396–408.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Black RE, Taylor CE, Arole S, *et al.*: **Comprehensive review of the evidence regarding the effectiveness of community-based primary health care in improving maternal, neonatal and child health: 8. summary and recommendations of the Expert Panel**. *J Glob Health*. 2017; **7**(1): 010908.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- PAHO: **Renewing Primary Health Care in the Americas: A Position Paper of the Pan American Health Organization/World Health Organization (PAHO/WHO)**. Washington, DC: Pan American Health Organization/World Health Organization, 2007.
[Reference Source](#)
- A renaissance in primary health care**. *Lancet*. 2008; **372**(9642): 863.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chan M: **Return to Alma-Ata**. *Lancet*. 2008; **372**(9642): 865–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lawn JE, Rohde J, Rifkin S, *et al.*: **Alma-Ata 30 years on: revolutionary, relevant, and time to revitalise**. *Lancet*. 2008; **372**(9642): 917–27.
[PubMed Abstract](#) | [Publisher Full Text](#)
- WHO, MOH of Kazakhstan, UNICEF: **Global Conference on Primary Health Care: 25-26 October 2018 – Astana, Kazakhstan**. 2018; (accessed 31 October 2018).
[Reference Source](#)
- WHO, UNICEF: **Declaration of Astana**. 2018; (accessed 31 October 2018).
[Reference Source](#)

Open Peer Review

Current Peer Review Status: ? ✓ ?

Version 1

Reviewer Report 07 January 2019

<https://doi.org/10.21956/gatesopenres.13929.r26827>

© 2019 Byass P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Peter Byass 

Epidemiology and Global Health, Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden

This article discusses developments in approaches and definitions to primary health care. I find it difficult to review against the label “research article” as it much more of a narrative review in structure and content, and indeed states explicitly “no data are associated with this article”.

The author chooses to make a sharp distinction from the outset between primary health care in low-income countries versus primary medical care in high-income countries, which I find somewhat dated and unhelpful. Of course it is true that any country’s primary care and economic development are inextricably interrelated. However, the majority of the 21st century world’s population actually lives in middle-income countries, which are not mentioned. Every country’s health system should be providing both preventive and curative care at the primary level, and indeed this is a fundamental vision of WHO’s current push towards “health for all” and “universal health coverage”, which are absolutely not only for low-income countries. However, the global reality is that there is a spectrum of countries on both health care and economic development axes, and it would have been more helpful for this article to have taken this global perspective as its basis, perhaps going on to extract how countries at the lower end of the development spectrum might most appropriately conceptualise primary care.

The range of different health care personnel that are needed to provide effective primary care in any situation are also rather unhelpfully dichotomised into “higher- and lower-level personnel”. This seems to go against the concept of primary care teams which have long been a feature of many health systems, for example in European countries, and perhaps reflects an over-emphasis on the very hierarchical model of the US health system.

Nevertheless there are important pointers here to positive ways in which primary care systems might develop in various settings.

There is a typo in the first major heading, which should read “light of 21st century evidence”.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Global health

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 04 January 2019

<https://doi.org/10.21956/gatesopenres.13929.r26828>

© 2019 Van Damme W et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Wim Van Damme 

Department of Public Health, Institute of Tropical Medicine Antwerp, Antwerp, Belgium

Juan Antonio Ricarte 

UHC Study Group, University of the Philippines, Manila, Philippines

Thank you for the opportunity to review this important article. Primary Health Care (PHC) indeed is, or should be, the foundation of any health system, and it is timely to revisit the definition of PHC in line with the transitions happening in the world over the past 40 years since the Alma-Ata declaration. The paper contributes to a much-needed discussion on the concept and definition of PHC; stressing the absolute primacy of essential services in contrast to the over-dominance of disease control programs and hospital services, which is observed in many countries.

Since its inception 40 years ago, the Alma-Ata vision for PHC, has often been interpreted in a restrictive way, as Perry rightly explains. But from the onset, there has also been a school of thought on PHC which embraced a comprehensive vision, close to the one proposed in this paper. This has been framed as a

clash between “selective PHC & comprehensive PHC”, or as “vertical vs. horizontal approaches”. The paper rightly tries to go beyond this “false dichotomy” and present an inclusive definition of PHC, acknowledging the importance of all components of an effective health system. It also shows that such vision is not only relevant for low- and middle-income countries, but for all societies.

The paper presents a good rationale for redefining PHC: given that technology is rapidly evolving and has become more accessible even in low-income contexts. It would be nice to give more examples on this (e.g. new low-cost diagnostics, telemedicine, e-health). The paper also presents relevant and concrete examples of how PHC services can be expanded at the local level, and how community participation should be fully part of PHC.

However, for this review, we also identify some points that could further strengthen the paper, or need some clarification:

- Relation between PHC & Universal Health Coverage (UHC), which has now become the dominant framing of health care;
- Relation between “continuous essential services” and “disease control programs”, pleading for “integration” is the dominant advocacy term, but “integration” can be interpreted in so many different ways that some more specifics are needed to clarify the What? Why? How? Additional explanation and examples might be needed on “strengthening the integration of PHC to vertical programs”. In addition to the examples on incentivizing HRH aligned to vertical programs, some PHC-vertical program linkages (e.g. alignment of resources/technology, referral systems, alignment of priorities) could be discussed.
- A number of terms used in the article don’t necessarily have universal definitions. For example:
 - A “health center” with surgical capabilities in one context, would be called a district hospital or infirmary in a different setting
 - The term “health provider” is used loosely, (i.e. “1 billion people have never seen a health provider” Vs. “Mothers are primary health care providers”).
- Primary health care is not synonymous to health care provided in a primary care facility. In comprehensive definitions of PHC, essential care in primary hospitals, such as C-sections, is part of the PHC spectrum. Perhaps an alternate approach would be to discuss in terms of essential primary health care services.

We look forward to the final publication.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: I, Wim Van Damme, am a member of the Advisory Board of the International Institute for Primary Health Care, Ethiopia (IIfPHC-E); and the author of the paper, Henry B Perry also is a member of IIfPHC-E's Advisory Board.

Reviewer Expertise: Public health, health policy, primary health care, disease control programs, health financing, global health initiatives, chronic disease care

We have read this submission. We believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 20 December 2018

<https://doi.org/10.21956/gatesopenres.13929.r26826>

© 2018 Hill P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Peter S. Hill 

School of Public Health, University of Queensland, Brisbane, Qld, Australia

Thank you for the opportunity to review this paper which revisits Primary Health Care (PHC) following the celebration of its 40 years in action. The proposed "expansion" is consistent with the spirit of PHC, in its local application, and the need to re-invest in the framing in dialogue with Universal Health Coverage.

There are two areas in which the paper could be enhanced: Firstly, in structural clarity. It would be useful to restate the original definition of PHC that the author is working with--there are more than one definition that are in circulation, but with the paper expanding it would be useful to be specific. Secondly, it would be useful to more overtly indicate which statement is the expanded definition. In the introduction there are five "components" that could represent the new definition; in the next section there are three "directions". It would be helpful if the relation between these was stated, and the actual "definition" more clearly marked. I also found the reiteration of "stronger" problematic in that it did not provide defined endpoints in the expanded activities--it becomes difficult to know at what point the new PHC is achieved. The second issue relates to the level of analysis--this is PHC in a centre/services/activities frame--there is also a higher level of systems analysis that could be approached that would locate these within a national (or universal?) frame. In this regard, reference to the WHR 2008 Primary Health Care now more than ever would be useful because of the reforms and reframing of PHC it advocated--and that would link well to this new visioning at a systems level.

I look forward to the final publication.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: global health systems; global health policy; PHC; UHC and the SDGs

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.
