



Dietary Vitamin E as a Protective Factor for Parkinson's Disease: Clinical and Experimental Evidence

Tommaso Schirinzi¹, Giuseppina Martella^{1,2}, Paola Imbriani^{1,2}, Giulia Di Lazzaro¹, Donatella Franco¹, Vito Luigi Colona¹, Mohammad Alwardat¹, Paola Sinibaldi Salimei³, Nicola Biagio Mercuri^{1,2}, Mariangela Pierantozzi¹ and Antonio Pisani^{1,2*}

¹ Department of Systems Medicine, University of Rome Tor Vergata, Rome, Italy, ² IRCCS Fondazione Santa Lucia, Rome, Italy, ³ Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, Italy

OPEN ACCESS

Edited by:

Giovanni Albani,
Istituto Auxologico Italiano (IRCCS),
Italy

Reviewed by:

Laura Avanzino,
Università di Genova, Italy
Graziella Madeo,
National Institutes of Health (NIH),
United States

*Correspondence:

Antonio Pisani
pisani@uniroma2.it

Specialty section:

This article was submitted to
Movement Disorders,
a section of the journal
Frontiers in Neurology

Received: 06 November 2018

Accepted: 05 February 2019

Published: 26 February 2019

Citation:

Schirinzi T, Martella G, Imbriani P, Di Lazzaro G, Franco D, Colona VL, Alwardat M, Sinibaldi Salimei P, Mercuri NB, Pierantozzi M and Pisani A (2019) Dietary Vitamin E as a Protective Factor for Parkinson's Disease: Clinical and Experimental Evidence. *Front. Neurol.* 10:148. doi: 10.3389/fneur.2019.00148

Effective disease-modifying treatments are an urgent need for Parkinson's disease (PD). A putative successful strategy is to counteract oxidative stress, not only with synthetic compounds, but also with natural agents or dietary choices. Vitamin E, in particular, is a powerful antioxidant, commonly found in vegetables and other components of the diet. In this work, we performed a questionnaire based case-control study on 100 PD patients and 100 healthy controls. The analysis showed that a higher dietary intake of Vitamin E was inversely associated with PD occurrence independently from age and gender (OR = 1.022; 95% CI = 0.999–1.045; $p < 0.05$), though unrelated to clinical severity. Then, in order to provide a mechanistic explanation for such observation, we tested the effects of Vitamin E and other alimentary antioxidants *in vitro*, by utilizing the homozygous PTEN-induced kinase 1 knockout (*PINK1*^{-/-}) mouse model of PD. *PINK1*^{-/-} mice exhibit peculiar alterations of synaptic plasticity at corticostriatal synapses, consisting in the loss of both long-term potentiation (LTP) and long-term depression (LTD), in the absence of overt neurodegeneration. Chronic administration of Vitamin E (alpha-tocopherol and the water-soluble analog trolox) fully restored corticostriatal synaptic plasticity in *PINK1*^{-/-} mice, suggestive of a specific protective action. Vitamin E might indeed compensate PINK1 haploinsufficiency and mitochondrial impairment, reverting some central steps of the pathogenic process. Altogether, both clinical and experimental findings suggest that Vitamin E could be a potential, useful agent for PD patients. These data, although preliminary, may encourage future confirmatory trials.

Keywords: Parkinson's disease, Vitamin E, antioxidant, neuroprotection, protective factors, diet, PINK1, synaptic plasticity

INTRODUCTION

Parkinson's disease (PD) is a common neurodegenerative disorder, idiopathic, and multifactorial, mainly due to the loss of dopaminergic neurons in the *Substantia Nigra pars compacta* (SNpc) and to excessive brain accumulation of α -synuclein positive cytoplasmic Lewy bodies (LB). PD causes a progressive and disabling syndrome, including motor and non-motor disturbances, which severely impair patients' quality of life. Therefore, effective disease modifying treatments represent an unmet clinical need (1–3).

Successful neuroprotection may imply a combined approach against the different partners of neurodegeneration (4). Since oxidative stress is a major player of the neurodegenerative process in PD, the modulation of redox balance has been extensively explored as a potential strategy to prevent neural death and disease progression. Indeed, different antioxidant agents are under investigation in several clinical trials (5–9). Besides synthetic compounds, an invaluable source of natural antioxidants is food, namely fruits and vegetables (10). Therefore, an antioxidant-rich diet could represent a viable option to boost antioxidant pathways counteracting neurodegeneration.

Multiple antioxidant species can be found in food, which operate differently in cellular metabolism. Vitamin E, in particular, is a powerful antioxidant, found in plants and abundant in many aliments consumed in our diet (11). Vitamin E family includes a number of lipophilic molecules (α -, β -, γ -, δ -tocopherols and α -, β -, γ -, δ -tocotrienols), whose antioxidant properties rely on lipoperoxyl radical scavenging activity (11). Their neuroprotective effects have been demonstrated in multiple experimental models; likewise, the reduced levels of these molecules in humans have been associated with the occurrence of neurodegenerative diseases (12, 13).

We hypothesized that a higher dietary intake of Vitamin E might protect from progressive neurodegeneration in PD. Therefore, we conducted a study including: (1) a retrospective assessment of dietary Vitamin E intake (VEI) in PD patients compared to healthy controls, aimed at determining if a different dietary VEI is associated with diverse clinical conditions; (2) an *in vitro* protocol in brain slices of a PD mouse model, aimed at evaluating the effects of Vitamin E on synaptic plasticity abnormalities, a peculiar endophenotype observed in distinct PD models. Specifically, we used homozygous PTEN-induced kinase 1 (*PINK1*) knockout mice (*PINK1*^{-/-}), an established model of subclinical PD, which might reflect the early phases of the disease. In this model, we previously observed a significant decrease in dopamine release, which is the major determinant of the loss of bidirectional synaptic plasticity at corticostriatal synapses. Indeed, both long-term potentiation (LTP) and long-term depression (LTD) are impaired in these mice, in the absence of overt neuronal degeneration, thereby representing an early pathophysiological event preceding cellular death (14–16).

METHODS

Case-Control Study Population

The study involved 200 consecutive subjects (100 PD patients and 100 sex/age matched controls), afferent to the Neurology Unit of Tor Vergata University Hospital (Rome, Italy). PD was diagnosed according to the United Kingdom PD Society Brain Bank criteria. Controls (CTL) were healthy subjects, without history of neurological diseases or neurological signs at clinical examination, enrolled among non-blood relatives of patients. Exclusion criteria were cognitive decline with Mini-Mental status Examination (MMSE) <25 (adjusted for age and educational level); gastrointestinal disorders and malabsorption; abdominal surgery; diabetes; obesity (BMI > 29); alcoholism; internal

failures (e.g., liver, heart); feeding problems; dietary restrictions; habit to taking vitamins integration. All the participants signed a written informed consent. The study was carried out according to the Declaration of Helsinki and was approved by the local ethical committee (Tor Vergata, Rome—Italy; number 98–09).

Assessment of VEI

All subjects underwent a structured *ad hoc* interview assessing dietary habits: the interview relied on a questionnaire, including explicative pictures to avoid misunderstanding. Subjects were asked how frequently each specific Vitamin E-rich aliment was habitually consumed in the preceding year (2 = more than once a week; 1 = at least once a month; 0 = never). Vitamin E-rich aliments' daily portions were named in a list including fresh fruits (e.g., kiwi, mango), dried fruits (e.g., almonds, walnuts), vegetable (e.g., spinach, broccoli), seeds (e.g., sunflower, pumpkin), oil (e.g., olive, sunflower), fish (e.g., bluefish, crayfish); (source: US Department of Agriculture, USDA (17)). Individual VEI was finally estimated by summing the products of each food's vitamin E content (mg) * the frequency of eating (0, 1, 2). Vitamin E content values were obtained from the Swedish Food Administration Database (18).

Statistical Analysis

The distribution of collected variables was preliminary examined with the Shapiro–Wilk test. Then, the non-normally distributed data were log-transformed to allow statistical analysis. Differences between the groups were tested by parametric (one-way ANOVA) or non-parametric (chi-square) tests, as appropriate. In addition, possible differences in the VEI depending on the H&Y stage and gender were tested by using the one-way ANOVA. The association between PD and VEI was assessed by means binomial logistic regression, adjusting the model for age and gender.

Experimental Electrophysiology on PD Mouse Model

Animal Model and Experimental Setting

Treatment and handling of animals were carried out in accordance with both the EC and Italian guidelines (86/609/EEC; DLS 116/1992, Directive 2010/63/EU; DLS/26 04/03/2014) and were further approved by the University of Rome Tor Vergata statute (n. 153/2001A) and by Animal Care and Use Committee of University of Rome “Tor Vergata.” Transgenic mice (8- to 10-weeks old) were generated as previously described (14).

Intracellular recordings were obtained from striatal neurons in a parasagittal brain slice (300 μ m) (9, 19, 20). A single slice was transferred in a recording chamber (35°C, 2–3 ml/min) and submerged in a continuously flowing Krebs' solution (35°C, 2–3 ml/min) bubbled with 95% O₂ and 5% CO₂. Krebs' solution was composed of (in mM): 126 NaCl, 2.5 KCl, 1.3 MgCl₂, 1.2 NaH₂PO₄, 2.4 CaCl₂, 10 glucose, and 18 NaHCO₃. Intracellular recording electrodes were filled with 2M KCl (30–60 M Ω). To evoke excitatory postsynaptic potentials (EPSPs), a bipolar electrode was placed in the white matter, in close proximity to the recording electrode or in layer VI of the cortex. Test stimuli were delivered at a frequency of 0.1 Hz in the presence

of 50 μM Picrotoxin to block GABA A-mediated responses. The pharmacological effects on EPSPs recorded from knockout mice (*PINK1*^{-/-}) were calculated as percentage of control amplitude in the wild-type (WT) or *PINK1*^{+/+} neuronal population.

For high-frequency stimulation (HFS, three trains 100 Hz, 3 s, 20 s apart), stimulus intensity was raised to reach threshold level. After HFS delivery, the amplitude of EPSPs was plotted over-time as percentage of the control EPSP. Magnesium was omitted from the medium for LTP induction (9, 19, 20).

Signals were recorded with an Axoclamp 2B amplifier (Axon Instruments, Foster City CA 94404, USA), displayed on a oscilloscope and stored on PC using Digidata 1,500 A and pClamp 10.6 (Axon Instruments, Molecular Devices, USA). Data were examined of line by clampfit 10.7 software (Axon Instruments, Molecular Devices, USA). After initial analysis, all data were elaborated by Origin Microcal 2016 (Adalta) software.

Treatments

Effects of Vitamin E (alpha-tocopherol and the water-soluble analog Trolox) were assessed in comparison to other antioxidant agents of alimentary origin, in order to test their action specificity. In particular, for our experiments we specifically selected: beta-carotene (21, 22), lycopene (22, 23), lutein (24), folic acid (25), ascorbic acid/Vitamin C (26), retinol/Vitamin A (27), Vitamin K1/phyloquinone, and Vitamin K2/menaquinones (28, 29).

The effects of drug treatments in *PINK1*^{-/-} mice were tested in two different conditions: (1) *ex-vivo*, by acute preincubation of parasagittal brain slices; (2) *in vivo*, by chronically administered intraperitoneal injections.

For acute treatment, a single slice was incubated, from 40 min before HFS induction and for the duration of the whole experiment (about 1 h), in a bath solution containing the drug dissolved in Krebs' solution. Selected compounds were used in bath at the respective dose of: alpha-tocopherol = 100 μM and Trolox = 100 μM (9); beta-carotene = 100 μM (30); lutein = 20 μM (24); lycopene = 5 μM (22, 23); folic acid = 100 μM (25); Vitamin A = 1 μM (27); Vitamin C = 1–3 mM (26); Vitamin K1 = 20 μM , and Vitamin K2 = 10 μM (28, 29).

For chronic treatments, all compounds were solved in ringer lactate and administered via intraperitoneal injections for 7 days consecutively (9). Dose treatment was: alpha-tocopherol = 100 mg/kg/7 days and Trolox = 5 mg/kg/7 days (9); beta-carotene = 2 mg/kg/7 days (31); lutein = 3 mg/kg/7 days (32); lycopene = 50 mg /kg/7 days (31); folic acid = 2 mg/kg/7 days (33); Vitamin A = 0.5 mg/kg/7 days (31); Vitamin C = 100 mg/kg/7 days (34); Vitamin K2 = 50 mg/kg/7 days (35); Vitamin K1 = 150 mg/kg/7 days (36).

Drug Source

Beta-carotene, lycopene, folic acid, lutein, folic acid, Vitamin A, Vitamin C, Vitamin K1 and K2, alpha-tocopherol, and Trolox were purchased from Sigma-Adrich, Italy. All the other drugs were purchased by Panreac Quimica (Spain).

TABLE 1 | Clinical-demographic parameters and VEI values of the study population.

		PD	CTL	Significance
Gender	(M/F)	55.2%	44.8%	46.9% 53.1% ns
Age (years)	Mean	63.3	60.5	
	St.dev.	8.5	10.6	
Log10 Age	Mean	1.80	1.77	ns
	St.dev.	0.06	0.08	
VEI (mg)	Mean	31.6	38.4	
	St.dev.	13.5	17.8	
Log10 VEI	Mean	1.45	1.54	$p < 0.05$
	St.dev.	0.21	0.21	
H&Y	Mean	2.4	–	
	St.dev.	0.7	–	

Statistical Analysis

Data are presented as mean \pm standard error of the mean (SEM). Statistical significance between pre and post HFS stimulation was evaluated using Student *T*-test. Percentage values were calculated for each individual experiment. An analysis of variance with the Tukey's *post-hoc* test was performed among the groups ($P < 0.05$; alpha = 0.01). Statistical significance was set at <0.05 .

RESULTS


Case-Control Study

Clinical-demographic parameters and the VEI of the study population are summarized in **Table 1**. PD and CTL were homogeneous in age and gender distribution; VEI was significantly higher in CTL (38.4 mg \pm 17.8) than PD (31.6 mg \pm 13.5; Statistical analysis was conducted on Log-transformed values, resulting $p < 0.05$). Conversely, VEI did not differ depending on the gender in both groups, neither among the stages of H&Y in PD patients. The binomial logistic regression showed that VEI was directly associated with CTL status, independently from age and gender (Odd Ratio, OR = 1.022; 95% CI = 0.999–1.045; $p < 0.05$).

Electrophysiology in PD Mouse Model

According to our previous findings (14, 16), HFS protocol performed on parasagittal slice preparation induced a robust LTD in MSNs recorded from WT mice (58.24 \pm 3.79% of control; $n = 16$ **Figure 1A**), whereas it failed to elicit LTD in *PINK1*^{-/-} mice (99.61 \pm 2.88% of control; $n = 22$ $p < 0.05$ *t*-test **Figure 1A**). After removal of magnesium from the bathing medium, HFS induced LTP in WT mice (168.26 \pm 5.21% of control; $n = 12$; *t*-test $p < 0.05$ **Figure 1B**). In *PINK1*^{-/-} mice, HFS also increased EPSPs compared to pre HFS (123.39 \pm 4.88% of control; $n = 12$; *t*-test $p < 0.05$, **Figure 1B**), but the magnitude was significantly lower than WT mice ($p < 0.05$ ANOVA), suggesting the impairment of this form of plasticity.

None of the other drugs, but Vitamin E, was able to rescue either LTD or LTP in both acute and chronic treatment



(>5 observations for each experimental condition; *T*-test, $p > 0.05$; **Table 2**). Specifically, Trolox (the water-soluble analog of Vitamin E) fully rescued both the forms of synaptic plasticity either after acute or chronic administration (LTD protocol: $55.88 \pm 6.01\%$ of control; $n = 16$; *t*-test $p < 0.05$ **Figure 1C**; LTP protocol: $175.30 \pm 4.88\%$ of control; $n = 10$; *t*-test $p < 0.05$ **Figure 1D**, **Table 2**); alpha-tocopherol restored in chronic conditions (intraperitoneal injection) (LTD protocol: $56.9 \pm 4.21\%$ of control; $n = 8$; *t*-test $p < 0.05$; LTP protocol: $177.2 \pm 6.36\%$ of control; $n = 6$; *t*-test $p < 0.05$, **Table 2**).

DISCUSSION

In this study, both the clinical retrospective analysis and our electrophysiological experiments demonstrate that Vitamin E might exert potential beneficial effects in PD.

The case-control analysis showed that dietary VEI is higher in healthy subjects than age/sex matched PD patients. Such a reduced intake in PD patients might suggest a lack of its putative protective action, independently from age and gender. Although consistent with other data from larger and prospective cohorts (17, 18), a number of limitations should be considered in the interpretation of the result, such as the sample size, the recall bias, and the absence of accurate measurement for the dietary intake. In fact, VEI was just approximately estimated by a retrospective *ad hoc* questionnaire, scoring how frequently the standard portions of aliments with Vitamin E higher content were assumed in the last year, and not precisely quantified. However, to prevent confounding factors due to the occurrence of the disease, we excluded from the study patients with alimentary restrictions (e.g., dysphagia), dementia or any concomitant condition affecting feeding behavior or

TABLE 2 | The table summarizes the effects of acute and chronic treatment of every single compound on corticostriatal synaptic plasticity.

Drug	Administration	Rescue LTD	Rescue LTP
Beta-carotene	Acute	NO	NO
	Chronic		
Lutein	Acute	NO	NO
	Chronic		
Lycopene	Acute	NO	NO
	Chronic		
Folic acid	Acute	NO	NO
	Chronic		
Vitamin A	Acute	NO	NO
	Chronic		
Vitamin C	Acute	NO	NO
	Chronic		
Vitamin K2	Acute	NO	NO
	Chronic		
Vitamin K1	Acute	NO	NO
	Chronic		
Alpha-tocopherol (Vitamin E)	Acute	NO	NO
	Chronic	YES	YES
Trolox (vitamin E)	Acute	YES	YES
	Chronic	YES	YES

Only alpha-tocopherol and Trolox resulted effective. Bold highlights drugs restoring synaptic plasticity.

intestinal absorption. Moreover, the homogeneous distribution of demographic features in the study population and the statistical methodology might have limited the influence of other potential confounding factors (e.g., sex/age-based dietary choices, sex/age-dependent differences in internal metabolism). In addition to confirm previous findings (17, 18), here we noticed that VEI did not correlate with severity of PD, assessed by H&Y score. Actually, the sample size and the exclusion from the model of other clinical determinants (e.g., disease duration, therapy) might represent a bias; indeed, it is possible that both levodopa dose and disease duration, which are usually higher in more advanced patients, may affect vitamin absorption (37), causing some deficiency unrelated to the alimentary habits. Also the occurrence and severity of constipation, which in turn might influence dietary choices, pharmacotherapy and intestinal function (38), has not been addressed in the study. Therefore, caution is required in the interpretation of our preliminary results. To this regard, it should be mentioned that other authors, also by utilizing recall-based questionnaires, excluded significant associations between alimentary VEI and PD (39, 40); these studies differed in sample size, but were performed out of Mediterranean area. Hence, we should consider regional diet as a further potential confounding factor. Certainly, prospective cohort studies, eventually supported by direct vitamin dosage, are necessary to assess the weight of VEI in PD pathogenesis.

The protective action of Vitamin E on PD has been further explored by using an experimental model of preclinical PD.

PINK1^{-/-} mice indeed exhibit the disruption of bidirectional plasticity at corticostriatal synapses, even in the absence of overt neurodegeneration. Several studies indicate this model as representative of a critical time-window of the disease in which a specific intervention may revert the pathophysiological cascade leading to symptoms onset, being thus appropriate to test the efficacy of disease-modifying strategies (3, 14–16). Our experiments show that the administration of Vitamin E (alpha-tocopherol and Trolox), but not other dietary antioxidant compounds (beta-carotene, lycopene, lutein, folic acid, ascorbic acid/Vitamin C, retinol/Vitamin A, Vitamin K1/phyloquinone, and Vitamin K2/menaquinones), was able to revert synaptic plasticity abnormalities in *PINK1*^{-/-} mice. *PINK1* haploinsufficiency precipitates mitochondrial functioning, impairing mitophagy, and above all, energy production under increased demand (41). This, in turn, accounts for the reduced synaptic vesicle release at dopaminergic terminals and the subsequent breakdown of corticostriatal synaptic plasticity (9, 14, 16). It is thus conceivable that Vitamin E, unlike other vitamins, specifically rescues striatal homeostasis and neurotransmission in *PINK1*^{-/-} mice, by enhancing mitochondrial metabolism (42, 43) and energy-dependent processes. Indeed, it has been recently demonstrated that Vitamin E, but not other antioxidants, such as Vitamin C, fully rescued longevity in a short-lived *Candida elegans gas-1(fc21)* model of respiratory chain complex I defect (44). Moreover, in other experimental models, Vitamin E resulted to be able to activate cellular pathways involved in antioxidant, detoxifying, and anti-inflammatory responses and to promote bioenergy at mitochondrial level (45, 46). Regarding *PINK1*^{-/-} rodents model, Shim and colleagues demonstrated that Trolox dramatically improved mitochondrial metabolism in *PINK1*-deficient dopaminergic cells, by increasing complex I and complex IV's activity (47). Because neurotransmitter release depends on mitochondrial bioenergetics (48), we hypothesize a recovery of a physiological dopaminergic transmission with the subsequent rescue of corticostriatal plasticity. However, a specific set of experiments is required to assess this issue in *PINK1*^{-/-} mice. Yet, the complex interactions between *PINK1*-mediated mitochondrial activities and Vitamin E-induced cellular reactions (49, 50) could then explain the inefficacy of other proved alimentary antioxidants in restoring corticostriatal synaptic plasticity.

Since mitochondrial dysfunction is critical also in pathogenesis of idiopathic PD (41, 51, 52), such a mechanism may justify a beneficial action of higher dietary VEI on PD in humans. Furthermore, Vitamin E seems to intervene on other pathogenic pathways of PD, such as lysosome metabolism (53), expanding potential restorative effects. Definitely, larger studies are mandatory to validate this hypothesis.

Regardless the limitations, our findings suggest a potential protective action of a Vitamin E rich diet. These data may indicate that Vitamin E represents a potential therapeutic target for disease-modifying treatments in PD. Therefore, diets including Vitamin E rich aliments could be an immediate option to reduce the risk of PD and other neurodegenerative

diseases (54), although specific confirmatory trials are necessary.

DATA AVAILABILITY

The datasets generated for this study are available on request to the corresponding author.

AUTHOR CONTRIBUTIONS

TS, GM, and AP conceived the study and wrote the manuscript. GD, VC, DF, and MA collected clinical data. GM and PI performed the experiments. TS and MP performed statistical

analysis. NM, MP, and PS contributed to interpretation of results and edited the manuscript.

ACKNOWLEDGMENTS

The study was partially funded by a Grant from Fondazione Baroni, Rome—Italy (to AP, 2016) and PRIN-MIUR 2015FNWP34 (to AP, 2015). We thank Mr. Massimo Tolu for his excellent technical support. We acknowledge the Master of Science in Physical Activity and Health Promotion and the Department of Biomedicine and Prevention, University of Rome Tor Vergata (Rome—Italy) for the support.

REFERENCES

- Schirinzi T, Martella G, Elia AD, Di Lazzaro G, Imbriani P, Madeo G, et al. Outlining a population (at risk) of Parkinson's disease: evidence from a case-control study. *Parkinsons Dis.* (2016) 2016:9646057. doi: 10.1155/2016/9646057
- Kalia LV, Lang AE. Parkinson's disease. *Lancet* (2015) 386:896–912. doi: 10.1016/S0140-6736(14)61393-3
- Schirinzi T, Madeo G, Martella G, Maltese M, Picconi B, Calabresi P, et al. Early synaptic dysfunction in Parkinson's disease: insights from animal models. *Mov Disord.* (2016) 31:802–13. doi: 10.1002/mds.26620
- Lang AE, Espay AJ. Disease modification in Parkinson's disease: current approaches, challenges, and future considerations. *Mov Disord.* (2018) 33:660–77. doi: 10.1002/mds.27360
- Jin H, Kanthasamy A, Ghosh A, Anantharam V, Kalyanaraman B, Kanthasamy AG. Mitochondria-targeted antioxidants for treatment of Parkinson's disease: preclinical and clinical outcomes. *Biochim Biophys Acta Mol Basis Dis.* (2014) 1842:1282–94. doi: 10.1016/j.bbadis.2013.09.007
- Surmeier DJ, Obeso JA, Halliday GM. Parkinson's disease is not simply a prion disorder. *J Neurosci.* (2017) 37:9799–807. doi: 10.1523/JNEUROSCI.1787-16.2017
- Crotty GF, Ascherio A, Schwarzschild MA. Targeting urate to reduce oxidative stress in Parkinson disease. *Exp Neurol.* (2017) 298(Pt B):210–24. doi: 10.1016/j.expneurol.2017.06.017
- Imbriani P, Schirinzi T, Meringolo M, Mercuri NB, Pisani A. Centrality of early synaptopathy in Parkinson's disease. *Front Neurol.* (2018) 9:103. doi: 10.3389/fneur.2018.00103
- Martella G, Madeo G, Maltese M, Vanni V, Puglisi F, Ferraro E, et al. Neurobiology of Disease Exposure to low-dose rotenone precipitates synaptic plasticity alterations in PINK1 heterozygous knockout mice. *Neurobiol Dis.* (2016) 91:21–36. doi: 10.1016/j.nbd.2015.12.020
- Thapa A, Carroll NJ. Molecular sciences dietary modulation of oxidative stress in Alzheimer's disease. *Int J Mol Sci.* (2017) 18:1583. doi: 10.3390/ijms18071583
- Jiang Q. Natural forms of vitamin E: metabolism, antioxidant, and anti-inflammatory activities and their role in disease prevention and therapy. *Free Radic Biol Med.* (2014) 72:76–90. doi: 10.1016/j.freeradbiomed.2014.03.035
- Kaneai N, Arai M, Takatsu H, Fukui K, Urano S. Vitamin E inhibits oxidative stress-induced denaturation of nerve terminal proteins involved in neurotransmission. *J Alzheimer's Dis.* (2012) 28:183–9. doi: 10.3233/JAD-2011-111133
- Ulatowski LM, Manor D. Vitamin E and neurodegeneration. *Neurobiol Dis.* (2015) 84:78–83. doi: 10.1016/j.nbd.2015.04.002
- Kitada T, Pisani A, Porter DR, Yamaguchi H, Tschertner A, Martella G, et al. Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice. *Proc Natl Acad Sci USA.* (2007) 104:11441–6. doi: 10.1073/pnas.0702717104
- Imbriani P, Sciamanna G, Santoro M, Schirinzi T, Pisani A. Promising rodent models in Parkinson's disease. *Parkinsonism Relat Disord.* (2018) 46:S10–4. doi: 10.1016/j.parkreldis.2017.07.027
- Madeo G, Schirinzi T, Martella G, Latagliata EC, Puglisi F, Shen J, et al. PINK1 heterozygous mutations induce subtle alterations in dopamine-dependent synaptic plasticity. *Mov Disord.* (2014) 29:41–53. doi: 10.1002/mds.25724
- Zhang SM, Hernán MA, Chen H, Spiegelman D, Willett WC, Ascherio A. Intakes of vitamins E and C, carotenoids, vitamin supplements, and PD risk. *Neurology* (2002) 59:1161–9.
- Yang F, Wolk A, Håkansson N, Pedersen NL, Wirdefeldt K. Dietary antioxidants and risk of Parkinson's disease in two population-based cohorts. *Mov Disord.* (2017) 32:1631–6. doi: 10.1002/mds.27120
- Martella G, Tassone A, Sciamanna G, Platania P, Cuomo D, Viscomi MT, et al. Impairment of bidirectional synaptic plasticity in the striatum of a mouse model of DYT1 dystonia: role of endogenous acetylcholine. *Brain* (2009) 132:2336–49. doi: 10.1093/brain/awp194
- Maltese M, Stanic J, Tassone A, Sciamanna G, Ponterio G, Vanni V, et al. Early structural and functional plasticity alterations in a susceptibility period of DYT1 dystonia mouse striatum. *Elife* (2018) 7:33331. doi: 10.7554/eLife.33331
- Chang C-S, Chang C-L, Lai G-H. Reactive oxygen species scavenging activities in a chemiluminescence model and neuroprotection in rat pheochromocytoma cells by astaxanthin, beta-carotene, and canthaxanthin. *Kaohsiung J Med Sci.* (2013) 29:412–21. doi: 10.1016/j.kjms.2012.12.002
- Di Tomo P, Canali R, Ciavardelli D, Di Silvestre S, De Marco A, Giardinelli A, et al. β -Carotene and lycopene affect endothelial response to TNF- α reducing nitro-oxidative stress and interaction with monocytes. *Mol Nutr Food Res.* (2012) 56:217–27. doi: 10.1002/mnfr.201100500
- Qu M, Jiang Z, Liao Y, Song Z, Nan X. Lycopene prevents amyloid [Beta]-induced mitochondrial oxidative stress and dysfunctions in cultured rat cortical neurons. *Neurochem Res.* (2016) 41:1354–64. doi: 10.1007/s11064-016-1837-9
- Li S-Y, Lo ACY. Lutein protects RGC-5 cells against hypoxia and oxidative stress. *Int J Mol Sci.* (2010) 11:2109–17. doi: 10.3390/ijms11052109
- Zhu M, Li B, Ma X, Huang C, Wu R, Zhu W, et al. Folic acid protected neural cells against aluminum-maltolate-induced apoptosis by preventing miR-19 downregulation. *Neurochem Res.* (2016) 41:2110–8. doi: 10.1007/s11064-016-1926-9
- Harrison FE, May JM. Vitamin C function in the brain: vital role of the ascorbate transporter SVCT2. *Free Radic Biol Med.* (2009) 46:719–30. doi: 10.1016/j.freeradbiomed.2008.12.018
- Kona SL, Shrestha A, Yi X, Joseph S, Barona HM, Martinez-Ceballos E. RAR β -dependent signaling represses neuronal differentiation in mouse ES cells. *Differentiation* (2017) 98:55–61. doi: 10.1016/j.diff.2017.11.002
- Sakaue M, Mori N, Okazaki M, Kadowaki E, Kaneko T, Hemmi N, et al. Vitamin K has the potential to protect neurons from methylmercury-induced cell death *in vitro*. *J Neurosci Res.* (2011) 89:1052–8. doi: 10.1002/jnr.22630

29. Nakajima M, Furukawa S, Hayashi K, Yamada A, Kawashima T, Hayashi Y. Age-dependent survival-promoting activity of vitamin K on cultured CNS neurons. *Brain Res Dev Brain Res.* (1993) 73:17–23.
30. Lin H-W, Chang T-J, Yang D-J, Chen Y-C, Wang M, Chang Y-Y. Regulation of virus-induced inflammatory response by β -carotene in RAW264.7 cells. *Food Chem.* (2012) 134:2169–75. doi: 10.1016/j.foodchem.2012.04.024
31. Glise D, Riondel J, Favier A. Comparative distribution of beta-carotene and lycopene after intraperitoneal administration in mice. *In Vivo* (1998) 12:447–54.
32. Li S-Y, Yang D, Fu ZJ, Woo T, Wong D, Lo ACY. Lutein enhances survival and reduces neuronal damage in a mouse model of ischemic stroke. *Neurobiol Dis.* (2012) 45:624–32. doi: 10.1016/j.nbd.2011.10.008
33. Gupta A, Puri V, Sharma R, Puri S. Folic acid induces acute renal failure (ARF) by enhancing renal prooxidant state. *Exp Toxicol Pathol.* (2012) 64:225–32. doi: 10.1016/j.etp.2010.08.010
34. Sahraei H, Aliabadi AA, Zarrindast M-R, Ghoshooni H, Nasiri A, Barzegari-Sorkheh AA, et al. Ascorbic acid antagonizes nicotine-induced place preference and behavioral sensitization in mice. *Eur J Pharmacol.* (2007) 560:42–8. doi: 10.1016/j.ejphar.2006.12.019
35. Onodera K, Shinoda H, Zushida K, Taki K, Kamei J. Antinociceptive effect induced by intraperitoneal administration of vitamin K2 (menatetrenone) in ICR mice. *Life Sci.* (2000) 68:91–7.
36. Sakamoto N, Kimura M, Hiraike H, Itokawa Y. Changes of phyloquinone and menaquinone-4 concentrations in rat liver after oral, intravenous and intraperitoneal administration. *Int J Vitam Nutr Res.* (1996) 66:322–8.
37. Rajabally YA, Martey J, Levodopa, vitamins, ageing and the neuropathy of Parkinson's disease. *J Neurol.* (2013) 260:2844–8. doi: 10.1007/s00415-013-7079-8
38. Barichella M, Cereda E, Cassani E, Pinelli G, Iorio L, Ferri V, et al. Dietary habits and neurological features of Parkinson's disease patients: implications for practice. *Clin Nutr.* (2017) 36:1054–61. doi: 10.1016/j.clnu.2016.06.020
39. Hughes KC, Gao X, Kim IY, Rimm EB, Wang M, Weisskopf MG, et al. Intake of antioxidant vitamins and risk of Parkinson's disease. *Mov Disord.* (2016) 31:1909–14. doi: 10.1002/mds.26819
40. Morens DM, Grandinetti A, Waslien CI, Park CB, Ross GW, White LR. Case-control study of idiopathic Parkinson's disease and dietary vitamin E intake. *Neurology* (1996) 46:1270–4.
41. Pickrell AM, Youle RJ. The roles of PINK1, parkin, and mitochondrial fidelity in Parkinson's disease. *Neuron* (2015) 85:257–73. doi: 10.1016/j.neuron.2014.12.007
42. Duan L, Li J, Ma P, Yang X, Xu S. Vitamin E antagonizes ozone-induced asthma exacerbation in Balb/c mice through the Nrf2 pathway. *Food Chem Toxicol.* (2017) 107:47–56. doi: 10.1016/j.fct.2017.06.025
43. Vineetha RC, Binu P, Arathi P, Nair RH. L-ascorbic acid and α -tocopherol attenuate arsenic trioxide-induced toxicity in H9c2 cardiomyocytes by the activation of Nrf2 and Bcl2 transcription factors. *Toxicol Mech Methods* (2018) 28:353–60. doi: 10.1080/15376516.2017.1422578
44. Polyak E, Ostrovsky J, Peng M, Dingley SD, Tsukikawa M, Kwon YJ, et al. N-acetylcysteine and vitamin E rescue animal longevity and cellular oxidative stress in pre-clinical models of mitochondrial complex I disease. *Mol Genet Metab.* (2018) 123:449–62. doi: 10.1016/j.ymgme.2018.02.013
45. Petrillo S, Piermarini E, Pastore A, Vasco G, Schirinzi T, Carrozzo R, et al. Nrf2-inducers counteract neurodegeneration in frataxin-silenced motor neurons: disclosing new therapeutic targets for friedreich's ataxia. *Int J Mol Sci.* (2017) 18:2173. doi: 10.3390/ijms18102173
46. Dinkova-Kostova AT, Abramov AY. The emerging role of Nrf2 in mitochondrial function. *Free Radic Biol Med.* (2015) 88:179–88. doi: 10.1016/j.freeradbiomed.2015.04.036
47. Shim JH, Yoon SH, Kim K-H, Han JY, Ha J-Y, Hyun DH, et al. The antioxidant Trolox helps recovery from the familial Parkinson's disease-specific mitochondrial deficits caused by PINK1- and DJ-1-deficiency in dopaminergic neuronal cells. *Mitochondrion* (2011) 11:707–15. doi: 10.1016/j.mito.2011.05.013
48. Madeo G, Martella G, Schirinzi T, Ponterio G, Shen J, Bonsi P, et al. Aberrant striatal synaptic plasticity in monogenic parkinsonisms. *Neuroscience* (2012) 211:126–35. doi: 10.1016/j.neuroscience.2011.07.065
49. Murata H, Takamatsu H, Liu S, Kataoka K, Huh N, Sakaguchi M. NRF2 regulates PINK1 expression under oxidative stress conditions. *PLoS ONE* (2015) 10:e0142438. doi: 10.1371/journal.pone.0142438
50. Buendia I, Michalska P, Navarro E, Gameiro I, Egea J, León R. Nrf2-ARE pathway: an emerging target against oxidative stress and neuroinflammation in neurodegenerative diseases. *Pharmacol Ther.* (2016) 157:84–104. doi: 10.1016/j.pharmthera.2015.11.003
51. Tufekci KU, Civi Bayin E, Genc S, Genc K. The Nrf2/are pathway: a promising target to counteract mitochondrial dysfunction in Parkinson's disease. *Parkinsons Dis.* (2011) 2011:314082. doi: 10.4061/2011/314082
52. Holmström KM, Kostov RV, Dinkova-Kostova AT. The multifaceted role of Nrf2 in mitochondrial function. *Curr Opin Toxicol.* (2017) 2:80–91. doi: 10.1016/j.cotox.2016.10.002
53. Marin T, Contreras P, Castro J, Chamorro D, Balboa E, Bosch-Morató M, et al. Vitamin E dietary supplementation improves neurological symptoms and decreases c-Abl/p73 activation in niemann-pick C mice. *Nutrients* (2014) 6:3000–17. doi: 10.3390/nu6083000
54. Barnard ND, Bush AI, Ceccarelli A, Cooper J, de Jager CA, Erickson KI, et al. Dietary and lifestyle guidelines for the prevention of Alzheimer's disease. *Neurobiol Aging* (2014) 35:S74–8. doi: 10.1016/j.neurobiolaging.2014.03.033

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Schirinzi, Martella, Imbriani, Di Lazzaro, Franco, Colona, Alwardat, Sinibaldi Salimei, Mercuri, Pierantozzi and Pisani. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.