



Thigh Ischemia-Reperfusion Model Does Not Accelerate Pulmonary VO_2 Kinetics at High Intensity Cycling Exercise

Lucas Helal^{1*}, Paulo Cesar do Nascimento Salvador², Ricardo Dantas de Lucas² and Luiz Guilherme Antonacci Guglielmo²

¹ Exercise Pathophysiology Laboratory, Hospital de Clínicas de Porto Alegre, Graduate Program in Cardiology and Cardiovascular Sciences, School of Medicine, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil,

² Physical Effort Laboratory, Graduate Program in Biodynamics and Human Performance, Universidade Federal de Santa Catarina, Florianópolis, Brazil

OPEN ACCESS

Edited by:

Keith Russell Brunt,
Dalhousie University, Canada

Reviewed by:

Joanna E. MacLean,
University of Alberta, Canada
Trevor James King,
University of Guelph, Canada

*Correspondence:

Lucas Helal
lucas.helal@ufrgs.br
orcid.org/0000-0002-6900-7185

Specialty section:

This article was submitted to
Respiratory Physiology,
a section of the journal
Frontiers in Physiology

Received: 20 April 2018

Accepted: 08 February 2019

Published: 25 February 2019

Citation:

Helal L, do Nascimento Salvador PC, de Lucas RD and Guglielmo LGA (2019) Thigh Ischemia-Reperfusion Model Does Not Accelerate Pulmonary VO_2 Kinetics at High Intensity Cycling Exercise. *Front. Physiol.* 10:160. doi: 10.3389/fphys.2019.00160

Background: We aimed to investigate the effect of a priming ischemia-reperfusion (IR) model on the kinetics of pulmonary oxygen uptake (VO_2) and cardiopulmonary parameters after high-intensity exercise. Our primary outcome was the overall VO_2 kinetics and secondary outcomes were heart rate (HR) and O_2 pulse kinetics. We hypothesized that the IR model would accelerate VO_2 and cardiopulmonary kinetics during the exercise.

Methods: 10 recreationally active men (25.7 ± 4.7 years; 79.3 ± 10.8 kg; 177 ± 5 cm; 44.5 ± 6.2 mL kg^{-1} min^{-1}) performed a maximal incremental ramp test and four constant load sessions at the midpoint between ventilatory threshold and VO_2 max on separate days: two without IR (CON) and two with IR (IR). The IR model consisted of a thigh bi-lateral occlusion for 15 min at a pressure of 250 mmHg, followed by 3 min off, before high-intensity exercise bouts.

Results: There were no significant differences for any VO_2 kinetics parameters (VO_2 base 1.08 ± 0.08 vs. 1.12 ± 0.06 L min^{-1} ; $P = 0.30$; $\tau = 50.1 \pm 7.0$ vs. 47.9 ± 6.4 s; $P = 0.47$), as well as for HR (MRT_{180s} 67.3 ± 6.0 vs. 71.3 ± 6.1 s; $P = 0.54$) and O_2 pulse kinetics (MRT_{180s} 40.9 ± 3.9 vs. 48.2 ± 5.6 s; $P = 0.31$) between IR and CON conditions, respectively.

Conclusion: We concluded that the priming IR model used in this study had no influence on VO_2 , HR, and O_2 pulse kinetics during high-intensity cycling exercise.

Keywords: oxygen uptake kinetics, ischemia-reperfusion, cardiopulmonary test, physical exercise, exercise physiology

INTRODUCTION

Pulmonary oxygen uptake (VO_2) kinetics plays a role in the oxidative metabolism during constant load exercise (Poole et al., 2008). Considering that the area above the curve is accepted as a representative surrogate of the required additional energy supplying – i.e., anaerobic energy sources (Margarita et al., 1963), any acute or chronic adjustments that could reduce the time to attain a steady state of oxygen uptake and improves the energy imbalance becomes relevant for aerobic exercise performance (Gastin, 2001) or health related outcomes (Poole and Jones, 2012).

Several methods have been used to acutely challenge the VO₂ control system and to enhance VO₂ kinetics, such as breathing hyperoxic gas (MacDonald et al., 1998, 2000); prior exercise (Germino et al., 1996; Burnley et al., 2001, 2002) a reactive hyperemia induced by ischemia-reperfusion (IR) (Walsh et al., 2002; Faisal et al., 2010) – Which have resulted in controversial findings. The IR model involves blocking arterial to the target site (usually through a cuff pressure device), for a given period (continuously or intermittently). Upon cuff release, blood flow is augmented well above resting values (Kooijman et al., 2008). The brief exposure of muscle tissue to a blood flow absence causes an immediate reduction of the local O₂ saturation, followed by a reduced adenosine triphosphate (ATP) and phosphocreatine (PCr) intramuscular concentrations, and an increase in adenosine diphosphate (ADP) and inorganic phosphate (Pi) intramuscular concentrations (Boushel et al., 1998). On the very end, an increase in vascular dilation (Enko et al., 2011), a potential increase in local blood flow, O₂ muscle saturation and cardiac function (Daly and Bondurant, 1962; Faisal et al., 2010) is expected, which may accelerate VO₂ kinetics by improving O₂ delivery and availability. Nonetheless, the blood flow restriction (e.g., increases in ADP, Pi, etc.) may affect the VO₂ time course through spin-off manifestations, likely improvements in the mitochondrial oxidation activity through the increases in ADP and Pi concentrations (Boushel et al., 1998; Walsh et al., 2002). Despite the physiological rationale, the IR model has received little attention to study VO₂ kinetics at the onset of exercise.

Few studies were conducted with contradictory and inconclusive findings, like an impairment (Faisal et al., 2010), no effect (Kido et al., 2015) or an improvement (Walsh et al., 2002) in the pulmonary VO₂ kinetics. At high intensity exercise, VO₂ kinetics appears to be sensitive to the fundamental component amplitude (A_p) (Walsh et al., 2002; Wilkerson et al., 2006), slow component amplitude (A_s) (MacDonald et al., 1998; Burnley et al., 2002; Walsh et al., 2002), time-constant of fundamental component (τ_p) (MacDonald et al., 1998) and the “overall” kinetics (Walsh et al., 2002; Tordi et al., 2003; Nascimento et al., 2015). Also, VO₂ kinetics is dependent on systemic and peripheral mechanisms, such as increased activity of oxidative enzymes, muscle blood flow and improved O₂ distribution in the exercise muscles (Burnley et al., 2002; Wilkerson et al., 2006; Faisal et al., 2010; Poole and Jones, 2012). Notwithstanding, there is a question that still provokes further investigation for the physiologists: is the VO₂ kinetics at the onset of exercise limited by muscle O₂ delivery or by an oxidative enzyme inertia? In that scenario the reactive hyperemia by occluding target members, then augmenting muscle blood flow for several minutes prior to a subsequent exercise bout (Faisal et al., 2010), with less pronounced changes in the metabolic environment (i.e., peripheral mechanisms) seems an innovative approach.

Finally, it has been suggested that O₂ delivery does not limit the phase II of VO₂ kinetics in healthy subjects during a high intensity exercise (Poole et al., 2008), at least in some instances. However, at some “tipping point,” an O₂ limitation could occur and VO₂ kinetics can be modified through alterations in O₂ availability. Specifically, high-intensity approaches the tipping

point, where VO₂ kinetics is O₂-delivery dependent and may potentially be more susceptible to the influence of blood-muscle O₂ influx (Poole and Jones, 2012). Thus, this study aims to investigate the effect of a priming IR protocol prior to high-intensity exercise on pulmonary VO₂ kinetics and cardiac parameters (heart rate and O₂ pulse). We hypothesized that a prior IR stimulus would accelerate VO₂ and cardiac related kinetics at the onset of high-intensity cycling.

MATERIALS AND METHODS

Participants

Ten recreationally active men (age 25 ± 5 years; body mass 79.3 ± 11 kg; height 177 ± 5 cm; VO₂ max 44.5 ± 6.2 mL kg⁻¹ min⁻¹) volunteered to participate in this study. Participants were classified as physically active by the American College of Sports Medicine (ACSM) criteria (cardiorespiratory exercise training for ≥ 30 min-days⁻¹ on ≥ 5 days-week⁻¹ for a total of ≥ 150 min-week⁻¹) (Garber et al., 2011). Some of them were involved in further recreational sport activities but not in any systematic physical exercise program. Participants were informed about risks and possible discomfort related to the protocol study. After that, a written informed consent was given before starting the study.

The subjects were instructed to avoid any intake of caffeine for 3 h, or alcohol and strenuous exercise in the 24 h preceding the test sessions. They were also instructed to arrive at the laboratory in a rested and fully hydrated state, and to eat the last meal up to 2 h prior to the experiment. All tests were conducted at the same time of day in controlled environmental laboratory conditions (19–22°C; 50–60% RH), to minimize the effects of diurnal biological variation on the results. All procedures were performed in accordance to the Declaration of Helsinki statement and were reviewed and approved by the Institutional Research Board (Ethics Committee for Research in Human Beings, Universidade Federal de Santa Catarina, IRB number 51045315.7.0000.0121).

Experimental Design

The subjects visited the laboratory on five occasions. In the first visit, participants underwent a maximal incremental cycling test to determine intensity parameters for the experimental protocol (Visit 1); the other four experimental visits were randomly assigned, and separated by at least 48 h.

The experimental trials consisted of 3 min of baseline cycling at 20 W plus 8 min at $\Delta 50\%$ intensity. Gas exchange was collected throughout exercise bouts to analyze VO₂, and O₂ pulse kinetics. Heart rate (HR) was measured during the entire exercise bouts as well. In two of the four trials, cycling exercise was preceded by 15 min bilateral thigh blood flow occlusion, whereas two trials without IR were conducted as control sessions (**Figure 1**).

VO₂ max, GET and $\Delta 50\%$ Determination

To determine physiological indexes for exercise sessions, we conducted the maximal incremental ramp test on a cycle ergometer (Lode Excalibur, Groningen, Netherlands). Participants exercised for 4 min at a constant load of 20 W

to stabilize gas measurements and then the exercise intensity increased systematically by 1 W every 2 s (rate of 30 W min⁻¹) until volitional exhaustion.

Oxygen uptake was measured breath-by-breath using a gas analyzer (Quark PFT Ergo, COSMED, Rome, Italy). The gas analyzer was calibrated immediately before each test using ambient air and certified alpha standard gasses containing 16.0% of oxygen and 5.0% of carbon dioxide. The turbine flowmeter was calibrated with a syringe of three liters (Quark PFT Ergo, COSMED, Rome, Italy). HR was continuously recorded by a HR monitor integrated into the gas analyzer system. Earlobe capillary blood lactate samples (25 μL) were taken before and immediately after the incremental exercise from the ear lobe and analyzed by an electrochemical method (YSI 2700 STAT, Yellow Springs, OH, United States).

Maximal oxygen uptake was defined by two criteria, depending on the presence of a VO₂ plateau. The VO₂ plateau was defined as any 60 s period in which the VO₂ stabilized without increases higher than 150 mL min⁻¹. In the absence of a VO₂ plateau, we considered the maximum oxygen uptake as VO₂ max, by the maximal observed value in a 15 s interval (Day et al., 2003). Maximal power output (W_{peak}), maximal heart rate (HR_{max}) and maximal lactate were considered as the highest values observed during exercise. Gas exchange threshold (GET) was calculated using the V-slope method (Beaver et al., 1986). The increase in VO₂ ventilatory equivalent (VE/VO₂) with no further increases in VCO₂ ventilatory equivalent (VE/VCO₂) was verified through visual inspection and used as a secondary criterion.

The Δ50% exercise intensity was calculated by defining GET-VO₂ and VO₂ max, and then adding 50% of the difference between them to GET-VO₂. After this, we made an interpolation using the linear regression model with the equation of a VO₂/intensity relationship (Souza et al., 2016), in order to determine the Δ50% workload.

VO₂, HR, and O₂ Pulse Kinetics

Each testing session consisted of a square-wave transition of 3 min at 20 W followed by 8 min at Δ50% intensity. Earlobe capillary blood lactate samples were collected 30 s before the baseline ([Lac]_{rest}), 30 s before exercise ([Lac]_{pre}) and immediately after the exercise ([Lac]_{post}). O₂ pulse was determined as the fraction between VO₂ and HR (Whipp et al., 1996), both interpolated by 1 s intervals after matching the data of the two identical sessions corresponding to IR and CON conditions.

Data Acquisition

Oxygen uptake and HR were recorded breath-by-breath during the entire exercise sessions (Quark PFT Ergo, COSMED, Rome, Italy) and were exported as raw data for filtering and analysis (OriginPro 7.0, United States). For VO₂, signals suggesting sighs and coughs were initially removed from each test. According to Lamarra et al. (1987), occasional breath values were excluded from analysis using three standard deviations from the local mean as the criterion. Breath-by-breath data were linearly interpolated by 1 s intervals and then time-aligned to the start of exercise. After this procedure, both data series of each condition

were matched and averaged in order to provide a unique profile for each exercise condition. The last filtering procedure was applied to reduce interpolated data to a 5 s stationary mean (Rossiter et al., 2005). The cardiopulmonary phase (i.e., first 20 s of exercise) was not included in the analysis (Rossiter et al., 2005).

For modeling the VO₂ profile, a non-linear regression technique was applied (Rossiter et al., 2002), where phase II and a slow component were fitted separately. The phase II was modeled from the end of the cardio-dynamic phase to the beginning of the slow component phase. The slow component phase was determined by four criteria: (1) the narrowest confidence interval for τ; (2) breakpoint and the systematic rising of phase II amplitude and time-constant with a decrease in TD; (3) breakpoint and a systematic rising of fitting R²; (4) visual inspection of residual plot, considering the end-phase by the farthest point from zero. The model was constrained in the VO₂ baseline (Barstow et al., 1996) to identify key parameters according to the Eq. 1:

Equation 1. Mono-exponential model for VO₂ kinetics.

$$VO_2(t) = VO_2 \text{ base} + Ap \left[1 - e^{-\frac{(t-TDp)}{\tau_p}} \right] \quad (1)$$

where VO₂ (t) represents VO₂ values for a given time t; VO₂ base is the average VO₂ of the last minute of 20 W load period; Ap is the amplitude of phase II; τ_p is the phase II time-constant; and TD is the phase II time delay.

The slow component amplitude was determined by the difference between the VO₂ end and the sum of the VO₂ baseline and Ap, according to Eq. 2:

Equation 2. VO₂ slow component calculation.

$$As = VO_2 \text{ end} - (VO_2 \text{ base} + Ap) \quad (2)$$

Where, As is the slow component amplitude; VO₂ end is the average VO₂ value over the last 15 s; and Ap is the phase II amplitude.

The mean response time (MRT), which describes an overall pattern of VO₂ kinetics for the exercise bout, was calculated by a mono-exponential model from the beginning to the end of the exercise, excluding the time delay (Barstow and Molé, 1991), according to Eq. 3. The same equation was used to fit HR and O₂ pulse kinetics at the onset of the exercise (Wilkinson et al., 2006), with the end-point fixed at 180 s in order to augment fitting quality (i.e., better r²), since the bi-exponential pattern of HR and O₂ pulse were not detected for all of the subjects:

Equation 3. Mean-response time (overall kinetics) for VO₂, HR and O₂ pulse.

$$VO_2(t) = VO_2 \text{ base} + Ap \left[1 - e^{-\frac{t}{\tau}} \right] \quad (3)$$

Ischemia-Reperfusion Protocol

For IR, a 15 min bilateral thigh occlusion was introduced by the application of 250 mmHg pressure simultaneously in each thigh of the participants while they rested in a supine position (Ger-Ar, São Paulo, Brazil). Occlusion cuffs, chosen to be 20% wider than the upper leg diameter, were placed at the inguinal leg site. After 15 min of inflated cuffs, they were released, and the exercise load

started following 3 min of 20 W cycling. Capillary blood lactate was taken from the ear lobe at rest and every 5 min until the end of the occlusion period ([Lac]_{rest}, [Lac]₅, [Lac]₁₀, and [Lac]₁₅). The occlusion time, mode and the time between cuff releasing and the beginning of exercise were determined based on a time-window where the blood flow remained augmented against the baseline conditions (Hampson and Piantadosi, 1988; Paganelli et al., 1989; Faisal et al., 2010). The period of time between the subject's displacement from the bed to the cycle ergometer, as well as the data acquisition procedures (cuff displacement, face mask, HR strap adjustment, and sitting) were also controlled, in order to ensure trial reliability.

Statistical Analysis

Descriptive data is presented as mean ± standard deviation. Firstly, we tested the normality of the distribution by the Shapiro-Wilk test ($n < 50$). For variance analysis, sphericity assumptions were tested by the Mauchly's test, and corrections were made by the Greenhouse-Geisser factor. For VO₂, HR, and O₂ pulse kinetics parameter comparison, the two-tailed Student *t*-test for paired samples was used. Log transformations were made when necessary. For lactate concentration analysis between-conditions, the two-way ANOVA for repeated measures (condition vs. time) was conducted. For comparisons of lactate concentrations during thigh occlusion (in supine position), we used a one-way ANOVA for repeated measures. The Bonferroni's *post hoc* analysis was used to estimate point-by-point differences after ANOVA. The sample size was calculated for a moderate effect size ($ES = 0.5$), with an α level of 0.05 and a statistical power of 80%, resulting in 10 subjects. VO₂, HR and O₂ pulse data were loaded into the Origin 8.0 software and the statistical analysis was completed using the SPSS 22.0 for Windows® and the GraphPad Prism for Mac® (Prism 7.0, United States). An α level of 0.05 was set to claim for statistical significance.

RESULTS

Physiological Indexes

Table 1 shows the physiological indexes derived from the maximal incremental test.

VO₂ Kinetics at $\Delta 50\%$ After Ischemia-Reperfusion Model (IRT) and Control (CONT) Condition

Table 2 shows VO₂ kinetics parameters obtained during the 8 min square wave exercise with or without previous limb IR protocol. In Figure 2, the overall exercise VO₂ pattern is illustrated by averaging all the subject's 5 s data points. VO₂ measured at the end of square-wave trials were 3.38 ± 0.11 L min⁻¹ and 3.41 ± 0.09 L min⁻¹ ($P = 0.39$) for CON and IR conditions, respectively, overestimating predicted VO₂ about $22.9\% \pm 0.3\%$ and $24.1 \pm 0.2\%$. There were no differences between the conditions for any VO₂ kinetics parameters (VO₂ base, A_p, A_s, VO₂ end, τ_p , TD_p e MRT). In addition, there were no interactional effects between the condition and time for

TABLE 1 | Physiological indexes obtained after maximal incremental test.

Variables	Mean ± SD	Minimum	Maximum
VO ₂ max (L min ⁻¹)	3.48 ± 0.40	2.70	4.06
Wmax (W)	320 ± 28	290	365
GET VO ₂ (L min ⁻¹)	2.03 ± 0.32	1.53	2.53
GET %VO ₂ max (%)	58.2 ± 5.20	48.20	63.60
GET W (W)	132 ± 24	110	170
GET %Wmax (%)	41.3 ± 5.40	37.60	53.10
$\Delta 50\%$ VO ₂ (L min ⁻¹)	2.75 ± 0.35	215	325
$\Delta 50\%$ %VO ₂ max (%)	79.1 ± 2.60	74.10	81.80
$\Delta 50\%$ W (W)	211 ± 28	163	249
$\Delta 50\%$ %Wmax (%)	65.8 ± 4.70	56.30	73

VO₂ max, maximal oxygen uptake; Wmax, maximal power output; GET, gas exchange threshold; $\Delta 50\%$, GET plus 50% of the difference between it and VO₂ max.

TABLE 2 | VO₂ kinetics parameters.

Variables	CON	IR	P
Baseline			
VO ₂ base (L min ⁻¹)	1.08 ± 0.08	1.12 ± 0.06	0.30
Primary component			
A _p (L min ⁻¹)	2.07 ± 0.64	2.07 ± 0.48	0.99
τ_p (s)	50.1 ± 7.04	47.9 ± 6.37	0.47
TD _p (s)	6.11 ± 1.75	4.40 ± 1.2	0.34
Slow component			
A _s (L min ⁻¹)	0.22 ± 0.19	0.21 ± 0.17	0.75
Overall kinetics			
A _t (L min ⁻¹)	3.38 ± 0.11	3.41 ± 0.09	0.39
MRT _{VO2} (s)	76.6 ± 6.6	75.6 ± 5.2	0.75

CON – exercise without previous IR; IR: exercise preceded by IR, VO₂ base: average VO₂ of the lasting 60 s of 20 W baseline load; A_p, primary component amplitude; A_s, slow component amplitude; A_t, overall amplitude; τ_p , primary component time-constant; TD_p, primary component time-delay; MRT, mean response time; $\alpha = 0.05$.

the capillary blood lactate concentrations ($P = 0.16$), as well as no effect of conditions (CON vs. IR) ($P = 0.20$) for [Lac]_{rest} (0.86 ± 0.32 and 1.16 ± 0.38 mmol L⁻¹), [Lac]_{pre} (1.04 ± 0.35 and 1.16 ± 0.44 mmol L⁻¹) and [Lac]_{final} (10.26 ± 1.62 and 9.43 ± 1.46 mmol L⁻¹); and for [Lac] during bilateral thigh blood flow occlusion ($P = 0.64$; Figure 3).

HR and O₂ Pulse Kinetics

Table 3 shows HR and O₂ pulse kinetics results. MRT and the amplitude values are related to the time-fixed first 180 s of the exercise. There were no differences for any analyzed variables.

DISCUSSION

Main Findings

The main aim of this study was to investigate the impact of a prior bilateral thigh IR model on pulmonary VO₂ kinetics during a cycling exercise at $\Delta 50\%$ intensity. The secondary aim was to investigate central control of the O₂ pathway via the






FIGURE 2 | VO₂ during high-intensity exercise in IR and CON conditions. Points are averaged in 5 s means along the exercise period (8 min). Each point is representative of the mean ± SD of all 10 studied subjects. Mean VO₂ baseline for CON: 1.08 ± 0.08; mean VO₂ baseline for IR: 1.12 ± 0.06.



kinetic responses or HR and estimations of stroke volume (via O₂ pulse) at the onset of exercise. Our major finding was that the proposed IR protocol did not modify VO₂ kinetics during

TABLE 3 | HR and O₂ kinetics.

Variables	CON	IR	P
HR			
HRbase (bpm)	93 ± 13	91 ± 11	0.57
A _{180s} (bpm)	61 ± 11	64 ± 9	0.10
MRT _{180s} (s)	67.25 ± 6.03	71.33 ± 6.08	0.54
O ₂ pulse			
(O ₂ pulse)base (mL bpm ⁻¹)	11.99 ± 0.81	12.29 ± 0.94	0.37
A _{180s} (mL bpm ⁻¹)	8.94 ± 0.48	9.96 ± 0.91	0.35
MRT _{180s} (s)	40.89 ± 3.85	48.15 ± 5.55	0.31

HR base: heart rate of the lasting 60 s of the 20 W baseline load; (O₂ pulse) base: O₂ pulse of the lasting 60 s of the 20 W baseline load; A_{180s}: fixed-time amplitude at 180 s after the onset of the exercise; MRT_{180s}: time-constant of the variables fixed at 180 s after the onset of the exercise.

high-intensity exercise, neither HR and O₂ pulse, rejecting our hypothesis that a previous blood flow occlusion protocol would accelerate pulmonary VO₂ kinetics in a high intensity exercise. Our data also suggest that the IR model did not accelerate any central variables kinetics at the onset of the exercise. To the best of our knowledge, this is the first study to report the effects of IR on HR and O₂ pulse kinetics during a high intensity cycling exercise.

In light of the previous studies in the literature related to IR model, the absence of effects is in accordance to the study of Kido et al. (2015), that did not find any acceleration of pulmonary VO₂ kinetics during a severe intensity exercise when preceded by a similar protocol (ischemic preconditioning – 3 × 5 min bilateral occlusion, with 300 mmHg of pressure, ending within 5 min before exercise), especially at the onset of the exercise (τ_p) and also for “overall” kinetics (MRT). Faisal et al. (2010) found a diminished muscle VO₂ when the exercise began following 3 min of recovery after 15 min of forearm ischemia during an upper arm heavy exercise model, although an enhancement of forearm blood flow was observed. On the other hand, the findings of Walsh et al. (2002) suggests that “overall” pulmonary VO₂ kinetics could be accelerated by previous thigh IR lasting 5 to 10 min using about 250 mmHg of pressure. However, the major difference between the Walsh et al. (2002) study to ours/others was the timing between the end of a blood flow occlusion and the beginning of exercise.

Whilst it has been suggested that the local muscular hyperemia could be accessed by up to 2–3 min after the cuff release (Mullen

et al., 2001; Faisal et al., 2010), we consider this potential as a feature to explain possible differences. In our study, we chose a 3 min time frame until the beginning of the exercise based on a study of Faisal et al. (2010). In addition, the baseline exercise before the $\Delta 50\%$ exercise, which is recommended and often used for pulmonary VO₂ kinetics experimental assessments (Rossiter et al., 2002; Poole and Jones, 2012), may have influenced muscle blood flow levels before $\Delta 50\%$ exercise as well.

A classic study of Daly and Bondurant (1962) provided accurate measurements of HR, cardiac output and stroke volume after 15 min of a 250 mmHg forearm blood flow occlusion, suggesting a sudden rise of those variables in the first 15 s of cuff releasing. Thus, the central parameter results, such as HR and stroke volume (O₂ pulse), need to be considered when evaluating VO₂ kinetics. Also, it is noteworthy to highlight that we did not find any effects of IR model on [Lac] during the occlusion period, neither any differences in [Lac] rest values and [Lac]_{pre}. Blood lactate has been suggested as a properly mediator of O₂ transport enhancement by vascular tissue dilation (Krustrup et al., 2001; Tordi et al., 2003), which could partially explain the VO₂ kinetics acceleration after priming exercise conditions (Wilkerson et al., 2004; Nascimento et al., 2016). We also observed a concomitant absence of change in [Lac] and central hemodynamic parameters following the IT intervention. This may have occurred due to a higher than expected O₂ delivery and availability during thigh occlusion, as well as a better mitochondrial oxidation caused by higher levels of ADP, Pi, and Cr, which would attenuate any changes in [Lac].

Limitations and Future Directions

We considered that our findings need to be interpreted in light of its limitations. Here we point out some future directions for research that investigators could further consider.

When defining our protocol, we considered three major factors: (1) the pressure cuff level; (2) the time period for occlusion; (3) and the time between the end of occlusion and the beginning of the exercise. In regard to the first, a 250 mmHg level was chosen because of its efficiency to completely occlude the femoral artery blood flow (Sharma et al., 2014) and because it has been successfully by others (Salvador et al., 2015). For the same reason, the occlusion time was chosen – i.e., 15 min of bilateral thigh IR, as it had been shown to trigger reactive hyperemia (Kooijman et al., 2008) and necessary metabolic disturbances to maximize (Sharma et al., 2014). Finally, the time between the end of the IR model and the beginning of the exercise was selected for two main reasons: firstly, a physiological rationale to augment the local blood flow, supported by previous experimental results (Faisal et al., 2010); and secondly a logistic reason that is related to the necessary time to strictly adhere to the entire necessary methodological procedures for the VO₂ kinetics assessments in cycle ergometry, such as the participants transposition from the supine to the upright position, moving to the ergometer, cuff deflation and equipment removal.

Our results reinforce the absence of the central limitation in the onset-exercise pulmonary VO₂ kinetics, and, strengthen

the evidence on this side of the discussion. Therefore, we recommend that further research limited to this aspect is unlikely to change the direction of the evidence by its robustness. However, investigators still interested in using IR to study the onset-exercise pulmonary or muscular VO₂ kinetics, and should consider a reduction in the time to the beginning of the exercise and to control the confounding factors necessary to reduce muscle inertia and rest muscle tension (Hughson and Morrissey, 1983) such as the 3 min of a baseline exercise.

CONCLUSION

We found that VO₂, HR, and O₂ pulse kinetics during a high-intensity cycling exercise was not accelerated when preceded by a 15 min bilateral thigh blood flow occlusion and 3 min off. These results have important implications in possible strategies to accelerate VO₂ kinetics and improve metabolic efficiency. Considering there were no effects of IR when preceded by the off-period between the blood flow release and the exercise starting, further research should use other methods to investigate whether VO₂ kinetics are limited by oxygen delivery or oxidative enzyme inertia.

ETHICS STATEMENT

IRB: Comitê de Ética em Pesquisa com Seres Humanos – Universidade Federal de Santa Catarina Procedures: we first submitted to IRB's appreciation our project with rationale, methods, risks, and benefits. Together, the Consent Form was approved and, before any enrollment with this study, the subjects needed to assign it. In the consent form, information regarding risk, benefits, study's rationale and relevance and their wrights to drop out of the study whenever they want was made available to them. All signed forms are still stored in the project's folder.

AUTHOR CONTRIBUTIONS

All authors designed the study and interpreted the data. LH and PS collected and analyzed the data. LH wrote the first draft. PS, RdL, and LG wrote the final draft.

FUNDING

LH and PS received Ph.D. funding support by the CAPES Foundation. LG is granted by the CNPq Council.

ACKNOWLEDGMENTS

We want to thank the Exercise Physiology Laboratory staff (LAEF/UFSC) for supporting data collection, Dr. Eurico Wilhelm for supporting the reporting development.

REFERENCES

- Barstow, T. J., Jones, A. M., Nguyen, P. H., and Casaburi, R. (1996). Influence of muscle fiber type and pedal frequency on oxygen uptake kinetics of heavy exercise. *J. Appl. Physiol.* 81, 1642–1650. doi: 10.1152/jappl.1996.81.4.1642
- Barstow, T. J., and Molé, P. A. (1991). Linear and nonlinear characteristics of oxygen uptake kinetics during heavy exercise. *J. Appl. Physiol.* 71, 2099–2106. doi: 10.1152/jappl.1991.71.6.2099
- Beaver, W. L., Wasserman, K., and Whipp, B. J. (1986). A new method for detecting anaerobic threshold by gas exchange. *J. Appl. Physiol.* 60, 2020–2027. doi: 10.1152/jappl.1986.60.6.2020
- Boushel, R., Pott, F., Madsen, P., Radegran, G., Nowak, M., Quistorff, B., et al. (1998). Muscle metabolism from near infrared spectroscopy during rhythmic handgrip in humans. *Eur. J. Appl. Physiol. Occup. Physiol.* 79, 41–48. doi: 10.1007/s004210050471
- Burnley, M., Doust, J. H., Carter, H., and Jones, A. M. (2001). Effects of prior exercise and recovery duration on oxygen uptake kinetics during heavy exercise in humans. *Exp. Physiol.* 86, 417–425. doi: 10.1113/eph8602122
- Burnley, M., Doust, J. H., and Jones, A. M. (2002). Effects of prior heavy exercise, prior sprint exercise and passive warming on oxygen uptake kinetics during heavy exercise in humans. *Eur. J. Appl. Physiol.* 87, 424–432. doi: 10.1007/s00421-002-0647-8
- Daly, W. J., and Bondurant, S. (1962). Effects of oxygen breathing on the heart rate, blood pressure, and cardiac index of normal men—resting, with reactive hyperemia, and after atropine. *J. Clin. Invest.* 41, 126–132. doi: 10.1172/JCI104454
- Day, J. R., Rossiter, H. B., Coats, E. M., Skasick, A., and Whipp, B. J. (2003). The maximally attainable VO₂ during exercise in humans: the peak vs. maximum issue. *J. Appl. Physiol.* 95, 1901–1907. doi: 10.1152/japplphysiol.00024.2003
- Enko, K., Nakamura, K., Yunoki, K., Miyoshi, T., Akagi, S., Yoshida, M., et al. (2011). Intermittent arm ischemia induces vasodilatation of the contralateral upper limb. *J. Physiol. Sci.* 61, 507–513. doi: 10.1007/s12576-011-0172-9
- Faisal, A., Dyson, K. S., and Hughson, R. L. (2010). Prolonged ischaemia impairs muscle blood flow and oxygen uptake dynamics during subsequent heavy exercise. *J. Physiol.* 588, 3785–3797. doi: 10.1113/jphysiol.2010.188698
- Garber, C. E., Blissmer, B., Deschenes, M. R., Franklin, B. A., Lamonte, M. J., Lee, I. M., et al. (2011). Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med. Sci. Sports Exerc.* 43, 1334–1359. doi: 10.1249/MSS.0b013e318213febf
- Gastin, P. B. (2001). Energy system interaction and relative contribution during maximal exercise. *Sports Med.* 31, 725–741. doi: 10.2165/00007256-200131100-00003
- Gerbino, A., Ward, S. A., and Whipp, B. J. (1996). Effects of prior exercise on pulmonary gas-exchange kinetics during high-intensity exercise in humans. *J. Appl. Physiol.* 80, 99–107. doi: 10.1152/jappl.1996.80.1.99
- Hampson, N. B., and Piantadosi, C. A. (1988). Near infrared monitoring of human skeletal muscle oxygenation during forearm ischemia. *J. Appl. Physiol.* 64, 2449–2457. doi: 10.1152/jappl.1988.64.6.2449
- Hughson, R. L., and Morrissey, M. A. (1983). Delayed kinetics of VO₂ in the transition from prior exercise. Evidence for O₂ transport limitation of VO₂ kinetics: a review. *Int. J. Sports Med.* 4, 31–39. doi: 10.1055/s-2008-1026013
- Kido, K., Suga, T., Tanaka, D., Honjo, T., Homma, T., Fujita, S., et al. (2015). Ischemic preconditioning accelerates muscle deoxygenation dynamics and enhances exercise endurance during the work-to-work test. *Physiol. Rep.* 3:e12395. doi: 10.14814/phy2.12395
- Kooijman, M., Thijssen, D. H. J., de Groot, P. C. E., Bleeker, M. W. P., van Kuppevelt, H. J. M., Green, D. J., et al. (2008). Flow-mediated dilatation in the superficial femoral artery is nitric oxide mediated in humans. *J. Physiol.* 586, 1137–1145. doi: 10.1113/jphysiol.2007.145722
- Krustrup, P., Gonzalez-Alonso, J., Quistorff, B., and Bangsbo, J. (2001). Muscle heat production and anaerobic energy turnover during repeated intense dynamic exercise in humans. *J. Physiol.* 536, 947–956. doi: 10.1111/j.1469-7793.2001.00947.x
- Lamarra, N., Whipp, B. J., Ward, S. A., and Wasserman, K. (1987). Effect of interbreath fluctuations on characterizing exercise gas exchange kinetics. *J. Appl. Physiol.* 62, 2003–2012. doi: 10.1152/jappl.1987.62.5.2003
- MacDonald, M. J., Shoemaker, J. K., Tschakovsky, M. E., and Hughson, R. L. (1998). Alveolar oxygen uptake and femoral artery blood flow dynamics in upright and supine leg exercise in humans. *J. Appl. Physiol.* 85, 1622–1628. doi: 10.1152/jappl.1998.85.5.1622
- MacDonald, M. J., Tarnopolsky, M. A., and Hughson, R. L. (2000). Effect of hyperoxia and hypoxia on leg blood flow and pulmonary and leg oxygen uptake at the onset of kicking exercise. *Can. J. Physiol. Pharmacol.* 78, 67–74. doi: 10.1139/cjpp-78-1-67
- Margaria, R., Cerretelli, P., Di Prampero, P. E., Massari, C., and Torelli, G. (1963). Kinetics and mechanism of oxygen debt contraction in man. *J. Appl. Physiol.* 18, 371–377. doi: 10.1152/jappl.1963.18.2.371
- Mullen, M. J., Kharbanda, R. K., Cross, J., Donald, A. E., Taylor, M., Vallance, P., et al. (2001). Heterogeneous nature of flow-mediated dilatation in human conduit arteries in vivo: relevance to endothelial dysfunction in hypercholesterolemia. *Circ. Res.* 88, 145–151. doi: 10.1161/01.RES.88.2.145
- Nascimento, P. C., de Aguiar, R. A., Teixeira, A. S., de Souza, K. M., de Lucas, R. D., Denadai, B. S., et al. (2016). Are the oxygen uptake and heart rate off-kinetics influenced by the intensity of prior exercise? *Respir. Physiol. Neurobiol.* 230, 60–67. doi: 10.1016/j.resp.2016.05.007
- Nascimento, P. C., de Lucas, R. D., Souza, K. M., De Aguiar, R. A., Denadai, B. S., and Guglielmo, L. G. A. (2015). The effect of prior exercise intensity on oxygen uptake kinetics during high-intensity running exercise in trained subjects. *Eur. J. Appl. Physiol.* 115, 147–156. doi: 10.1007/s00421-014-3000-0
- Paganelli, W., Pendergast, D. R., Kones, J., and Cerretelli, P. (1989). The effect of decreased muscle energy stores on the VO₂ kinetics at the onset of exercise. *Eur. J. Appl. Physiol. Occup. Physiol.* 59, 312–316.
- Poole, D. C., Barstow, T. J., McDonough, P., and Jones, A. M. (2008). Control of oxygen uptake during exercise. *Med. Sci. Sports Exerc.* 40, 462–474. doi: 10.1249/MSS.0b013e31815ef29b
- Poole, D. C., and Jones, A. M. (2012). Oxygen uptake kinetics. *Compr. Physiol.* 2, 933–996. doi: 10.1002/cphy.c100072
- Rossiter, H., Howe, F., and Ward, S. (2005). “Intramuscular phosphate and pulmonary VO₂ kinetics during exercise,” in *Oxygen Uptake Kinetics in Sport, Exercise and Medicine*, eds D. Poole and A. M. Jones (London: Routledge), 154–184.
- Rossiter, H. B., Ward, S. A., Howe, F. A., Kowalchuk, J. M., Griffiths, J. R., and Whipp, B. J. (2002). Dynamics of intramuscular 31P-MRS Pi peak splitting and the slow components of PCr and O₂ uptake during exercise. *J. Appl. Physiol.* 93, 2059–2069. doi: 10.1152/japplphysiol.00446.2002
- Salvador, A. F., De Aguiar, R. A., Lisboa, F. D., Pereira, K. L., Cruz, R. S., and Caputo, F. (2015). Ischemic preconditioning and exercise performance: a systematic review and meta-analysis. *Int. J. Sports Physiol. Perform.* 11, 4–14. doi: 10.1123/ijsp.2015-0204
- Sharma, V., Cunniffe, B., Verma, A. P., Cardinale, M., and Yellon, D. (2014). Characterization of acute ischemia-related physiological responses associated with remote ischemic preconditioning: a randomized controlled, crossover human study. *Physiol. Rep.* 2:e12200. doi: 10.14814/phy2.12200
- Souza, K. M., De Lucas, R. D., Helal, L., Guglielmo, L. G. A., Greco, C. C., and Denadai, B. S. (2016). Agreement analysis between critical power and intensity corresponding to 50% Δ in cycling exercise. *Rev. Bras. Cineantropom. Desempenho Hum.* 18, 197–206. doi: 10.5007/1980-0037.2016v18n2p197
- Tordi, N., Perrey, S., Harvey, A., and Hughson, R. L. (2003). Oxygen uptake kinetics during two bouts of heavy cycling separated by fatiguing sprint exercise in humans. *J. Appl. Physiol.* 94, 533–541. doi: 10.1152/japplphysiol.00532.2002
- Walsh, M. L., Takahashi, A., Endo, M., Miura, A., and Fukuba, Y. (2002). Effects of ischaemia on subsequent exercise-induced oxygen uptake kinetics in healthy adult humans. *Exp. Physiol.* 87, 227–235. doi: 10.1113/eph8702262
- Whipp, B. J., Higgenbotham, M. B., and Cobb, F. C. (1996). Estimating exercise stroke volume from asymptotic oxygen pulse in humans. *J. Appl. Physiol.* 81, 2674–2679. doi: 10.1152/jappl.1996.81.6.2674
- Wilkerson, D. P., Berger, N. J., and Jones, A. M. (2006). Influence of hyperoxia on pulmonary O₂ uptake kinetics following the onset of exercise in humans. *Respir. Physiol. Neurobiol.* 153, 92–106. doi: 10.1016/j.resp.2005.09.006

Wilkerson, D. P., Koppo, K., Barstow, T. J., and Jones, A. M. (2004). Effect of prior multiple-sprint exercise on pulmonary O₂ uptake kinetics following the onset of perimaximal exercise. *J. Appl. Physiol.* 97, 1227–1236. doi: 10.1152/jappphysiol.01325.2003

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Helal, do Nascimento Salvador, de Lucas and Guglielmo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.