



# Inhibitory effect of microalgae and cyanobacteria extracts on influenza virus replication and neuraminidase activity

Thauane Silva<sup>1</sup>, Paulo S. Salomon<sup>2</sup>, Lidilhone Hamerski<sup>3</sup>, Juline Walter<sup>4</sup>, Rafael B. Menezes<sup>2</sup>, José Edson Siqueira<sup>3</sup>, Aline Santos<sup>1</sup>, Jéssica Aparecida Moraes Santos<sup>1</sup>, Natália Ferme<sup>1</sup>, Thaise Guimarães<sup>1</sup>, Giovana O. Fistarol<sup>2</sup>, Paulo I. Hargreaves<sup>2</sup>, Cristiane Thompson<sup>4</sup>, Fabiano Thompson<sup>4</sup>, Thiago Moreno Souza<sup>5,6</sup>, Marilda Siqueira<sup>1</sup> and Milene Miranda<sup>1</sup>

<sup>1</sup>Laboratório de Vírus Respiratórios e do Sarampo, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil

<sup>2</sup>Laboratório de Fitoplâncton Marinho, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>3</sup>Instituto de Pesquisas de Produtos Naturais, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>4</sup>Laboratório de Microbiologia Marinha, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>5</sup>Laboratório de Imunofarmacologia, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil

<sup>6</sup>Centro de Desenvolvimento Tecnológico em Saúde, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil

## ABSTRACT

**Background.** The influenza virus can cause seasonal infections with mild to severe symptoms, circulating worldwide, and it can affect people in any age group. Therefore, this infection is a serious public health problem that causes severe illness and death in high-risk populations. Every year, 0.5% of the world's population is infected by this pathogen. This percentage can increase up to ten times during pandemics. Influenza vaccination is the most effective way to prevent disease. In addition, anti-influenza drugs are essential for prophylactic and therapeutic interventions. The oseltamivir (OST, a neuraminidase inhibitor) is the primary antiviral used in clinics during outbreaks. However, OST resistant viruses may emerge naturally or due to antiviral pressure, with a prevalence of 1–2% worldwide. Thus, the search for new anti-influenza drugs is extremely important. Currently, several groups have been developing studies describing the biotechnological potential of microalgae and cyanobacteria, including antiviral activity of their extracts. In Brazil, this potential is poorly known and explored.

**Methods.** With the aim of increasing the knowledge on this topic, 38 extracts from microalgae and cyanobacteria isolated from marine and freshwater biomes in Brazil were tested against: cellular toxicity; OST-sensitive and resistant influenza replications; and neuraminidase activity.

**Results.** For this purpose, Madin-Darby Canine Kidney (MDCK)-infected cells were treated with 200 µg/mL of each extract. A total of 17 extracts (45%) inhibited influenza A replication, with seven of them resulting in more than 80% inhibition. Moreover, functional assays performed with viral neuraminidase revealed two extracts (from *Leptolyngbya* sp. and Chlorellaceae) with IC<sub>50</sub> mean < 210 µg/mL for influenza A and B, and also OST-sensitive and resistant strains. Furthermore, MDCK cells exposed to 1 mg/mL of all the extracts showed viability higher than 80%.

Submitted 23 April 2018

Accepted 10 September 2018

Published 26 October 2018

Corresponding author

Milene Miranda,  
mmiranda@ioc.fiocruz.br,  
milenediasmiranda@gmail.com

Academic editor

Peirong Jiao

Additional Information and  
Declarations can be found on  
page 13

DOI 10.7717/peerj.5716

© Copyright  
2018 Silva et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Discussion.** Our results suggest that extracts of microalgae and cyanobacteria have promising anti-influenza properties. Further chemical investigation should be conducted to isolate the active compounds for the development of new anti-influenza drugs. The data generated contribute to the knowledge of the biotechnological potential of Brazilian biomes that are still little explored for this purpose.

**Subjects** Marine Biology, Virology, Freshwater Biology

**Keywords** Cyanobacteria, Microalgae, Neuraminidase inhibition, Anti-influenza extracts, OST-sensitive and resistant influenza viruses

## INTRODUCTION

Lower acute respiratory infections (ARIs) are a persistent and pervasive public health problem, since they constitute one of the main causes of morbidity and mortality, with greater burden of disease worldwide than human immunodeficiency virus infection, malaria, cancer, or heart attacks (*Mizgerd, 2008; Pichon, Lina & Josset, 2017*).

World Health Organization (WHO) data indicate influenza A viruses as the main viral agents causing ARI, and therefore of great epidemiological importance. Seasonal influenza epidemics, currently involving influenza A(H3N2), A(H1N1)pdm09 and B (Yamagata (Yam) and Victoria (Vic) lineages), affect 10–20% of the human population each year (*WHO, 2018b*). According to new estimates by the United States Centers for Disease Control and Prevention, up to 650,000 deaths annually are associated with respiratory diseases from seasonal influenza (*WHO, 2017*). Although influenza B only infects humans, influenza A infects both humans and animals. Furthermore, humans can be infected with avian, swine and other zoonotic influenza viruses, such as A(H5N1), A(H7N9), and A(H9N2) subtypes (*WHO, 2018a*).

The influenza A virus belongs to *Orthomyxoviridae* family. The genomic single stranded RNA (RNAss) is octa-segmented, negative-sense, surrounded by a helical capsid with externally lipoprotein envelope, in which glycoproteins hemagglutinin (HA) and neuraminidase (NA) are inserted (*Camp et al., 2013*). Frequently, minor modifications in these envelope proteins may alter the affinity of vaccine antibodies or inactivate them, preventing recognition of the virus by the immune system, causing repetitive influenza outbreaks worldwide. However, in rare moments the combination of the eight genomic segments (reassortment) can occur, such as between influenza animal and human subtypes. This event, named antigenic shift, can result in strains capable of causing large regional or global pandemic outbreaks (*Zhu, Wang & Wang, 2017*).

The primary method of prevention is annual vaccination. Antiviral medications for prevention and treatment of influenza are an important adjunct to vaccines, especially for at-risk groups, including young children, older people, pregnant women and people with certain health conditions (*Del Giudice & Rappuoli, 2015; Rotrosen & Neuzil, 2017*).

The most important class of antiviral recommended for the control of influenza epidemics and eventual pandemics is the Neuraminidase Inhibitors (NAIs), particularly oseltamivir (OST) and zanamivir (ZAN). These compounds are active against all influenza A

subtypes and the two major influenza B lineages. Thus, the emergence of NAIs resistance could be a major clinical concern. Although most currently circulating influenza A and B strains are susceptible to NAIs, the pressure imposed by OST has led to the selection of OST-resistant mutants, with a prevalence of 1–2% in different countries ([Dixit et al., 2013](#); [Lopes e Souza et al., 2015](#); [Souza et al., 2011](#)). The OST-resistant strains with compensatory mutations may arise in an independent fashion, with samples being identified in different states of Brazil and in other countries ([Lopes e Souza et al., 2015](#)). Reports have shown single or multiple substitutions or deletions in the NA gene, which can promote a phenotype cross-resistance to the two main NAIs (oseltamivir and zanamivir) used in clinics, mostly in immunocompromised individuals ([Abed & Boivin, 2017](#)). Systematic circulation of these viral strains may jeopardise the use of the first line of anti-influenza drugs in the future. Thus, the search for new anti-influenza compounds is pivotal for public health.

Marine and freshwater natural products have been reported to contain different biological activities, which allow them to provide health and cosmetic benefits, such as antioxidant, anti-inflammatory, anticancer and antiviral activity, including against some respiratory viruses ([Chen et al., 2016](#); [Hasui et al., 1995](#); [Khalid et al., 2017](#); [Mayer & Hamann, 2004](#); [Mayer et al., 2007](#); [Mayer et al., 2017](#); [Mendes Gda et al., 2010](#)). Microalgae and cyanobacteria are huge natural sources of high-value compounds with health-promoting properties. These marine and freshwater organisms are a rich source of bioactive compounds such as vitamins, proteins with essential amino acids, polysaccharides, fatty acids, minerals, enzymes, fiber and photosynthetic pigments (carotenoids and chlorophylls) ([Montalvao et al., 2016](#)). The carotenoids contain a great compound range of different classes that are biosynthesized by condensation of isoprene units in (micro)algae and cyanobacteria ([Gong & Bassi, 2016](#); [Hynstova et al., 2018](#); [Sathasivam & Ki, 2018](#)).

The aim of this study is to evaluate the antiviral activity of natural extracts against influenza A and B, oseltamivir-sensitive and resistant strains and lineages. The 38 extracts were obtained from microalgae and cyanobacteria cultures isolated from fresh (Paraguaçu River and Lençóis Maranhenses Lakes) and marine water (Abrolhos Bank and Guanabara Bay) biomes in Brazil. We evaluated these extracts for cellular toxicity, influenza replication and neuraminidase activity. The extracts presented low cytotoxicity, and 17 of them inhibited influenza replication in more than 80% *in vitro*, with two inhibiting neuraminidase activity from influenza A and B OST-sensitive and resistant. These data contribute to the knowledge of the biotechnological potential of Brazilian biomes, which are underexplored for antiviral research and bioprospecting.

## MATERIALS AND METHODS

### Origin of microalgae and cyanobacteria strains

Microalgae and cyanobacteria strains used in this study belong to the Culture Collection of Microalgae at UFRJ (CCMR, Marine Phytoplankton Laboratory, Federal University of Rio de Janeiro). These strains were isolated by single-cell sorting in a flow cytometer ([Fistarol et al., 2018](#)) and are maintained in culture by successive transfers to fresh liquid medium at 3–6 week intervals, depending on the species. All 26 strains used here are

originated from fresh (Paraguçu River and Shallow Lake System of Lençóis Maranhenses) and marine water (Guanabara Bay estuary and Abrolhos coral reefs) biomes in Brazil (Table 1). Taxonomic identification of the strains was based on morphology and, for most of them, also on DNA sequencing (SSU rDNA) (Fistarol *et al.*, 2018). The field experiments were approved by Ministério do Meio Ambiente (MMA), Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) and Sistema de Autorização e Informação em Biodiversidade (SISBIO), approval number: 35854-2. The access to genetic heritage was approved by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), approval number: 010339/2014-0; and Sistema Nacional de Gestão do Patrimônio Genético e do Conhecimento Tradicional Associado (SisGen), approval number: A03D5C4.

### Microalgae and cyanobacteria cultivation

Microalgae and cyanobacteria were grown in autoclaved sterile medium at  $26 \pm 1$  °C, with a photon flux of ca.  $300 \mu\text{moles photon m}^{-2} \text{s}^{-1}$  and 16 h light: 8 h dark photoperiod. Freshwater strains were grown in ASM-1 medium whereas marine strains were grown in f/2 medium (Gorham *et al.*, 1964; Guillard & Lorenzen, 1972; Reynolds & Jaworski, 1978). Cultures were grown in glass balloons with 300–500 mL of culture medium for 3–6 weeks, depending on the growth rate of each strain. The cultures were stored at  $-80$  °C until extracts production.

### Extracts production

All the microalgae and cyanobacteria cultures (cells and culture medium) were frozen, thawed and filtered through cheesecloth. The filtrate was extracted three times with ethyl acetate (added ratio 1:1 of volume) and three times with n-butanol (added ratio 1:1 of volume). The organic layers were concentrated under reduced pressure to dryness. A total of 38 extracts were produced, 26 in ethyl acetate (EtOAc) and 12 in n-butanol (n-BuOH). The crude extracts were stored at  $-20$  °C until analysis. The extracts from one to 12 strains were obtained in the first batch with both solvents. After the screening test for anti-influenza activity, the second batch of extracts was produced from 13 to 26 strains, and they were solely obtained with ethyl acetate. Dried extracts were resuspended in dimethyl sulfoxide (DMSO) in a final concentration of 100 mg/mL for the *in vitro* tests. All the extracts were analyzed by TLC, HPLC, and Mass Spectrometry (MS). However, we could not identify any compounds, but the chemical profiles of the active extracts were preserved. Thus, the procedure must be repeated on a larger scale for the more active anti-influenza extracts. The chromatograms and mass spectra for extracts 5 and 21 are provided in the File S1 (S01).

### Cells and virus

Madin-Darby canine kidney (MDCK) cells (London line) were kindly donated by the Centers for Disease Control and Prevention (CDC), Influenza Reagent Resources (IRR) (FR-58). These cells were used for cytotoxicity analysis, viral growth and experimental assays with influenza virus infection. These cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco, Waltham, MA, USA) supplemented with 10% of fetal bovine serum (FBS; Gibco) and 100 U/mL penicillin and 100 mg/mL streptomycin (Sigma-Aldrich,

**Table 1** Extracts identification and collection specifications.

Extract identification	Taxon	Source	Collection point	Latitude	Longitude	Collection data
1	Unidentified filamentous green algae	Freshwater	Lençóis Maranhenses Lakes	2°32'11"S	41°51'10"W	August, 2012
2	<i>Staurastrum</i> sp.	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
3	Unidentified filamentous green algae	Freshwater	Paraguaçu River	12°57'8.48"S	41°16'39.29"W	November, 2012
4	<i>Scenedesmus</i> sp.	Freshwater	Lençóis Maranhenses Lakes	12°50'25.91"S	41°19'26.52"W	November, 2012
5	<i>Leptolyngbya</i> sp. <sup>a</sup>	Marine	Abrolhos Bank	17°59'52.8"S	38°40'15.6"W	October, 2013
6	Unidentified coccoid green algae	Freshwater	Paraguaçu River	12°57'8.48"S	41°16'39.29"W	November, 2012
7	Unidentified coccoid green algae	Freshwater	Lençóis Maranhenses Lakes	12°50'25.91"S	41°19'26.52"W	November, 2012
8	<i>Scenedesmus abundans</i>	Freshwater	Lençóis Maranhenses Lakes	2°32'11"S	41°51'10"W	August, 2012
9	<i>Scenedesmus vacuolatus</i>	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
10	<i>Chlamydomonas angulosa</i>	Freshwater	Lençóis Maranhenses Lakes	2°38'50"S	42°49'43"W	August, 2012
11	Chlorellaceae	Freshwater	Lençóis Maranhenses Lakes	2°38'50"S	42°49'43"W	August, 2012
12	Desmidiaceae	Freshwater	Lençóis Maranhenses Lakes	2°35'08"S	42°48'03"W	August, 2012
13	<i>Chlamydomonas</i> sp.	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
14	<i>Selenastrum</i> sp.	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
15	Scenedesmaceae	Freshwater	Lençóis Maranhenses Lakes	2°33'42"S	42°51'49"W	August, 2012
16	Unidentified coccoid green algae	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
17	<i>Leptolyngbya</i> sp. <sup>a</sup>	Marine	Abrolhos Bank	12°50'25.91"S	41°19'26.52"W	November, 2012
18	<i>Desmodesmus armatus</i>	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
19	<i>Symbiodinium</i> sp.	Marine	Abrolhos Bank	17°57'32.7"S	38°30'20.3"W	March, 2012
20	<i>Chattonella</i> sp.	Marine	Guanabara Bay	22°50'01.0"S	43°12'29.0"W	January, 2016
21	Chlorellaceae	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
22	<i>Desmodesmus perforatus</i>	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
23	Chlorellaceae	Freshwater	Lençóis Maranhenses Lakes	2°32'11"S	41°51'10"W	August, 2012
24	Desmidiaceae	Freshwater	Lençóis Maranhenses Lakes	2°30'10"S	42°51'50"W	August, 2012
25	<i>Romeria</i> sp. <sup>a</sup>	Marine	Abrolhos Bank	18°00'13.0"S	39°14'57.0"W	February, 2014
26	<i>Nanofrustulum shiloi</i>	Marine	Abrolhos Bank	17°57'07.0"S	39°13'11.0"W	February, 2013

**Notes.**<sup>a</sup>Cyanobacteria strains.

St. Louis, MO, USA). Cells were cultured at 37 °C in 5% CO<sub>2</sub> atmosphere ([Szretter, Balish & Katz, 2006](#)).

The viruses were grown and titrated according to WHO manual for the laboratory diagnosis and virological surveillance of influenza ([WHO, 2011](#)). The virus titration was performed by 50% Tissue Culture Infectious Dose (TCID<sub>50</sub>) assay. We analyzed the inhibitory effect of the extracts against influenza A strains and B lineages, OST-sensitive

(Wild type—WT) and resistant (mutant): A/California/04/2009 (A(H1N1)pdm09-WT); A/Perth/261/2009 (A(H1N1)pdm09-H275Y); A/Switzerland/9715293/2013 (A(H3N2)-WT); B/Phuket/3073/2013 Yam (WT); B/Brisbane/60/2008 Vic (WT); B/Perth/211/2001 Yam (WT) and B/Perth/211/2001 Yam (D197E). All the viruses were kindly provided by the International Society for Influenza and other Respiratory Viruses Diseases-Antiviral Group (isirv-AVG) and CDC –IRR.

### Cytotoxicity assay

One hundred microliters of  $2.0 \times 10^4$  MDCK cells were seeded into the 96-well culture plates (flat bottom) and grown for 24 h at 37 °C in a 5% CO<sub>2</sub> atmosphere, then, the extracts were added with a final concentration of 1 mg/mL in DMEM. Control cells were treated with 1.0% DMSO, which did not affect the growth of the cells. The extracts were diluted in culture medium DMEM, the DMSO final concentration was equal to 1.0% (v/v). After 48 h of cells incubation with extracts or vehicle (DMSO), a freshly prepared XTT (Sigma) solution was added, as specified by the manufacturer's instructions. The XTT is a colorless or slightly yellow compound that is reduced to formazan (bright orange) by mitochondrial dehydrogenases of viable cells, as described before (Roehm et al., 1991; Scudiero et al., 1988). The cell viability was calculated by the comparison between absorbance (A475 nm–A660 nm) from treated (A) and untreated (B) cells using the formula:  $A/B \times 100$ . Tests were carried out performing three replicates ( $n = 3$ ).

### Screening of extracts on viral replication

Initially, we performed a screening to evaluate the inhibitory effect of the extracts. MDCK cells ( $2.0 \times 10^5$  cells/well) were seeded in 24-well tissue culture plates and incubated overnight. Then, the monolayers of cells were infected with A(H1N1)pdm09-WT or A(H3N2)-WT at 400 TCID<sub>50</sub> per well during 1 h at 37 °C and 5% CO<sub>2</sub>. After, the viral suspension was removed and the monolayers were treated with 200 µg/mL of each extract. There is no established protocol or concentration for the anti-influenza screening of crude extracts of natural products. There are papers that use 2 mg/ml as the maximum concentration, while others use 100 µg/ml (Ding et al., 2017; Ghoke et al., 2018; Rajasekaran et al., 2013; Shoji et al., 2017; Sood et al., 2012). At 48 h post-infection (hpi), the supernatant was harvested to quantify the influenza titer. For this purpose, the supernatants were diluted 1:5 in NA-Star buffer and the neuraminidase activity was analyzed in triplicate by the NA-Star™ assay kit (Life Technologies, Carlsbad, CA, USA), according to the manufacturer's instructions. The antiviral activity of the extracts was calculated with respect to virus control only.

### EC<sub>50</sub> determination

Half maximal effective concentration was determined for the extract that inhibited influenza replication in more than 80%. MDCK cells were seeded and infected as described above. After 1 hpi, the viral suspension was removed and the monolayers were treated with different extract concentrations (400 – 12.5 µg/mL). Oseltamivir carboxylate (100 – 3.2 µM) (OST-car; kindly donated by Hoffman-La Roche Inc., Basel, Switzerland) was used as a control. For influenza titration, the supernatants were harvested at 48 hpi, and the



influenza virus was titrated by TCID<sub>50</sub>/mL using MDCK cells ( $5 \times 10^4$  cells/well in 96-well plates) through the Reed and Muench method ([Reed & Mounch, 1938](#); [Sacramento et al., 2015](#)). Non-linear regression of the dose response curves was performed to determine the 50% inhibitory effect on viral replication (EC<sub>50</sub>) for the extracts and reference compound (OST-car).

### Functional antiviral assay—IC<sub>50</sub> determination

For extracts with high influenza inhibition (80%) in the screening test, we performed a functional antiviral assay to determine the concentration required to inhibit 50% of the NA enzymatic activity (IC<sub>50</sub>). Wild-type and resistant strains of influenza A and B were tested against different concentrations of extracts (400 – 12.5 or 200 – 6.25 µg/mL), OST-car (1,000 – 0.01 nM) was used as control. This assay was done using the NA-Star<sup>TM</sup> assay kit (Life Technologies, USA), according to the manufacturer's instructions ([Lopes e Souza et al., 2015](#); [Souza et al., 2011](#); [Souza et al., 2013](#)). The IC<sub>50</sub> was calculated using a non-linear regression.

### Statistical analysis

The dose–response curves used to calculate the pharmacological parameter values were generated using Excel 2010 for Windows software (Microsoft) ([Souza et al., 2007](#)). All of the experiments were performed at least three times, and the results are displayed as the mean or the mean ± standard error (SEM).

## RESULTS

### Low cytotoxicity of the extracts

The cells viability was the first analysis conducted, extracts with high cell toxicity *in vitro* were disregarded from the antiviral analysis. All the extracts produced were tested at 1 mg/mL in MDCK cell cultures. As control, we exposed MDCK to DMSO in the same proportion as we used in the extracts test, 1.0% V/V in culture medium, which did not affect the cells viability. The results showed that redox mitochondrial activity of the tested cells with extracts was inhibited at most by 20% after 48 h of exposure. Approximately 60% of extracts in ethyl acetate (EtOAc) and n-butanol (n-BuOH) solvent showed 90% of cell viability ([Table 2](#)).

### The extracts inhibit influenza replication in more than 80%

After cytotoxicity analysis, we tested anti-influenza activity with all the extracts produced, EtOAc and n-BuOH. For this purpose, MDCK-infected cells were treated with 200 µg/mL of each extract. After 48 hpi, the supernatants were harvested and the influenza neuraminidase activity was analyzed. We observed that 17 extracts (16 in EtOAc and 1 in n-BuOH) inhibited the replication of influenza A strains, A(H1N1)pdm09-WT and A(H3N2)-WT. Seven of them (41%) resulted in more than 80% of inhibition ([Table 2](#)). Just one n-BuOH extract from filamentous green algae was capable of inhibiting influenza replication, but at low levels. Despite this, an EtOAc extract obtained from another filamentous green alga was able to produce satisfactory inhibitory effects ([Table 2](#)). All

**Table 2** Percentage of cell viability and influenza inhibition (screening) after exposure to extracts in ethyl acetate (EtOAc) or n-butanol (n-BuOH) solvent.

Extract	Taxon	Cell viability (%)		Screening (%)			
		EtOAc	n-BuOH	A(H1N1)pdm09-WT		A(H3N2)-WT	
				EtOAc	n-BuOH	EtOAc	n-BuOH
1	Unidentified filamentous green algae	80 ± 1	80 ± 2	87 ± 2	NI	82 ± 2	NI
2	<i>Staurastrum</i> sp.	90 ± 2	90 ± 2	85 ± 2	NI	82 ± 1	NI
3	Unidentified filamentous green algae	80 ± 1	80 ± 2	NI	30 ± 4	NI	20 ± 2
4	<i>Scenedesmus</i> sp.	90 ± 2	90 ± 2	83 ± 3	NI	80 ± 3	NI
5	<i>Leptolyngbya</i> sp. <sup>a</sup>	90 ± 2	90 ± 1	90 ± 2	NI	85 ± 2	NI
6	Unidentified coccoid green algae	90 ± 3	90 ± 1	73 ± 2	NI	70 ± 4	NI
7	Unidentified coccoid green algae	90 ± 1	90 ± 2	NI	NI	NI	NI
8	<i>Scenedesmus abundans</i>	90 ± 3	90 ± 2	70 ± 4	NI	75 ± 1	NI
9	<i>Scenedesmus vacuolatus</i>	80 ± 2	80 ± 2	NI	NI	NI	NI
10	<i>Chlamydomonas angulosa</i>	90 ± 1	90 ± 1	NI	NI	NI	NI
11	Chlorellaceae	80 ± 3	80 ± 4	NI	NI	NI	NI
12	Desmidiaceae	80 ± 1	80 ± 2	60 ± 2	NI	56 ± 2	NI
13	<i>Chlamydomonas</i> sp.	90 ± 3	NP	NI	NP	NI	NP
14	<i>Selenastrum</i> sp.	80 ± 2	NP	NI	NP	NI	NP
15	Scenedesmaceae	90 ± 3	NP	NI	NP	NI	NP
16	Unidentified coccoid green algae	90 ± 4	NP	NI	NP	NI	NP
17	<i>Leptolyngbya</i> sp. <sup>a</sup>	90 ± 3	NP	57 ± 2	NP	55 ± 2	NP
18	<i>Desmodesmus armatus</i>	80 ± 3	NP	94 ± 2	NP	90 ± 2	NP
19	<i>Symbiodinium</i> sp.	80 ± 2	NP	59 ± 4	NP	62 ± 4	NP
20	<i>Chattonella</i> sp.	90 ± 1	NP	70 ± 5	NP	65 ± 2	NP
21	Chlorellaceae	90 ± 4	NP	99 ± 1	NP	95 ± 1	NP
22	<i>Desmodesmus perforates</i>	80 ± 3	NP	58 ± 3	NP	50 ± 4	NP
23	Chlorellaceae	90 ± 3	NP	46 ± 2	NP	38 ± 4	NP
24	Desmidiaceae	90 ± 1	NP	84 ± 1	NP	82 ± 2	NP
25	<i>Romeria</i> sp. <sup>a</sup>	80 ± 3	NP	NI	NP	NI	NP
26	<i>Nanofrustulum shiloi</i>	90 ± 3	NP	66 ± 2	NP	60 ± 2	NP

**Notes.**<sup>a</sup>Cyanobacteria strains.

NP, not produced; NI, no inhibition.

DMSO was used as a control at 0.01% V/V, and the cell viability remained at 100%.

the EtOAc extract from Desmidiaceae family (*Staurastrum* sp.) and four out of six EtOAc extract from Scenedesmaceae family (*Scenedesmus* sp. and *Desmodesmus* sp.) were capable of inhibiting influenza replication with different magnitudes (Table 2). Besides these, extract 21, obtained from Chlorellaceae green microalgae presented the greatest capacity to inhibit both influenza replications. However, others from the same family had limited or null capacity against this virus (Table 2).

**EC<sub>50</sub> determination and SI evaluation**

We determined the EC<sub>50</sub> for seven marine extracts that *inhibited* in more than 80% the replication of both influenza A(H1N1)pdm09-WT and A(H3N2)-WT in the screening



**Table 3** The EC<sub>50</sub>, CC<sub>50</sub> and SI values from extracts that inhibited influenza A replication higher than 80% in screening assay.

Extract	Taxon	EC <sub>50</sub> <sup>a</sup> (μg/mL)		CC <sub>50</sub> <sup>b</sup> (μg/mL)	SI <sup>c</sup>	
		A(H1N1)pdm09-WT	A(H3N2)-WT		A(H1N1)pdm09-WT	A(H3N2)-WT
1	Unidentified filamentous green algae	130	150	>1,000	>7.7	>6.7
2	<i>Staurastrum</i> sp.	70	90	>1,000	>14.3	>11.1
4	<i>Scenedesmus</i> sp.	130	130	>1,000	>7.7	>7.7
5	<i>Leptolyngbya</i> sp. <sup>d</sup>	80	85	>1,000	>12.5	>11.8
18	<i>Desmodesmus armatus</i>	55	60	>1,000	>18.4	>16.7
21	Chlorellaceae	30	40	>1,000	>33.3	>25.0
24	Desmidiaceae	50	55	>1,000	>20.0	>18.2
OST-car		0.012 μM	0.032 μM	2,132 μM	177,7	66,63

**Notes.**

<sup>a</sup>EC<sub>50</sub>, the concentration required to reduced inhibition of viral infection-induced cytopathogenicity by 50%. Values represent the mean of duplicate samples from three independent experiments.

<sup>b</sup>CC<sub>50</sub>, the concentration required to reduced normal, non-infected cell viability by 50%. Values represent the mean of duplicate samples from three independent experiments.

<sup>c</sup>SI, selective index is determined by the ratio between CC<sub>50</sub> and EC<sub>50</sub>.

<sup>d</sup>Cyanobacteria strain.

OST-car, OST carboxylate.

assay. The selective index (SI) is determined by the ratio between CC<sub>50</sub> and EC<sub>50</sub>, and represents the relative effectiveness of the investigational product in inhibiting viral replication compared to inducing cell death. The EC<sub>50</sub> of five extracts was less than 100 μg/mL with SI higher than 10.0 (Table 3). The oseltamivir carboxylate (OST-car) was used as positive control with 177,667 and 66,625 SI for A(H1N1)pdm09-WT and A(H3N2)-WT, respectively. The SI values between a pool of molecules (extracts) and a molecule chemically defined (OST) are incomparable.

### Neuraminidase inhibition and IC<sub>50</sub> determination

We performed a functional assay to analyze neuraminidase activity with all the seven extracts that inhibited influenza infection in more than 80%. IC<sub>50</sub> is the concentration required to inhibit 50% of the influenza NA enzymatic activity. Thus, lower IC<sub>50</sub> values imply in higher NA inhibition activity. Two of them, extracts from the cyanobacteria *Leptolyngbya* sp. (5) and microalga Chlorellaceae (21) inhibited influenza A NA activity with IC<sub>50</sub> lower than 210 and 20 μg/mL, respectively (Table 4 and Figs. 1A–1D). The OST-car was used as control (Table 4 and Figs. 1E and 1F). The HPLC and mass spectrum of extracts 5 and 21 are shown S01-1 and S01-2, respectively. The mass spectrum of extract 5 (S01-3) indicated the presence of chlorinated compounds m/z = 214.9850 and 232.9946. On the other hand, the mass spectrum of extract 21 (S01-4) exhibited signals that indicated the presence of peptides, terpenes and/or alkaloids.

### EC<sub>50</sub> and IC<sub>50</sub> determination to influenza A and B OST-sensitive and resistant viruses

The OST is the main antiviral used in clinics against Influenza A and B. The pressure imposed by this drug has led to the selection of resistant mutants (Dixit et al., 2013; Lopes e Souza et al., 2015; Souza et al., 2011). Thus, the search for new anti-influenza compounds

**Table 4** The EC<sub>50</sub> and IC<sub>50</sub> values from extracts for OST-sensitive and resistant influenza A and B viruses.

Influenza Virus strain	EC <sub>50</sub> <sup>a</sup> (μg/mL)		IC <sub>50</sub> <sup>b</sup> (μg/mL)		
	<i>Leptolyngbya</i> sp. (extract 5)	Chlorellaceae (extract 21)	<i>Leptolyngbya</i> sp. (extract 5)	Chlorellaceae (extract 21)	OST-car (nM)
A(H1N1)pdm09-WT	80	30	149 ± 5	16 ± 2	0.1 ± 0.2
A(H3N2)-WT	85	40	162 ± 3	14 ± 3	2.1 ± 0.3
A/Perth/261/2009 (A(H1N1)pdm09H275Y)	168	50	208 ± 4	72 ± 3	112 ± 2
B/Phuket/3073/2013 Yam (WT)	215	70	207 ± 5	68 ± 2	2.8 ± 1
B/Brisbane/60/2008 Vic (WT)	230	75	199 ± 4	41 ± 2	3 ± 0.5
B/Perth/211/2001 Yam (WT)	200	100	179 ± 3	30 ± 5	5.0 ± 1
B/Perth/211/2001 Yam (D197E)	260	120	184 ± 5	42 ± 4	10 ± 2

**Notes.**

<sup>a</sup>EC<sub>50</sub>, the concentration required to reduced inhibition of viral infection-induced cytopathogenicity by 50%. Values represent the mean of duplicate samples from three independent experiments.

<sup>b</sup>IC<sub>50</sub>, the concentration required to inhibit 50% of the NA enzymatic activity. Values represent the mean of duplicate samples from three independent experiments ± standard error.

OST-car, OST carboxylate.


that inhibits both types of influenza and their resistant strains or lineages is extremely important. Therefore, we tested extracts from *Leptolyngbya* sp. (5) and Chlorellaceae (21) against replication and NA activity of influenza A OST-resistant (Table 4 and Figs. 1B and 1D) and B OST-sensitive and resistant (Table 4 and Figs. 1A–1D) viruses. The OST-car was used as control (Table 4 and Figs. 1E and 1F). Both extracts inhibited influenza replication and NA activity with low EC<sub>50</sub> and IC<sub>50</sub>, including for OST-resistant strains and lineages (Table 4 and Fig. 1). The raw data of the IC<sub>50</sub> calculations are provided in S02.

## DISCUSSION

Influenza virus is the most important pathogen that causes acute lower respiratory infections (WHO, 2018b). The OST, a neuraminidase inhibitor, is the main anti-influenza used in clinics, but strains resistant to this molecule have been in clinical treatments reported worldwide, mandating the development of novel therapeutics (Dixit et al., 2013; Lopes e Souza et al., 2015; Souza et al., 2011).

Microalgae and cyanobacteria are huge natural sources of compounds of high nutritional and medicinal value. These marine and freshwater organisms are a rich source of bioactive compounds such as vitamins, proteins with essential amino acids, polysaccharides, fatty acids, minerals, photosynthetic pigments, enzymes, and fiber (Gogineni & Hamann, 2018; Montalvao et al., 2016).

Cyanobacteria are a promising yet underexplored source for novel natural products, including antiviral compounds. More detailed screening studies have identified antiviral activities on sulfoglycolipids and lectins (Lopes et al., 2011; Mundt et al., 1997; Niedermeyer, 2015; Sharaf et al., 2010; Zainuddin et al., 2002). However, chemically different groups of compounds in cyanobacteria include alkaloids, lipopeptides, macrolides, and others (Kiuru et al., 2014). Several studies have been conducted to test microalgae compounds against pathogenic human viruses. Antiviral compounds extracted from microalgae are mainly polysaccharides. Polyunsaturated aldehydes, carotenoid, terpenes, and alkaloids are other



**Figure 1** *Leptolyngbya sp.* and *Chlorellaceae* extracts inhibit influenza A and B OST-sensitive and -resistant viruses. The NA activity of the influenza A and B OST-sensitive and -resistant viruses were measured in the presence of different concentrations of extracts using a chemiluminescent substrate, NA-star kit (Life Technologies, Carlsbad, CA, USA). (A and B) Extract 5 (400 – 12.5  $\mu\text{g/mL}$ ). (C and D) Extract 21 (200 – 6.25  $\mu\text{g/mL}$ ). (E and F) OST carboxylate (1,000 – 0.01 nM), the reference compound was used as control. The results were obtained in relative luminescence units (RLU), but were converted to a percentage of the control for normalization of the data displayed. These experiments were performed three times and the means were expressed in the graphs.

Full-size [DOI: 10.7717/peerj.5716/fig-1](https://doi.org/10.7717/peerj.5716/fig-1)

bioactive molecules produced by microalgae (Falaise et al., 2016; Martinez Andrade et al., 2018; Sathasivam & Ki, 2018).

Thus, we tested anti-influenza activity of extracts derived from microalgae and cyanobacteria collected from Paraguaçu River, Shallow Lake System of Lençóis Maranhenses, Guanabara Bay estuary and Abrolhos coral reefs.

In screening assay we observed that seven extracts were able to inhibit seasonal influenza A replication viruses in more than 80%, five of them with SI higher than 10. Aqueous and methanolic extracts of cultured cyanobacteria of *Microcystis* and *Spirulina* genus inhibit influenza replication with SI comparable to ethyl acetate extract from *Leptolyngbya*

sp. (cyanobacteria) (Chen *et al.*, 2016; Zainuddin *et al.*, 2002). Besides that, one sulfated polysaccharide purified from the marine microalga, *Gyrodinium impudium*, inhibits influenza A infection with SI higher than 200, but not influenza B (Kim *et al.*, 2012).

In addition, *Trichodesmium erythraeum* (Cyanobacterial) aplysiatoxin-related compounds debromoaplysiatoxin and 3-methoxydebromoaplysiatoxin displayed anti-CHIKV effects at concentrations that resulted in minimal cytotoxicity (Gupta *et al.*, 2014). Activity against influenza A has been found for the cyclic depsipeptides named ichthyopeptins A and B isolated from cyanobacterium *Microcystis ichthyolabe* (Zainuddin *et al.*, 2007). *Nostoc ellipsosporum* (Cyanobacterial) lectin named Cyanovirin-N (CV-N) is a potent anti-influenza A as well as anti-influenza B. The CV-N binds directly to the viruses, inactivating them (Singh *et al.*, 2017). The green algae and land plants form a monophyletic lineage named chlorophytes, the first is an ancestor of the second (Delaux *et al.*, 2015). Therefore, it is possible that they present similar primary and secondary metabolites (Cheynier *et al.*, 2013; Iwai & Yokono, 2017). A broad spectrum of plant extracts inhibit the influenza replication, including extracts from ethyl acetate solvent, with SI comparable to that observed in extracts 2, 4, 18, 21 and 24, from green microalgae *Staurastrum* sp., *Scenedesmus* sp., *Desmodesmus armatus*, Chlorellaceae and Desmidiaceae, respectively (Hour *et al.*, 2013; Wu *et al.*, 2010).

The Chlorellaceae family, mainly *Chlorella* genus, produce a broad spectrum of carotenoids with different anti-angiogenic activities, cardioprotective, anti-cancer, anti-diabetic, anti-inflammatory, anti-oxidant and others (Sathasivam & Ki, 2018). Furthermore, retinoids, products of carotenoid metabolism, inhibit mumps virus replication preventing the infection of healthy cells by induction of a retinoid inducible gene I (RIG-I), retinoic acid receptor (RAR) and interferon (Soye *et al.*, 2013). Interferon-induced transmembrane proteins (IFITMs) inhibit infection of diverse enveloped viruses, such as West Nile, dengue, Zika and influenza by blocking virus-endosome fusion (Brass *et al.*, 2009; Desai *et al.*, 2014; Mesquita *et al.*, 2014; Savidis *et al.*, 2016). Thus, it is possible that the Chlorellaceae (microalgal) extract leads to the increase of this cell restrictive protein, suggesting a mechanism of action for the influenza replication inhibitory effect.

In addition, two extracts from *Leptolyngbya* sp. (cyanobacteria) and Chlorellaceae family (microalgae) inhibited the replication and neuraminidase activity of influenza A and B viruses, including OST-sensitive and resistant strains and lineages. These results were similar to that observed for a water extract of *Taxodium distichum* (Cupressaceae family), which presented excellent efficacy against both influenza viruses, particularly OST-resistant clinical isolates and swine-origin influenza strains (Hsieh *et al.*, 2016). These data indicate that microalgae and cyanobacteria are a promising source of anti-influenza natural products.

As mentioned before, there are several cellular factors described as restricting the replication of influenza virus (Villalon-Letelier *et al.*, 2017). Furthermore, despite the inhibitory effect against neuraminidase activity, the extracts may have molecules capable of inhibiting different stages of the viral replicative cycle, and/or they can activate host cell restriction factors, but these effects will still be investigated. Thus, microalgae and cyanobacteria are important marine and freshwater microorganisms for bioprospecting in

antiviral research. The microorganisms evaluated in this study were collected in Brazil, but they are not exclusively Brazilian. It may be relevant to compare the metabolites production of the same taxonomic groups collected in different parts of the world.

## CONCLUSIONS

In this study we tested anti-influenza activity of extracts derived by microalgae and cyanobacteria collected from Paraguaçu River, Shallow Lake System of Lençóis Maranhenses, Guanabara Bay estuary and Abrolhos coral reefs. We found seven extracts, in ethyl acetate solvent, that inhibit seasonal influenza A and B replication in MDCK cells in more than 80%. Two extracts from *Leptolyngbya* sp. (cyanobacteria) and Chlorellaceae family (microalgae) inhibited influenza A and B replication and neuraminidase activity, from OST-sensitive and resistance strains and lineages. Thus, these organisms are important for bioprospecting in antiviral research.

## ACKNOWLEDGEMENTS

Great thanks to the International Society for Influenza and other Respiratory Viruses Diseases-Antiviral Group (isrv-AVG), the Centers for Disease Control and Prevention (CDC), Influenza Reagent Resources (IRR) and Hoffman-La Roche Inc., Basel, Switzerland for donating materials to conduct this investigation.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) and Instituto Oswaldo Cruz - Fiocruz. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:  
Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).  
Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ).  
Instituto Oswaldo Cruz - Fiocruz.

### Competing Interests

Fabiano Thompson is an Academic Editor for PeerJ.

### Author Contributions

- Thauane Silva performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Paulo S. Salomon and Lidilhone Hamerski conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

- Juline Walter performed the experiments, prepared figures and/or tables, approved the final draft.
- Rafael B. Menezes, José Edson Siqueira, Aline Santos, Jéssica Aparecida Moraes Santos, Natália Ferme, Thaise Guimarães, Giovana O. Fistarol and Paulo I. Hargreaves performed the experiments, approved the final draft.
- Cristiane Thompson conceived and designed the experiments and contributed reagents/materials/analysis tools.
- Fabiano Thompson and Marilda Siqueira conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Thiago Moreno Souza conceived and designed the experiments, analyzed the data, approved the final draft.
- Milene Miranda conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

The field experiments were approved by Ministério do Meio Ambiente (MMA), Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) and Sistema de Autorização e Informação em Biodiversidade (SISBIO), approval number: 35854-2. The access to genetic heritage was approved by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), approval number: 010339/2014-0; and Sistema Nacional de Gestão do Patrimônio Genético e do Conhecimento Tradicional Associado (SisGen), approval number: A03D5C4.

### Data Availability

The following information was supplied regarding data availability:

The raw data are provided in the [Supplemental Files](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5716#supplemental-information>.

## REFERENCES

- Abed Y, Boivin G. 2017.** A review of clinical influenza A and B infections with reduced susceptibility to both Oseltamivir and Zanamivir. *Open Forum Infectious Diseases* 4(3):ofx105 DOI 10.1093/ofid/ofx105.
- Brass AL, Huang IC, Benita Y, John SP, Krishnan MN, Feeley EM, Ryan BJ, Weyer JL, Van der Weyden L, Fikrig E, Adams DJ, Xavier RJ, Farzan M, Elledge SJ. 2009.** The IFITM proteins mediate cellular resistance to influenza A H1N1 virus, West Nile virus, and dengue virus. *Cell* 139:1243–1254 DOI 10.1016/j.cell.2009.12.017.



- Camp JV, Chu YK, Chung DH, McAllister RC, Adcock RS, Gerlach RL, Wiemken TL, Peyrani P, Ramirez JA, Summersgill JT, Jonsson CB. 2013. Phenotypic differences in virulence and immune response in closely related clinical isolates of influenza A 2009 H1N1 pandemic viruses in mice. *PLOS ONE* 8:e56602 DOI 10.1371/journal.pone.0056602.
- Chen YH, Chang GK, Kuo SM, Huang SY, Hu IC, Lo YL, Shih SR. 2016. Well-tolerated Spirulina extract inhibits influenza virus replication and reduces virus-induced mortality. *Scientific Reports* 6:24253 DOI 10.1038/srep24253.
- Cheyrier V, Comte G, Davies KM, Lattanzio V, Martens S. 2013. Plant phenolics: recent advances on their biosynthesis, genetics, and ecophysiology. *Plant Physiology and Biochemistry* 72:1–20 DOI 10.1016/j.plaphy.2013.05.009.
- Del Giudice G, Rappuoli R. 2015. Inactivated and adjuvanted influenza vaccines. *Current Topics in Microbiology and Immunology* 386:151–180 DOI 10.1007/82\_2014\_406.
- Delaux PM, Radhakrishnan GV, Jayaraman D, Cheema J, Malbreil M, Volkening JD, Sekimoto H, Nishiyama T, Melkonian M, Pokorny L, Rothfels CJ, Sederoff HW, Stevenson DW, Surek B, Zhang Y, Sussman MR, Dunand C, Morris RJ, Roux C, Wong GK, Oldroyd GE, Ane JM. 2015. Algal ancestor of land plants was preadapted for symbiosis. *Proceedings of the National Academy of Sciences of the United States of America* 112:13390–13395 DOI 10.1073/pnas.1515426112.
- Desai TM, Marin M, Chin CR, Savidis G, Brass AL, Melikyan GB. 2014. IFITM3 restricts influenza A virus entry by blocking the formation of fusion pores following virus-endosome hemifusion. *PLOS Pathogens* 10:e1004048 DOI 10.1371/journal.ppat.1004048.
- Ding Y, Zeng L, Li R, Chen Q, Zhou B, Cheng PL, Yutao W, Zheng J, Yang Z, Zhang FCP. 2017. The Chinese prescription lianhuapingwen capsule exerts anti-influenza activity through the inhibition of viral propagation and impacts immune function. *BMC Complementary and Alternative Medicine* 17:130 DOI 10.1186/s12906-017-1585-7.
- Dixit R, Khandaker G, Ilgoutz S, Rashid H, Booy R. 2013. Emergence of oseltamivir resistance: control and management of influenza before, during and after the pandemic. *Infectious Disorders Drug Targets* 13:34–45 DOI 10.2174/18715265112129990006.
- Falaise C, Francois C, Travers MA, Morga B, Haure J, Tremblay R, Turcotte F, Pasetto P, Gastineau R, Hardivillier Y, Leignel V, Mouget JLCP. 2016. Antimicrobial compounds from Eukaryotic Microalgae against human Pathogens and diseases in aquaculture. *Marine Drugs* 14(9):159 DOI 10.3390/md14090159.
- Fistarol GO, Hargreaves P, Walter JM, Viana TV, Brito CL, Gomes P, Rezende CE, Gregoracci G, Rua C, Thompson CC, Thompson FL, Salomon PS. 2018. Rapid isolation of culturable microalgae from a tropical shallow lake system. *Journal of Applied Phycology* 30:1807–1819 DOI 10.1007/s10811-018-1404-7.
- Ghoke SS, Sood R, Kumar N, Pateriya AK, Bhatia S, Mishra A, Dixit R, Singh VK, Desai DN, Kulkarni DD, Dimri U, Singh VPCP. 2018. Evaluation of antiviral activity of *Ocimum sanctum* and *Acacia arabica* leaves extracts against H9N2 virus using

- embryonated chicken egg model. *BMC Complementary and Alternative Medicine* **18**:174 DOI [10.1186/s12906-018-2238-1](https://doi.org/10.1186/s12906-018-2238-1).
- Gogineni V, Hamann MTCPCN. 2018.** Marine natural product peptides with therapeutic potential: chemistry, biosynthesis, and pharmacology. *Biochimica et Biophysica Acta/General Subjects* **1862**:81–196 DOI [10.1016/j.bbagen.2017.08.014](https://doi.org/10.1016/j.bbagen.2017.08.014).
- Gong M, Bassi A. 2016.** Carotenoids from microalgae: a review of recent developments. *Biotechnology Advances* **34**:1396–1412 DOI [10.1016/j.biotechadv.2016.10.005](https://doi.org/10.1016/j.biotechadv.2016.10.005).
- Gorham P, McLachlan J, Hammer U, Kim W. 1964.** Isolation and culture of toxic strains of *Anabaena flos-aquae* (Lyngb.) de BrÃ©b. *Verhandlungen des Internationalen Verein Limnologie* **15**:796–804.
- Guillard R, Lorenzen C. 1972.** Yellow-green algae with chlorophyllide c. *Journal of Phycology* **8**:10–14 DOI [10.1111/j.0022-3646.1972.00010.x](https://doi.org/10.1111/j.0022-3646.1972.00010.x).
- Gupta DK, Kaur P, Leong ST, Tan LT, Prinsep MR, Chu JJCP. 2014.** Anti-Chikungunya viral activities of aplysiatoxin-related compounds from the marine cyanobacterium *Trichodesmium erythraeum*. *Marine Drugs* **12**:115–127 DOI [10.3390/md12010115](https://doi.org/10.3390/md12010115).
- Hasui M, Matsuda M, Okutani K, Shigeta S. 1995.** *In vitro* antiviral activities of sulfated polysaccharides from a marine microalga (*Cochlodinium polykrikoides*) against human immunodeficiency virus and other enveloped viruses. *International Journal of Biological Macromolecules* **17**:293–297 DOI [10.1016/0141-8130\(95\)98157-T](https://doi.org/10.1016/0141-8130(95)98157-T).
- Hour MJ, Huang SH, Chang CY, Lin YK, Wang CY, Chang YS, Lin CW. 2013.** Baicalein, Ethyl Acetate, and Chloroform extracts of *Scutellaria baicalensis* inhibit the Neuraminidase activity of Pandemic 2009 H1N1 and seasonal influenza A viruses. *Evidence-based Complementary and Alternative Medicine* **2013**:750803 DOI [10.1155/2013/750803](https://doi.org/10.1155/2013/750803).
- Hsieh CF, Chen YL, Lin CF, Ho JY, Huang CH, Chiu CH, Hsieh PW, Horng JT. 2016.** An extract from *Taxodium distichum* targets hemagglutinin- and neuraminidase-related activities of influenza virus *in vitro*. *Scientific Reports* **6**:36015 DOI [10.1038/srep36015](https://doi.org/10.1038/srep36015).
- Hynstova V, Sterbova D, Klejdus B, Hedbavny J, Huska D, Adam V. 2018.** Separation, identification and quantification of carotenoids and chlorophylls in dietary supplements containing *Chlorella vulgaris* and *Spirulina platensis* using High Performance Thin Layer Chromatography. *Journal of Pharmaceutical and Biomedical Analysis* **148**:108–118 DOI [10.1016/j.jpba.2017.09.018](https://doi.org/10.1016/j.jpba.2017.09.018).
- Iwai M, Yokono M. 2017.** Light-harvesting antenna complexes in the moss *Physcomitrella patens*: implications for the evolutionary transition from green algae to land plants. *Current Opinion in Plant Biology* **37**:94–101 DOI [10.1016/j.pbi.2017.04.002](https://doi.org/10.1016/j.pbi.2017.04.002).
- Khalid M, Khalid N, Ahmed I, Hanif R, Ismail M, Janjua H. 2017.** Comparative studies of three novel fresh water microalgae strains for synthesis of silver nanoparticles: Insights of characterization, antibacterial, cytotoxicity and antiviral activities.
- Kim M, Yim JH, Kim SY, Kim HS, Lee WG, Kim SJ, Kang PS, Lee CK. 2012.** *In vitro* inhibition of influenza A virus infection by marine microalga-derived sulfated

polysaccharide p-KG03. *Antiviral Research* **93**:253–259

DOI [10.1016/j.antiviral.2011.12.006](https://doi.org/10.1016/j.antiviral.2011.12.006).

**Kiuru P, D’Auria MV, Muller CD, Tammela P, Vuorela H, Yli-Kauhaluoma J. 2014.**

Exploring marine resources for bioactive compounds. *Planta Medica* **80**:1234–1246

DOI [10.1055/s-0034-1383001](https://doi.org/10.1055/s-0034-1383001).

**Lopes e Souza TM, Fintelman-Rodrigues N, Resende PC, Mesquita M, Gregianini**

**TS, Bozza FA, Pecego AC, Fernandes SB, Cury AL, Riediger IN, Siqueira MM.**

**2015.** Oseltamivir-resistant influenza A(H1N1)pdm2009 strains found in Brazil are endowed with permissive mutations, which compensate the loss of fitness imposed by antiviral resistance. *Memorias do Instituto Oswaldo Cruz* **110**:101–105

DOI [10.1590/0074-02760140330](https://doi.org/10.1590/0074-02760140330).

**Lopes VR, Schmidtke M, Helena Fernandes M, Martins R, Vasconcelos V. 2011.**

Cytotoxicity in L929 fibroblasts and inhibition of herpes simplex virus type 1 Kupka by estuarine cyanobacteria extracts. *Toxicology In Vitro* **25**:944–950

DOI [10.1016/j.tiv.2011.03.003](https://doi.org/10.1016/j.tiv.2011.03.003).

**Martinez Andrade KA, Lauritano C, Romano G, Ianora ACP. 2018.** Marine microalgae with anti-cancer properties. *Marine Drugs* **16**(5):165 DOI [10.3390/md16050165](https://doi.org/10.3390/md16050165).

**Mayer AM, Hamann MT. 2004.** Marine pharmacology in 2000: marine compounds with antibacterial, anticoagulant, antifungal, anti-inflammatory, antimalarial, antiplatelet, antituberculosis, and antiviral activities; affecting the cardiovascular, immune, and nervous systems and other miscellaneous mechanisms of action.

*Marine Biotechnology* **6**:37–52 DOI [10.1007/s10126-003-0007-7](https://doi.org/10.1007/s10126-003-0007-7).

**Mayer AM, Rodriguez AD, Berlinck RG, Hamann MT. 2007.** Marine pharmacology in 2003-4: marine compounds with anthelmintic antibacterial, anticoagulant, antifungal, anti-inflammatory, antimalarial, antiplatelet, antiprotozoal, antituberculosis, and antiviral activities; affecting the cardiovascular, immune and nervous systems, and other miscellaneous mechanisms of action. *Comparative Biochemistry and Physiology—Part C: Toxicology & Pharmacology* **145**:553–581 DOI [10.1016/j.cbpc.2007.01.015](https://doi.org/10.1016/j.cbpc.2007.01.015).

**Mayer AMS, Rodriguez AD, Tagliatalata-Scafati O, Fusetani N. 2017.** Marine pharmacology in 2012-2013: marine compounds with antibacterial, antidiabetic, antifungal, anti-inflammatory, antiprotozoal, antituberculosis, and antiviral activities; affecting the immune and nervous systems, and other miscellaneous mechanisms of action. *Marine Drugs* **15**(9):273 DOI [10.3390/md15090273](https://doi.org/10.3390/md15090273).

**Mendes Gda S, Soares AR, Martins FO, Albuquerque MC, Costa SS, Yoneshigue-Valentin Y, Gestinari LM, Santos N, Romanos MT. 2010.** Antiviral activity of the green marine alga *Ulva fasciata* on the replication of human metapneumovirus. *Revista do Instituto de Medicina Tropical de Sao Paulo* **52**:3–10 DOI [10.1590/S0036-46652010000100001](https://doi.org/10.1590/S0036-46652010000100001).

**Mesquita M, Fintelman-Rodrigues N, Sacramento CQ, Abrantes JL, Costa E, Temerozo JR, Siqueira MM, Bou-Habib DC, Souza TM. 2014.** HIV-1 and its gp120 inhibits the influenza A(H1N1)pdm09 life cycle in an IFITM3-dependent fashion. *PLOS ONE* **9**:e101056 DOI [10.1371/journal.pone.0101056](https://doi.org/10.1371/journal.pone.0101056).

- Mizgerd JP. 2008.** Acute lower respiratory tract infection. *New England Journal of Medicine* 358:716–727 DOI 10.1056/NEJMra074111.
- Montalvao S, Demirel Z, Devi P, Lombardi V, Hongisto V, Perala M, Hattara J, Imamoglu E, Tilvi SS, Turan G, Dalay MC, Tammela P. 2016.** Large-scale bio-prospecting of cyanobacteria, micro- and macroalgae from the Aegean Sea. *New Biotechnology* 33:399–406 DOI 10.1016/j.nbt.2016.02.002.
- Mundt S, Nowotny A, Mentel R, Lesnau A, Lindequist U. 1997.** Antiviral activity of the cyanobacterium *Microcystis aeruginosa* SPH 01.
- Niedermeyer TH. 2015.** Anti-infective natural products from Cyanobacteria. *Planta Medica* 81:1309–1325 DOI 10.1055/s-0035-1546055.
- Pichon M, Lina B, Josset L. 2017.** Impact of the respiratory microbiome on host responses to respiratory viral infection. *Vaccine* 5(4):40 DOI 10.3390/vaccines5040040.
- Rajasekaran D, Palombo EA, Chia Yeo T, Lim Siok Ley D, Lee Tu C, Malherbe F, Grollo LCP. 2013.** Identification of traditional medicinal plant extracts with novel anti-influenza activity. *PLOS ONE* 8:e79293 DOI 10.1371/journal.pone.0079293.
- Reed J, Mounch M. 1938.** An endpoint assay to evaluate the 50% tissue cytopathic effect. *Journal of General Virology* 5:25–29.
- Reynolds C, Jaworski G. 1978.** Enumeration of natural *Microcystis* populations. *Brazilian Journal of Phycology* 13:269–277.
- Roehm NW, Rodgers GH, Hatfield SM, Glasebrook AL. 1991.** An improved colorimetric assay for cell proliferation and viability utilizing the tetrazolium salt XTT. *Journal of Immunological Methods* 142:257–265 DOI 10.1016/0022-1759(91)90114-U.
- Rotrosen ET, Neuzil KM. 2017.** Influenza: a global perspective. *Pediatric Clinics of North America* 64:911–936 DOI 10.1016/j.pcl.2017.03.007.
- Sacramento CQ, Marttorelli A, Fintelman-Rodrigues N, De Freitas CS, De Melo GR, Rocha ME, Kaiser CR, Rodrigues KF, Da Costa GL, Alves CM, Santos-Filho O, Barbosa JP, Souza TM. 2015.** Aureonitol, a fungi-derived Tetrahydrofuran, inhibits influenza replication by targeting its surface Glycoprotein Hemagglutinin. *PLOS ONE* 10:e0139236 DOI 10.1371/journal.pone.0139236.
- Sathasivam R, Ki JS. 2018.** A review of the biological activities of microalgal carotenoids and their potential Use in healthcare and cosmetic industries. *Marine Drugs* 16(1):26 DOI 10.3390/md16010026.
- Savidis G, Perreira JM, Portmann JM, Meraner P, Guo Z, Green S, Brass AL. 2016.** The IFITMs inhibit Zika virus replication. *Cell Reports* 15:2323–2330 DOI 10.1016/j.celrep.2016.05.074.
- Scudiero DA, Shoemaker RH, Paull KD, Monks A, Tierney S, Nofziger TH, Currens MJ, Seniff D, Boyd MR. 1988.** Evaluation of a soluble tetrazolium/formazan assay for cell growth and drug sensitivity in culture using human and other tumor cell lines. *Cancer Research* 48:4827–4833.
- Sharaf M, Amara A, Aboul-Enein A, Helmi S, Ballot A, Astani A, Schnitzler P. 2010.** Molecular authentication and characterization of the antiherpetic activity of the cyanobacterium *Arthrospira fusiformis*. *Pharmazie* 65:132–136.

- Shoji M, Woo SY, Masuda A, Win NN, Ngwe H, Takahashi E, Kido H, Morita H, Ito T, Kuzuhara T. 2017.** Anti-influenza virus activity of extracts from the stems of *Jatropha multifida* Linn. collected in Myanmar. *BMC Complementary and Alternative Medicine* 17:96 DOI 10.1186/s12906-017-1612-8.
- Singh RS, Walia AK, Khattar JS, Singh DP, Kennedy JF. 2017.** Cyanobacterial lectins characteristics and their role as antiviral agents. *International Journal of Biological Macromolecules* 102:475–496 DOI 10.1016/j.ijbiomac.2017.04.041.
- Sood R, Swarup D, Bhatia S, Kulkarni DD, Dey S, Saini M, Dubey SC. 2012.** Antiviral activity of crude extracts of *Eugenia jambolana* Lam. against highly pathogenic avian influenza (H5N1) virus. *Indian Journal of Experimental Biology* 50:179–186.
- Souza TM, Abrantes JL, De AER, Leite Fontes CF, Frugulhetti IC. 2007.** The alkaloid 4-methylaaptamine isolated from the sponge *Aaptos aaptos* impairs Herpes simplex virus type 1 penetration and immediate-early protein synthesis. *Planta Medica* 73:200–205 DOI 10.1055/s-2007-967109.
- Souza TM, Mesquita M, Resende P, Machado V, Gregianini TS, Fernandes SB, Oliveira IC, Rosa Mdo C, Marinelli R, De Azeredo-Lima CH, Motta FC, Aguiar-Oliveira Mde L, Siqueira MM. 2011.** Antiviral resistance surveillance for influenza A virus in Brazil: investigation on 2009 pandemic influenza A (H1N1) resistance to oseltamivir. *Diagnostic Microbiology and Infectious Disease* 71:98–99 DOI 10.1016/j.diagmicrobio.2011.05.006.
- Souza TM, Resende PC, Fintelman-Rodrigues N, Gregianini TS, Ikuta N, Fernandes SB, Cury AL, Rosa Mdo C, Siqueira MM. 2013.** Detection of oseltamivir-resistant pandemic influenza A(H1N1)pdm2009 in Brazil: can community transmission be ruled out? *PLOS ONE* 8:e80081 DOI 10.1371/journal.pone.0080081.
- Soye KJ, Trottier C, Di Lenardo TZ, Restori KH, Reichman L, Miller Jr WH, Ward BJ. 2013.** *In vitro* inhibition of mumps virus by retinoids. *Virology Journal* 10:337 DOI 10.1186/1743-422x-10-337.
- Szretter KJ, Balish AL, Katz JM. 2006.** Influenza: propagation, quantification, and storage. *Current Protocols in Microbiology* 15:Unit 15G.1.1–15G.1.22 DOI 10.1002/0471729256.mc15g01s3.
- Villalon-Letelier F, Brooks AG, Saunders PM, Londrigan SL, Reading PC. 2017.** Host cell restriction factors that limit influenza A infection. *Viruses* 9(12):376 DOI 10.3390/v9120376.
- World Health Organization (WHO). 2011.** Manual for the laboratory diagnosis and virological surveillance of influenza. World Health Organization. Available at [http://www.who.int/influenza/gisrs\\_laboratory/manual\\_diagnosis\\_surveillance\\_influenza/en/](http://www.who.int/influenza/gisrs_laboratory/manual_diagnosis_surveillance_influenza/en/).
- World Health Organization (WHO). 2017.** Up to 650,000 people die of respiratory diseases linked to seasonal flu each year. Available at <http://www.who.int/mediacentre/news/releases/2017/seasonal-flu/en/>.
- World Health Organization (WHO). 2018a.** Influenza (Avian and other zoonotic). Available at [http://www.who.int/mediacentre/factsheets/avian\\_influenza/en/](http://www.who.int/mediacentre/factsheets/avian_influenza/en/).

- World Health Organization (WHO). 2018b.** Influenza (Seasonal). Available at <http://www.who.int/mediacentre/factsheets/fs211/en/>.
- Wu Q, Yu C, Yan Y, Chen J, Zhang C, Wen X. 2010.** Antiviral flavonoids from *Mosla scabra*. *Fitoterapia* **81**:429–433 DOI [10.1016/j.fitote.2009.12.005](https://doi.org/10.1016/j.fitote.2009.12.005).
- Zainuddin EN, Mentel R, Wray V, Jansen R, Nimtz M, Lalk M, Mundt S. 2007.** Cyclic depsipeptides, ichthyopeptins A and B, from *Microcystis ichthyoblabe*. *Journal of Natural Products* **70**:1084–1088 DOI [10.1021/np060303s](https://doi.org/10.1021/np060303s).
- Zainuddin EN, Mundt S, Wegner U, Mentel R. 2002.** Cyanobacteria a potential source of antiviral substances against influenza virus. *Medical Microbiology and Immunology* **191**:181–182 DOI [10.1007/s00430-002-0142-1](https://doi.org/10.1007/s00430-002-0142-1).
- Zhu W, Wang C, Wang BZ. 2017.** From variation of influenza viral proteins to vaccine development. *International Journal of Molecular Sciences* **18**(7):1554 DOI [10.3390/ijms18071554](https://doi.org/10.3390/ijms18071554).