



# Effects of graded concentrations of supplemental selenium on selenium concentrations in tissues and prediction equations for estimating dietary selenium intake in pigs

Ah Reum Son<sup>1,\*</sup>, Jin-young Jeong<sup>2,\*</sup>, Kyu Ree Park<sup>1</sup>, Minseok Kim<sup>2,5</sup>, Sung Dae Lee<sup>2</sup>, Ji-Hyock Yoo<sup>3</sup>, Yoon-Jung Do<sup>4</sup>, Kondreddy Eswar Reddy<sup>2</sup> and Hyun-Jeong Lee<sup>2</sup>

<sup>1</sup>Department of Animal Science and Technology, Konkuk University, Seoul, Republic of Korea

<sup>2</sup>Animal Nutritional Physiology Team, National Institute of Animal Science, Rural Development Administration, Wanju-gun, Republic of Korea

<sup>3</sup>Department of Agro-food Safety, National Institute of Agricultural Science, Rural Development Administration, Wanju-gun, Republic of Korea

<sup>4</sup>Division of Animal Diseases & Health, National Institute of Animal Science, Rural Development Administration, Wanju-gun, Republic of Korea

<sup>5</sup>Current affiliation: Department of Animal Science, College of Agriculture and Life Sciences, Chonnam National University, Gwangju, Republic of Korea

\*These authors contributed equally to this work.

## ABSTRACT

The experiment was conducted to determine the effects of graded dietary selenium (Se) on organ weight and Se concentrations in tissues and to develop equations for estimating dietary Se intake in pigs. Sixteen barrows (initial body weight =  $30.0 \pm 2.6$ ) were allotted to four dietary treatments including graded Se supplementations with 0, 1, 5, and 50 mg/kg of diet. The experimental diets fed to the pigs for 30 d, and then the pigs were euthanized, and the organs, muscle, and urine samples were collected. The hair and blood samples of pigs were collected on d 15 and 30. Equations were developed for predicting daily Se intake using the Se concentration in plasma, hair, liver, kidneys, muscle, or urine. For graded dietary Se concentrations, linear and quadratic effects on the final body weight, weight and relative weight of liver and kidneys were not observed. The Se concentration in plasma, hair, liver, kidneys, muscle, and urine were linearly and quadratically increased as dietary Se concentration increased ( $P < 0.001$ ). The dietary Se concentration was positively correlated with the Se concentrations in the plasma, organs, muscle, and urine ( $r > 0.81$ ,  $P < 0.001$ ). The equations for estimating dietary Se intake using the Se concentration in the plasma, hair, or organ as an independent variable were significant ( $P < 0.05$ ). In conclusion, the dietary Se concentration was well reflected in the Se concentration in the plasma, hair, liver, kidneys, and urine. The Se concentration in the plasma, hair, liver, and kidneys can be used as an independent variable for estimating the Se intake.

Submitted 6 April 2018  
Accepted 19 September 2018  
Published 19 October 2018

Corresponding authors  
Kondreddy Eswar Reddy,  
[dreswar4u@gmail.com](mailto:dreswar4u@gmail.com),  
[dreswar4uk@rediffmail.com](mailto:dreswar4uk@rediffmail.com)  
Hyun-Jeong Lee, [hyunj68@korea.kr](mailto:hyunj68@korea.kr)

Academic editor  
Joao Rocha

Additional Information and  
Declarations can be found on  
page 11

DOI 10.7717/peerj.5791

 Copyright  
2018 Son et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

**Subjects** Agricultural Science, Toxicology

**Keywords** Selenium, Swine, Accumulation, Organ

## INTRODUCTION

Selenium (Se) is one of essential trace minerals, and it plays important biochemical roles as a component of selenoproteins to prevent oxidative damages in the body (*Barbosa et al., 2017*). Furthermore, as part of selenoproteins, the Se modulates physiological processes related with the immune system and thyroid metabolism for pigs (*Mahan, 2001; Dalgaard et al., 2018*). The Se deficiency in pigs induces sudden death, reduced reproduction and milk production, and impaired immune response (*National Research Council, 2012*). Positive effects of supplemental dietary Se on meat quality and antioxidative status for pigs have been reported (*Calvo et al., 2017; Schwarz et al., 2017*). However, excessive consumption of Se occurs toxicity in animals. For the pigs, generally 5 to 25 ppm of Se in diets causes a chronic selenosis. Major symptoms of selenosis are reduced growth rate and feed intake, hair loss, hoof lesion, vomiting, and incoordination (*Mahan, 2001; National Research Council, 2005*).

For Se, safety margin between adequate amount and overconsumption is narrow in humans and other mammals, so it has become more important to estimate the Se intake (*Reis et al., 2017; Ullah et al., 2018*). In addition, deficient or excessive Se intake depends on regions because the Se concentration in vegetables can be largely affected by Se concentration in soil (*Reis et al., 2017*). Natural and anthropogenic sources of Se including bedrocks, volcanic activity, Se-fertilizers, and coal burning influence the Se concentration in soil (*Winkel et al., 2015*), and this subsequently can affect the Se concentration in plants and animal body. Moreover, livestock products have relatively high Se concentration compared with fruits and cereals (*Choi et al., 2009; Ullah et al., 2018*). It has been reported that Se concentration in organs and tissues of pigs increased as dietary Se concentrations increased (*Goehring et al., 1984; Kim & Mahan, 2001a; Kim & Mahan, 2001b; Mateo et al., 2007*). In EU and USA, supplementation of Se in complete feeds for swine is regulated at a level not to exceed 0.2 and 0.3 mg/kg, respectively. Regulatory limits for the Se in swine diets can be determined based on accumulation rate of dietary Se to edible portion of pigs. However, studies on relationship between dietary Se intake and concentration in tissues of pigs are limited.

Therefore, the objectives of this experiment were (1) to determine the effects of graded supplemental Se concentrations on body and organ weights and the Se concentrations in the plasma, hair, soft tissues (liver, kidneys, and muscle), and urine; and (2) to generate equations for estimating daily Se intake using the Se concentrations in the plasma, hair, soft tissues, or urine as an independent variable for pigs.

## MATERIALS AND METHODS

The protocols used for the animal experimental procedures were reviewed and approved by the Institutional Animal Care and Use Committee of the National Institute of Animal Science, South Korea No. 2015-147 of 29 May 2015. The experimental procedure of current study was performed as described in our previous study (*Reddy et al., 2017*).

**Table 1** Ingredients and chemical compositions of control diet (as-fed basis).

Item	Control diet
Ingredients (%)	
Ground corn	58.56
Soybean meal (46% crude protein )	14.00
Extruded soybean meal	12.00
Whey powder (12% crude protein)	7.00
Fish meal	3.45
Soybean oil	1.60
L-Lysine·hydrochloride (78%)	0.43
DL-Methionine (99%)	0.14
L-Threonine (99%)	0.12
Monocalcium phosphate	1.08
Ground limestone	0.60
Choline chloride (50%)	0.20
Sodium chloride	0.32
Vitamin-trace mineral premix <sup>a</sup>	0.50
Calculated nutrients (%)	
Metabolizable energy (kcal/kg)	3,444
Crude fiber	2.29
Crude protein	20.78
Lysine	1.47
Methionine	0.49
Calcium	0.75
Phosphorus	0.45
Selenium <sup>b</sup> (mg/kg)	0.13

**Notes.**

<sup>a</sup>Provided the following quantities per kg of complete diet: vitamin A 11,000 IU; vitamin D<sub>3</sub> 1,500 IU; vitamin E 44.1 IU; vitamin K<sub>3</sub> 4.0 mg; vitamin B<sub>1</sub> 1.4 mg; vitamin B<sub>2</sub> 5.22 mg; vitamin B<sub>5</sub> 20.0 mg; vitamin B<sub>12</sub> 0.01 mg; Niacin 26.0 mg; Pantothenic acid 14 mg; Folic acid 0.8 mg; Biotin 44 µg; Fe 100.0 mg as iron sulfate; Cu 16.50 mg as copper sulfate; Zn 90.0 mg as zinc sulfate; Mn 35.0 mg as manganese sulfate; I 0.30 mg as calcium iodate.

<sup>b</sup>Calculated value from *National Research Council (2012)* and *Prabhu (2015)*.

**Animals, diets, and experimental design**

A total of 16 barrows with a mean initial body weight (BW) 30 kg (standard deviation = 2.6) was randomly assigned to four dietary treatments. A control diet without dietary Se supplementation was prepared mainly based on corn and soybean meal ([Table 1](#)). The control diet was formulated to meet or exceed the nutrient requirements estimates (*National Research Council, 2012*). All pigs were fed the control diet during the experimental period. The Se as a sodium selenite put into a soft capsule, and then the encapsulated Se was supplemented to additional three dietary treatment groups along with each meal (1, 5, or 50 mg/kg of diet). The encapsulated Se was mixed with each meal, and an experimenter checked and confirmed that the pigs completely consumed their feed allowance and capsule at each feeding time. An experimental period lasted for 30 d, and the daily feed allowance of each pigs was 1 kg and 2 kg for first and last 15 d, respectively. The pigs were fed two equal meals daily, and freely accessed to water.

**Table 2** Final body weight (BW) and weight of liver and kidneys for pigs fed the diets containing graded concentrations of supplemental selenium (Se).<sup>a</sup>

Item	Supplemental Se, mg/kg of diet				SEM <sup>b</sup>	P-value	
	0	1	5	50		Linear	Quadratic
Initial BW, kg	29.0	28.8	29.4	29.9	–	–	–
Final BW, kg	52.0	49.7	52.3	52.1	2.0	0.724	0.698
Weight, g							
Liver	1,672	1,658	1,870	1,772	89	0.599	0.100
Kidney	135	141	160	160	9	0.141	0.082
Relative weight to BW, %							
Liver	3.24	3.34	3.57	3.41	0.16	0.776	0.165
Kidney	0.26	0.28	0.31	0.31	0.02	0.186	0.142

**Notes.**<sup>a</sup>Each least squares mean represents four observations.<sup>b</sup>SEM, standard error of the means.**Table 3** Selenium (Se) concentrations in plasma, hair, liver, kidneys, muscle, and urine for pigs fed the diets containing graded concentrations of supplemental Se.<sup>a</sup>


Item	Supplemental Se, mg/kg of diet				SEM <sup>b</sup>	P-value	
	0	1	5	50		Linear	Quadratic
Plasma, mg/L							
d 15	0.25	0.32	1.96	3.52	0.16	<0.001	<0.001
d 30	0.18	0.35	2.10	4.25	0.19	<0.001	<0.001
Hair, mg/kg							
d 15	0.42	0.45	1.06	2.42	0.08	<0.001	<0.001
d 30	0.41	0.46	1.47	5.19	0.05	<0.001	<0.001
Liver, mg/kg	9.62	16.2	41.9	83.9	2.22	<0.001	<0.001
Kidneys, mg/kg	18.5	37.3	62.8	114.9	2.4	<0.001	<0.001
Muscle, mg/kg	2.62	3.52	4.57	5.78	0.15	<0.001	<0.001
Urine, mg/L	0.06	0.19	1.01	1.47	0.08	<0.001	<0.001

**Notes.**<sup>a</sup>Each least squares mean represents four observations.<sup>b</sup>SEM, standard error of the means.**Sampling and processing**

On d 0 and 30, all pigs were individually weighed, and blood and hair were collected from each pig. The blood was collected from jugular vein, and then samples were taken into metal free heparinized collection tubes (BD Vacutainer<sup>®</sup>, Cat. No. 366480, Franklin Lakes, NJ, USA), and immediately centrifuged at 4 °C for 15 min at 2,500 rpm. The plasma samples were stored at –20 °C for until analysis. At the end of experiment, all pigs were euthanized by an anesthetic overdose with the combination of barbiturates and pentobarbital. After slaughter, the liver, both kidneys, and thigh muscle samples were collected, and then immediately frozen by using the liquid nitrogen, and stored at –80 °C. From the urinary bladder, the urine samples were collected by using syringe, and were stored at –20 °C for until analysis.


**Table 4** Correlation coefficients ( $r$ ) between supplemental selenium (Se) concentration and final body weight (BW), liver and kidneys weights relative to BW, and Se concentration in blood, hair, liver, kidneys, muscle, and urine.

Item	Supplemental Se concentrations	
	$r$	$P$ -value
Final BW (kg)	0.100	0.713
Liver weight relative to BW	0.077	0.777
Kidneys weight relative to BW	0.343	0.194
Se concentration		
Plasma (mg/L)		
d 15	0.884	<0.001
d 30	0.909	<0.001
Hair (mg/kg)		
d 15	0.960	<0.001
d 30	0.990	<0.001
Liver (mg/kg)	0.936	<0.001
Kidneys (mg/kg)	0.929	<0.001
Muscle (mg/kg)	0.840	<0.001
Urine (mg/L)	0.810	<0.001



**Figure 1** Linear and quadratic regression equations for estimating daily selenium (Se) intake (mg/d) based on the Se concentration in plasma (mg/L) on d 15 and 30. Each data point represents least squares mean of four observations.  $r^2$ , coefficient of determination; RMSE, root mean square of error.

Full-size [DOI: 10.7717/peerj.5791/fig-1](https://doi.org/10.7717/peerj.5791/fig-1)



**Figure 2** Linear and quadratic regression equations for estimating daily selenium (Se) intake (mg/d) based on the Se concentration in hair (mg/kg) on d 15 and 30. Each data point represents least squares mean of four observations.  $r^2$ , coefficient of determination; RMSE, root mean square of error.

Full-size [DOI: 10.7717/peerj.5791/fig-2](https://doi.org/10.7717/peerj.5791/fig-2)


### Sample digestion procedure and elemental analysis

The liver, kidneys, and muscle samples were dried at  $-105\text{ }^{\circ}\text{C}$  until constant weight is reached (Linden, Olsson & Oskarsson, 1999), and then finely ground before analysis. For the hair samples, a preconditioning process was performed according to a previous study (Chaturvedi et al., 2014). The amounts of samples used for each analysis were: 1 mL of plasma, 0.1 g of hair, 0.2 g of dried liver, kidneys, or muscle, and 1.5 mL of urine. Each sample was analyzed for the Se concentration in triplicate. Each digestion procedure included three blank samples.

The procedure of digestion and analysis was based on a previous study (Ashoka et al., 2009). A 2.5 mL of concentrated  $\text{HNO}_3$  and 0.5 mL of concentrated  $\text{HCl}$  were added to the samples placed in a glass tube with cap, and the tubes kept in a water bath at  $85\text{ }^{\circ}\text{C}$ . Three hours later, the samples were cooled at room temperature. After cooling, the samples were diluted to 50 mL using a 2%  $\text{HNO}_3$  solution (Cubadda, Raggi & Coni, 2006). The digested samples were analyzed for the Se concentration using an ICP-MS (Agilent 7500, Santa Clara, CA, USA).

### Statistical analysis

The MIXED procedure of SAS (SAS Institute Inc., Cary, NC, USA) were used to analyze experimental data. The model included the dietary treatment as an independent variable.



**Figure 3** Linear and quadratic regression equations for estimating daily selenium (Se) intake (mg/d) based on Se concentration in dried liver (mg/kg). Each data point represents least squares mean of four observations.  $r^2$ , coefficient of determination; RMSE, root mean square of error.


Full-size  DOI: [10.7717/peerj.5791/fig-3](https://doi.org/10.7717/peerj.5791/fig-3)

Linear and quadratic effects of dietary Se concentrations were tested using the orthogonal polynomial contrast. To generate the contrast coefficients for unequally spaced dietary Se concentrations, the IML procedure of SAS was used. The CORR procedure of SAS was used to determine the correlation between dietary Se concentration and Se concentrations of plasma, hair, soft tissues (liver, kidneys, and muscle), and urine. The REG and NLIN procedures of SAS were used to develop linear and quadratic equations for predicting daily Se intake of the pigs. The experimental unit was the pig, and level of statistical significance (alpha level) was set at 0.05.

## RESULTS

The final BW of pigs was not affected by dietary Se concentration (Table 2). The linear and quadratic effects on the weight and relative weight of liver and kidneys with increasing dietary Se concentration were not observed. The Se concentrations in plasma, hair, liver, kidneys, muscle, and urine were linearly and quadratically increased with increasing dietary Se concentration ( $P < 0.001$ ; Table 3).

The correlations between the dietary Se and final BW or relative weight to BW of organ were not observed (Table 4). There were positive correlations between the dietary Se concentration and the Se concentrations in the plasma, hair liver, kidneys, muscle, and urine ( $r > 0.81$ ,  $P < 0.001$ ). Using the Se concentration in plasma, hair, liver, kidneys, muscle, or



**Figure 4** Linear regression equation for estimating daily selenium (Se) intake (mg/d) based on the Se concentration in dried kidneys (mg/kg). Each data point represents least squares mean of four observations.  $r^2$ , coefficient of determination; RMSE, root mean square of error.


Full-size DOI: 10.7717/peerj.5791/fig-4

urine as an independent variable, the linear and quadratic equations for estimating daily Se intake were developed. The linear and quadratic equations using the Se concentration in plasma or hair were significant ( $P < 0.01$ ; Figs. 1 and 2). The quadratic equations based on the Se concentration in the dried liver or kidneys had significance ( $P < 0.05$ ; Figs. 3 and 4) and high  $r^2$  (0.998). There was no significance for the prediction equations developed based on the Se concentration in the dried muscle or urine (Figs. 5 and 6).

## DISCUSSION

The dietary Se concentration did not affect the final BW of pigs in the present study. This result agreed with previous studies (Mahan, Cline & Richert, 1999; Mateo et al., 2007). In contrast, it was reported that the dietary Se cause decreased growth performance in the pigs (Goehring et al., 1984; Mahan & Moxon, 1984; Kim & Mahan, 2001a; Kim & Mahan, 2001b). The influence of dietary Se can be affected by source of Se, age, physical status, feeding duration, and dietary Se concentration (National Research Council, 2005). In previous studies reported decreased growth performance for pigs fed diets containing excessive Se, experimental period lasted 5 to 12 wk, and the pigs had ad libitum access to the experimental diets (Goehring et al., 1984; Mahan & Moxon, 1984; Kim & Mahan, 2001a; Kim & Mahan, 2001b). However, in the current study, the daily feed allowance of pigs was restricted at 1 or 2 kg during the 30 d of experimental period. In addition, some previous






**Figure 5** Linear and quadratic regression equations for estimating daily selenium (Se) intake (mg/d) based on the Se concentration in dried muscle (mg/kg). Each data point represents least squares mean of four observations.  $r^2$ , coefficient of determination; RMSE, root mean square of error.

Full-size  DOI: [10.7717/peerj.5791/fig-5](https://doi.org/10.7717/peerj.5791/fig-5)

studies used weanling pigs (Goehring *et al.*, 1984; Mahan & Moxon, 1984) which can be more sensitive to Se compared with older pigs. For these reasons, there were differences in effects of dietary Se on the growth performance between the studies.

The Se concentrations in the plasma, hair, soft tissues, and urine increased as the dietary Se level increased in the present study. The increased Se concentration in the plasma with supplementary Se was consistent with results of previous studies fed either sodium selenite or organic Se to the pigs (Kim & Mahan, 2001a; Zhan *et al.*, 2007; Zhou *et al.*, 2009). Normal Se concentration in serum of pig ranges from 0.08 to 0.15 mg/L (Mahan, 2001). However, the Se concentration in plasma for pigs fed the control diet in the present study was slightly greater than the previous studies (0.22 vs. 0.06 to 0.12 mg/L; Kim & Mahan, 2001a; Zhan *et al.*, 2007; Zhou *et al.*, 2009).

The normal Se concentration in the tissues of pigs is kidney > liver > glandular tissue > muscle in declining order of abundance (Mahan, 2001). In the current study, the kidney and liver had also higher Se concentration regardless of dietary Se level. This result indicated that ingested Se was mainly retained the kidney and liver for the pigs. In the previous studies (Mahan & Moxon, 1984; Kim & Mahan, 2001a; Kim & Mahan, 2001b; Mahan *et al.*, 2005), relatively high Se concentration in the kidney and liver compared with other tissues was also observed in pigs fed diets with or without Se supplementation. In this study, the hair collected on d 30 had higher Se concentration rather than that collected



**Figure 6** Linear and quadratic regression equations for estimating daily selenium (Se) intake (mg/d) based on the Se concentration in urine (mg/L). Each data point represents least squares mean of four observations.  $r^2$ , coefficient of determination; RMSE, root mean square of error.

Full-size DOI: [10.7717/peerj.5791/fig-6](https://doi.org/10.7717/peerj.5791/fig-6)

on d 15. It was also reported that the Se concentration in the hair increased as feeding period of dietary Se was extended from 4 to 12 wk for both inorganic and organic Se supplementation (Kim & Mahan, 2001b). Although the feeding period of dietary Se was shorter compared with the literature, the result may confirm that the Se concentration in the hair is affected by the feeding duration of Se-containing diet.

In the current study, the prediction equations for estimating daily Se intake based on the Se concentration in the plasma, hair, soft tissues, or urine were developed. A high correlation of dietary Se to various tissues (loin, liver, heart, and hair) Se concentration was reported (Mahan et al., 2005). The coefficient of determination ( $r^2$ ) of prediction equations developed in present study was greater than 0.83 for the plasma, hair, liver, and kidney. This result may indicate that the Se concentration in the plasma, hair, liver, and kidney is used as an independent variable for estimating daily Se intake for the pigs. Moreover, the Se concentration in the plasma and hair can be used as an antemortem biomarker of dietary status for the pigs.

In conclusion, the dietary Se concentration was highly correlated with Se concentration in the plasma, hair, liver, kidney, muscle, and urine. The Se concentration in the plasma, hair, liver, and kidneys can be used as good indicators for estimating the Se intake. The prediction equations for estimating Se intake developed in present study may be useful to determine regulatory limits for Se concentration in swine diets.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was carried out with the support of the “Research Program for Agriculture Science and Technology Development (Project No. PJ01093202),” National Institute of Animal Sciences, Rural Development Administration, Republic of Korea. This paper was supported by the KU Research Professor Program of Konkuk University. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Research Program for Agriculture Science and Technology Development: PJ01093202.  
National Institute of Animal Sciences.  
Rural Development Administration.  
KU Research Professor Program of Konkuk University.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Ah Reum Son and Jin-young Jeong authored or reviewed drafts of the paper.
- Kyu Ree Park analyzed the data, prepared figures and/or tables.
- Minseok Kim and Yoon-Jung Do reviewed drafts of the paper.
- Sung Dae Lee conceived and designed the experiments, reviewed drafts of the paper.
- Ji-Hyock Yoo performed chemical analysis.
- Kondreddy Eswar Reddy conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Hyun-Jeong Lee conceived and designed the experiments, contributed reagents/materials/analysis tools, approved the final draft, reviewed drafts of the paper.

### Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The protocols used for the animal experimental procedures were reviewed and approved by the Institutional Animal Care and Use Committee of the National Institute of Animal Science, South Korea No. 2015-147 of 29 May 2015.

### Data Availability

The following information was supplied regarding data availability:

The raw data are provided in a [Data S1](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5791#supplemental-information>.

## REFERENCES

- Ashoka S, Peake BM, Bremner G, Hageman KJ, Reid MR. 2009. Comparison of digestion methods for ICP-MS determination of trace elements in fish tissues. *Analytica Chimica Acta* 653:191–199 DOI 10.1016/j.aca.2009.09.025.
- Barbosa NV, Nogueira CW, Nogara PB, De Bem AF, Aschner M, Rocha JBT. 2017. Organoselenium compounds as mimics of selenoproteins and thiol modifier agents. *Metallomics* 9:1703–1734 DOI 10.1039/C7MT00083A.
- Calvo L, Segura J, Toldrá F, Flores M, Rodríguez AI, López-Bote CJ, Rey AI. 2017. Meat quality, free fatty acid concentration, and oxidative stability of pork from animals fed diets containing different sources of selenium. *Food Science and Technology International* 23:716–728 DOI 10.1177/1082013217718964.
- Chaturvedi R, Banerjee S, Chattopadhyay P, Bhattacharjee CR, Raul P, Borah K. 2014. High iron accumulation in hair and nail of people living in iron affected areas of Assam, India. *Ecotoxicology Environmental Safety* 110:216–220 DOI 10.1016/j.ecoenv.2014.08.028.
- Choi Y, Kim J, Lee H-S, Kim C, Hwang IK, Park HK, Oh C-H. 2009. Selenium content in representative Korean foods. *Journal of Food Composition and Analysis* 22:117–122 DOI 10.1016/j.jfca.2008.11.009.
- Cubadda F, Raggi A, Coni E. 2006. Element fingerprinting of marine organisms by dynamic reaction cell inductively coupled plasma mass spectrometry. *Analytical and Bioanalytical Chemistry* 384:887–896 DOI 10.1007/s00216-005-0256-6.
- Dalgaard TS, Briens M, Engberg RM, Lauridsen C. 2018. The influence of selenium and selenoproteins on immune responses of poultry and pigs. *Animal Feed Science and Technology* 238:73–83 DOI 10.1016/j.anifeedsci.2018.01.020.
- Goehring TB, Palmer IS, Olson OE, Libal GW, Wahlstrom RC. 1984. Toxic effects of selenium on growing swine fed corn-soybean meal diets. *Journal Animal Science* 59:733–737 DOI 10.2527/jas1984.593733x.
- Kim YY, Mahan DC. 2001a. Comparative effects of high dietary levels of organic and inorganic selenium on selenium toxicity of growing-finishing pigs. *Journal Animal Science* 79:942–948 DOI 10.2527/2001.794942x.
- Kim YY, Mahan DC. 2001b. Effect of dietary selenium source, level, and pig hair color on various selenium indices. *Journal Animal Science* 79:949–955 DOI 10.2527/2001.794949x.
- Linden A, Olsson IM, Oskarsson A. 1999. Cadmium levels in feed components and kidneys of growing/finishing pigs. *Journal AOAC International* 82:1288–1297.
- Mahan DC. 2001. Selenium and vitamin E in swine nutrition. In: Lewis AJ, Southern LL, eds. *Swine nutrition*. Second edition. New York: CRC Press, LLC, 281–314.
- Mahan DC, Brendemuhl JH, Carter SD, Chiba LI, Crenshaw TD, Cromwell GL, Dove CR, Harper AF, Hill GM, Hollis GR, Kim SW, Lindemann MD, Maxwell CV, Miller PS, Nelssen JL, Richert BT, Southern LL, Stahly TS, Stein HH, Van Heugten E, Yen JT. 2005. Comparison of dietary selenium fed to grower-finisher pigs from various

- regions of the United States on resulting tissue Se and loin mineral concentrations. *Journal of Animal Science* **83**:852–857 DOI [10.2527/2005.834852x](https://doi.org/10.2527/2005.834852x).
- Mahan DC, Cline TR, Richert B. 1999.** Effects of dietary levels of selenium-enriched yeast and sodium selenite as selenium sources fed to growing-finishing pigs on performance, tissue selenium, serum glutathione peroxidase activity, carcass characteristics, and loin quality. *Journal of Animal Science* **77**:2172–2179 DOI [10.2527/1999.7782172x](https://doi.org/10.2527/1999.7782172x).
- Mahan DC, Moxon AL. 1984.** Effect of inorganic selenium supplementation on selenosis in postweaning swine. *Journal of Animal Science* **58**:1216–1221 DOI [10.2527/jas1984.5851216x](https://doi.org/10.2527/jas1984.5851216x).
- Mateo RD, Spallholz JE, Elder R, Yoon I, Kim SW. 2007.** Efficacy of dietary selenium sources on growth and carcass characteristics of growing-finishing pigs fed diets containing high endogenous selenium. *Journal of Animal Science* **85**:1177–1183 DOI [10.2527/jas.2006-067](https://doi.org/10.2527/jas.2006-067).
- National Research Council. 2005.** *Mineral tolerance of animals*. Second edition. Washington, D.C.: The National Academies Press.
- National Research Council. 2012.** *Nutrition requirements of swine*. Eleventh edition. Washington, D.C.: The National Academy Press.
- Prabhu PAJ. 2015.** Minerals in fish: does the source matter? In: Aquaculture and fisheries. PhD thesis, Wageningen University, Wageningen, 19.
- Reddy KE, Park KR, Lee SD, Yoo J-H, Son AR, Lee H-J. 2017.** Effects of graded concentrations of supplemental lead on lead concentrations in tissues of pigs and prediction equations for estimating dietary lead intake. *PeerJ* **5**:e3936 DOI [10.7717/peerj.3936](https://doi.org/10.7717/peerj.3936).
- Reis AR, El-Ramady H, Santos EF, Gratão PL, Schomburg L. 2017.** Overview of selenium deficiency and toxicity worldwide: affected areas, selenium-related health issues, and case studies. In: Pilon-Smits EAH, Winkel LHE, Lin ZQ, eds. *Selenium in plants: molecular, physiological, ecological and evolutionary aspects*. Cham: Springer International Publishing, 209–230.
- Schwarz C, Ebner KM, Furtner F, Duller S, Wetscherek W, Wernert W, Kandler W, Schedle K. 2017.** Influence of high inorganic selenium and manganese diets for fattening pigs on oxidative stability and pork quality parameters. *Animal* **11**:345–353 DOI [10.1017/S1751731116001518](https://doi.org/10.1017/S1751731116001518).
- Ullah H, Liu G, Yousaf B, Ali MU, Abbas Q, Munir MAM, Mian MM. 2018.** Developmental selenium exposure and health risk in daily foodstuffs: a systematic review and meta-analysis. *Ecotoxicology Environmental Safety* **149**:291–306 DOI [10.1016/j.ecoenv.2017.11.056](https://doi.org/10.1016/j.ecoenv.2017.11.056).
- Winkel LHE, Vriens B, Jones GD, Schneider LS, Pilon-Smits E, Bañuelos GS. 2015.** Selenium cycling across soil-plant-atmosphere interfaces: a critical review. *Nutrients* **7**:4199–4239 DOI [10.3390/nu7064199](https://doi.org/10.3390/nu7064199).
- Zhan XA, Wang M, Zhao RQ, Li WF, Xu ZR. 2007.** Effects of different selenium source on selenium distribution, loin quality and antioxidant status in finishing pigs. *Animal Feed Science and Technology* **132**:202–211 DOI [10.1016/j.anifeedsci.2006.03.020](https://doi.org/10.1016/j.anifeedsci.2006.03.020).

**Zhou JC, Zhao H, Li JG, Xia XJ, Wang KN, Zhang YJ, Liu Y, Zhao Y, Lei XG. 2009.**  
Selenoprotein gene expression in thyroid and pituitary of young pigs is not affected  
by dietary selenium deficiency or excess. *The Journal of Nutrition* **139**:1061–1066  
[DOI 10.3945/jn.109.104901](https://doi.org/10.3945/jn.109.104901).