



# Exploring the mechanistic link between corticosterone and insulin-like growth factor-1 in a wild passerine bird

Zsófia Tóth<sup>1</sup>, Jenny Q. Ouyang<sup>2</sup> and Ádám Z. Lendvai<sup>1,3</sup>

<sup>1</sup> Department of Evolutionary Zoology and Human Biology, University of Debrecen, Debrecen, Hungary

<sup>2</sup> Department of Biology, University of Nevada, Reno, NV, United States of America

<sup>3</sup> Department of Geology, University of Babeş-Bolyai, Cluj-Napoca, Romania

## ABSTRACT

**Background.** Physiological regulators of life history trade-offs need to be responsive to sudden changes of resource availability. When homeostasis is challenged by unpredictable stressors, vertebrates respond through a set of physiological reactions, which can promote organismal survival. Glucocorticoids have been traditionally recognized as one of the main regulators of the physiological stress response, but the role of an evolutionarily more conserved pathway, the hypothalamic-pituitary-somatotropic (HPS) axis producing insulin-like growth factor-1 (IGF-1) has received much less attention. Although IGF-1 is known to affect several life history traits, little is known about its role in the physiological stress response and it has never been studied directly in adult wild animals.

**Methods.** In this study, we combined field observations with a controlled experiment to investigate how circulating levels of IGF-1 change in response to stress and whether this change is due to concomitant change in glucocorticoids in a free-living songbird, the bearded reedling *Panurus biarmicus*. We used a standard capture-restraint protocol in field observation, in which we took first and second (stress induced: 15 minutes later) samples. In a follow-up experiment, we used a minimally invasive oral corticosterone manipulation.

**Results.** We showed that corticosterone levels significantly increased while IGF-1 levels significantly decreased during capture and handling stress. However, change in corticosterone levels were not related to change in IGF-1 levels. We found that experimentally elevated corticosterone levels did not affect IGF-1 levels.

**Discussion.** Our results are the first to highlight that circulating IGF-1 levels are responsive to stress independently from glucocorticoids and suggest that the HPS axis is an autonomous physiological pathway that may play an important role as regulator of life-history decisions.

**Subjects** Ecology, Zoology

**Keywords** Glucocorticoid, IGF-1, Stress response, *Panurus biarmicus*, HPS axis, Life-history trade offs

## INTRODUCTION

Resource allocation trade-offs are central to the evolution of life-histories. Physiological mediators of such trade-offs need to monitor resource availability and transmit a signal

Submitted 6 June 2018

Accepted 15 October 2018

Published 14 December 2018

Corresponding authors

Zsófia Tóth, [hylidae91@gmail.com](mailto:hylidae91@gmail.com)

Ádám Z. Lendvai,

[az.lendvai@gmail.com](mailto:az.lendvai@gmail.com)

Academic editor

George Bentley

Additional Information and  
Declarations can be found on  
page 14

DOI 10.7717/peerj.5936

 Copyright

2018 Tóth et al.

Distributed under

Creative Commons CC-BY 4.0

**OPEN ACCESS**

to relevant parts of the organism to adjust energy expenditure in face of environmental variation. Possible candidates of such key life-history regulatory mechanisms must therefore integrate information from both the external and internal environment and be responsive to changes in resource availability (*Harshman & Zera, 2007*). One way to investigate whether a given physiological mechanism could have such life-history regulatory functions is to assess whether it fulfils these requirements of information processing, integration and responsiveness. A useful framework to analyse these questions is when individuals are exposed to stressors. This approach is biologically and ecologically relevant because in order to successfully reproduce and survive, all organisms must be able to cope with environmental challenges. An organism can display a stress response when challenged by unpredictable, noxious changes in the environment, such as the attack of a predator, an infection, inclement weather, or food shortage (*Wingfield et al., 1998; Kitaysky, Wingfield & Piatt, 1999; Romero, Reed & Wingfield, 2000; Hawlena & Schmitz, 2010*). Mounting an appropriate stress response requires a dramatic reorganization of resource allocation. Understanding this response provides great insight into life-history decisions and resource allocation mechanisms.

An important regulator of the stress response is the endocrine system, which plays a central role in regulating the adjustment of morphology, physiology and behaviour to deal with current conditions (*Ricklefs & Wikelski, 2002; Flatt & Heyland, 2011; Wingfield & Boonstra, 2013*). Hormones function as integrators that process information from the environment and orchestrate multiple processes simultaneously to maximise survival and reproduction under the given circumstances (*Wingfield et al., 1998; Martin et al., 2011*). Although some pathways have been clearly represented in the study of the stress response (e.g., HPA axis—hypothalamic-pituitary-adrenal axis), other, more evolutionarily conserved pathways are understudied even though they can play a key role in the evolution of the endocrine system.

The HPA cascade begins at the brain that perceives the stressor and launches a series of downstream hormonal changes. At the endpoint of this axis, the adrenal cortex secrete increased amounts of glucocorticoids into the bloodstream, which stimulate energy production via gluconeogenesis. However, this pathway appeared during the evolution of vertebrates, by which period robust regulation of homeostasis and resource allocation had already evolved (*Stoks, 2001; Baker Michael, 2003; Bijlsma & Loeschcke, 2005; Pauwels, Stoks & De Meester, 2005; Hawlena & Schmitz, 2010*). A more conserved endocrine pathway is the insulin/insulin-like signalling pathway (the IIS pathway), which is present in all animals and regulates resource allocation and stress resistance (*Broughton et al., 2005; Harshman & Zera, 2007; Dantzer & Swanson, 2012*). In vertebrates, the IIS pathway is integrated into the Hypothalamic-Pituitary-Somatotropic (HPS) axis. As part of this hormonal cascade, growth hormone (GH) stimulates the secretion of insulin-like growth factor-1 (IGF-1), primarily from the liver (*Roith, Scavo & Butler, 2001*).

IGF-1 is an evolutionarily highly conserved nutrient sensing hormone that has pleiotropic effects influencing key life-history traits and major life-history trade-offs among growth, reproduction and lifespan (*Dantzer & Swanson, 2012*). IGF-1 is negatively related to lifespan (*Holzenberger et al., 2003; Lewin et al., 2016*), but has a positive effect on

growth, sexual maturation and reproduction (Crain et al., 1995; Yakar et al., 1999; Pine et al., 2006; Flatt et al., 2008; Sparkman, Vleck & Bronikowski, 2009; Lewin et al., 2016; Lodjak, Mägi & Tilgar, 2014; Swanson & Dantzer, 2014). IGF-1 therefore may mediate the trade-off between longevity and reproduction because reduced IGF-1 signalling has been shown to increase lifespan and the expression of genes involved in stress resistance while increased IGF-1 signalling is necessary for reproduction (Holzenberger et al., 2003; Harshman & Zera, 2007; Dantzer & Swanson, 2012; Lewin et al., 2016).

Although the role of IGF-1 has been established in resource allocation, its role in coping with stressful situations and the crosstalk between the HPA and the HPS axes remain poorly understood. It has been shown that IGF-1 levels change under nutritional and handling stress. For instance, in response to 5 min of restraint, circulating IGF-1 levels decreased by 21% within 60 min in Yorkshire pigs *Sus crofa domesticus* and remained suppressed for up to 150 min (Farmer et al., 1991). A more recent study in pigs also found that restraint stress caused a drop in circulating IGF-1 levels and affected other components of the IIS pathway, suggesting that the IGF-system represents a physiologically relevant biomarker of stress response (Wirthgen et al., 2017). Similarly, IGF-1 levels significantly decreased due to short-term (15 min) and long term (24 h) confinement stress in sunshine bass (hybrid of *Morone saxatilis* and *M. chrysops*), in Atlantic salmon *Salmo salar* and in rainbow trout *Oncorhynchus mykiss* (Wilkinson et al., 2006; Davis & Peterson, 2006).

Whether the decrease in IGF-1 levels is a direct consequence of exposure to stressors on the HPS axis or is due to the stress-induced activity of the HPA axis is controversial (Unterman et al., 1993; Dell et al., 1999; Davis & Peterson, 2006). On the one hand, it is well established that glucocorticoids initiate and orchestrate the emergency life-history stage within minutes to hours from the appearance of the stressor in temperate bird species (Breuner, Greenberg & Wingfield, 1998; Wingfield et al., 1998; Romero & Remage-Healey, 2000; Lohmus, Sundström & Moore, 2006), and we also know that glucocorticoids are interconnected with the somatotrophic axis (Dell et al., 1999). For instance, glucocorticoids play a role in the embryonic development of the HPS axis by affecting GH gene expression, IGF-1 transcription (Dell et al., 1999; Bossis & Porter, 2003; Reindl & Sheridan, 2012). Experimental studies have also shown that exogenous glucocorticoids cause lower IGF-1 levels in rats (Gayan-Ramirez et al., 1999), chicken (Leili & Scanes, 1998) and fish (Kajimura et al., 2003; Peterson & Small, 2005). On the other hand, these effects seem to operate only at high glucocorticoid concentrations and/or at a prolonged exposure. For example, although Bossis & Porter (2003) found that glucocorticoids mediate the embryonic development of the HPS axis, hormone treatment for at least 8 h was necessary to detect a significant increase in GH gene expression. Similarly, while a pharmacological increase in cortisol resulted in a decrease of IGF-1 levels in tilapia *Oreochromis mossambicus* 24–48 h post injection (Kajimura et al., 2003), a more moderate, short-term elevation in cortisol did not affect IGF-1 levels in sunshine bass (Davis & Peterson, 2006).

While some of these laboratory, aquacultural and agricultural studies suggest a relationship between stress, glucocorticoids and IGF-1, our knowledge remains very limited about how stressors affect IGF-1 levels in free-living organisms. Importantly, we do not know how the HPA and HPS axes are linked mechanistically. Birds are particularly

interesting model systems for studying the IIS, because their metabolism is faster yet their lifespan is longer than similar-sized mammals (Costantini, 2008), and differences in how the IIS is regulated compared to the most studied mammalian models might provide a deeper understanding of the evolution of this pathway (Holmes & Ottinger, 2003; Dantzer & Swanson, 2012). Although the IGF-system has been extensively studied in poultry (reviewed in McMurtry, 1998) almost nothing is known about its regulation in free-living birds (notable exceptions are Lodjak, Mägi & Tilgar, 2014; Lodjak, Tilgar & Mägi, 2016; Lodjak et al., 2017)

While interest in the ecological and evolutionary relevance of IGF-1 has recently increased (Sparkman, Vleck & Bronikowski, 2009; Sparkman et al., 2010; Palacios, Sparkman & Bronikowski, 2012; Lodjak, Mägi & Tilgar, 2014; Reding et al., 2016; Lodjak, Tilgar & Mägi, 2016; Lodjak et al., 2017), to the best of our knowledge no study has directly tested the effects of stressors on the activity of the HPS axis or investigated the mechanistic link between the HPA and HPS axes in any free-living organisms. Therefore, we aimed at answering whether: (1) acute stress affects plasma levels of IGF-1 and whether (2) glucocorticoids directly affect circulating IGF-1 levels. First, we hypothesized that acute stress will affect the activity of the HPS axis; therefore, we predicted that circulating IGF-1 levels will decrease in response to short-term acute stress. Second, to investigate the crosstalk between the HPA and the HPS axes, we experimentally increased corticosterone levels using a minimally invasive technique. We predicted that if the HPS axis receives direct input from glucocorticoids, then increased glucocorticoid levels will down-regulate IGF-1 levels. Alternatively, if the HPS axis responds to external stressors directly, then we predicted that short-term elevation of circulating glucocorticoids will not affect IGF-1 levels. To answer these questions, we studied free-living bearded reedlings *Panurus biarmicus*, a small (~14 g), sexually dimorphic, resident songbird common throughout wetlands of Eurasia.

## MATERIAL AND METHODS

### Study animal and field study

We captured 17 wintering free-living bearded reedlings *Panurus biarmicus* (Linnaeus, 1758) in Hungary at Virágoskúti-halastó (N47.6518, E21.3589) between September 2015 and January 2016. We caught the birds with continuously observed mist-nets and subjected them to a standardized capture-handling-restraint protocol (Wingfield, 1994). The first blood sample (50~100  $\mu$ l) was taken as soon as possible after the bird hit the net (mean handling time: 4:58 min; range: 2–10 min). The initial handling times contained one high outlier, inclusion or removal of this point did not affect our results qualitatively. The handling time was not detectably related to either corticosterone levels ( $t = 1.276$ ,  $p = 0.223$ ) or IGF-1 levels ( $t = -0.785$ ,  $p = 0.445$ ). The bird was then placed into an opaque cloth bag and the collection of a subsequent blood sample (50–100  $\mu$ l) was started 15 min after the initial capture (completed mean time: 17:51 min; range: 15–19 min). We chose to take the second blood sample after 15 min because we were interested in the short-term acute stress-response, and we wanted to minimize the possible downstream effects of corticosterone on IGF-1 levels. The total volume of the two blood samples was

under 140  $\mu$ l that met the recommendation of [Owen \(2011\)](#). The study was approved by the Institutional Animal Care and Use Committee at the University of Debrecen (DEMAB/19-6/2015) and the regional government agency (HBB/17/00870-3/2015).

## **Experimental study**

### ***Housing protocol***

Twenty-one wintering bearded reedlings were captured with mist-nets in Hungary at Hortobágy-Halastó (N47.6211, E21.0757) between 18 of October and 16 of November in 2016. After capture, they were housed in an outdoor aviary at the Botanical Garden of the University of Debrecen where they were kept for 4 months and acclimated to captivity. Four weeks prior to the experiment, the birds were transferred to individual cages measuring 25  $\times$  25  $\times$  25 cm. All sides of the cages (except for the front) were made of a non-transparent board (OSB); therefore, birds were kept in visual but not acoustic isolation. Individual cages were separated by a non-transparent removable divider. Because bearded reedlings are very social, live in flocks and maintain strong pair bonds throughout the year ([Lovász, Fenyvesi & Gyurácz, 2017](#); [Griggio & Hoi, 2011](#)), long-term individual separation may be perceived as stressful for the birds (as found in other social species, e.g., [Remage-Healey, Adkins-Regan & Romero, 2003](#)). To avoid such additional stressors, before the experiment, birds were kept in pairs, by removing the divider between adjacent cages. Food (a mixture of apple, carrot, quark, cracked dried fish, dried *Gammarus* sp., cracked corn, cracked dry cat food, a commercial soft food mixture for birds and mealworms) and water were available *ad libitum* at all times.

### ***Experimental design***

Corticosterone levels were manipulated orally using a minimally invasive technique described by [Breuner, Greenberg & Wingfield \(1998\)](#), in which corticosterone dissolved in peanut oil was injected into mealworms *Tenebrio molitor*. We used a randomized block design with two doses of exogenous corticosterone and a control manipulation for each block.

The day before experimental day, we removed the food from the birds 1.5 h before sunset and the mobile dividers were inserted into the cages to keep the birds individually. Water was still available *ad libitum*. The next morning (between 8:00–9:00, to avoid daily variation in hormone levels and to standardize the duration of the food removal), the experimenter quietly entered the room (the cages were oriented in a way that the birds could not see the door) and gave one mealworm to the selected bird through a small hole covered by a semi-transparent layer at the back of the cages, so that the bird could not see the experimenter, but we could observe the birds and record the time when they consumed the mealworm (mean time of ingestion was 39 s). We paid particular attention to avoid any visual or acoustic contacts with the birds. Fifteen minutes after the bird consumed the mealworm, the bird was captured through a backdoor at the cages and a single blood sample ( $\sim$ 70  $\mu$ l) was taken as soon as possible. After blood sampling, body mass (to the nearest 0.1 g) of the sampled individuals was also recorded and they were released back to their cage. The exact times when we entered the room, when the mealworm was given, when the bird ate the mealworm, when we caught the birds in the cages and when blood

sampling was completed were all recorded. Handling time was defined as the time between when we opened the door of individual cages and blood sample collected. Total procedural time was defined as the time elapsed between the experimenter entering the room and when blood sampling was completed. Treatments were carried out in blocks, so that 3 birds in a block got the mealworms subsequently (with approximately 1 min staggering). Two blocks were sampled in a morning. After the treatments and sampling, the birds received the usual ad libitum bird chow and fresh water and were left undisturbed for the rest of the day. After we processed all birds, the experiment was repeated one week later, in which each individual received a different treatment than in the first trial, hence all birds received two different treatments. The treatments for the second trial were randomized again within blocks and birds were randomly assigned to the blocks.

### **Mealworm injection**

Mealworms were injected with 20  $\mu$ l of peanut oil (VWR catalogue number: ACRO416855000) containing one of the following concentrations of corticosterone (Sigma catalogue number: NET399250UC): (1) control, no corticosterone; (2) low corticosterone, 0.2 mg/ml (4  $\mu$ g corticosterone + 16  $\mu$ l peanut oil); and (3) 0.5 mg/ml corticosterone concentration (10  $\mu$ g corticosterone + 10  $\mu$ l peanut oil). We made a stock solution for every concentration before we started the experiment. Hereafter we used those stock solution for injection. After thoroughly vortexing the solution, we injected it into mealworms with a 1 ml syringe using a 26G-needle. The chosen corticosterone concentrations were based on previous studies and were calculated as dose per body mass. In red-eyed vireos *Vireo olivaceus* (12–16 g), plasma corticosterone concentrations were elevated with a 0.2 mg/ml concentration corticosterone solution (which corresponds to 0.28  $\mu$ g corticosterone per 1g of body mass) (Löhmus, Sundström & Moore, 2006). In Gambel's white crowned sparrows *Zonotrichia leucophrys gambelii* (25–28 g) a low (0.2 mg/ml–0.15  $\mu$ g/g) and a high (1 mg/ml–0.7  $\mu$ g/g) dose were used, but the low dose did not elevate significantly the plasma corticosterone levels (Breuner, Greenberg & Wingfield, 1998). Nestling Zebra-finches *Taeniopygia guttata* (6–15 g) received a dose of 0.25 mg/ml (1.19  $\mu$ g/g) corticosterone, which resulted in a significant increase in circulating corticosterone levels (Spencer & Verhulst, 2008). Therefore, our manipulations correspond to a dose of 0.25  $\mu$ g/g (low) and 0.62  $\mu$ g/g (high).

### **Blood sampling**

We took blood samples by puncturing the brachial vein with a 26G-needle and collecting blood in heparinized capillary tubes. Samples were kept on ice until transferring them to the lab (1–7 h in the field and max. 1 h in the experiment). Samples were centrifuged at 2,200 g for 10 min and the plasma was removed with a Hamilton syringe. We divided the plasma into 2 aliquots, one for IGF-1 (15  $\mu$ l) and one for corticosterone (15  $\mu$ l). We stored the samples at  $-20^{\circ}\text{C}$  until assayed for corticosterone by radioimmunoassay (RIA) and assayed for IGF-1 by enzyme-linked immunosorbent assay (ELISA).



## Hormone assays

Plasma IGF-1 levels were measured in duplicates by a commercial avian ELISA kit (catalogue: cIGF1ELISA, lot number D00035) from IBT GmbH, Germany. The assay was developed to measure chicken IGF-1, and the amino acid sequence of IGF-1 is identical in chicken and in *P. biarmicus* (ÁZ Lendvai et al., 2018, unpublished data), therefore this assay was expected to perform well in our study species. Serial dilutions of a plasma pool of *P. biarmicus* were parallel of the standard curve. IGF-1 was separated from its binding proteins using an acidic extraction in accordance with the manufacturer's instructions. The final concentrations were determined colourimetrically measuring the absorbance at 450 nm using a Tecan F50 microplate reader. In this assay, we also included chicken *Gallus gallus* plasma samples as a reference, and the obtained IGF-1 concentrations for the chicken samples ( $375.2 \pm 69.4$  SE ng/ml) were an order of magnitude higher than what we had expected based on the literature (30–50 ng/ml, [Ballard et al., 1990](#)). Therefore we measured known concentrations of an international IGF-1 gold standard (WHO/NIBSC 02/254, a product used in different laboratories to calibrate IGF-1 values; [Chanson et al., 2016](#)) and recalibrated all concentrations against this standard. IGF-1 concentrations of the recalibrated chicken reference samples (53.0 ng/ml) were similar to published results and were in agreement with an in-house ELISA developed in our laboratory (ÁZ Lendvai et al., 2018, unpublished data). Therefore, we used the recalibrated values in our analyses. Note however, that the recalibration only affects the absolute values reported, since the concentrations obtained originally would yield the same results, albeit on a different scale. Minimal detection limit was 1.5 ng/ml, and none of the samples fell below this limit. Intra-assay coefficient of variation was 3.9%, inter-assay CV was 5.7%.

Total corticosterone from plasma samples was quantified through direct radioimmunoassay ([Lendvai, Bókonyi & Chastel, 2011](#)). We extracted corticosterone from plasma using diethyl-ether, and extracts were reconstituted in phosphate-buffered saline. We let the samples incubate overnight at 4 °C. We added ~10 K dpm of 3H-Cort (Catalogue number: NET399250UC, lot number: B00025; Perkin Elmer, Waltham, MA, USA), antiserum (Sigma C8784-100ST, lot number: 092M4784) and phosphate-buffered saline. We let the solution sit overnight at 4 °C. Dextran-coated charcoal was added to separate corticosterone bound to antibodies. After centrifugation, the radioactivity of the bound fraction was counted in a liquid scintillation counter (QuantaSmart). All samples were processed in one assay (intra-assay CV: 3.5%).

## Statistical analyses

We analysed our data in R statistical environment, R version 3.3.2 ([R Core Team, 2017](#)). We fitted linear mixed models with function 'lmer' from package lme4 ([Bates et al., 2015](#); version: 1.1-13), and we used stepwise backward model selection to find the best fitting model. Degrees of freedom for linear mixed models were calculated using the Satterthwaite approximation and corresponding p-values were obtained using the package lmerTest ([Kunetsova, Brockhoff & Christensen, 2016](#); version: 2.0-33).

To test the effects of handling stress on IGF-1 and corticosterone, we used generalized linear mixed models with individual as a random intercept. In the initial model, IGF-1 was

the dependent variable, handling (first or second sample) and its two-way interaction with body mass and sex were the explanatory variables, and ring number (as individual identity) was the random factor. The same initial model structure was used to model corticosterone levels. We also analysed the relationship between IGF-1 levels and corticosterone levels in the first sample with a linear model controlling for body mass. Next, we calculated the stress-induced change in both corticosterone and IGF-1 levels (values from the first sample subtracted from the second samples), and analysed whether the magnitude of change between these two hormones were related in a linear model. In this analysis, the dependent variable was the change in IGF-1 levels and the explanatory variable was the change in corticosterone levels, while controlling for body mass and sex.

The experimental data were analysed using linear mixed models. Here, we considered treatment as a three-level factor (high corticosterone, low corticosterone and control). In the initial model, IGF-1 or corticosterone was the dependent variable, treatment and its two-way interaction with sex and body mass were the explanatory variables. The treatment blocks and the weeks of the experiment were also included as fixed effects. In both models we controlled for the handling time and total procedural time.

## RESULTS

### Field samples

The 15 min capture-restraint stress induced a significant increase in corticosterone levels (Fig. 1A, Table 1), and body mass was positively related to corticosterone levels in the first sample (Table 1). Sex and the two-way interaction between stress and sex or body mass did not affect corticosterone levels (Table 1). Handling stress caused a significant decrease in IGF-1 levels (Fig. 1B, Table 2), and IGF-1 levels were higher in males (Table 2). Body mass was not related to IGF-1 levels in the first sample. Body mass and the two way interaction between stress and sex or body mass did not affect the IGF-1 levels (Table 2).


We did not find a significant relationship between IGF-1 and corticosterone levels in the first samples ( $t = 0.786$ ,  $p = 0.450$ ). The changes in IGF-1 and in corticosterone levels induced by the handling stress were also unrelated, even after controlling for body mass or sex ( $p$ 's  $> 0.9$ ). One bird showed an unusual pattern in the corticosterone data, in which corticosterone decreased in response to handling stress. Removing this data point did not alter any of our conclusions.

### Corticosterone manipulation

The body mass of the captive birds did not change during the two experimental weeks, and IGF-1 levels, corticosterone levels, block or sex were all unrelated to body mass ( $p$ 's  $> 0.5$ ). The experimental week or the treatment blocks did not affect corticosterone or IGF-1 levels ( $p$ 's  $> 0.2$ ).

Both low and high dose of treatment significantly increased corticosterone levels compared to the control treatment (Fig. 2A, Table 3). Other effects, including sex, mass and the two-way interaction of treatment with sex and mass, handling time and total procedural time did not influence corticosterone levels (Table 3). Although the manipulation was successful in creating differences in circulating corticosterone levels, the





**Figure 1** Effects of capture handling stress on corticosterone (A) and IGF-1 (B) levels. Capture handling stress causes (A) a significant increase in circulating corticosterone levels ( $n = 16$ ) and (B) a significant decrease in circulating IGF-1 levels ( $n = 17$ ) in free-living bearded reedlings (*Panurus biarmicus*). Squares denote the individual IGF-1 and corticosterone concentrations, and lines between the squares connect the first and second samples from the same individual. The dots and arrows beside the individual points represent the mean and standard error values of corticosterone (A) or IGF-1 (B) levels.

Full-size [DOI: 10.7717/peerj.5936/fig-1](https://doi.org/10.7717/peerj.5936/fig-1)

treatment did not affect IGF-1 levels (Fig. 2B, Table 4). Furthermore, sex, mass and the two-way interaction of treatment with sex and mass, handling time and total procedural time did not affect IGF-1 levels during the experiment (Table 4).

## DISCUSSION

We explored the mechanistic link between the HPA and HPS axes using a free-living songbird, resulting in two key findings. First, we found that in response to a standardized stressor, circulating IGF-1 levels decreased in wild bearded reedlings within 15 min. To our knowledge, this is the first study that reports such an effect for any free-living species. Second, we found that experimentally elevated corticosterone levels did not result in a decrease in IGF-1 levels during the same time frame as we found in the field. This result suggests that the somatotrophic axis may respond to environmental stimuli independently from the HPA axis and may be part of the adaptive physiological coping mechanisms used to maintain or restore homeostasis in stressful situations.

**Table 1 Results from the final linear mixed model after backward elimination for corticosterone in the field study.** Parameter estimates of variables affecting circulating corticosterone levels in free-living bearded reedlings (*Panurus biarmicus*). Results are from the final linear mixed-effects model after stepwise backward elimination of non-significant effects. The initial model structure was: Corticosterone  $\sim$  Handling  $\times$  (Sex + Mass). The terms excluded during model selection with the associated  $p$ -values in the model before elimination are shown below the table.

	Estimate	Std. Error	df	t-value	p-value
Intercept	-22.207	7.701	10.843	-2.884	0.015
Handling (stress)	2.619	0.570	10.664	4.593	<0.001
Mass	1.678	0.500	10.833	3.356	0.006

**Notes.**

Terms excluded: Handling\*Sex  $p = 0.458$ , Sex  $p = 0.401$ , Handling\*Mass  $p = 0.096$ .


**Table 2 Results from the final linear mixed model after the backward elimination for IGF-1 in the field study.** Parameter estimates of variables affecting circulating corticosterone levels in free-living bearded reedlings (*Panurus biarmicus*). Results are from the final linear mixed-effects model after stepwise backward elimination of non-significant effects. The initial model structure was: IGF-1  $\sim$  Handling  $\times$  (Sex + Mass). The terms excluded during model selection with the associated  $p$ -values in the model before elimination are shown below the table.

	Estimate	Std. Error	df	t-value	p-value
Intercept	55.666	3.217	13.809	17.306	<0.001
Handling (stress)	-6.698	1.173	13.235	-5.709	<0.001
Sex (males)	14.181	3.880	13.001	3.655	0.003

**Notes.**

Terms excluded: Handling\*Mass  $p = 0.644$ , Handling\*Sex  $p = 0.507$ , Mass  $p = 0.313$ .

Increases in plasma corticosterone levels in response to stress have been previously reported in many species. Our data obtained in bearded reedlings in the field is consistent with these findings, showing that corticosterone levels increase in response to capture/restraint stress and suggest that the birds perceived the procedure as stressful. Although many studies collect the stress-induced blood sample at 30 min (*Wingfield, Vleck & Moore, 1992; Remage-Healey & Romero, 2001; Buehler et al., 2008*), we chose to reduce the restraint period to study the effects of short-term acute stress on IGF-1 levels, while minimizing the potential confounding downstream effects of corticosterone. The effects of glucocorticoids are mainly genomic, which typically act over an hour, but require at least 15 min (reviewed in *Haller, Mikics & Makara, 2008*). Some short-term direct effects of corticosterone have been also demonstrated, such as effects on RNA synthesis (reviewed in *Haller, Mikics & Makara, 2008*). Therefore, by choosing a shorter restraint period, we aimed at decreasing the time during which the organism may have been exposed to the physiological effects of elevated glucocorticoids (*Buehler et al., 2008*). Although the effects of stress on IGF-1 levels in free-living organisms have not been reported before, this finding is consistent with previous studies in captive animals (*Farmer et al., 1991; Wilkinson et al., 2006; Davis & Peterson, 2006; Wirthgen et al., 2017*). However, our findings do not support the conclusion of earlier avian studies (*Lodjak, Mägi & Tilgar, 2014; Lodjak, Tilgar & Mägi, 2016; Lodjak et al., 2017*), in which handling time was reported to be unrelated to IGF-1 levels, although in nestlings. Such effects may be age-specific. For example,



**Figure 2** Effects of dietary corticosterone treatment on circulating (A) corticosterone ( $n = 42$ ) and (B) IGF-1 levels ( $n = 42$ ). Corticosterone levels were significantly higher in the low corticosterone ( $n = 14$ ) and the high corticosterone ( $n = 14$ ) group compared with the control group ( $n = 14$ ), although the IGF-1 levels did not differ between the treatment groups. We used 21 individuals in the experiment, and every bird received two different treatments.

Full-size DOI: [10.7717/peerj.5936/fig-2](https://doi.org/10.7717/peerj.5936/fig-2)

gilthead seabream *Sparus aurata* differed in expression of IGF-1 and IGF-1R mRNA levels during ontogeny (Perrot *et al.*, 1999). In a study on Korean native ogol chicken, circulating IGF-1 levels gradually increased during the post-hatching period (Yun *et al.*, 2005). The stress-responsiveness of IGF-1 may also vary with development, and may explain why Lodjak, Mägi & Tilgar (2014) and Lodjak, Tilgar & Mägi (2016) did not find that IGF-1 was affected by handling. Furthermore, stress-responsiveness of IGF-1 may also be species-specific, although we also found that capture-restraint stress caused a decrease in IGF-1 levels in adult free-living house sparrows *Passer domesticus* (CI Vágási *et al.*, 2018, unpublished data). Therefore, we suggest that further studies of IGF-1 levels should take into account the potentially confounding effect of additional stressors.

The decrease in IGF-1 under stress is consistent with the allostatic concept of the stress response (McEwen & Wingfield, 2003). Under stressful situations, the organism has to re-establish homeostasis and in order to do so, it has to suppress energetically costly anabolic processes and reinforce those behavioural and physiological processes that promote immediate survival (Wingfield *et al.*, 1998). IGF-1 is a prime regulator of anabolic processes and antagonistic of the catabolic effects of glucocorticoids, therefore the decrease

**Table 3 Results from the final linear mixed model on the effects of experimental treatment on corticosterone.** Parameter estimates of variables affecting circulating corticosterone levels after oral administration of corticosterone in bearded reedlings (*Panurus biarmicus*). Results are from the final linear mixed-effects model after stepwise backward elimination of non-significant effects. The initial model structure was: Corticosterone  $\sim$  Handling time + Procedural time + Treatment  $\times$  (Sex + Mass). The terms excluded during model selection with the associated *p*-values in the model before elimination are shown below the table.

	Estimate	Std. Error	df	t-value	p-value
Intercept	5.660	4.550	28.750	1.244	0.223
Treatment (low)	18.260	6.100	33.790	2.993	0.005
Treatment (high)	43.100	6.100	33.790	7.006	<0.001

**Notes.**

Terms excluded: Treatment (Low)\*Mass *p* = 0.685, Treatment (High)\*Mass *p* = 0.958, Treatment (Low)\*Sex *p* = 0.543, Treatment (High)\*Sex *p* = 0.578, Sex *p* = 0.530, Total procedural time *p* = 0.267, Handling time *p* = 0.338, Mass *p* = 0.249.

**Table 4 Results from the final linear mixed model testing the effects of experimental treatment on IGF-1 levels.** Parameter estimates of variables affecting circulating IGF-1 levels after oral administration of corticosterone in bearded reedlings (*Panurus biarmicus*). Results are from the final linear mixed-effects model after stepwise backward elimination of non-significant effects. Treatment was part of the experimental design, so were kept in the final model, despite being not-significant. The initial model structure was: IGF-1  $\sim$  Handling time + Procedural time + Treatment  $\times$  (Sex + Mass). The terms excluded during model selection with the associated *p*-values in the model before elimination are shown below the table.

	Estimate	Std. Error	df	t-value	p-value
Intercept	52.865	2.900	5.670	18.228	<0.001
Treatment (low)	-0.932	3.862	38.000	-0.241	0.810
Treatment (high)	0.709	3.862	38.000	0.184	0.855

**Notes.**

Terms excluded: Handling time *p* = 0.934, Treatment (Low)\*Mass *p* = 0.911, Treatment (High)\*Mass *p* = 0.729, Mass *p* = 0.620, Total procedural time *p* = 0.502, Treatment (Low)\*Sex *p* = 0.795, Treatment (High)\*Sex *p* = 0.052, Sex *p* = 0.561.

of IGF-1 under acute stress is consistent with its role as one of the physiological mechanisms responsible for maintaining homeostasis. For instance, in a previous study in mice, IGF-1 levels decreased markedly in food restricted animals and the individuals started to lose weight (O'Sullivan *et al.*, 1989). However, experimental IGF-1 administration during starvation reduced the rate of weight loss through the inhibition of the catabolic processes (O'Sullivan *et al.*, 1989). The sudden drop of IGF-1 levels in response to the stressor suggests that this physiological change prepares the animal for the metabolic challenges faced by slowing down anabolic processes and permitting the catabolic effects of glucocorticoids.

In light of these results, we expected that higher glucocorticoid stress responses would be associated with the largest decrease in IGF-1 levels. However, despite the opposite direction of change in the two hormones, neither levels in the first sample nor the magnitude of corticosterone and IGF-1 stress responses were related in the field study at the individual level: birds with the strongest corticosterone increase were not the ones that decreased their IGF-1 levels the most, and vice versa. Furthermore, we did not find any relationship between those parameters if we controlled for body mass. In order to test the relationship between the two hormones more thoroughly, we carried out an experiment in which we manipulated corticosterone in a minimally invasive manner.

Our dietary hormone treatment increased the circulating levels of corticosterone, while control birds did not show a marked increase in corticosterone levels over the course of the study. These results suggest that similarly to previous studies ([Breuner, Greenberg & Wingfield, 1998](#); [Löhmus, Sundström & Moore, 2006](#); [Spencer & Verhulst, 2008](#)), our oral hormone treatment was successful. Corticosterone concentrations after ingesting the mealworm were significantly higher in both the low and the high dose group compared to the controls, albeit with large individual variation. Despite this rapid increase in corticosterone levels in the absence of a physical stressor, IGF-1 levels in our treated birds remained at the level of the controls, with minimal effect sizes; therefore, we can be confident that the treatment did not affect IGF-1 secretion. Our results are similar to those reported in the sunshine bass, in which confinement stress for 15 min resulted in a decrease in IGF-1 levels, but the dietary hormone treatment did not affect plasma IGF-1 concentrations ([Davis & Peterson, 2006](#)).

These results suggest that the HPA and HPS axes are not linked downstream at the endpoint of these hormonal cascades, but the crosstalk between these pathways happens at the hypothalamic-pituitary level. We argue that this is the reason behind the discrepancy between the conclusions of studies using physiological and pharmacological doses (see above). Dexamethasone, a powerful glucocorticoid agonist, which is known for having a strong negative feedback on the HPA axis has also been shown to decrease plasma IGF-1 concentration in chickens ([Leili & Scanes, 1998](#)), which supports the notion that integration of the HPA and HPS axes is operating at higher regulatory levels.

According to a recent study performed on great tit nestlings, the relationship between corticosterone and IGF-1 varies with the nutritional condition of the individuals. [Lodjak, Tilgar & Mägi \(2016\)](#) found that pre-fledging plasma IGF-1 levels of nestlings in good condition (from broods that were experimentally reduced) were positively related to feather corticosterone (an integrated measure of corticosterone over several days during the development), whereas the association between IGF-1 and feather corticosterone levels was negative in nestlings with lower nutritional condition (from enlarged broods). In control broods however, there were no association between the two hormones. In our captive study, the birds had *ad libitum* food availability and they were all in good condition; therefore, one possible explanation for the absence of the relationship between IGF-1 and corticosterone may be that birds in good condition can afford to tolerate higher glucocorticoid concentrations without decreasing IGF-1 levels. In line with this possibility, biomedical studies have shown that IGF-1 diminishes the protein catabolic effects of glucocorticoids only in normally fed but not in starved subjects ([Botfield, Ross & Hinds, 1997](#)).

## CONCLUSIONS

In this study, we showed for the first time that IGF-1 levels decrease in response to stress in a free-living songbird, and that the magnitude of this response is not related to the glucocorticoid stress response. Furthermore, an experimental increase in corticosterone did not affect circulating IGF-1 levels. While glucocorticoids may still have non-linear or

permissive effects on IGF-1 regulation, our results suggest that the HPA and HPS axes are both stress responsive and are not tightly co-regulated at their downstream endpoints. These results raise the possibility that the interaction between IGF-1 and corticosterone may modulate the adaptive response of organisms in stressful situations. Investigations of the relationship between glucocorticoids and fitness remain equivocal, with some studies showing positive, negative and also no relationship (reviewed in [Bonier et al., 2009](#)). The lack of a general glucocorticoid-fitness relationship has been suggested to be a result of the flexibility and environmentally context dependent nature of glucocorticoids ([Bonier & Martin, 2016](#)). Our results showing that IGF-1 levels are responsive to stress independently from glucocorticoids suggest that the HPS axis is an autonomous physiological cascade that may be also involved in the mediation of life history decisions and affect fitness components ([Harshman & Zera, 2007](#); [Dantzer & Swanson, 2012](#); [Lewin et al., 2016](#)). IGF-1 is an evolutionary ancient regulatory hormone with a primary role to provide an organism-wide internal signal about resource availability, and may alter the function of glucocorticoids. If overall resource availability is high (as was the case in our captive study), then IGF-1 levels can act as a physiological buffer against the adverse effects of increased corticosterone levels. However, when the central nervous system receives input from the environment challenging the organism, it may require the reallocation of resources, which is reflected in decreased IGF-1 levels. Therefore, circulating IGF-1 levels can be an important biological indicator of individual internal state and a useful parameter to investigate in the study of life-history decisions.

## ACKNOWLEDGEMENTS

We thank Gyula Ölveczki, Janka Péntzes and Anna Anita Rác for excellent help in the fieldwork and the experiment. Scott MacDougall-Shackleton and an anonymous reviewer provided constructive comments on an earlier version of the manuscript. We are grateful for the Hortobágy National Park to make the work possible.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

The present study was financially supported by the Hungarian Scientific Research Fund (OTKA K113108), the National Development, Research and Innovation Fund (TÉT 15-1-2016-0044), the European Union and the European Social Fund (EFOP-3.6.1-16-2016-00022), and the Hungarian Ministry of National Resources—National Talent Program (NTP-EFÖ-P-15-041). During manuscript preparation, Jenny Q. Ouyang was supported by NSF OIA-1738594 and NIH P20 GM103650 and Ádám Z. Lendvai by the Romanian Ministry of Education (PN-III-P4-ID-PCE-2016-0572). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:



Hungarian Scientific Research Fund: OTKA K113108.  
National Development, Research and Innovation Fund: TÉT 15-1-2016-0044.  
European Union and the European Social Fund: EFOP-3.6.1-16-2016-00022.  
Hungarian Ministry of National Resources—National Talent Program: NTP-EFÖ-P-15-041.  
NSF: OIA-1738594.  
NIH: P20 GM103650.  
Romanian Ministry of Education: PN-III-P4-ID-PCE-2016-0572.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Zsófia Tóth conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Jenny Q. Ouyang conceived and designed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Ádám Z. Lendvai conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

### Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Regional government agency (licence no HBB/17/00870-3/2015).

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Field experiments were approved by regional government agency (licence no HBB/17/00870-3/2015).

### Data Availability

The following information was supplied regarding data availability:

The raw data are provided in the [Supplemental Files](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5936#supplemental-information>.

## REFERENCES

- Baker Michael E. 2003.** Evolution of adrenal and sex steroid action in vertebrates: a ligand-based mechanism for complexity. *BioEssays* 25:396–400  
[DOI 10.1002/bies.10252](https://doi.org/10.1002/bies.10252).

- Ballard FJ, Johnson RJ, Owens PC, Francis GL, Upton FM, McMurtry JP, Wallace JC. 1990.** Chicken insulin-like growth factor-I: amino acid sequence, radioimmunoassay, and plasma levels between strains and during growth. *General and Comparative Endocrinology* **79**:459–468 DOI [10.1016/0016-6480\(90\)90076-X](https://doi.org/10.1016/0016-6480(90)90076-X).
- Bates D, Mächler M, Bolker B, Walker S. 2015.** Fitting linear mixed-effects models using lme4. *Journal of Statistical Softwares* **67**:1–48 DOI [10.18637/jss.v067.i01](https://doi.org/10.18637/jss.v067.i01).
- Bijlsma R, Loeschcke V. 2005.** Environmental stress, adaptation and evolution: an overview. *Journal of Evolutionary Biology* **18**:744–749 DOI [10.1111/j.1420-9101.2005.00962.x](https://doi.org/10.1111/j.1420-9101.2005.00962.x).
- Bonier F, Martin PR. 2016.** How can we estimate natural selection on endocrine traits? Lessons from evolutionary biology. *Proceedings of the Royal Society B: Biological Sciences* **283**:20161887 DOI [10.1098/rspb.2016.1887](https://doi.org/10.1098/rspb.2016.1887).
- Bonier F, Martin PR, Moore IT, Wingfield JC. 2009.** Do baseline glucocorticoids predict fitness? *Trends in Ecology & Evolution* **24**:634–642 DOI [10.1016/j.tree.2009.04.013](https://doi.org/10.1016/j.tree.2009.04.013).
- Bossis I, Porter TE. 2003.** Evaluation of glucocorticoid-induced growth hormone gene expression in chicken embryonic pituitary cells using a novel in situ mRNA quantitation method. *Molecular and Cellular Endocrinology* **201**:13–23 DOI [10.1016/S0303-7207\(03\)00006-6](https://doi.org/10.1016/S0303-7207(03)00006-6).
- Botfield C, Ross RJM, Hinds CJ. 1997.** The role of IGFs in catabolism. *Baillière's Clinical Endocrinology and Metabolism* **11**:679–697 DOI [10.1016/S0950-351X\(97\)80958-5](https://doi.org/10.1016/S0950-351X(97)80958-5).
- Breuner CW, Greenberg AL, Wingfield JC. 1998.** Noninvasive corticosterone treatment rapidly increases activity in Gambel's white-crowned sparrows (*Zonotrichia leucophrys gambelii*). *General and Comparative Endocrinology* **111**:386–394 DOI [10.1006/gcen.1998.7128](https://doi.org/10.1006/gcen.1998.7128).
- Broughton SJ, Piper MDW, Ikeya T, Bass TM, Jacobson J, Driege Y, Martinez P, Hafen E, Withers DJ, Leivers SJ, Partridge L. 2005.** Longer lifespan, altered metabolism, and stress resistance in *Drosophila* from ablation of cells making insulin-like ligands. *Proceedings of the National Academy of Sciences of the United States of America* **102**:3105–3110 DOI [10.1073/pnas.0405775102](https://doi.org/10.1073/pnas.0405775102).
- Buehler DM, Bhola N, Barjaktarov D, Goymann W, Schwabl I, Tieleman BI, Piersma T. 2008.** Constitutive immune function responds more slowly to handling stress than corticosterone in a shorebird. *Physiological and Biochemical Zoology* **81**:673–681 DOI [10.1086/588591](https://doi.org/10.1086/588591).
- Chanson P, Arnoux A, Mayromati M, Brailly-Tabard S, Massart C, Young J, Piketty ML, Sourberbielle JC. 2016.** Reference values for IGF-1 serum concentrations: comparison of six immunoassays. *The Journal of Clinical Endocrinology & Metabolism* **101**:3450–3458 DOI [10.1210/jc.2016-1257](https://doi.org/10.1210/jc.2016-1257).
- Costantini D. 2008.** Oxidative stress in ecology and evolution: lessons from avian studies. *Ecology Letters* **11**:1238–1251 DOI [10.1111/j.1461-0248.2008.01246.x](https://doi.org/10.1111/j.1461-0248.2008.01246.x).
- Crain DA, Bolten AB, Bjørndal KA, Guillerte Jr LJ, Gross TS. 1995.** Size-dependent, sex-dependent, and seasonal changes in insulin-like growth factor I in the loggerhead

- sea turtle (*Caretta caretta*). *General and Comparative Endocrinology* **98**:219–226  
DOI [10.1006/gcen.1995.1063](https://doi.org/10.1006/gcen.1995.1063).
- Dantzer B, Swanson EM. 2012.** Mediation of vertebrate life histories via insulin-like growth factor-1. *Biological Reviews* **87**:414–429  
DOI [10.1111/j.1469-185X.2011.00204.x](https://doi.org/10.1111/j.1469-185X.2011.00204.x).
- Davis KB, Peterson BC. 2006.** The effect of temperature, stress, and cortisol on plasma IGF-I and IGFBPs in sunshine bass. *General and Comparative Endocrinology* **149**:219–225 DOI [10.1016/j.ygcen.2006.05.009](https://doi.org/10.1016/j.ygcen.2006.05.009).
- Dell G, Ward A, Shokrai A, Madej A, Engström W. 1999.** Regulation of the IGF system by glucocorticoids. *Zoological Science* **16**:377–385  
DOI [10.2108/zsj.16.377](https://doi.org/10.2108/zsj.16.377).
- Farmer C, Dubreuil P, Couture Y, Brazeau P, Petitclerc D. 1991.** Hormonal changes following an acute stress in control and somatostatin-immunized pigs. *Domestic Animal Endocrinology* **8**:527–536 DOI [10.1016/0739-7240\(91\)90022-C](https://doi.org/10.1016/0739-7240(91)90022-C).
- Flatt T, Heyland A. 2011.** *Mechanisms of life history evolution: the genetics and physiology of life history traits and trade-Offs*. Oxford: OUP.
- Flatt T, Min K-J, D’Alterio C, Villa-Cuesta E, Cumbers J, Lehmann R, Jones DL, Tatar M. 2008.** Drosophila germ-line modulation of insulin signaling and lifespan. *Proceedings of the National Academy of Sciences of the United States of America* **105**:6368–6373 DOI [10.1073/pnas.0709128105](https://doi.org/10.1073/pnas.0709128105).
- Gayán-Ramírez G, Vanderhoydonc F, Verhoeven G, Decramer M. 1999.** Acute treatment with corticosteroids decreases IGF-1 and IGF-2 expression in the rat diaphragm and gastrocnemius. *American Journal of Respiratory and Critical Care Medicine* **159**:283–289 DOI [10.1164/ajrccm.159.1.9803021](https://doi.org/10.1164/ajrccm.159.1.9803021).
- Griggio M, Hoi H. 2011.** An experiment on the function of the long-term pair bond period in the socially monogamous bearded reedling. *Animal Behaviour* **82**:1329–1325  
DOI [10.1016/j.anbehav.2011.09.016](https://doi.org/10.1016/j.anbehav.2011.09.016).
- Haller J, Mikics É, Makara GB. 2008.** The effects of non-genomic glucocorticoid mechanisms on bodily functions and the central neural system. A critical evaluation of findings. *Frontiers in Neuroendocrinology* **29**:273–291  
DOI [10.1016/j.yfrne.2007.10.004](https://doi.org/10.1016/j.yfrne.2007.10.004).
- Harshman LG, Zera AJ. 2007.** The cost of reproduction: the devil in the details. *Trends in Ecology & Evolution* **22**:80–86 DOI [10.1016/j.tree.2006.10.008](https://doi.org/10.1016/j.tree.2006.10.008).
- Hawlina D, Schmitz OJ. 2010.** Physiological stress as a fundamental mechanism linking predation to ecosystem functioning. *The American Naturalist* **176**:537–556  
DOI [10.1086/656495](https://doi.org/10.1086/656495).
- Holmes DJ, Ottinger MA. 2003.** Birds as long-lived animal models for the study of aging. *Experimental Gerontology* **38**:1365–1375  
DOI [10.1016/j.exger.2003.10.018](https://doi.org/10.1016/j.exger.2003.10.018).
- Holzenberger M, Dupont J, Ducos B, Leneuve P, Géloën A, Even PC, Cervera P, Le Bouc Y. 2003.** IGF-1 receptor regulates lifespan and resistance to oxidative stress in mice. *Nature* **421**:182–187 DOI [10.1038/nature01298](https://doi.org/10.1038/nature01298).

- Kajimura S, Hirano T, Visitacion N, Moriyama S, Aida K, Grau EG. 2003.** Dual mode of cortisol action on GH/IGF-I/IGF binding proteins in the tilapia, *Oreochromis mossambicus*. *Journal of Endocrinology* **178**:91–99  
DOI [10.1677/joe.0.1780091](https://doi.org/10.1677/joe.0.1780091).
- Kitaysky AS, Wingfield JC, Piatt JF. 1999.** Dynamics of food availability, body condition and physiological stress response in breeding Black-legged Kittiwakes. *Functional Ecology* **13**:577–584 DOI [10.1046/j.1365-2435.1999.00352.x](https://doi.org/10.1046/j.1365-2435.1999.00352.x).
- Kunetsova A, Brockhoff PB, Christensen RHB. 2016.** lmerTest: Tests in linear mixed effects models. R package version 2.0-33. Available at <https://CRAN.R-project.org/package=lmerTest>.
- Leili S, Scanes CG. 1998.** The effects of glucocorticoids (Dexamethasone) on insulin-like growth factor-I, IGF-binding proteins, and growth in chickens. *Proceedings of the Society for Experimental Biology and Medicine* **218**:329–333  
DOI [10.3181/00379727-218-44300](https://doi.org/10.3181/00379727-218-44300).
- Lendvai ÁZ, Bókony V, Chastel O. 2011.** Coping with novelty and stress in free-living house sparrows. *Journal of Experimental Biology* **214**:821–828  
DOI [10.1242/jeb.047712](https://doi.org/10.1242/jeb.047712).
- Lewin N, Swanson EM, Williams BL, Holekamp KE. 2016.** Juvenile concentrations of IGF-1 predict life-history trade-offs in a wild mammal. *Functional Ecology* **31**:894–902 DOI [10.1111/1365-2435.12808](https://doi.org/10.1111/1365-2435.12808).
- Lodjak J, Mägi M, Sild E, Mänd R. 2017.** Causal link between insulin-like growth factor 1 and growth in nestlings of a wild passerine bird. *Functional Ecology* **31**:184–191  
DOI [10.1111/1365-2435.12679](https://doi.org/10.1111/1365-2435.12679).
- Lodjak J, Mägi M, Tilgar V. 2014.** Insulin-like growth factor 1 and growth rate in nestlings of a wild passerine bird. *Functional Ecology* **28**:159–166  
DOI [10.1111/1365-2435.12164](https://doi.org/10.1111/1365-2435.12164).
- Lodjak J, Tilgar V, Mägi M. 2016.** Does the interaction between glucocorticoids and insulin-like growth factor 1 predict nestling fitness in a wild passerine? *General and Comparative Endocrinology* **225**:149–154 DOI [10.1016/j.yggen.2015.10.016](https://doi.org/10.1016/j.yggen.2015.10.016).
- Löhmus M, Sundström LF, Moore FR. 2006.** Non-invasive corticosterone treatment changes foraging intensity in red-eyed vireos *Vireo olivaceus*. *Journal of Avian Biology* **37**:523–526 DOI [10.1111/j.0908-8857.2006.03733.x](https://doi.org/10.1111/j.0908-8857.2006.03733.x).
- Lovász L, Fenyvesi L, Gyurác J. 2017.** Contribution to the study of pair formation strategy of the Bearded Reedling (*Panurus biarmicus*). *North-Western Journal of Zoology* **13**:297–302 Article number: e161609.
- Martin LB, Liebl AL, Trotter JH, Richards CL, McCoy K, McCoy MW. 2011.** Integrator networks: illuminating the black box linking genotype and phenotype. *Integrative and Comparative Biology* **51**:514–527 DOI [10.1093/icb/ucr049](https://doi.org/10.1093/icb/ucr049).
- McEwen BS, Wingfield JC. 2003.** The concept of allostasis in biology and biomedicine. *Hormones and Behavior* **43**:2–15 DOI [10.1016/S0018-506X\(02\)00024-7](https://doi.org/10.1016/S0018-506X(02)00024-7).
- McMurtry JP. 1998.** Nutritional and developmental roles of insulin-like growth factors in poultry. *The Journal of Nutrition* **128**:302S–305S DOI [10.1093/jn/128.2.302S](https://doi.org/10.1093/jn/128.2.302S).

- O'Sullivan U, Gluckman PD, Breier BH, Woodall S, Siddiqui RA, McCUTCHEON SN. 1989. Insulin-like growth factor-1 (igf-1) in mice reduces weight loss during starvation. *Endocrinology* 125:2793–2794 DOI 10.1210/endo-125-5-2793.
- Owen JC. 2011. Collecting, processing, and storing avian blood: a review. *Journal of Field Ornithology* 82:339–354 DOI 10.1111/j.1557-9263.2011.00338.x.
- Palacios MG, Sparkman AM, Bronikowski AM. 2012. Corticosterone and pace of life in two life-history ecotypes of the garter snake *Thamnophis elegans*. *General and Comparative Endocrinology* 175:443–448 DOI 10.1016/j.ygcn.2011.11.042.
- Pauwels K, Stoks R, De Meester L. 2005. Coping with predator stress: interclonal differences in induction of heat-shock proteins in the water flea *Daphnia magna*. *Journal of Evolutionary Biology* 18:867–872 DOI 10.1111/j.1420-9101.2005.00890.x.
- Perrot V, Moiseeva EB, Gozes Y, Chan SJ, Ingleton P, Funkenstein B. 1999. Ontogeny of the insulin-like growth factor system (IGF-I, IGF-II, and IGF-1R) in gilthead seabream (*Sparus aurata*): expression and cellular localization. *General and Comparative Endocrinology* 116:445–460 DOI 10.1006/gc.1999.7337.
- Peterson BC, Small BC. 2005. Effects of exogenous cortisol on the GH/IGF-I/IGFBP network in channel catfish. *Domestic Animal Endocrinology* 28:391–404 DOI 10.1016/j.domaniend.2005.01.003.
- Pine MD, Hiney JK, Dearth RK, Bratton GR, Dees WL. 2006. IGF-1 administration to prepubertal female rats can overcome delayed puberty caused by maternal Pb exposure. *Reproductive Toxicology* 21:104–109 DOI 10.1016/j.reprotox.2005.07.003.
- R Core Team. 2017. R: The R Project for Statistical Computing. R foundation for Statistical Computing, Vienna, Austria. Available at <https://www.r-project.org>.
- Reding DM, Addis EA, Palacios MG, Schwartz TS, Bronikowski AM. 2016. Insulin-like signaling (IIS) responses to temperature, genetic background, and growth variation in garter snakes with divergent life histories. *General and Comparative Endocrinology* 233:88–99 DOI 10.1016/j.ygcn.2016.05.018.
- Reindl KM, Sheridan MA. 2012. Peripheral regulation of the growth hormone-insulin-like growth factor system in fish and other vertebrates. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 163:231–245 DOI 10.1016/j.cbpa.2012.08.003.
- Remage-Healey L, Adkins-Regan E, Romero LM. 2003. Behavioral and adrenocortical responses to mate separation and reunion in the zebra finch. *Hormones and Behavior* 43:108–114 DOI 10.1016/S0018-506X(02)00012-0.
- Remage-Healey L, Romero LM. 2001. Corticosterone and insulin interact to regulate glucose and triglyceride levels during stress in a bird. *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology* 281:R994–R1003 DOI 10.1152/ajpregu.2001.281.3.R994.
- Ricklefs RE, Wikelski M. 2002. The physiology/life-history nexus. *Trends in Ecology & Evolution* 17:462–468 DOI 10.1016/S0169-5347(02)02578-8.

- Roith DL, Scavo L, Butler A. 2001.** What is the role of circulating IGF-I? *Trends in Endocrinology & Metabolism* **12**:48–52 DOI [10.1016/S1043-2760\(00\)00349-0](https://doi.org/10.1016/S1043-2760(00)00349-0).
- Romero LM, Reed JM, Wingfield JC. 2000.** Effects of weather on corticosterone responses in wild free-living passerine birds. *General and Comparative Endocrinology* **118**:113–122 DOI [10.1006/gcen.1999.7446](https://doi.org/10.1006/gcen.1999.7446).
- Romero LM, Ramage-Healey L. 2000.** Daily and seasonal variation in response to stress in captive starlings (*Sturnus vulgaris*): corticosterone. *General and Comparative Endocrinology* **119**:52–59 DOI [10.1006/gcen.2000.7491](https://doi.org/10.1006/gcen.2000.7491).
- Sparkman AM, Byars D, Ford NB, Bronikowski AM. 2010.** The role of insulin-like growth factor-1 (IGF-1) in growth and reproduction in female brown house snakes (*Lamprophis fuliginosus*). *General and Comparative Endocrinology* **168**:408–414 DOI [10.1016/j.ygcen.2010.05.006](https://doi.org/10.1016/j.ygcen.2010.05.006).
- Sparkman AM, Vleck CM, Bronikowski AM. 2009.** Evolutionary ecology of endocrine-mediated life-history variation in the garter snake *Thamnophis elegans*. *Ecology* **90**:720–728 DOI [10.1890/08-0850.1](https://doi.org/10.1890/08-0850.1).
- Spencer KA, Verhulst S. 2008.** Post-natal exposure to corticosterone affects standard metabolic rate in the zebra finch (*Taeniopygia guttata*). *General and Comparative Endocrinology* **159**:250–256 DOI [10.1016/j.ygcen.2008.09.007](https://doi.org/10.1016/j.ygcen.2008.09.007).
- Stoks R. 2001.** Food stress and predator-induced stress shape developmental performance in a damselfly. *Oecologia* **127**:222–229 DOI [10.1007/s004420000595](https://doi.org/10.1007/s004420000595).
- Swanson EM, Dantzer B. 2014.** Insulin-like growth factor-1 is associated with life-history variation across Mammalia. *Proceedings of the Royal Society of London B* **281**:2013–2458 DOI [10.1098/rspb.2013.2458](https://doi.org/10.1098/rspb.2013.2458).
- Unterman TG, Jentel JJ, Oehler DT, Lacson RG, Hofert JF. 1993.** Effects of glucocorticoids on circulating levels and hepatic expression of insulin-like growth factor (IGF)-binding proteins and IGF-I in the adrenalectomized streptozotocin-diabetic rat. *Endocrinology* **133**:2531–2539 DOI [10.1210/endo.133.6.7694841](https://doi.org/10.1210/endo.133.6.7694841).
- Wilkinson RJ, Porter M, Woolcott H, Longland R, Carragher JF. 2006.** Effects of aquaculture related stressors and nutritional restriction on circulating growth factors (GH, IGF-I and IGF-II) in Atlantic salmon and rainbow trout. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **145**:214–224 DOI [10.1016/j.cbpa.2006.06.010](https://doi.org/10.1016/j.cbpa.2006.06.010).
- Wingfield JC. 1994.** Modulation of the adrenocortical response to stress in birds. *Perspectives in Comparative Endocrinology* 520–528.
- Wingfield JC, Boonstra R. 2013.** Ecological processes and the ecology of stress: the impacts of abiotic environmental factors. *Functional Ecology* **27**:37–44 DOI [10.1111/1365-2435.12039](https://doi.org/10.1111/1365-2435.12039).
- Wingfield JC, Maney DL, Breuner CW, Jacobs JD, Lynn S, Ramenofsky M, Richardson RD. 1998.** Ecological bases of hormone—behavior interactions: the emergency life history stage. *American Zoologist* **38**:191–206 DOI [10.1093/icb/38.1.191](https://doi.org/10.1093/icb/38.1.191).
- Wingfield JC, Vleck CM, Moore MC. 1992.** Seasonal changes of the adrenocortical response to stress in birds of the Sonoran desert. *Journal of Experimental Zoology* **264**:419–428 DOI [10.1002/jez.1402640407](https://doi.org/10.1002/jez.1402640407).



- Wirthgen E, Kunze M, Goumon S, Walz C, Höflich C, Spitschak M, Brenmoehl J, Kanitz E, Tuchscherer M, Otten W, Gimsa U, Schön P, Manteuffel C, Tuchscherer A, Pfuhl R, Metges CC, Stabenow B, Erdmann S, Schluricke K, Faucitano L, Hoeflich A. 2017.** Interference of stress with the somatotropic axis in pigs—lights on new biomarkers. *Scientific Reports* 7:12055  
[DOI 10.1038/s41598-017-11521-5](https://doi.org/10.1038/s41598-017-11521-5).
- Yakar S, Liu J-L, Stannard B, Butler A, Accili D, Sauer B, LeRoith D. 1999.** Normal growth and development in the absence of hepatic insulin-like growth factor I. *Proceedings of the National Academy of Sciences of the United States of America* 96:7324–7329 [DOI 10.1073/pnas.96.13.7324](https://doi.org/10.1073/pnas.96.13.7324).
- Yun JS, Seo DS, Kim WK, Ko Y. 2005.** Expression and relationship of the insulin-like growth factor system with posthatch growth in the Korean Native Ogol chicken. *Poultry Science* 84:83–90 [DOI 10.1093/ps/84.1.83](https://doi.org/10.1093/ps/84.1.83).