



# Serum Antioxidative Enzymes Levels and Oxidative Stress Products in Children and Adolescents with Type I Diabetes Mellitus

Özlem Akgün, Nilgün Selçuk Duru, Murat Eevli

University of Health Sciences, İstanbul Haseki Training and Research Hospital, Clinic of Pediatrics, İstanbul, Turkey

## ABSTRACT

**Aim:** Type I diabetes mellitus (T1DM) is an oxidative stress condition in addition to being a chronic metabolic disease. In this study, our aim is to investigate the activity of antioxidative enzymes and the products of oxidative stress in children and adolescents with T1DM and compare the findings with those in healthy control subjects.

**Materials and Methods:** The study enrolled 41 children and adolescents with T1DM (mean age 11.4±3.3 years; 21 female, 20 male) and 25 healthy subjects (mean age 11.3±3.1 years; 8 female, 17 male) with a similar age and gender distribution. Serum samples were obtained to detect the antioxidative enzymes of paraoxonase (PON), arylesterase (ARE), oxidation degradation products of malondialdehyde (MDA) and also zinc which acts as an antioxidant.

**Results:** We found a significant decrease in PON activity and zinc levels in diabetics compared to the healthy controls ( $p=0.021$ ;  $p<0.001$ , respectively). Zinc was negatively correlated to hemoglobin A1c ( $r=-0.317$ ,  $p=0.049$ ). MDA and ARE did not show a significant difference in the T1DM patients compared to the healthy subjects.

**Conclusion:** Zinc level and PON activity were lower in diabetic children and adolescents. Further studies with larger samples are required to confirm their roles in the following and prognosis of T1DM.

**Keywords:** Antioxidant, oxidative stress, paraoxonase, Type I diabetes mellitus, zinc

## Introduction

Type I diabetes mellitus (T1DM) is the most common metabolic disorder resulting in the destruction of insulin producing pancreatic  $\beta$ -cells by lymphocytic infiltration (1). Genetic predisposition is important in the development of T1DM (2). In addition, some environmental factors such as viral infection, vaccines, low levels of vitamin D and dietary factors during infancy may trigger the development of T1DM in those individuals with genetic susceptibility (2). Some

metabolic and physiologic processes lead to reactive oxygen species (ROS) in the body. ROS are highly reactive molecules derived from the reduction of oxygen and can be harmful to some cell structures such as carbohydrates, nucleic acids, lipids and proteins (3). Their elimination is provided by the antioxidant defence system (4). Oxidative stress is the loss of balance between prooxidant and antioxidant systems (5). Oxidative stress may play a role in the pathogenesis of human diseases. Many studies have investigated the relationship between oxidative stress parameters and various

## Address for Correspondence

Nilgün Selçuk Duru MD, University of Health Sciences, İstanbul Haseki Training and Research Hospital, Clinic of Pediatrics, İstanbul, Turkey  
Phone: +90 532 256 40 10 E-mail: nilgundurdu@yahoo.com ORCID ID: [orcid.org/0000-0001-9105-0529](http://orcid.org/0000-0001-9105-0529)

The study originated from a thesis and has been previously presented at the 4<sup>th</sup> Çocuk Dostları Kongresi, March-2016, İstanbul/Turkey.

Received: 12.01.2018 Accepted: 11.04.2018

©Copyright 2018 by Ege University Faculty of Medicine, Department of Pediatrics and Ege Children's Foundation  
The Journal of Pediatric Research, published by Galenos Publishing House.

diseases such as some cancers, cardiovascular disease, Type II diabetes, cataracts and aging (5-9). Oxidative stress is believed to play a role both in the initial pathology of diabetes and in the development of vascular complications during the course of the disease (10-12). It can cause irreversible damage to the  $\beta$ -cells of the pancreatic islets (13). As a result, diabetic patients are susceptible to developing atherosclerotic cardiovascular diseases at early ages compared to healthy subjects (12). Many antioxidants are produced in the body to prevent the harmful effects of these oxidants (1). This study measured paraoxonase (PON) and arylesterase (ARE) activities as antioxidants, and the level of malondialdehyde (MDA), an end product of lipid peroxidation and the level of zinc which is a trace element acting as antioxidant in children with T1DM and also healthy control subjects. We aimed to compare them between groups and to investigate whether these parameters are associated with metabolic control, gender and diabetes duration.

## Materials and Methods

### Study Groups

The patient group consisted of 41 children and adolescents with T1DM (mean age  $11.4 \pm 3.3$  years, 20 males and 21 females). The patients were diagnosed according to criteria provided by American Diabetes Association (14) and the presence of positive autoimmune antibodies. Exclusion criteria were as follows: other systemic diseases, abnormal renal/hepatic biochemical values or macrovascular complications. The control group consisted of 25 healthy subjects (mean age  $11.3 \pm 3.1$  years; 8 female, 17 male). The study was approved by the University of Health Sciences, İstanbul Haseki Training and Research Hospital Local Ethics Committee (approval number: 48-11/10/2013). Written informed consent was obtained from each child included in the study or their parents before enrolling in the study. All patients had been treated with fast- and longacting insulin therapy from the onset of the disease. Data about the duration of illness and onset of the disease in those children with T1DM were obtained from the parents. Biochemical parameters such as glycosylated hemoglobin A1c (HbA1c) levels were determined in each child. The patients were also divided into subgroups according to their gender, glycaemic control (optimal and suboptimal glycaemic control:  $<9\%$ ; poor glycaemic control:  $\geq 9\%$ ) (12) and duration of disease ( $\leq 1$  year;  $>1$  year).

### Biochemical Analysis

Venous blood samples were collected after overnight fasting and were centrifuged at 2.000 rpm for 20 minutes; serum samples were stored at  $-70$  °C until assayed. PON and ARE activities were measured by an enzyme-linked immunosorbent assay (ELISA) technique using an ELISA kit (Eastbiopharm, USA). The PON activity was determined using

paraoxon as the substrate and measured by increases in the absorbance at 412 nm due to the formation of 4-nitrophenol. ARE activity was determined by measuring the rates of phenyl acetate and paraoxon hydrolysis at 548 nm. MDA was analyzed by a spectrophotometric method. MDA was measured using thiobarbituric acid (TBA) reaction substance production in the following manner. 50  $\mu$ L of sample was added to 750  $\mu$ L of acetic acid (20%), 100  $\mu$ L SDS (8.1%), 750  $\mu$ L TBA and 350  $\mu$ L distilled water. The mixture was heated at 100 °C for 45 min. Then, 0.5 mL of distilled water and 2.5 mL of butanol-pyridine 15:1 were added to the mixture and incubated. Then, the absorbance at 532 nm was determined. Zinc was detected by a spectrophotometric method. HbA1c was analyzed using ion-exchange high performance liquid chromatography (Adams A1c, Arkray).

### Statistical Analysis

SPSS (Statistical Package for the Social Science) 15.0 for Windows was used for the statistical analysis. Qualitative data are presented as counts and percentages. The association between qualitative variables was assessed using a chi-square test. Quantitative data are presented as mean  $\pm$  standard deviation for normally distributed data or otherwise as median and interquartile range. Student's t-test for independent samples was used to check for differences between two independent groups of normally distributed data and also by the Mann-Whitney U test. Spearman correlation coefficient was used to assess the relations between quantitative variables not following a normal distribution.  $P < 0.05$  was considered as statistically significant.

## Results

The study group comprised of 41 children and adolescents that were diagnosed with T1DM. Some markers related to oxidative stress were compared with a known control group of 25 healthy children and adolescents.

The mean age in the group was  $11.4 \pm 3.3$  years (range 3.5-18 years) and was similar to the control group  $11.3 \pm 3.1$  years (range 6-16 years) ( $p=0.912$ ). There were 21 female/20 male (51.2% female/48.8% male) in the T1DM group and 8 female/17 male (32.0% female/68.0% male) in the control group. The gender distribution was similar in both groups ( $p=0.127$ ). The main characteristics of the groups in the study are summarized in Table I. The average time from diagnosis of T1DM to participating in this study was  $34.0 \pm 49.2$  months (range 0-192 months). The average HbA1c value in the patients with T1DM was  $11.0 \pm 2.5\%$  (range 6.8-15.3%). While 28 of the children with T1DM had poor metabolic control, 13 patients had good metabolic control (Table I).

Regarding PON activities and zinc levels, we found statistically significant lower values for the diabetics compared to the controls ( $p < 0.001$ ,  $p=0.021$ , respectively). Also, ARE activity was lower in the diabetics vs. the controls,

**Table I.** Demographic features and biochemical data of the diabetic and control groups

Parameters	Type I diabetics	Control group	p value
Age (years)	11.4±3.3 (3.5-18)	11.3±3.1 (6-16)	0.912
Gender (females, %)	21 (51.2)	8 (32.0)	0.127
Duration of diabetes (month)	34.0±49.2 (0-192)	-	-
Long-acting insulin (U/day)	15.1±8.7 (4-36)	-	-
Fast-acting insulin (U/day)	20.2±11.7 (6-45)	-	-
HbA1c (%)	11.0±2.5 (6.8-15)	-	-
Zinc (mcg/dL)	103.9±51.9 (0.98-278)	163.8±40.6 (88-244.6)	<0.001**
Malondialdehyde (nmol/L)	108.1±166.5 (18-502)	123.7±181.3 (18-502)	0.402
Paraoxonase (ng/mL)	10.60±12.05 (3.6-37.65)	13.82±13.87 (3.3-37.65)	0.021*
Arylesterase (ng/mL)	0.19±0.32 (0.06-1.18)	0.28±0.47 (0.06-1.98)	0.177

\*p<0.05, \*\*p<0.001, Data are mean ± standard deviation, HbA1c: Hemoglobin A1c

**Table II.** Biochemical parameters comparison between diabetic children Type I diabetes mellitus and controls according to gender

		Type I diabetics	p value	Control group	p value
		Mean ± SD		Mean ± SD	
Zinc (mcg/dL)	Female	88.73±32.33	0.095	158.28±41.40	0.651
	Male	119.90±63.62		166.40±41.28	
Malondialdehyde (nmol/L)	Female	101.11±171.32	0.151	159.93±213.49	0.380
	Male	115.43±165.53		106.69±168.63	
Paraoxonase (ng/mL)	Female	10.55±12.28	0.314	19.11±16.27	0.539
	Male	10.65±12.12		11.33±12.34	
Arylesterase (ng/mL)	Female	0.22±0.37	0.449	0.39±0.67	0.793
	Male	0.17±0.25		0.22±0.36	

Data are mean ± SD, SD: Standard deviation

but not statistically significant (p=0.177). MDA levels of the diabetic patients were not statistically significant different from those of the controls (p=0.402) (Table I). We compared the parameters measured in the diabetic patients and the control group according to gender (Table II). We did not observe any significant difference between female and male children for PON, ARE activities, MDA and zinc levels (Table II).

We stratified the patients according to disease duration above and below one year. We did not observe any significant difference between children having a diabetes duration above one year and those below one year for PON, ARE, MDA and zinc (Table III). We compared the parameters measured in the diabetic group according to metabolic control. There was not a statically significant difference between children with poor and good metabolic control for PON, ARE, MDA and zinc (Table IV). A negative correlation was observed between zinc and HbA1c in children with T1DM (Rho=-0.317, p=0.049).

**Table III.** Biochemical parameters in Type I diabetes mellitus patients according to diabetes duration ≤1 year and >1 year

	Duration of diabetes		p value
	≤1 year	>1 year	
	Mean ± SD	Mean ± SD	
Zinc (mcg/dL)	107.45±51.03	101.37±55.03	0.655
Malondialdehyde (nmol/L)	95.64±161.72	124.71±177.30	0.613
Paraoxonase (ng/mL)	9.83±12.23	11.54±12.40	0.714
Arylesterase (ng/mL)	0.17±0.29	0.22±0.35	0.924

Data are mean ± SD, SD: Standard deviation

**Table IV.** Biochemical parameters in Type I diabetes mellitus patients according to hemoglobin A1c levels <9% and ≥9%

	HbA1c <9%	HbA1c ≥9%	
	Mean ± SD	Mean ± SD	p value
Zinc (mcg/dL)	89.24±38.51	111.11±57.20	0.206
Malondialdehyde (nmol/L)	101.20±157.64	116.82±176.84	0.813
Paraoxonase (ng/mL)	9.16±11.11	11.54±12.86	0.309
Arylesterase (ng/mL)	0.08±0.04	0.25±0.37	0.472

Data are mean ± SD, SD: Standard deviation, HbA1c: Hemoglobin A1c

**Table V.** Correlations of zinc, malondialdehyde, paraoxonase and arylesterase with age, duration of diabetes and hemoglobin A1c in children with Type I diabetes mellitus

		Age	Duration of T1DM	HbA1c	Zinc
Zinc (mcg/dL)	rho	-0.193	-0.087	<b>-0.317*</b>	-
	p	<b>0.126</b>	0.591	<b>0.049*</b>	-
Malondialdehyde (nmol/L)	rho	<b>0.299*</b>	0.211	0.051	-0.108
	p	0.016*	0.192	0.758	0.390
Paraoxonase (ng/mL)	rho	0.201	-0.003	0.057	0.231
	p	0.112	0.987	0.732	0.062
Arylesterase (ng/mL)	rho	<b>0.335*</b>	0.264	0.185	0.170
	p	<b>0.007*</b>	0.100	0.259	0.172

\*p<0.05, HbA1c: Hemoglobin A1c, T1DM: Type I diabetes mellitus

Age was positively correlated with ARE and MDA in the diabetic subjects (Rho=0.335, p=0.007; rho=0.299, p=0.016 respectively) (Table V).

## Discussion

Free oxygen radicals interact with cellular components such as proteins, lipids and nucleic acids and start lipid peroxidation (15). In an organism, production of free oxygen radicals and antioxidant defence mechanisms are in balance, and as long as this oxidative balance is kept, oxidative stress cannot damage the organism (1,15). Diabetes mellitus is associated with an endogenous inflammatory process and oxidative stress (1,10,13,16,17). The destruction of insulin producing β cells in T1DM patients elevates the plasma sugar level (18). It is a thought that high glucose levels

trigger oxidative stress and increase ROS in diabetics (18). In addition, the balance between oxidative and antioxidant processes is sensitive to the plasma glucose level (19). Therefore, an increase in ROS is generally accompanied by a decrease in antioxidant defence in T1DM patients (18,19). Prolonged oxidative stress may be associated with chronic complications of diabetes. As a result, diabetic patients are predisposed to atherosclerosis beginning at an early age (20). Most of the studies addressing these mechanisms were performed with diabetic adult patients (17,20-22). Therefore, the present study aimed to evaluate the biochemical markers of oxidative stress in children with T1DM. The end product of lipid peroxidation, MDA is an important marker of oxidative stress (23). High MDA levels were showed in diabetics (21-24). In some studies, statistically significantly higher levels were reported for patients with poor metabolic control than in patients with suboptimal and optimal metabolic control (17,21,23,25). These studies suggested that high glucose levels lead to lipid peroxidation and consequently to increased MDA (25). Erciyas et al. (23) proposed that the elevated MDA levels in children with T1DM with poor metabolic control may lead to vascular complications. Also, they recommended that MDA should be added to the routine laboratory evaluations in the follow-up of these patients (23). In contrast to these studies, Reis et al. (20) reported low MDA levels in patients with T1DM. In our study, MDA was similar between children with T1DM and the healthy controls. Also, MDA was not different between the groups in terms of disease duration, glycaemic control and gender. The reason for the different results in the studies is probably that many different enzymes and proteins play a role in oxidative stress. Enzymes with important functions in the fight against free radicals are known as antioxidants. PON and ARE have antioxidant and antiatherogenic effects. They are encoded by the same gene (26). Although PON shows polymorphic change, ARE does not show a genetic polymorphic change (26). There are studies indicating that PON polymorphism is a genetic predisposition to the complications of diabetes (27,28). Also, although the two enzymes have different natural substrates, the PON has the ability to hydrolyse phenylacetate, the natural substrate of ARE. PON prevents lipid oxidation which plays an important role in the development of micro- and macrovascular disease (12,27). Studies showed that PON activity was statistically significantly lower in patients with Type I diabetes compared to control groups (12,28,29). Craciun et al. (12) did not observe a correlation between PON activity and HbA1c in children with T1DM. In our study as in that of Craciun et al., (12) PON activity in the patient group was statistically significantly lower than in the controls but its correlation with HbA1c was not observed. Although ARE activity was lower than the control group; this difference

was not statistically significant. This result can be attributed to the fact that ARE activity is weaker than PON activity. However, the reason for decreased PON activity in patients with T1DM observed in our study as with studies of other investigators is still not fully understood (29). A possible explanation could be a modification of the enzyme's active centre affected by the glycation process. Even though we did not observe a negative correlation between the HbA1c value and PON activity, we speculated that lower PON activity in Type I diabetic patients could be the result of chronic hyperglycemia. Răchișan et al. (30) showed lower activities of PON and ARE in girls with T1DM than boys with ARE. There was no difference in PON and ARE activities in terms of gender in our study. In a study using a diabetic rat model, MDA and blood glucose were reduced in rats with T1DM treated with curcumin but superoxide dismutase and insulin increased (13). Curcumin is a kind of spice extensively used in Asian countries. It has antioxidant and anti-inflammatory effects (13). The antioxidant treatment is thought to improve beta-cell dysfunction, but the results are uncertain (31). Zinc is an essential element for the storage, secretion and action of insulin (31). In addition, it is a key co-factor of many antioxidant enzymes and also helps decrease the effects of inflammatory substances and oxidative stress (31). Zinc stimulates the synthesis of metallothionein, which cleanses hydroxyl radicals (31). Zinc transportation to insulin vesicles is facilitated by ZnT8 which is a transmembrane protein (31,32). Antibodies against ZnT8 are produced in patients with T1DM (33). A study from Sweden showed that low zinc in drinking water is associated with the risk of developing Type I diabetes during childhood (33). Lin et al. (34) did not observe a significant difference in zinc levels between diabetics and controls. As opposed to this study, serum zinc levels in our study were significantly lower in the diabetic patient group than in the control group. The reason for decreased zinc levels in our study is not clear. But a negative correlation with HbA1c was observed in this study. This result suggests that the decrease in zinc may be due to hyperglycemia. The small sample size was the main limitation of this study.

## Conclusion

PON, ARE activities and zinc levels were lower in children and adolescents with T1DM, but the decrease in the ARE activity was not statistically significant. In addition, a negative correlation was observed between zinc and HbA1c. Our results showed that the antioxidant defence systems decreased in children with T1DM. We conclude that antioxidant enzymes should be at normal levels to prevent or delay the complications of Type I diabetes in children, so we suggest that children with T1DM should adopt more physical activity, a healthier diet and less stressful lifestyle.

## Acknowledgements

The authors thank for the financial support provided by University of Health Sciences Haseki Training and Research Hospital, İstanbul, Turkey, Project No: 15

## Ethics

**Ethics Committee Approval:** The study was approved by the University of Health Sciences Haseki Training and Research Hospital Local Ethics Committee (approval number: 48-11/10/2013).

**Informed Consent:** Consent form was filled out by all participants.

**Peer-review:** Externally peer-reviewed.

## Authorship Contributions

Surgical and Medical Practices: Ö.A., N.S.D., Concept: N.S.D., Design: N.S.D., Data Collection or Processing: Ö.A., N.S.D., Analysis or Interpretation: Ö.A., N.S.D., M.E., Literature Search: Ö.A., N.S.D., Writing: N.S.D.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study received no financial support.

## References

1. Castro-Correia C, Maia ML, Norberto S, et al. Can Antioxidative Status Be Involved in Type 1 Diabetes? *J Clin Med Res* 2017;9:998-1001.
2. Tomita T. Apoptosis of pancreatic  $\beta$ -cells in Type 1 diabetes. *Bosn J Basic Med Sci* 2017;17:183-93.
3. Beckhauser TF, Francis-Oliveira J, De Pasquale R. Reactive Oxygen Species: Physiological and Physiopathological Effects on Synaptic Plasticity. *J Exp Neurosci* 2016;10(Suppl 1):23-48.
4. Apostolova N, Victor VM. Molecular strategies for targeting antioxidants to mitochondria: therapeutic implications. *Antioxid Redox Signal* 2015;22:686-729.
5. Samoylenko A, Hossain JA, Mennerich D, Kellokumpu S, Hiltunen JK, Kietzmann T. Nutritional countermeasures targeting reactive oxygen species in cancer: from mechanisms to biomarkers and clinical evidence. *Antioxid Redox Signal* 2013;19:2157-96.
6. Chang D, Zhang X, Rong S, et al. Serum antioxidative enzymes levels and oxidative stress products in age-related cataract patients. *Oxid Med Cell Longev* 2013;2013:587826.
7. Skibska B, Goraca A. The protective effect of lipoic acid on selected cardiovascular diseases caused by age-related oxidative stress. *Oxid Med Cell Longev* 2015;2015:313021.
8. Coudriet GM, Delmastro-Greenwood MM, Previte DM, et al. Treatment with a Catalytic Superoxide Dismutase (SOD) Mimetic Improves Liver Steatosis, Insulin Sensitivity, and Inflammation in Obesity-Induced Type 2 Diabetes. *Antioxidants (Basel)* 2017;6:E85.
9. Kubo E, Chhunchha B, Singh P, Sasaki H, Singh DP. Sulforaphane reactivates cellular antioxidant defense by inducing Nrf2/ARE/Prdx6 activity during aging and oxidative stress. *Sci Rep* 2017;7:14130.
10. Barseem N, Elsamalehy M. Gene Polymorphisms of Glutathione S-Transferase T1/M1 in Egyptian Children and Adolescents with

- Type 1 Diabetes Mellitus. *J Clin Res Pediatr Endocrinol* 2017;9:138-43.
11. Aral CA, Nalbantoğlu Ö, Nur BG, Altunsoy M, Aral K. Metabolic control and periodontal treatment decreases elevated oxidative stress in the early phases of type 1 diabetes onset. *Arch Oral Biol.* 2017;82:115-20.
  12. Craciun EC, Leucuta DC, Rusu RL, David BA, Cret V, Dronca E. Paraoxonase-1 activities in children and adolescents with type 1 diabetes mellitus. *Acta Biochim Pol* 2016;63:511-5.
  13. Xie Z, Wu B, Shen G, Li X, Wu Q. Curcumin alleviates liver oxidative stress in type 1 diabetic rats. *Mol Med Rep* 2018;17:103-8.
  14. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2012;35(Suppl 1):64-71.
  15. Altan N, Dinçel A, Koca C. Diabetes mellitus and oxidative stress. *Turk J Biochem* 2006;31:51-6.
  16. Kaneto H, Katakami N, Kawamori D, et al. Involvement of oxidative stress in the pathogenesis of diabetes. *Antioxid Redox Signal* 2007;9:355-66.
  17. Lin CC, Huang HH, Hu CW, et al. Trace elements, oxidative stress and glycemic control in young people with type 1 diabetes mellitus. *J Trace Elem Med Biol* 2014;28:18-22.
  18. Fatima N, Faisal SM, Zubair S, et al. Role of Pro-Inflammatory Cytokines and Biochemical Markers in the Pathogenesis of Type 1 Diabetes: Correlation with Age and Glycemic Condition in Diabetic Human Subjects. *PLoS One* 2016;11:e0161548.
  19. Menon V, Ram M, Dorn J, et al. Oxidative stress and glucose levels in a population-based sample. *Diabet Med* 2004;21:1346-52.
  20. Reis JS, Veloso CA, Volpe CM, et al. Soluble RAGE and malondialdehyde in type 1 diabetes patients without chronic complications during the course of the disease. *Diab Vasc Dis Res* 2012;9:309-14.
  21. Matteucci E, Giampietro O. Oxidative stress in families of type 1 diabetic patients. *Diabetes Care* 2000;23:1182-6.
  22. Koca C, Altan N, Dincel AS, Kosova F. Oxidative stress and serum leptin levels in patients with type 1 and 2 diabetes mellitus. *Turk J Biochem* 2008;6:99-107.
  23. Erciyas F, Taneli F, Arslan B, Uslu Y. Glycemic control, oxidative stress, and lipid profile in children with type 1 diabetes mellitus. *Arch Med Res* 2004;35:134-40.
  24. Abou-Seif MA, Youssef AA. Evaluation of some biochemical changes in diabetic patients. *Clin Chim Acta* 2004;346:161-70.
  25. Mishra N, Singh N. Blood viscosity, lipid profile, and lipid peroxidation in type-1 diabetic patients with good and poor glycemic control. *N Am J Med Sci* 2013;5:562-6.
  26. Kurban S, Akpınar Z, Mehmetoğlu İ. Investigation of serum paraoxonase and arylesterase activities and oxidative stress in patients with multiple sclerosis. *Genel Tip Derg* 2010;20:13-7.
  27. Hofer SE, Bennetts B, Chan AK, et al. Association between PON1 polymorphisms, PON activity and diabetes complications. *J Diabetes Complications* 2006;20:322-8.
  28. Fekih O, Triki S, Rejeb J, et al. Paraoxonase 1 polymorphisms (L55M and Q192R) as a genetic marker of diabetic nephropathy in youth with type 1 diabetes. *Endokrynol Pol* 2017;68:35-41.
  29. Wegner M, Pioruńska-Stolzmann M, Araszkievicz A, Zozulińska-Ziótkiewicz D, Wierusz-Wysocka B. Evaluation of paraoxonase 1 arylesterase activity and lipid peroxide levels in patients with type 1 diabetes. *Pol Arch Med Wewn* 2011;121:448-54.
  30. Râchişan AL, David BA, Căinap S, Miu N, Andreica M, Samaşca G. Immunological manifestations in type I diabetic children. *Roum Arch Microbiol Immunol* 2012;71:95-9.
  31. Gerber PA, Rutter GA. The Role of Oxidative Stress and Hypoxia in Pancreatic Beta-Cell Dysfunction in Diabetes Mellitus. *Antioxid Redox Signal* 2017;26:501-18.
  32. Elmaoğulları S, Uçaktürk SA, Elbeg Ş, et al. Prevalence of ZnT8 Antibody in Turkish Children and Adolescents with New Onset Type 1 Diabetes. *J Clin Res Pediatr Endocrinol* 2018;10:108-12.
  33. Samuelsson U, Oikarinen S, Hyöty H, Ludvigsson J. Low zinc in drinking water is associated with the risk of type 1 diabetes in children. *Pediatr Diabetes* 2011;12:156-64.
  34. Lin CC, Tsweng GJ, Lee CF, Chen BH, Huang YL. Magnesium, zinc, and chromium levels in children, adolescents, and young adults with type 1 diabetes. *Clin Nutr* 2016;35:880-4.