



The Prevalence of Colistin Resistant Strains and Antibiotic Resistance Gene Profiles in Funan River, China

Hongmei Tuot[†], Yanxian Yang[†], Xi Tao, Dan Liu, Yunxia Li, Xianjun Xie, Ping Li, Ju Gu, Linghan Kong, Rong Xiang, Changwei Lei, Hongning Wang and Anyun Zhang*

Animal Disease Prevention and Food Safety Key Laboratory of Sichuan Province, Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education, College of Life Sciences, Sichuan University, Chengdu, China

OPEN ACCESS

Edited by:

Gilberto Igrejas,
Universidade de Trás-os-Montes e
Alto Douro, Portugal

Reviewed by:

Magdalena Nüesch-Inderbinen,
University of Zurich, Switzerland
Roger Stephan,
University of Zurich, Switzerland

*Correspondence:

Anyun Zhang
zhanganyun@scu.edu.cn

[†] These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Antimicrobials, Resistance
and Chemotherapy,
a section of the journal
Frontiers in Microbiology

Received: 03 August 2018

Accepted: 29 November 2018

Published: 18 December 2018

Citation:

Tuo H, Yang Y, Tao X, Liu D, Li Y,
Xie X, Li P, Gu J, Kong L, Xiang R,
Lei C, Wang H and Zhang A (2018)
The Prevalence of Colistin Resistant
Strains and Antibiotic Resistance
Gene Profiles in Funan River, China.
Front. Microbiol. 9:3094.
doi: 10.3389/fmicb.2018.03094

Anthropogenic activities near urban rivers may have significantly increased the acquisition and dissemination of antibiotic resistance. In this study, we investigated the prevalence of colistin resistant strains in the Funan River in Chengdu, China. A total of 18 *mcr-1*-positive isolates (17 *Escherichia coli* and 1 *Enterobacter cloacae*) and 6 *mcr-3*-positive isolates (2 *Aeromonas veronii* and 4 *Aeromonas hydrophila*) were detected, while *mcr-2*, *mcr-4* and *mcr-5* genes were not detected in any isolates. To further explore the overall antibiotic resistance in the Funan River, water samples were assayed for the presence of 15 antibiotic resistance genes (ARGs) and class 1 integrons gene (*int1*). Nine genes, *sul1*, *sul2*, *int1*, *aac(6′)-Ib-cr*, *bla_{CTX-M}*, *tetM*, *ermB*, *qnrS*, and *aph(3′)-IIIa* were found at high frequencies (70–100%) of the water samples. It is worth noting that *mcr-1*, *bla_{KPC}*, *bla_{NDM}* and *vanA* genes were also found in water samples, the genes that have been rarely reported in natural river systems. The absolute abundance of selected antibiotic resistance genes [*sul1*, *aac(6′)-Ib-cr*, *ermB*, *bla_{CTX-M}*, *mcr-1*, and *tetM*] ranged from 0 to 6.0 (log₁₀ GC/mL) in water samples, as determined by quantitative polymerase chain reaction (qPCR). The *sul1*, *aac(6′)-Ib-cr*, and *ermB* genes exhibited the highest absolute abundances, with 5.8, 5.8, and 6.0 log₁₀ GC/mL, respectively. The absolute abundances of six antibiotic resistance genes were highest near a residential sewage outlet. The findings indicated that the discharge of resident sewage might contribute to the dissemination of antibiotic resistant genes in this urban river. The observed high levels of these genes reflect the serious degree of antibiotic resistant pollution in the Funan River, which might present a threat to public health.

Keywords: colistin, antibiotic resistance, *mcr-1*, *mcr-3*, urban river, quantitative polymerase chain reaction

INTRODUCTION

Multi-drug resistant (MDR) Gram-negative pathogens are resistant to almost all antibiotics, including cephalosporins, quinolones, aminoglycosides and carbapenems, making treatment difficult. Colistin is considered the last line of defense against MDR Gram-negative pathogens, playing an important role in the treatment of severe bacterial infections (Zavascki et al., 2007). Unfortunately, the recent emergence of plasmid-mediated colistin resistance genes in carbapenem-resistant *Enterobacteriaceae* presents a serious new threat to human health. The plasmid-mediated colistin resistance gene *mcr-1* was first discovered Liu et al. (2016). Soon afterward, another mobile

phosphoethanolamine transferase gene, termed *mcr-2*, was discovered in porcine and bovine *Escherichia coli* isolates in Belgium (Xavier et al., 2016). Recently, Yin et al. (2017) discovered a novel *mcr* subtype, *mcr-3*, encoded on an IncI2 plasmid in an *E. coli* isolated from a pig in China. The *mcr-4* and *mcr-5* genes were detected in Europe almost simultaneously (Borowiak et al., 2017; Carattoli et al., 2017). Although there have been numerous reports of colistin resistance genes in animals and humans, fewer studies have focused on *mcr*-bearing isolates from aquatic environments.

Due to the continual release of antibiotic residues and antibiotic resistant bacteria (ARB) into the environment from hospitals, livestock facilities, and sewage treatment plants (STP), antibiotic resistant genes (ARGs) are regarded as environmental contaminants (Pruden et al., 2006; Zurfluh et al., 2017). The occurrence and dissemination of antibiotic resistance in pathogenic and zoonotic bacteria pose a potential threat to human health (Rosenberg Goldstein et al., 2012; Neyra et al., 2014). Moreover, an increasing number of bacteria are resistant to multiple antibiotics, and are able to transfer their resistant determinants among different bacterial species and genera in aquatic environments (Akinbowale et al., 2006). Urban rivers may provide an ideal setting for the acquisition and dissemination of antibiotic resistance because they are frequently impacted by anthropogenic activities. Although antibiotic resistance is a major and developing public health concern, the surveillance of this phenomenon in urban rivers is remarkably limited.

The Funan River, a major urban river in Chengdu used for agricultural activities (e.g., irrigation and cultivation) as well as recreational activities (e.g., swimming and fishing), was used as the model in this study to analyze the magnitude of antibiotic resistance in urban rivers.

The objectives of this study were: (1) to determine the prevalence of colistin resistance strains in the Funan River; (2) to investigate the MDR phenotypes and genotypes of isolated colistin resistant strains; (3) to screen for resistance determinants, including *sul1*, *sul2*, *bla_{CTX-M}*, *bla_{VIM}*, *bla_{KPC}*, *bla_{NDM}*, *qnrS*, *aac(6′)-Ib-cr*, *vanA*, *mecA*, *ermB*, *ermF*, *tetM*, *aph(3′)-IIIa*, and *mcr-1*, and the class 1 integron gene (*int1*) in water samples from the Funan River.

MATERIALS AND METHODS

Sampling of River Water

To investigate the prevalence of colistin resistant strains, 30 water samples (2 L) were collected from the Funan River near densely populated areas in September 2017. To further explore the antibiotic resistance of bacteria throughout the Funan River, 10 water samples (2 L) were collected from representative locations along the river (Figure 1). The representative locations included river intersections, streams near parks, and sewage outlets near residential areas, the hospital, and the municipal wastewater treatment plant (WWTP). The site near the residential sewage outlet is designated RWW and the sample near the municipal wastewater treatment plant is designated WWTP. Sites P1, P2,

and P3 are close to various parks and HWW1 and HWW2 are close to the hospital sewage outlet. Site RI is located adjacent to the intersection of a tributary and the mainstream of the river. Sites UWP and DWP are upstream and downstream of Wetland Park, respectively. Water samples were collected from each site, immediately placed on ice, and transported to the laboratory within 4 h. The samples were then maintained at 4°C until investigation.


Bacterial Isolation

A total of 30 water samples were concentrated by vacuum filtration through 0.22 μm filter membranes. The membranes were washed and the collected material was suspended in 10 ml of sterile PBS. A volume of 1 ml thereof was added to 9 ml of Brain Heart Infusion (BHI) broth with polymyxin B at a final concentration of 4 μg/mL. After incubation at 37°C overnight, 100 μl culture samples were streaked onto MacConkey agar plates. Fifty colonies were picked from each MacConkey agar plates and subsequently grown in BHI broth with 4 μg/mL polymyxin B for 18–24 h. Isolates were screened for the presence of *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4*, and *mcr-5* by PCR. Next, *mcr*-positive isolates were purified by subculturing. The *mcr*-positive isolates were identified using 16S rRNA gene sequencing and the BD Phoenix-100 Automated Microbiology System (BD Diagnostic Systems, Sparks, NV, United States).

Antimicrobial Resistance Testing and Detection of *mcr*-Positive Strains Genotype

The minimum inhibitory concentration (MIC) of colistin was determined by broth microdilution. The antimicrobial susceptibility was interpreted according to the guidelines of the European Committee on Antimicrobial Susceptibility Testing (EUCAST) version 6.0 (EUCAST, 2017). Fourteen antimicrobial agents were tested: ampicillin (AMP, 10 μg), amoxicillin/clavulanic acid (AMC, 20/10 μg), cefotaxime (CTX, 30 μg), ceftriaxone (CRO, 30 μg), ceftazidime (CAZ, 30 μg), ceftiofloxacin (FOX, 30 μg), imipenem (IPM, 10 μg), ertapenem (ETP, 10 μg), aztreonam (ATM, 30 μg), ciprofloxacin (CIP, 5 μg), fosfomycin (FOS, 50 μg), tetracycline (TE, 30 μg), amikacin (AK, 30 μg) and trimethoprim/sulfamethoxazole (SXT, 1.25/23.75 μg). Antimicrobial susceptibility was determined by the agar disk diffusion method. Isolates were classified as susceptible, intermediate, or resistant using the breakpoints specified by the Clinical and Laboratory Standards Institute (CLSI) (CLSI, 2016). *Escherichia coli* ATCC 25922 was used as the quality control strain.

After DNA extraction using the TIANamp bacteria DNA kit (TIANGEN, China), the isolates were screened for the presence of 21 antibiotic resistance genes (*bla_{KPC}*, *bla_{OXA-48}*, *bla_{NDM}*, *bla_{VIM}*, *bla_{IMP}*, *bla_{SHV}*, *bla_{TEM}*, *bla_{CTX-M-1}*, *bla_{CTX-M-9}*, *fosA3*, *qnrB*, *qnrS*, *floR*, *oqxAB*, *sul1*, *sul2*, *tetM*, *tetA*, *aac(6′)-Ib-cr*, *rmtA*, and *rmtB*) (Berendonk et al., 2015; Zheng et al., 2015; Liu et al., 2016), and the



primers and PCR conditions used are listed in **Table 1**. Negative and positive controls for PCR of each gene were utilized.

Total DNA Extraction and Detection of ARGs

To further explore the extent of antibiotic resistance throughout the Funan River, water samples were collected from 10 locations (**Figure 1**). Total DNA was extracted using the Water DNA kit (OMEGA, United States) from the bacteria sample trapped by 0.22 μm pore filter (2 L samples). Standard PCR performed as listed in **Table 1** was used to detect 15 ARGs (*sul1*, *sul2*, *bla_{CTX-M}*, *bla_{VIM}*, *bla_{KPC}*, *bla_{NDM}*, *qnrS*, *aac(6′)-Ib-cr*, *vanA*, *mecA*, *ermB*, *ermF*, *tetM*, *aph(3′)-IIIa* and *mcr-1*) and the class 1 integron gene (*int11*). Negative and positive controls were used for each set of PCR primers. PCR amplification reactions were conducted in 20 μl volumes containing 1 \times PCR Master Mix (Tsingke, China), 1.0 μl template DNA, and 0.5 μM of each primer. After amplification, 5 μl samples of the PCR products were loaded

on a 1.0% agarose gel containing GoldView, and separated electrophoretically in 1 \times TAE buffer at 120 V for 20 min and visualized.

Quantitative Polymerase Chain Reaction

To compare the abundance of ARGs for different sampling sites, the gene copy numbers of the *sul1*, *aac(6′)-Ib-cr*, *ermB*, *bla_{CTX-M}*, and *tetM* genes were quantified using qPCR assays. These genes confer resistance to five major classes of antibiotics: sulphonamides, aminoglycosides, macrolides, β -lactams, and tetracyclines. The levels of *mcr-1* and 16S rRNA genes were also quantified. To quantitate the amounts of these genes, the levels were compared to the levels in standard samples prepared from plasmids containing these specific genes, as described previously (Chen and Zhang, 2013). The standard samples were diluted to yield a series of 10-fold concentrations and were subsequently used to generate qPCR standard curves. The R^2 values were higher than 0.990 for all standard curves. The 20 μl qPCR mixtures contained 10 μL of SYBR premix Ex TaqTM (TaKaRa, Dalian, China), 0.5 μM of each forward and reverse primer, and 1 μl of template DNA. The final volume was adjusted to

TABLE 1 | Standard primer pairs used in this study.

Target genes	Sequence (5' → 3')	Amplicon size(bp)	Reference
<i>mcr-1</i>	CGGTCAGTCCGTTTGTTC CTTGGTCGGTCTGTA GGG	350	Liu et al., 2016
<i>mcr-2</i>	TGGTACAGCCCCTTTATT GCTTGAGATTGGGTATGA	1617	Xavier et al., 2016
<i>mcr-3</i>	TTGGCACTGTATTTGCATT TTAACGAAATTGGCTGGAACA	542	Yin et al., 2017
<i>mcr-4</i>	ATTGGGATAGTCGCCTTTTT TTACAGCCAGAATCATTATCA	487	Carattoli et al., 2017
<i>mcr-5</i>	ATGCGGTTGTCTGCATTATC TCATTGTGGTTGCCTTTTCTG	1644	Borowiak et al., 2017
<i>bla_{KPC}</i>	ATGTCACTGTATCGCCGTC TTRACTGCCCGTTGACGCC	902	Zheng et al., 2015
<i>bla_{OXA-48}</i>	TTGGTGCCATCGATTATCGG GAGCACTTCTTTGTGATGGC	744	Zheng et al., 2015
<i>bla_{NDM}</i>	ATGGAATTGCCCAATATTATGCAC TCAGCGCAGCTTGTCCGGC	813	Zheng et al., 2015
<i>bla_{VIM}</i>	TTTGGTCGCATATCGCAACG CCATTCAGCCAGATCGGCAT	500	Zheng et al., 2015
<i>bla_{IMP}</i>	GTTTATGTTACATACWTCG GGTTTAAAYAAAACAACCAC	432	Zheng et al., 2015
<i>bla_{SHV}</i>	ATTTGTGCGTCTTACTCGC TTTATGGCGTTACCTTTGACC	861	Zheng et al., 2015
<i>bla_{TEM}</i>	ATGAGTATCAACATTTCCCGTG TTACCAATGCTTAATCAGTGAG	861	Zheng et al., 2015
<i>bla_{CTX-M}</i>	TTTGCGATGTGCGAGTACCAGTAA CGATATCGTTGGTGGTGCCATA	759	Zheng et al., 2015
<i>bla_{CTX-M-1}</i>	AAAAATCACTGCGCCAGTTC AGCTTATTCATCGCCACGTT	415	Zheng et al., 2015
<i>bla_{CTX-M-9}</i>	CAAAGAGAGTGCAACGGATG ATTGAAAGCGTTCATCACC	205	Zheng et al., 2015
<i>fosA3</i>	GCGTCAAGCCTGGCATT GCCGTGAGGTCGAGAAA	282	Hou et al., 2012
<i>qnrB</i>	GATCGTGAAAGCCAGAAAGG ACGATGCCTGGTAGTTGTCC	469	Wang et al., 2017
<i>qnrS</i>	ACGACATTCGTCAACTGCAA TAAATTGGCACCCGTAGGC	540	Wang et al., 2017
<i>oqxAB</i>	CCCTGGACCGCACATAAAG AAAGAACAAGATTCACCGCAAC	1140	Wang et al., 2017
<i>sul1</i>	ATGGTGACGGTGTTCGGCATTCTG CTAGGCATGATCTAACCCCTCGGTC	840	Hur et al., 2011
<i>sul2</i>	GAATAAATCGCTCATTTTTCCGG CGAATTCCTGCGGTTCTTTCCAGC	810	Hur et al., 2011
<i>tetM</i>	AGTGGAGCGATTACAGAA CATATGTCCTGGCGTGTCTA	158	Adefisoye and Okoh, 2016
<i>tetA</i>	GCTACATCCTGCTTGCCCTC CATAGATCGCCGTGAAGAGG	210	Adefisoye and Okoh, 2016
<i>aac(6')-Ib-cr</i>	TTGCGATGCTCTATGAGTGGCTA CTCGAATGCCTGGCGTGTTT	482	Eftekhari and Seyedpour, 2015
<i>rmtA</i>	CTAGCGTCCATCCTTTCCCTC TTGCTTCCATGCCCTTGCC	635	Wang et al., 2017
<i>rmtB</i>	GCTTTCTGCGGGCGATGTAA ATGCAATGCCGCGCTCGTAT	173	Wang et al., 2017

(Continued)

TABLE 1 | Continued

Target genes	Sequence (5'→3')	Amplicon size(bp)	Reference
<i>floR</i>	GTCGAGAAATCCCATGAGTTCA CAGACAGGATACCGACATTCAC	1645	Cloekaert et al., 2000
<i>int1</i>	GGGTCAAGGATCTGGATTTCCG ACATGCGTGAAATCATCGTCCG	484	Mazel et al., 2000
<i>vanA</i>	AATACTGTTTGGGGTTGCTC TTTTCCGGCTCGACTTCCT	734	Kafil and Asgharzadeh, 2014
<i>mecA</i>	TGGTATGTGGAAGTTAGATTGGGAT CTAATCTCATATGTGTTCTGTATTGGC	155	Paterson et al., 2012
<i>ermB</i>	GATACCGTTTACGAAATTGG GAATCGAGACTTGAGTGTGC	364	Zhang et al., 2016
<i>ermF</i>	CGACACAGCTTTGGTTGAAC GGACCTACCTCATAGACAAG	309	Zhang et al., 2016
<i>aph(3')-IIIa</i>	GCC GAT GTG GAT TGC GAA AA GCT TGA TCC CCA GTA AGT CA	269	Udo and Dashti, 2000

20 µl by addition of DNase-free water. The IQTM5 real-time PCR system was employed for amplification and quantification, using the following protocol: 30 s at 95°C, 40 cycles of 5 s at 95°C, 30 s at the annealing temperature, and extension for another 30 s at 72°C. For detection, simultaneous fluorescence signal was scanned at 72°C, followed by a melt curve stage with temperature ramping from 65 to 95°C. Details of the qPCR primers of the target genes and the annealing temperatures are given in Table 2. The method design was adopted from prior research (Thornton and Basu, 2011). The copy numbers of the selected ARGs were normalized against the 16S rRNA gene copy number. Therefore, the copy number unit is described as copies/16S.

Statistical Analysis

Statistical analysis was performed using SPSS 17.0 (IBM, United States). One-Way ANOVA was employed to analyze

the results and values of $P < 0.05$ were considered statistically significant.

RESULTS AND DISCUSSION

The Prevalence of *mcr*-Positive Isolates in the Funan River

The screening of 1500 isolates for *mcr* yielded a total of 24 *mcr*-positive isolates. They included 18 *mcr-1* positive isolates (17 *Escherichia coli* and 1 *Enterobacter cloacae*) and 6 *mcr-3* positive isolates (2 *Aeromonas veronii* and 4 *Aeromonas hydrophila*). *mcr-2*, *mcr-4*, or *mcr-5* were not observed in any of the isolates.

Many reports have described the presence in *mcr-1* in animal- and human- derived *Enterobacteriaceae* isolates isolated worldwide (Du et al., 2016; Liu et al., 2016; Malhotra-Kumar et al., 2016; Shen et al., 2016), but only two previous studies identified *mcr-1* in waterborne *Enterobacteriaceae*. One study reported

TABLE 2 | Quantitative polymerase chain reaction primer pairs used in this study.

Target genes	Sequence (5'→3')	Amplicon size(bp)	Annealing temperatures (°C)	Reference
<i>sul1</i>	CACCGGAAACATCGCTGCA AAGTTCCGCCGCAAGGCT	158	60	Luo et al., 2010
<i>aac(6')-Ib-cr</i>	GTTTCTTCTTCCCACCATCC AGTCCGTCACCTCCATACATTG	103	60	Yang et al., 2018
<i>ermB</i>	CACCGAACACTAGGGTTGC TGTGGTATGGCGGGTAAGT	129	55	This study
<i>bla_{CTX-M}</i>	CAGATTCGGTTCGCTTTCAC GCAAATACTTTATCGTGCTGATG	103	55	Yang et al., 2018
<i>mcr-1</i>	CATCGCGGACAATCTCGG AAATCAACACAGGCTTTAGCAC	116	56	Yang et al., 2017
<i>tetM</i>	TTCAGGTTTACTCGGTTCA GAAGTTAAATAGTGTCTTGGAG	106	55	This study
16S rRNA	CGGTGAATACGTTTCYCGG GGWTACCTGTACGACTT	128	55	Suzuki et al., 2000

detection of the *mcr-1* gene in 1 out of 74 *Enterobacteriaceae* isolated from 21 rivers and lakes in Switzerland that produced extended spectrum β -lactamases (ESBLs) (Zurfluh et al., 2016). In a separate study, similar to our results, Zhou et al. (2017) isolated 23 *mcr-1*-positive isolates from environmental water sources in Hangzhou, indicating that *mcr-1*-carrying *Enterobacteriaceae* may be common in lakes and rivers in China. Data addressing the prevalence of *mcr-3* is limited. Recently, a novel *mcr* variant, *mcr-3*, was first discovered on an IncI2 plasmid from a strain of *E. coli* isolated from a pig in China (Yin et al., 2017). Since then, *mcr-3*-positive strains have been identified in humans and food (Ling et al., 2017; Liu L. et al., 2017). Worryingly, *mcr-3* has been detected on the chromosome of *Aeromonas veronii*, and these chromosomally encoded *mcr-3* determinants can become plasmid-bound and transferable (Cabello et al., 2017; Ling et al., 2017). Recently, Shen et al. (2018a) presented evidence that *mcr* determinants originated from aquatic environments, including *mcr-3* harboring *Aeromonas* spp. Because *Aeromonas* species are prevalent in aquatic environments, the occurrence of colistin resistant isolates in urban rivers is of great concern as these

strains may contribute to the potential dissemination of *mcr* determinants.

Antimicrobial Resistance Phenotypes and Genotypes of *mcr-1* and *mcr-3*-Positive Strains

As shown in Table 3, we next analyzed the antimicrobial resistance phenotypes and genotypes of the isolated *mcr-1* and *mcr-3* positive strains, and found 21 (87.5%) multidrug resistance isolates. The antimicrobial resistance testing showed that all isolates were resistant to colistin (MIC \geq 4 μ g/mL). Of the other antimicrobials tested, the most frequent resistance was to CTX (75%, 18 isolates), followed by CAZ (50%, 12 isolates), AMP (50%, 12 isolates), CRO (45.8%, 11 isolates), ATM (45.8%, 11 isolates), SXT (41.7%, 10 isolates), FOS (29.2%, 7 isolates), TE (25%, 6 isolates), AK (20.8%, 5 isolates), CIP (20.8%, 5 isolates), IPM (16.7%, 4 isolates), FOX (12.5%, 3 isolates), AMC (12.5%, 3 isolates), and ETP (4.2%, 1 isolate). The high occurrence of ESBL producers is worrisome, and corresponds to Zurfluh et al. (2013)

TABLE 3 | The antimicrobial resistance genotypes, phenotypes and MIC values of colistin of *mcr-1* and *mcr-3* positive strains.

Isolates	Species	Antibiotic resistant genes	Antimicrobial resistance phenotypes ^a	MIC values of colistin (μ g/mL)
E22	<i>Escherichia coli</i>	<i>mcr-1</i>	CTX, CAZ, AMP, ATM	16
E23	<i>Escherichia coli</i>	<i>mcr-1</i>	CTX, CAZ	16
E24	<i>Escherichia coli</i>	<i>mcr-1</i>	CTX, CAZ	16
E25	<i>Escherichia coli</i>	<i>mcr-1</i>	CTX, CAZ, ATM	16
E26	<i>Escherichia coli</i>	<i>mcr-1</i>	CTX, CAZ, ATM, AK	16
E27	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>sul2</i>	CRO, ATM, SXT	16
E28	<i>Escherichia coli</i>	<i>mcr-1</i>	CTX, CRO, CAZ, ATM, AK	16
E29	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>bla</i> _{CTX-M-9} , <i>fosA3</i> , <i>qnrS</i> , <i>floR</i> , <i>oqxAB</i> , <i>sul1</i> , <i>sul2</i> , <i>tetA</i> , <i>aac(6')-lb-cr</i>	CTX, CRO, AMP, SXT, CIP	16
E30	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>sul1</i> , <i>tetA</i>	CRO, FOX, ATM, SXT, FOS, TE	16
E31	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>floR</i> , <i>sul2</i> , <i>tetM</i>	CTX, CRO, CAZ, SXT	16
E32	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>floR</i> , <i>sul2</i> , <i>tetM</i>	CTX, CRO, CAZ, SXT	16
E33	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>floR</i> , <i>sul2</i> , <i>tetM</i>	CTX, CRO, CAZ, SXT	16
E34	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>bla</i> _{TEM} , <i>bla</i> _{CTX-M-9}	CTX, CRO, CAZ, AMP	16
E35	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>bla</i> _{TEM} , <i>bla</i> _{CTX-M-9} , <i>floR</i> , <i>oqxAB</i> , <i>sul1</i> , <i>sul2</i> , <i>tetM</i> , <i>tetA</i> , <i>aac(6')-lb-cr</i> , <i>rmtB</i>	CTX, CRO, FOX, AMP, ATM, SXT, TE, AK, CIP	16
E36	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>bla</i> _{TEM} , <i>bla</i> _{CTX-M-9} , <i>fosA3</i> , <i>oqxAB</i> , <i>sul1</i> , <i>sul2</i>	CTX, CRO, AMP, ATM, SXT, FOS, CIP	16
E38	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>tetM</i> , <i>tetA</i>	CTX, CAZ, AMP, ATM, TE	16
E39	<i>Escherichia coli</i>	<i>mcr-1</i> , <i>qnrS</i> , <i>tetA</i>	CTX, ATM, AMC, TE, FOS, CIP	8
E37	<i>Enterobacter cloacae</i>	<i>mcr-1</i> , <i>floR</i> , <i>sul2</i> , <i>rmtA</i> , <i>rmtB</i>	CTX, FOX, AMP, AMC, SXT, AK, IPM	16
A4	<i>Aeromonas veronii</i>	<i>mcr-3</i> , <i>bla</i> _{SHV} , <i>sul1</i>	CTX, IPM	4
A19	<i>Aeromonas hydrophila</i>	<i>mcr-3</i> , <i>bla</i> _{TEM} , <i>bla</i> _{CTX-M-9} , <i>qnrB</i> , <i>sul1</i> , <i>sul2</i> , <i>tetA</i>	CTX, CRO, CAZ, AMP, ATM, AMC, FOS, TE, IPM	16
A48	<i>Aeromonas hydrophila</i>	<i>mcr-3</i> , <i>sul1</i> , <i>rmtA</i> , <i>rmtB</i>	AMP, FOS	8
A49	<i>Aeromonas hydrophila</i>	<i>mcr-3</i> , <i>sul1</i> , <i>sul2</i> , <i>rmtA</i> , <i>rmtB</i>	AMP, FOS, AK	8
A52	<i>Aeromonas hydrophila</i>	<i>mcr-3</i> , <i>qnrS</i> , <i>floR</i> , <i>sul1</i> , <i>tetA</i>	AMP, FOS, TE CIP	4
A54	<i>Aeromonas veronii</i>	<i>mcr-3</i> , <i>sul1</i>	AMP, SXT, IPM, ETP	4

^aCTX, cefotaxime; CRO, ceftriaxone; CAZ, ceftazidime; FOX, ceftoxitin; AMP, ampicillin; ATM, aztreonam; AMC: amoxicillin-clavulanic acid; SXT, trimethoprim-sulfamethoxazole; FOS, fosfomicin; TE, tetracycline; AK, amikacin; CIP, ciprofloxacin; IPM, imipenem; ETP, ertapenem.

who found 74 ESBL-producing isolates from 21 (36.2%) of 58 rivers and lakes, and all showed the multidrug resistance phenotype. In another study, 70% of fluoroquinolone resistant *E. coli* isolated from an urban river showed resistance to three or more classes of antibiotics (Zurfluh et al., 2014). The widespread distribution of MDR bacteria suggested serious drug-resistant pollution in river water. In this study, cephalosporin resistant strains were found most frequently, which may be related to the extensive use of cephalosporins for clinical and veterinary purposes. Overall, high usage has led to increased occurrence and wide distribution of ESBLs in bacteria (Bradford, 2001; Bonnet, 2004).


The *mcr-1* and *mcr-3* positive isolates were next assayed for the presence of other ARGs. The *bla_{SHV}*, *bla_{TEM}* and *bla_{CTX-M-9}* genes were detected in 1 (4.2%), 4 (16.7%), and 5 (20.8%) isolates, respectively. None of the isolates were positive for *bla_{KPC}*, *bla_{OXA-48}*, *bla_{NDM}*, *bla_{VIM}*, *bla_{IMP}* or *bla_{CTX-M-1}*. Fifteen (62.5%) of isolates contain sulphonamide resistance genes (*sul1* in 5 isolates, *sul2* in 5 isolates, and *sul1/sul2* combined in 5 isolates). Some isolates contained genes encoding tetracycline resistance, with 20.8% and 29.2% positive for *tetM* and *tetA* genes, respectively. Some isolates contained genes encoding fluoroquinolone resistance genes, *qnrB*, *qnrS*, and *oqxAB*, which were detected in 1(4.2%), 3(12.5%), and 3(12.5%) isolates, respectively. Genes associated with aminoglycoside resistance, *aac(6′)-Ib-cr*, *rmtA*, and *rmtB*, were amplified in 2 (8.3%), 3 (12.5%), and 4 (16.7%) isolates, respectively. The *floR* gene was detected in 7 (29.2%) isolates and the *fosA3* gene was identified in 2 (8.3%) isolates. According to a recent report, 77.3% of *mcr-1*-positive *E. coli* (34/44) carried at least 1 ESBL gene, and several isolates carried 3 or more ESBL genes (Wu et al., 2018). Furthermore, *bla_{CTX-M-9}* was one of the most

prevalent genes among the identified ESBL genes in China (Liu et al., 2015). Consistent with previous reports, sulphonamides and tetracycline resistance genes are the most abundant ARGs in rivers (Yang et al., 2018). We identified two strains (E29 and E36) that carried *mcr-1*, *fosA3*, and *bla_{CTX-M-9}* genes from river samples (Table 3). The *mcr-1*, *fosA3*, and ESBLs genes were previously identified in *E. coli* isolated from animal and food samples (Liu X. et al., 2017; Lupo et al., 2018), and the presence of these multidrug-resistant strains in urban river may present a serious threat to public health.

Prevalence of Antibiotic Resistance Genes in the Funan River

In this study, the prevalence of ARGs in water samples was investigated by sampling various sites along the Funan River. The *sul1*, *qnrS*, *tetM*, and *int11* genes were detected in samples from all 10 sampling sites (100%). Additionally, *aac(6′)-Ib-cr*, *sul2*, *aph(3′)-IIIa*, *ermB*, and *bla_{CTX-M}* were detected at high rates of 90%, 90%, 90%, 80% and 70%, respectively. Many studies have reported the presence of these genes in aquatic environments (Hu et al., 2008; D’Costa et al., 2011; van Hoek et al., 2011; Lin et al., 2015; Makowska et al., 2016). Interestingly, the *aph(3′)-IIIa* gene has rarely been reported in river water microorganisms, but has been reported in clinical specimens (Tuhina et al., 2016). The detection of the *aph(3′)-IIIa* gene was high in this study, suggesting contamination of the Funan River with resistant bacteria carrying the *aph(3′)-IIIa* gene.

Genes conferring resistance to the last line of antibiotics, including *mcr-1*, *bla_{NDM}*, *bla_{KPC}* and *vanA* genes, were detected at rates of 30%, 20%, 10%, and 10%, respectively. *bla_{VIM}* was not detected at any site. The *mcr-1* gene was detected in 30%



of samples, suggesting the Funan River could act as a reservoir for the *mcr-1* gene. The *bla*_{NDM}, *bla*_{KPC} and *vanA* genes were detected near the WWTP (Figure 1). Although *mcr-1* is found frequently in human and animal settings, there is only limited data for urban rivers (Marathe et al., 2017; Ovejero et al., 2017; Yang et al., 2017). Similarly, Marathe et al. detected *bla*_{NDM} and *bla*_{KPC} genes in the sediments of an Indian river (Marathe et al., 2017). Although a *bla*_{VIM} positive carbapenem-resistant strain was isolated from a river in Switzerland (Zurfluh et al., 2013), here is a lack of data on *bla*_{VIM} in the non-clinical environment. The *vanA* gene is associated with vancomycin resistance and has been found in wastewater biofilms and in drinking water biofilms in Mainz (Schwartz et al., 2003). Although these genes have rarely been identified in natural aquatic environments, given the dangerous infections that can arise from ARB (and which subsequently create intractable challenges for clinical treatment), further observation of the prevalence of these genes in aquatic environments is required.

Abundance of ARGs

Concerning the absolute abundance of ARGs in the Funan River, ARGs were detected at levels that ranged from 0 to 6.0 log₁₀ GC/mL (Figure 2). The *sul1*, *aac(6′)-Ib-cr*, and *ermB* genes were the dominant ARGs in the Funan River with mean absolute abundances of 4.8, 4.1, and 3.4 log₁₀ GC/mL, respectively. The *sul1* gene exhibited the most prominent average abundance in water samples. Previous studies reported that *sul1* is abundant in numerous water areas, including the Tordera River Basin (Proia et al., 2016) and the Haihe River (Luo et al., 2010). Although the *mcr-1* gene was not detected in water samples at some sites, three sites (RWW, HWW1, and HWW2) displayed 2.0–2.7 log₁₀ GC/mL. Notably, the highest detected level of *mcr-1* (2.7 log₁₀ GC/mL) was higher than that in previous reports about the Haihe river (2.6 log₁₀ GC/mL) (Yang et al., 2017). The absence of *mcr-1* in some samples may indicate that no *mcr-1* positive strains were present in the water samples or that the levels of *mcr-1* were below the detection limit. Site RWW is located near the residential sewage outlet, suggesting the presence of *mcr-1* was related to human activity. Consistently, *mcr-1* was detected at HWW1 and HWW2, adjacent to the hospital sewage outlets, suggesting the spread of *mcr-1* from hospitals to urban river, although colistin is not used widely in human medicine. The *mcr-1* abundance at RWW (2.7 log₁₀ GC/mL) was slightly higher than that at HWW1 (2.6 log₁₀ GC/mL) and at HWW2 (2.3 log₁₀ GC/mL). Similarly, the prevalence of *mcr-1*-positive *E. coli* from healthy individuals (0.7–6.2%) is higher than the prevalence for inpatients (0.4–2.9%) (Shen et al., 2018b). It is striking that *mcr* is the only gene that was absent from sites other than RWW and HWW. The reasons for high rate of fecal carriage of *mcr* in humans in China may reflect the rapid emergence of plasmid-encoded *mcr-1* within many MDR *E. coli* carried by humans and also be related to the significant diversity and genetic flexibility of MGEs harboring *mcr-1* (Zhong et al., 2018).

At RWW, RI, and WWTP, the absolute abundances of certain ARGs (*sul1*, *aac(6′)-Ib-cr*, and *ermB*) were significantly higher than those at other sampling sites ($P < 0.05$). At P3 and DWP, the absolute abundances of most ARGs were significantly

lower than the levels detected at the other sites ($P < 0.05$). RWW was associated with the highest absolute abundance of the six ARGs (*mcr-1*, *sul1*, *aac(6′)-Ib-cr*, *ermB*, *bla*_{CTX-M}, and *tetM*) (Figure 2). Samples near the wastewater treatment plant (WWTP) and densely populated areas exhibited a relatively greater content of resistant genes. Wastewater discharge may contribute to the spread of ARGs into the environment, thereby affecting the bacterial communities of the receiving river (Marti et al., 2013; Xu et al., 2015). Our results indicate that human activities influence the dissemination of resistance genes in the Funan River. Remarkably, the absolute abundances of most ARGs were low at the DWP sampling point, located downstream of the wetland park. This is consistent with a decrease in the ARGs levels of the effluents from a constructed wetland with a free surface flow (Liu et al., 2014).

As shown in Figure 2, the relative abundances of each ARG are only partly correlated with their absolute abundance. That is, although the absolute abundances of most ARGs at RWW, RI and WWTP were relatively high, their relative abundances were comparatively low. These differences may be related to the differences in the proportion of resistant bacteria to total bacteria at each site (Tao et al., 2014).

CONCLUSION

This study describes 18 *mcr-1*-positive strains and 6 *mcr-3*-positive strains isolated from the Funan River, of which 87.5% were found to be MDR. The *sul1*, *sul2*, *int11*, *aac(6′)-Ib-cr*, *bla*_{CTX-M}, *tetM*, *ermB*, *qnrS* and *aph(3′)-IIIa* genes were abundant in the Funan River. Interestingly, the *mcr-1*, *bla*_{KPC}, *bla*_{NDM}, and *vanA* genes were detected, although these four resistance genes have rarely been found in natural river systems. Notably, the *mcr-1* gene was detected at a rate of 30%. Our results suggest urban activities may increase the prevalence of antibiotic resistance genes and demonstrate the current presence of drug-resistance pollution in the Funan River. The processes by which the dissemination of ARGs occurs in urban rivers should be the focus of future studies.

AUTHOR CONTRIBUTIONS

AZ designed the study. HT, DL, XX, and PL carried out the sampling work. HT, YY, XT, and JG performed the experiments. AZ, HT, RX, LK, and CL analyzed the data. AZ, HT, YL, and HW drafted the manuscript. All authors have read and approved the final manuscript.

FUNDING

This work was supported by the National Key Research and Development Program of China (Grant No. 2018YFD0500300), the General Program of National Natural Science Foundation of China (Grant No. 31572548), the Applied Basic Research Program in Sichuan Province (2018JY0572), and the Science & Technology Pillar Program in Sichuan Province (2018HH0027).

REFERENCES

- Adefisoye, M. A., and Okoh, A. I. (2016). Identification and antimicrobial resistance prevalence of pathogenic *Escherichia coli* strains from treated wastewater effluents in Eastern Cape, South Africa. *Microbiologyopen* 5, 143–151. doi: 10.1002/mbio.3.319
- Akinbowale, O. L., Peng, H., and Barton, M. D. (2006). Antimicrobial resistance in bacteria isolated from aquaculture sources in Australia. *J. Appl. Microbiol.* 100, 1103–1113. doi: 10.1111/j.1365-2672.2006.02812.x
- Berendonk, T. U., Manaia, C. M., Merlin, C., Fatta-Kassinos, D., Cytryn, E., Walsh, F., et al. (2015). Tackling antibiotic resistance: the environmental framework. *Nat. Rev. Microbiol.* 13, 310–317. doi: 10.1038/nrmicro3439
- Bonnet, R. (2004). Growing group of extended-spectrum-lactamases: the CTX-M Enzymes. *Antimicrob. Agents Chemother.* 48, 1–14. doi: 10.1128/AAC.48.1.1-14.2004
- Borowiak, M., Fischer, J., Hammerl, J. A., Hendriksen, R. S., Szabo, I., and Malorny, B. (2017). Identification of a novel transposon-associated phosphoethanolamine transferase gene, *mcr-5*, conferring colistin resistance in d-tartrate fermenting *Salmonella enterica* subsp. *enterica* serovar *Paratyphi B*. *J. Antimicrob. Chemother.* 72, 3317–3324. doi: 10.1093/jac/dkx327
- Bradford, P. A. (2001). Extended-spectrum beta-lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clin. Microbiol. Rev.* 14, 933–951. doi: 10.1128/CMR.14.4.933-951.2001
- Cabello, F., Tomova, A., Ivanova, L., and Godfrey, H. (2017). Aquaculture and *mcr* colistin resistance determinants. *mBio* 8:e01229-17. doi: 10.1128/mBio.01229-17
- Carattoli, A., Villa, L., Feudi, C., Curcio, L., Orsini, S., Luppi, A., et al. (2017). Novel plasmid-mediated colistin resistance *mcr-4* gene in *Salmonella* and *Escherichia coli*, Italy 2013, Spain and Belgium, 2015 to 2016. *Euro. Surveill.* 22:30589. doi: 10.2807/1560-7917.ES.2017.22.31.30589
- Chen, H., and Zhang, M. (2013). Occurrence and removal of antibiotic resistance genes in municipal wastewater and rural domestic sewage treatment systems in eastern China. *Environ. Int.* 55, 9–14. doi: 10.1016/j.envint.2013.01.019
- Clockaert, A., Baucheron, S., Flaujac, G., Schwarz, S., Kehrenberg, C., Martel, J. L., et al. (2000). Plasmid-mediated florfenicol resistance encoded by the *floR* gene in *Escherichia coli* isolated from cattle. *Antimicrob. Agents Chemother.* 44, 2858–2860. doi: 10.1128/AAC.44.10.2858-2860.2000
- CLSI (2016). *Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Sixth Informational Supplement M100S*. Wayne, PA: CLSI.
- D'Costa, V. M., King, C. E., Kalan, L., Morar, M., Sung, W. W., Schwarz, C., et al. (2011). Antibiotic resistance is ancient. *Nature* 477, 457–461. doi: 10.1038/nature10388
- Du, H., Chen, L., Tang, Y. W., and Kreiswirth, B. N. (2016). Emergence of the *mcr-1* colistin resistance gene in carbapenem-resistant *Enterobacteriaceae*. *Lancet. Infect. Dis.* 16, 287–288. doi: 10.1016/S1473-3099(16)00056-56
- Eftekhari, F., and Seyedpour, S. M. (2015). Prevalence of *qnr* and *aac(6)-Ib-cr* genes in clinical isolates of *Klebsiella pneumoniae* from imam hussein hospital in Tehran. *Iran. J. Med. Sci.* 40, 515–521.
- EUCAST (2017). *European Committee on Antimicrobial Susceptibility Testing Clinical Breakpoints – Bacteria (v. 7.0)*. Available at: http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_7.1_Breakpoint_Tables.pdf
- Hou, J., Huang, X., Deng, Y., He, L., Yang, T., Zeng, Z., et al. (2012). Dissemination of the fosfomycin resistance gene *fosA3* with CTX-M beta-lactamase genes and *rmtB* carried on IncFII plasmids among *Escherichia coli* isolates from pets in China. *Antimicrob. Agents Chemother.* 56, 2135–2138. doi: 10.1128/AAC.05104-11
- Hu, J., Shi, J., Chang, H., Li, D., Yang, M., and Kamagata, Y. (2008). Phenotyping and genotyping of antibiotic-resistant *Escherichia coli* isolated from a natural river basin. *Environ. Sci. Technol.* 42, 3415–3420. doi: 10.1021/es7026746
- Hur, J., Kim, J. H., Park, J. H., Lee, Y. J., and Lee, J. H. (2011). Molecular and virulence characteristics of multi-drug resistant *Salmonella enteritidis* strains isolated from poultry. *Vet. J.* 189, 306–311. doi: 10.1016/j.tvjl.2010.07.017
- Kafil, H. S., and Asgharzadeh, M. (2014). Vancomycin-resistant *Enterococcus faecium* and *Enterococcus faecalis* isolated from education hospital of Iran. *Maedica* 9, 323–327.
- Lin, L., Yuan, K., Liang, X., Chen, X., Zhao, Z., Yang, Y., et al. (2015). Occurrences and distribution of sulfonamide and tetracycline resistance genes in the Yangtze River Estuary and nearby coastal area. *Mar. Pollut. Bull.* 100, 304–310. doi: 10.1016/j.marpolbul.2015.08.036
- Ling, Z., Yin, W., Li, H., Zhang, Q., Wang, X., Wang, Z., et al. (2017). Chromosome-mediated *mcr-3* variants in *Aeromonas veronii* from chicken meat. *Antimicrob. Agents Chemother.* 61:e01272-17. doi: 10.1128/AAC.01272-17
- Liu, L., He, D., Lv, L., Liu, W., Chen, X., Zeng, Z., et al. (2015). *bla*_{CTX-M-1/9/1} hybrid genes may have been generated from *bla*_{CTX-M-15} on an IncI2 plasmid. *Antimicrob. Agents Chemother.* 59, 4464–4470. doi: 10.1128/AAC.00501-15
- Liu, L., Liu, Y. H., Wang, Z., Liu, C. X., Huang, X., and Zhu, G. F. (2014). Behavior of tetracycline and sulfamethazine with corresponding resistance genes from swine wastewater in pilot-scale constructed wetlands. *J. Hazard. Mater.* 278, 304–310. doi: 10.1016/j.jhazmat.2014.06.015
- Liu, X., Li, R., Zheng, Z., Chen, K., Xie, M., Chan, E. W., et al. (2017). Molecular characterization of *Escherichia coli* isolates carrying *mcr-1*, *fosA3*, and extended-spectrum-beta-lactamase genes from food samples in China. *Antimicrob. Agents Chemother.* 61:e00064-17. doi: 10.1128/AAC.00064-17
- Liu, L., Feng, Y., Zhang, X., McNally, A., and Zong, Z. (2017). New variant of *mcr-3* in an extensively drug-resistant *Escherichia coli* clinical isolate carrying *mcr-1* and *bla*_{NDM-5}. *Antimicrob. Agents Chemother.* 61, e1757–e1717. doi: 10.1128/AAC.01757-17
- Liu, Y. Y., Wang, Y., Walsh, T. R., Yi, L. X., Zhang, R., Spencer, J., et al. (2016). Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect. Dis.* 16, 161–168. doi: 10.1016/S1473-3099(15)00424-7
- Luo, Y., Mao, D., Rysz, M., Zhou, Q., Zhang, H., Xu, L., et al. (2010). Trends in antibiotic resistance genes occurrence in the Haihe river, China. *Environ. Sci. Technol.* 44, 7220–7225. doi: 10.1021/es100233w
- Lupo, A., Saras, E., Madec, J. Y., and Haenni, M. (2018). Emergence of *bla*_{CTX-M-55} associated with *fosA*, *rmtB* and *mcr* gene variants in *Escherichia coli* from various animal species in France. *J. Antimicrob. Chemother.* 73, 867–872. doi: 10.1093/jac/dkx489
- Makowska, N., Koczura, R., and Mokracka, J. (2016). Class I integrase, sulfonamide and tetracycline resistance genes in wastewater treatment plant and surface water. *Chemosphere* 144, 1665–1673. doi: 10.1016/j.chemosphere.2015.10.044
- Malhotra-Kumar, S., Xavier, B. B., Das, A. J., Lammens, C., Hoang, H. T. T., Pham, N. T., et al. (2016). Colistin-resistant *Escherichia coli* harbouring *mcr-1* isolated from food animals in Hanoi. *Vietnam. Lancet. Infect. Dis.* 16, 286–287. doi: 10.1016/S1473-3099(16)00014-1
- Marathe, N. P., Pal, C., Gaikwad, S. S., Jonsson, V., Kristiansson, E., and Larsson, D. G. J. (2017). Untreated urban waste contaminates Indian river sediments with resistance genes to last resort antibiotics. *Water Res.* 124, 388–397. doi: 10.1016/j.watres.2017.07.060
- Marti, E., Jofre, J., and Balcazar, J. L. (2013). Prevalence of antibiotic resistance genes and bacterial community composition in a river influenced by a wastewater treatment plant. *PLoS One* 8:e78906. doi: 10.1371/journal.pone.0078906
- Mazel, D., Dychinco, B., Webb, V. A., and Davies, J. (2000). Antibiotic resistance in the ECOR collection: integrons and identification of a novel *aad* gene. *Antimicrob. Agents Chemother.* 44, 1568–1574. doi: 10.1128/AAC.44.6.1568-1574.2000
- Neyra, R. C., Frisancho, J. A., Rinsky, J. L., Resnick, C., Carroll, K. C., Rule, A. M., et al. (2014). Multidrug-resistant and methicillin-resistant *Staphylococcus aureus* (MRSA) in hog slaughter and processing plant workers and their community in North Carolina (USA). *Environ. Health Perspect.* 122, 471–477. doi: 10.1289/ehp.1306741
- Ovejero, C. M., Delgado-Blas, J. F., Calero-Caceres, W., Muniesa, M., and Gonzalez-Zorn, B. (2017). Spread of *mcr-1*-carrying *Enterobacteriaceae* in

- sewage water from Spain. *J. Antimicrob. Chemother.* 72, 1050–1053. doi: 10.1093/jac/dkw533
- Paterson, G. K., Larsen, A. R., Robb, A., Edwards, G. E., Pennycott, T. W., Foster, G., et al. (2012). The newly described *mecA* homologue, *mecALGA251*, is present in methicillin-resistant *Staphylococcus aureus* isolates from a diverse range of host species. *J. Antimicrob. Chemother.* 67, 2809–2813. doi: 10.1093/jac/dks329
- Proia, L., von Schiller, D., Sánchez-Melsió, A., Sabater, S., Borrego, C. M., Rodríguez-Mozaz, S., et al. (2016). Occurrence and persistence of antibiotic resistance genes in river biofilms after wastewater inputs in small rivers. *Environ. Pollut.* 210, 121–128. doi: 10.1016/j.envpol.2015.11.035
- Pruden, A., Pei, R., Storteboom, H., and Carlson, K. H. (2006). Antibiotic resistance genes as emerging contaminants: studies in Northern Colorado. *Environ. Sci. Technol.* 40, 7445–7450. doi: 10.1021/es060413l
- Rosenberg Goldstein, R. E., Micallef, S. A., Gibbs, S. G., Davis, J. A., He, X., George, A., et al. (2012). Methicillin-resistant *Staphylococcus aureus* (MRSA) detected at four U.S. wastewater treatment plants. *Environ. Health Perspect.* 120, 1551–1558. doi: 10.1289/ehp.1205436
- Schwartz, T., Kohnen, W., Jansen, B., and Obst, U. (2003). Detection of antibiotic-resistant bacteria and their resistance genes in wastewater, surface water, and drinking water biofilms. *FEMS. Microbiol. Ecol.* 43, 325–335. doi: 10.1111/j.1574-6941.2003.tb01073.x
- Shen, Y., Yin, W., Liu, D., Shen, J., and Wang, Y. (2018a). Reply to Cabello et al., “Aquaculture and *mcr* colistin resistance determinants. *mBio.* 9:e01629-18. doi: 10.1128/mBio.01629-18
- Shen, Y., Zhou, H., Xu, J., Wang, Y., Zhang, Q., Walsh, T., et al. (2018b). Anthropogenic and environmental factors associated with high incidence of *mcr-1* carriage in humans across China. *Nat. Microbiol.* 3, 1054–1062. doi: 10.1038/s41564-018-0205-8
- Shen, Z., Wang, Y., Shen, Y., Shen, J., and Wu, C. (2016). Early emergence of *mcr-1* in *Escherichia coli* from food-producing animals. *Lancet. Infect. Dis.* 16:293. doi: 10.1016/S1473-3099(16)00061-X
- Suzuki, M. T., Taylor, L. T., and DeLong, E. F. (2000). Quantitative analysis of small-subunit rRNA genes in mixed microbial populations via 5'-nuclease assays. *Appl. Environ. Microbiol.* 66, 4605–4614. doi: 10.1128/AEM.66.11.4605-4614.2000
- Tao, C. W., Hsu, B. M., Ji, W. T., Hsu, T. K., Kao, P. M., Hsu, C. P., et al. (2014). Evaluation of five antibiotic resistance genes in wastewater treatment systems of swine farms by real-time PCR. *Sci. Total Environ.* 496, 116–121. doi: 10.1016/j.scitotenv.2014.07.024
- Thornton, B., and Basu, C. (2011). Real-time PCR (qPCR) primer design using free online software. *Biochem. Mol. Biol. Educ.* 39, 145–154. doi: 10.1002/bmb.20461
- Tuhina, B., Anupurba, S., and Karuna, T. (2016). Emergence of antimicrobial resistance and virulence factors among the unusual species of *enterococci*, from North India. *Indian. J. Pathol. Microbiol.* 59, 50–55. doi: 10.4103/0377-4929.174795
- Udo, E. E., and Dashti, A. A. (2000). Detection of genes encoding aminoglycoside-modifying enzymes in *staphylococci* by polymerase chain reaction and dot blot hybridization. *Int. J. Antimicrob. Agents* 13, 273–279. doi: 10.1016/S0924-8579(99)00124-7
- van Hoek, A. H., Mevius, D., Guerra, B., Mullany, P., Roberts, A. P., and Aarts, H. J. (2011). Acquired antibiotic resistance genes: an overview. *Front. Microbiol.* 2:203. doi: 10.3389/fmicb.2011.00203
- Wang, Y., Zhang, A., Yang, Y., Lei, C., Jiang, W., Liu, B., et al. (2017). Emergence of *Salmonella enterica* serovar Indiana and California isolates with concurrent resistance to cefotaxime, amikacin and ciprofloxacin from chickens in China. *Int. J. Food Microbiol.* 262, 23–30. doi: 10.1016/j.ijfoodmicro.2017.09.012
- Wu, C., Wang, Y., Shi, X., Wang, S., Ren, H., Shen, Z., et al. (2018). Rapid rise of the ESBL and *mcr-1* genes in *Escherichia coli* of chicken origin in China, 2008–2014. *Emerg. Microbes Infect.* 7:30. doi: 10.1038/s41426-018-0033-1
- Xavier, B. B., Lammens, C., Ruhul, R., Kumar-Singh, S., Butaye, P., Goossens, H., et al. (2016). Identification of a novel plasmid-mediated colistin-resistance gene, *mcr-2*, in *Escherichia coli*, Belgium, June 2016. *Euro. Surveill.* 21, doi: 10.2807/1560-7917
- Xu, J., Xu, Y., Wang, H., Guo, C., Qiu, H., He, Y., et al. (2015). Occurrence of antibiotics and antibiotic resistance genes in a sewage treatment plant and its effluent-receiving river. *Chemosphere* 119, 1379–1385. doi: 10.1016/j.chemosphere.2014.02.040
- Yang, D., Qiu, Z., Shen, Z., Zhao, H., Jin, M., Li, H., et al. (2017). The occurrence of the colistin resistance gene *mcr-1* in the Haihe river (China). *Int. J. Environ. Res. Public Health* 14:E576. doi: 10.3390/ijerph14060576
- Yang, Y., Shi, W., Lu, S. Y., Liu, J., Liang, H., Yang, Y., et al. (2018). Prevalence of antibiotic resistance genes in bacteriophage DNA fraction from Funan River water in Sichuan, China. *Sci. Total Environ.* 626, 835–841. doi: 10.1016/j.scitotenv.2018.01.148
- Yin, W., Li, H., Shen, Y., Liu, Z., Wang, S., Shen, Z., et al. (2017). Novel Plasmid-mediated colistin resistance gene *mcr-3* in *Escherichia coli*. *mBio* 8:e00543-17. doi: 10.1128/mBio.00543-17
- Zavascki, A. P., Goldani, L. Z., Li, J., and Nation, R. L. (2007). Polymyxin B for the treatment of multidrug-resistant pathogens: a critical review. *J. Antimicrob. Chemother.* 60, 1206–1215. doi: 10.1093/jac/dkm357
- Zhang, J., Chen, M., Sui, Q., Wang, R., Tong, J., and Wei, Y. (2016). Fate of antibiotic resistance genes and its drivers during anaerobic co-digestion of food waste and sewage sludge based on microwave pretreatment. *Bioresour. Technol.* 217, 28–36. doi: 10.1016/j.biortech.2016.02.140
- Zheng, B., Zhang, J., Ji, J., Fang, Y., Shen, P., Ying, C., et al. (2015). Emergence of *Raoultella ornithinolytica* coproducing IMP-4 and KPC-2 carbapenemases in China. *Antimicrob. Agents Chemother.* 59, 7086–7089. doi: 10.1128/AAC.01363-15
- Zhong, L., Phan, H., Shen, C., Vihta, K., Sheppard, A., Huang, X., et al. (2018). High rates of human fecal carriage of *mcr-1*-positive multidrug-resistant *Enterobacteriaceae* emerge in china in association with successful plasmid families. *Clin. Infect. Dis.* 66, 676–685. doi: 10.1093/cid/cix885
- Zhou, H. W., Zhang, T., Ma, J. H., Fang, Y., Wang, H. Y., Huang, Z. X., et al. (2017). Occurrence of plasmid- and chromosome-carried *mcr-1* in waterborne *Enterobacteriaceae* in China. *Antimicrob. Agents Chemother.* 61:e00017-e17. doi: 10.1128/AAC.00017-17
- Zurfluh, K., Abgottspon, H., Hächler, H., Nüesch-Inderbinen, M., and Stephan, R. (2014). Quinolone resistance mechanisms among extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli* isolated from rivers and lakes in Switzerland. *PLoS One* 9:e95864. doi: 10.1371/journal.pone.0095864
- Zurfluh, K., Bagutti, C., Brodmann, P., Alt, M., Schulze, J., Fanning, S., et al. (2017). Wastewater is a reservoir for clinically relevant carbapenemase and 16S rRNA methylase producing *Enterobacteriaceae*. *Int. J. Antimicrob. Agents* 50, 436–440. doi: 10.1016/j.ijantimicag.2017.04.017
- Zurfluh, K., Hächler, H., Nüesch-Inderbinen, M., and Stephan, R. (2013). Characteristics of extended-spectrum beta-lactamase- and carbapenemase-producing *Enterobacteriaceae* isolates from rivers and lakes in Switzerland. *Appl. Environ. Microbiol.* 79, 3021–3026. doi: 10.1128/AEM.00054-13
- Zurfluh, K., Poirel, L., Nordmann, P., Nüesch-Inderbinen, M., Hächler, H., and Stephan, R. (2016). Occurrence of the Plasmid-borne *mcr-1* colistin resistance gene in Extended-Spectrum-beta-Lactamase-producing *Enterobacteriaceae* in river water and imported vegetable samples in Switzerland. *Antimicrob. Agents Chemother.* 60, 2594–2595. doi: 10.1128/AAC.00066-16

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Tuo, Yang, Tao, Liu, Li, Xie, Li, Gu, Kong, Xiang, Lei, Wang and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.