



Brain Networks Reorganization During Maturation and Healthy Aging-Emphases for Resilience

Gabriel Gonzalez-Escamilla^{1†}, Muthuraman Muthuraman^{1†}, Venkata C. Chirumamilla¹, Johannes Vogt^{2†} and Sergiu Groppa^{1*†}

¹ Department of Neurology, University Medical Center, Johannes Gutenberg University Mainz, Mainz, Germany, ² Institute for Microscopic Anatomy and Neurobiology, University Medical Center, Johannes Gutenberg University Mainz, Mainz, Germany

OPEN ACCESS

Edited by:

Patrik Roser,
Psychiatrische Dienste Aargau AG,
Switzerland

Reviewed by:

Leandro Da Costa Lane Valiengo,
Universidade de São Paulo, Brazil
Raluca Stetcu,
Spiru Haret University, Romania

*Correspondence:

Sergiu Groppa
segroppa@uni-mainz.de

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Public Mental Health,
a section of the journal
Frontiers in Psychiatry

Received: 06 April 2018

Accepted: 29 October 2018

Published: 21 November 2018

Citation:

Gonzalez-Escamilla G,
Muthuraman M, Chirumamilla VC,
Vogt J and Groppa S (2018) Brain
Networks Reorganization During
Maturation and Healthy
Aging-Emphases for Resilience.
Front. Psychiatry 9:601.
doi: 10.3389/fpsy.2018.00601

Maturation and aging are important life periods that are linked to drastic brain reorganization processes which are essential for mental health. However, the development of generalized theories for delimiting physiological and pathological brain remodeling through life periods linked to healthy states and resilience on one side or mental dysfunction on the other remains a challenge. Furthermore, important processes of preservation and compensation of brain function occur continuously in the cerebral brain networks and drive physiological responses to life events. Here, we review research on brain reorganization processes across the lifespan, demonstrating brain circuits remodeling at the structural and functional level that support mental health and are parallelized by physiological trajectories during maturation and healthy aging. We show evidence that aberrations leading to mental disorders result from the specific alterations of cerebral networks and their pathological dynamics leading to distinct excitability patterns. We discuss how these series of large-scale responses of brain circuits can be viewed as protective or malfunctioning mechanisms for the maintenance of mental health and resilience.

Keywords: resilience, lifespan, brain networks, brain reorganization, health maintenance

INTRODUCTION

Aging is related to alterations of cognitive functioning accompanied by structural and functional brain reorganization (1, 2). Maintained cognitive function late in life is generally achieved by the integrated communication of specific brain regions (3, 4). The functional and structural reorganization of brain circuits occur continuously during the lifespan and play an essential role for preserving brain health (5–7). Hence, abnormal cognitive function may build upon specific alterations of brain networks and their dynamic responses to life events or physiological processes during maturation or aging (8–10). An exact understanding of structural and functional longitudinal properties and a precise characterization of the tissue properties are crucial for modeling the long-term processes and to distinguish between healthy and disease-specific alterations. Hence, modeling interregional connectivity and specific reorganization of cerebral networks topology is likely to promote our understanding of underlying mechanisms of mental health and resilience to life events (11).

Functional connectivity patterns can be obtained from the temporal correlations of spontaneous neurophysiological signal fluctuations between brain regions either by electroencephalography

(EEG) or functional magnetic resonance imaging (fMRI), and then analyzed within a graph-theoretical framework (7, 12, 13). On the other hand, the structural network connectivity can be accessed by pure structural MRI measures like cortical thickness and volume or by white matter fiber tracts obtained through tractography between predefined regions of interest (6, 14, 15) as shown in **Figure 1**. These network fingerprints are predictive measures of disease-related clinical symptoms (6, 7) or therapy outcomes (16).

In this review, we present existing evidence for a profound understanding of lifespan-related reorganization processes that can be related to protective mechanisms that help our brain to cope with age-related situations and reduce the burden for brain alterations linked to mental disorders. Unraveling complementary functional and structural fingerprints should give important insight on inter-individual courses. On the basis of our recent results on the importance of the cerebral networks in disease outcome (6, 13, 16–19), we discuss the impact of the structural gray matter tissue integrity and reorganization of normal appearing white matter and evolving functional adaptations for clinical phenotypes. Non-invasive structural and fMRI characterization of the neuronal circuits will be discussed from a longitudinal perspective.

We hypothesize that (i) cerebral networks in resilient subjects with preserved mental health despite traumatic events are characterized by a reorganization of the gray and white matter compartments with a strengthening of distinct regional connectivity patterns and preserved structural integrity in the key anatomical regions (prefrontal cortex, hippocampus and corpus callosum); (ii) these brain circuits remodeling processes are partially mirrored in age-related reorganization during maturation or healthy aging; (iii) neurocognitive and clinical impairment of mental health is associated with exhaustion of network compensation, which manifests in divergent lifespan patterns of network reorganization or a breakdown of functional responses; (iv) structural and functional lateralization patterns together with inter-hemispheric connections and the connectivity fingerprints in the above mentioned networks together with their functional interactions have a large impact on network compensation and thus inter-individual mental status. The overall aim of this work is to identify distinct network connectivity and integrity patterns reflecting compensation processes for global network functioning. We focus first on maturation processes and healthy aging and draw parallels to mechanisms of resilience behavior.


AGE-RELATED BRAIN REMODELING AND EXCITABILITY DYNAMICS DURING MATURATION

Accurate synaptic transmission is a fundamental requirement for normal brain function (20); signaling alterations at the excitatory synapse leading to cortical hyperexcitability have been related to psychiatric disorders such as schizophrenia (21–24). However, to what extent alterations in brain maturation leading to cortical miswiring and subsequent cortical hyperexcitability

contribute to psychiatric disorders has not been fully elucidated. During brain development and maturation, neuronal activity is an important regulator of cortical connectivity. Experimental data suggests that disrupted neuronal activity during circuit maturation results in a failure of the refining of the circuit, inducing miswiring and increased network hyperexcitability due to alterations of the postsynaptic compartment (25, 26). In line with this, recent data shows that during juvenile brain development neuronal activity is needed for the proper formation of interhemispheric connections, while inhibition of neuronal activity resulted in decreased neuronal connectivity (27). In outgrowing axons, increased neuronal activity and proper connectivity was suggested to depend on axonal Ca^{2+} -signaling leading to activation of the CaMKK/CaMKI alpha cascade, thereby supporting axonal outgrowth (28). However, although neuronal activity was shown to be also important for subcortical-cortical projections, like the thalamo-cortical fibers (29), recent data suggests that pathological increased cortical excitability during juvenile brain development affects the formation of cortical connections leading to decreased cortical connectivity (30). Thus, neuronal activity during brain maturation has to be balanced: too low or too high neuronal activity levels are associated with cortical miswiring and cortical network hyperexcitability and may lead to psychiatric disorders at further adult ages. Indeed, alteration of cortical excitation/inhibition (E/I) balance and subsequent alteration of cortico-cortical modulation was shown to lead to increased cortical gamma oscillations (31, 32). Moreover, recent human data has shown that schizophrenia patients display increased spontaneous gamma activity during auditory steady-state stimulation reflecting a disrupted E/I balance (33).

A second critical period of cortical restructuring, which is present in distinct and interrelated connectivity development and cortical regions activity shaping, occurs during adolescence (34). Twin studies suggest that these cortical growth trajectories are determined by different sets of genes, which are active in interconnected brain subregions (35). Further studies revealed that regional alterations in the gray matter properties occurred in specific brain networks, which are relevant for the development of psychiatric disorders. Reorganization of these brain networks in adolescence is suggested to result in a particular vulnerability for psychiatric disorders (36, 37). Indeed, analyses in patients with childhood-onset schizophrenia have identified an abnormal pattern of cortical growth in the cingulo-fronto-temporal area of these patients, and suggest a specific impact of genetic systems in these neuroanatomical modules affecting their connectivity (38). Interestingly, network-specific alterations, which increase vulnerability to brain disorders, are not restricted to developmental periods, but have also been found to be present in different neurodegenerative disorders (39). These data suggest that developmental disturbances during adolescence, leading to increased vulnerability to psychiatric and neurodegenerative disorders rely on network-driven alterations of specific brain networks.

The idea that altered neuronal connectivity during brain maturation may lead to psychiatric disorders is supported by longitudinal studies focusing on delayed development



of brain connectivity. Adolescents with childhood-onset of schizophrenia, as well as their clinically unaffected siblings, showed reduced structural integrity and connectivity deficits in the left occipito-temporal areas (40). Although the maturation deficits of cortical connectivity have been shown to become normal with age in patients' siblings (41), this is not the case for the adolescents with childhood-onset schizophrenia (42). These findings support the hypothesis that maturation of cortical connectivity is an important factor for resilience to psychiatric disorders, in which alterations in cortical connectivity at certain life periods, as present in adolescents, may affect resilience toward psychiatric disorders. This is further supported by a recent study which correlated the microarchitecture of corpus callosum to the ability of individuals, who were exposed to high stress, to resist mental disorders (43). Here, young adolescents (mean age: 14.4 ± 1.31 years) with a high resilience to psychiatric disorders displayed higher fractional anisotropy (FA) values in the anterior corpus callosum when compared to adolescents at-risk for mental disorders or with controls (43).

In addition, findings showing altered morphological connectivity following abnormal adolescent brain maturation and associated with cortical hyperexcitability have taken a central role for the current view on the development of psychiatric disorders (24). This is in line with findings that early cortical hyperexcitability has a deleterious effect on brain development leading to sequelae later in life. For instance, Dube and co-workers (44) have shown that stress during sensitive early life periods led to cortical hyperexcitability at later life periods, where 57% of the individuals with early life stress developed epileptic seizures (44). However, and despite the immediate drastic effects like epileptiform discharges (5), continued and sustained cortical hyperexcitability may lead to psychiatric disorders, as described in animal models for autism phenotypes (45), or may contribute to the pathological cascade of events that contribute to the development of Alzheimer's disease (46). Interestingly, not only generalized cortical excitability, but hyperexcitability of specific brain regions was shown to be involved in fear reactions and reduced extinction suggesting an important link between neuronal excitability of specific brain regions like the dentate gyrus in the pathology of post-traumatic stress disorder. In sum, proper maturation from the synaptic level up to the cortical circuit level assuring correct neurotransmission and cortical connectivity is a prerequisite for proper resilient behavior at adult ages, while alterations in E/I balance and in cortico-cortical connectivity may lead to psychiatric disorders.

BRAIN REORGANIZATION AND NETWORK COMPENSATION DURING AGING

From early life through adulthood the brain is constantly changing; later in life the physiological aging process (i.e., free from neurodegeneration) is associated with modification of intrinsic neuronal excitability, together with functional and structural connectivity reorganization. These processes are typically associated with preservation or rather decline in performance across several cognitive domains. In elderly

people, preserved function is thought to be underpinned by compensatory mechanisms (47), in which proximal or distal brain regions to those that decline over time because of natural aging, are recruited to maintain function (48).

From the functional perspective, age-associated adaptations of intrinsic neuronal excitability have been related with changes in cellular micro-architecture (i.e., membrane ion channels, receptors and vesicle fusion processes) and its molecular signaling (49, 50). Although the mechanisms of action are not yet fully understood, numerous evidence has shown involvement of ion-gated channels (e.g., voltage-gated Ca^{2+} channels and mechano-gated K^{+} channels) in mediating the loss of plasticity in neurons, making neurons more susceptible to deleterious processes such as oxidative stress (51). As an example, previous studies have shown increased loss of dopamine synthesis with healthy aging in the striatal system (52), which is particularly vulnerable to oxidative stress (53, 54). At this stage in life, the rate of dopamine loss is more prominent than the regional loss of gray matter tissue (52) and is related to cognitive performance (55), hence suggesting that this transition could be initiated by changes in the ion channels. However, further research is still needed in order to unveil the importance and to better characterize the impact of molecular changes related to the widespread functional and structural brain changes associated with healthy aging.

At the macro level, extra-cephalic electrophysiological recordings have repeatedly evidenced age-related alterations of oscillatory activity across distributed portions of the cortex. Since a loss of approximately 10% of all neocortical neurons over the lifespan occurs (56), alterations of local activity can be explained by an impaired synchronization of neuronal activity in specific frequency ranges. One possible cause of impaired synchronous firing activity can be an alteration of the cortico-cortical neuromodularity mechanisms or imbalances in the subcortico-cortical circuits, directly impacting spontaneous neuronal firing rates and changing the E/I balance. This appears to be the case for high frequency bands, for instance beta (13–30 Hz) and gamma (30–45 Hz) bands, in which increased power has been reported with increasing age (57), whereas decreases in lower alpha (8–10.5 Hz) and a slowing of peak alpha frequency appear with aging (58). Here, age-related interhemispheric asymmetry in power has been related to an increased excitability within the subcortico-cortical circuits (57), which is interpreted as a compensatory mechanism. However, a less consistent scenario emerges for frequencies in the slow wave range, delta (1–4 Hz) and theta (4–7.5 Hz) bands, in which both increases and decreases have been reported (59–62). Notably, apart from increased functional activity (47, 63), reduction of brain lateralization has been described as an important age-related mechanism for compensatory functions (64). However, this view considers only detrimental aspects of aging toward pathology and ignores a causal dynamical relationship with neural adaptation (i.e., plasticity) to life events (e.g., education, intellectual engagement and daily activities) during maturation.

On the structural part, although annual decreases on the order of 0.2–0.5% are well documented (65), MRI-derived morphometric measures (gray matter volume and thickness)

have demonstrated heterogeneous effects of aging, in which despite occurrence of disseminated atrophy across the whole brain, changes vary from region to region and tissue type, particularly over the cortical mantle (65, 66). Of all cortical regions, the frontal and parietal cortices appear the most susceptible to age-related changes (65, 67–70), with accumulating evidence showing increased involvement of the temporal regions (65). At the subcortical level, the hippocampus, caudate nucleus and cerebellum are the most age-susceptible regions (67), whereas involvement of regions belonging to the limbic system appear only limited (71–74). How these changes participate or guide aging processes is still unknown. For instance, reduction of the integrity in the prefrontal cortex is related with functional hyper-activation of the same region during task performance, suggesting the existence of compensatory mechanisms that support the maintenance of cognitive function (1, 75). Anatomical brain lateralization also seems to be nontrivial during aging, since for example, a trend for faster gray matter loss of the left prefrontal cortex relative to the right one has been described (76). Although this finding has not been consistently reported (77), individuals with smaller left than right hemispheric structural integrity are more likely to report cognitive deficits (78). Furthermore, converging evidence exists that an aging-related asymmetric loss of integrity in the parietal and temporal cortices is associated with cognitive functioning (79).

Examination of the white matter tissue has pointed to reduced microstructural integrity in the fiber tracts of the frontal and parietal lobes, as well as in the corpus callosum in elderly persons (80–82). White matter alterations are associated with decrements in cognitive performance, speed of processing, memory and executive functions (82). Moreover, age-related metabolic decreases in the middle and superior temporal cortex, albeit less pronounced than in frontal regions, have been related to white matter disturbances in the long fronto-temporo-occipital association pathways (83). Although these changes are likely to be explained by changes in myelin content (84), due to its influence on signal conduction (85), the effects of age on myelin are complex; even though some reduction in myelin sheaths is observed with age, the myelin is continuously produced throughout life, but perhaps in an uncontrolled or dysfunctional manner (86). Recent advances in MRI methods have successfully ascertained the *in-vivo* assessment of myelin content, showing a negative correlation with aging in the white matter (87, 88) and also to a lesser extent the gray matter (89). Notably, separation of myelin effects from cerebrovascular alterations is not easy using MRI approaches (90), because focal age-related anatomical changes in the white matter, e.g., white matter lesions, can also result from changes in blood pressure (91).

IMPACT OF DIVERGENT PATTERNS OF NETWORK REORGANIZATION FOR RESILIENCE AND SUSTAINED MENTAL HEALTH

Detection of altered brain circuits in the comparison between healthy subjects and patients only partially captures the neurobiological complexity of reorganization processes

associated with maturation and aging and their influence on the development of pathological trajectories (92). The modern, network view of the human brain envisions circuits that are not only shaped by interactions (connections) between their constituent elements (brain regions), but also by their complex topological organization and temporal dynamics. These factors mainly determine the differentiation between physiological vs. pathological processes and hold the key to describing reorganization associated with resilience and sustained mental health (11). Hence, the efficient organization of the brain networks results from delicate balancing of the opposing requirements for information integration and segregation, allowing effective complex cognitive and perceptual functions essential for mental health (93, 94).

Functional connectivity during resting state (i.e., individuals are relaxed and awake, but not engaged in task-directed cognition) has been increasingly applied to investigate the reorganization of the brain across the lifespan, showing that the network topology has a tendency to become randomly organized with increasing age, losing its efficient organization. Such dynamics seem to be good predictors of the individual transition from young to middle age (95), where the information flows from and to the frontal and parietal regions have a primordial role for physiological preservation of mental function.

Interestingly, despite the amount of research showing that chronic exposure to high levels of stress is associated with increased susceptibility to anxiety and other mental disorders (96–98), there is compelling evidence that more graded exposure to stress might reduce such vulnerabilities and promote resilience (99) or may even positively influence several cognitive domains such as memory functions influencing cerebral networks encompassing the hippocampus and amygdala (100). Of note, the stress responses mediated by the amygdala are regulated by the medial prefrontal cortex and their coupling (101–103); thus, stress exposure impairs prefrontal cortex-mediated cognitive functions and switches the control of stress behavior and emotion to interconnected brain circuits (104). These results support the hypothesis of compensation (105), establishing that the recruitment of secondary networks is a mediator of the relationship between structural brain damage and memory or targeted attacks (13). Hence, failure in the compensatory processes related to stress could reduce the ability of the brain circuits to compensate insults, increasing the rate at which functioning is impaired by new challenges (106). Indeed, sustained activation of the circuits involved in coping with stress situations could pass from being an adaptive or compensatory response, to lead to impairments in learning, memory, and the ability to regulate future stress responses (107, 108) and increase the vulnerability to a range of mental disorders over a lifetime (109). On the contrary, forms of early enrichment could induce an accurate heterotypic adjustment at molecular, synaptic and brain circuits levels that could strengthen resilience behavior (99, 110).

In some clinical conditions, such as depression or post-traumatic stress disorder (PTSD), an impaired structural network with abnormal hippocampal integrity and diminished function was described and considered as disease landmarks (111, 112). Smaller hippocampal volumes have been as well attested in

women with major depression and related to experiences of childhood trauma, while depressed individuals without similar trauma events had hippocampal volumes similar to healthy controls (113). Moreover, the unexposed twins of PTSD patients show a similar degree of hippocampal decrease, but without clinical implications (114). Hence, decreased hippocampal volume in these patients cannot be considered a mere disease outcome, but may be a pre-existing risk factor and could be related to early and continued exposure to aversive situations (115).

A further example largely associated with aging is that of Alzheimer's disease (AD), in which a continuum exists comprising a preclinical stage, a symptomatic prodementia stage known as mild cognitive impairment (MCI), and the final stage of dementia (116). The molecular hallmarks of MCI subjects who progress to AD show positive biomarkers of amyloid- β (A β) and tau-related neural injury (117, 118). Periphery biomarkers, such as lower levels of cerebrospinal fluid (CSF) A β , indicate increased accumulation in the brain, whereas increased CSF tau levels indicate damaged neuronal microtubules, clearly evidencing synaptic dysfunction (i.e., desynchronization and hypersynchronization) due to AD (119) that negatively impacts synaptic plasticity and causes synaptic loss, which in turn leads to impairment of neural networks involved in memory and cognition.


Studying network organization patterns in such conditions offers the advantage of addressing developmental trajectories, with causal and longitudinal interpretation and offering the possibility to differentiate primary brain circuits alterations from secondary wide-spread function loss and to highlight

the involvement of particular brain regions as network nodes or connectivity dynamics as active processes (120). In this sense, network compensation as an adaptive mechanism for resilience is not only of use to explain stress coping and to closely track and predict preserved mental health (7), but can be also illustratively studied to monitor adaptation and compensation that occur during the lifespan (69, 121). While age-related decreased connections from and to the frontal regions and increased connections to the posterior (parietal) modules have been described (122) these have been closely related with maintained cognitive function (123, 124). Moreover, variations in network efficiency predict memory function across different life periods and are suggested to reflect differences in information processing between association and sensorimotor systems (125). This all points to the fact that the adjustment of network organization is responsible for buffering alterations in cognitive ability (126, 127), and the loss of optimal topological organization is associated with disease development and impaired brain function (128, 129). Hence, the capacity to react to threats is built into specific brain circuits whose development is influenced by multiple experiences and present differential susceptibility during the lifespan.

BRAIN NETWORKS FINGERPRINTS OF REORGANIZATION DURING LIFESPAN AND RESILIENCE

The dynamics of brain reorganization during lifespan partly mirror structural (130) and functional networks behavior (131, 132), increasing efficiency during maturation through young adulthood until reaching a peak at about 30–40 years of age, that acts as an inflection point, after which the efficiency of brain circuits starts to decrease and physiological aging begins. More broadly, the structural integrity of the brain, as measured by the brain parenchymal fraction (BPF) presents a similar trend across the lifespan (133), whereas the cognitive ability increases throughout young adulthood and decreases in older adulthood (134). Variations in the trajectories are expected in relation to biological and environmental factors, such as genetics, lifestyle, education, socio-cultural background, exercise engagements and learning.

Despite the above mentioned age-related lifespan brain reorganization processes that explain links from structure to cognitive and mental function (1, 2), whether all these changes result from modifications in earlier or later life periods remains an open question. More importantly, which compensation patterns occur at different life points and how brain circuits can compensate for detrimental events is only partly understood (135–137). This is a central point of resilience research, which postulates that active processes of brain reorganization play an essential role for preserved mental function (11), where the connectivity patterns shaping excitability regulation mechanisms are principally involved in sustaining mental health and physiological trajectories of cognitive function (138). Accordingly, we propose an integrative model as shown in **Figure 2**.



To confront the increased endogenous challenges (i.e., changes to neural anatomy and physiology), as well as exogenous challenges (i.e., those brought about by traumatic events or by changes to the environment), brain circuits must present tremendous abilities to flexibly adapt. Maintained mental health can be generally achieved by the integrated communication between frontal and parietal brain regions (3, 4). Within this view, reorganization principles of the brain's topological architecture, beyond being mere compensatory mechanisms (47, 75), could be thought of as manifestations of neural plasticity, synaptic adaptation and reorganization of information flows (69, 139–142). This hypothesis is based on the fact that age-related functional and structural reorganization processes spatially correspond to evolving patterns of activation, and that these structure-function patterns are tightly associated with cognitive performance (1, 143). More supporting evidence shows that adults who do not show age-related adjustment in EEG theta-alpha power are more likely to exhibit cognitive deficits than those who adapt (62). Accordingly, the process of resilience cannot be restricted to early life but should operate throughout the entire lifespan.

Recent studies have emphasized that the integral topological architecture of the brain networks, i.e., modular organization, supports increased cognitive demands (144–146) and that the reorganization of its patterns are tightly related to aging (122, 125, 147). How this process of increased modularization of brain networks is related to mental health or resilience is not yet clear but should be a matter of future studies. Hence, adaptation of interregional connectivity and specific changes in network topology are likely to be contributing mechanisms to healthy aging (147, 148). This phenomenon is characterized by modification within different brain subsystems (122), which contain regions critical for several cognitive functions and/or are particularly sensitive to disease development. Specifically, the prefrontal, anterior and posterior cingulate cortices, as well as the precuneus and the inferior parietal lobe have been consistently shown to present age-related changes in connectivity relevant to preserved mental function and resilience (3, 4).

Common patterns of functional and structural network reorganization have been reported at different time scales (149), mainly involving reduced connectivity in the fronto-parietal or default mode networks which seem to be the normal adaptation to healthy aging. On the contrary, although the temporal cortex shows a similar trajectory with age as other regions (150, 151) the changes to this region show signs of neurodegeneration (152, 153). These findings pinpoint the possibility of specific regional contributions to differentiate healthy from abnormal aging trajectories, and opens the possibility that failures in age-related network reorganization predispose the brain to the development of the so-called disconnection syndromes.

In the same line, specific network topology patterns have already been related to the disease course and clinical progression in AD (154) and Parkinson's disease (16), multiple sclerosis (6, 15), schizophrenia (155, 156), depression (157) and PTSD (158, 159). These observations emphasize that the network adaptations are not merely a consequence of pathological

alterations, but should be seen as integrative processes of functional and structural alterations and compensation for optimal network functioning (123), which act until the set-point at which the network performance cannot be maintained and compensation abates (7). Furthermore, patterns of network reorganization are linked to plasticity in the normal brain (160) and with maintenance of function despite continuous damage (161, 162). This suggests that the correct adaptation of distinct brain circuits may be the key mechanism underlying resilience (as in our proposed model shown in **Figure 2**), where parallel processes of lifespan-related reorganization in brain circuits can be drawn, and used to improve our understanding, under an holistic framework, of the interrelation between physiological and pathological developments and experienced life events (7).

CONCLUSION

Clearly, brain networks development and reorganization across the human lifespan are active and continuous processes, which allow the emergence of resilience mechanisms across the entire lifespan. Within these networks, the frontal and parietal regions are certainly modulators of maintained health, where the intrinsic dynamics and excitability patterns of these regions and their connections, for instance to the temporal lobe and hippocampus, are able to reduce vulnerability and risk for damage. Hence, correct adaptation of these connectivity patterns could be the key to healthy aging and diminish the risk for developing neurological and neuropsychiatric disease at different stages across the lifespan. Future research should aim for multivariate full-brain investigations using not only larger study populations, also with the help of combined biomarkers at the micro-level, such as blood markers or genetic profiling, and at macro-level, e.g., brain imaging and network analyses; all together with longitudinal designs in order to better capture the full dynamics of lifespan modifications.

AUTHOR CONTRIBUTIONS

GG-E, MM, JV, and SG made substantial contributions to the conception and design of the work; performed the literature search, drafted parts of the manuscript and revised it critically. GG-E, and MM prepared the first version and revised subsequent versions of the manuscript. JV and SG made substantial contributions to the interpretation of data. VC made substantial contributions to the interpretation of data; he made the figures and revised the manuscript critically. GG-E, MM, VC, JV, and SG give their final approval of the version to be published and agree to be responsible for all aspects of the work so that questions about the accuracy or integrity of any part of the work are adequately investigated and solved.

ACKNOWLEDGMENTS

The authors would like to thank Cheryl Ernest for proofreading the manuscript. This work was supported by a grant from the German Research Council (DFG; CRC-1193).

REFERENCES

- Grady C. The cognitive neuroscience of ageing. *Nat Rev Neurosci.* (2012) 13:491–505. doi: 10.1038/nrn3256
- Song J, Birn RM, Boly M, Meier TB, Nair VA, Meyerand ME, et al. Age-related reorganizational changes in modularity and functional connectivity of human brain networks. *Brain Connect.* (2014) 4:662–76. doi: 10.1089/brain.2014.0286
- Andrews-Hanna JR, Snyder AZ, Vincent JL, Lustig C, Head D, Raichle ME, et al. Disruption of large-scale brain systems in advanced aging. *Neuron* (2007) 56:924–35. doi: 10.1016/j.neuron.2007.10.038
- Sambataro F, Murty VP, Callicott JH, Tan HY, Das S, Weinberger DR, et al. Age-related alterations in default mode network: impact on working memory performance. *Neurobiol Aging* (2010) 31:839–52. doi: 10.1016/j.neurobiolaging.2008.05.022
- Groppa S, Siebner HR, Kurth C, Stephani U, Siniatchkin M. Abnormal response of motor cortex to photic stimulation in idiopathic generalized epilepsy. *Epilepsia* (2008) 49:2022–9. doi: 10.1111/j.1528-1167.2008.01709.x
- Fleischer V, Groger A, Koirala N, Drobny A, Muthuraman M, Kolber P, et al. Increased structural white and grey matter network connectivity compensates for functional decline in early multiple sclerosis. *Mult Scler.* (2017) 23:432–41. doi: 10.1177/1352458516651503
- Fleischer V, Radetz A, Ciolac D, Muthuraman M, Gonzalez-Escamilla G, Zipp F, et al. Graph theoretical framework of brain networks in multiple sclerosis: a review of concepts. *Neuroscience* (2017) S0306-4522(17)30761-3. doi: 10.1016/j.neuroscience.2017.10.033. [Epub ahead of print].
- Van Den Heuvel MP, Sporns O. Network hubs in the human brain. *Trends Cogn Sci.* (2013) 17:683–96. doi: 10.1016/j.tics.2013.09.012
- Stam CJ. Modern network science of neurological disorders. *Nat Rev Neurosci.* (2014) 15:683–95. doi: 10.1038/nrn3801
- Medaglia JD, Pasqualetti F, Hamilton RH, Thompson-Schill SL, Bassett DS. Brain and cognitive reserve: translation via network control theory. *Neurosci Biobehav Rev.* (2017) 75:53–64. doi: 10.1016/j.neubiorev.2017.01.016
- Kalisch R, Baker DG, Basten U, Boks MP, Bonanno GA, Brummelman E, et al. The resilience framework as a strategy to combat stress-related disorders. *Nat Hum Behav.* (2017) 1:784–90. doi: 10.1038/s41562-017-0200-8
- Chirumamilla VC, Fleischer V, Drobny A, Anjum T, Muthuraman M, Zipp F, et al. Functional connectivity analysis using whole brain and regional network metrics in MS patients. *Conf Proc IEEE Eng Med Biol Soc.* (2016) 2016:4039–42. doi: 10.1109/EMBC.2016.7591613
- Drobny A, Yuen KS, Muthuraman M, Reitz SC, Fleischer V, Klein J, et al. Changes in brain functional connectivity patterns are driven by an individual lesion in MS: a resting-state fMRI study. *Brain Imaging Behav.* (2016) 10:1117–26. doi: 10.1007/s11682-015-9476-3
- He Y, Chen ZJ, Evans AC. Small-world anatomical networks in the human brain revealed by cortical thickness from MRI. *Cereb Cortex* (2007) 17:2407–19. doi: 10.1093/cercor/bhl149
- Muthuraman M, Fleischer V, Kolber P, Luessi F, Zipp F, Groppa S. Structural Brain Network Characteristics Can Differentiate CIS from Early RRMS. *Front Neurosci.* (2016) 10:14. doi: 10.3389/fnins.2016.00014
- Koirala N, Fleischer V, Glaser M, Zeuner KE, Deuschl G, Volkmann J, et al. Frontal lobe connectivity and network community characteristics are associated with the outcome of subthalamic nucleus deep brain stimulation in patients with parkinson's disease. *Brain Topogr.* (2018) 31:311–21. doi: 10.1007/s10548-017-0597-4
- Groppa S, Herzog J, Falk D, Riedel C, Deuschl G, Volkmann J. Physiological and anatomical decomposition of subthalamic neurostimulation effects in essential tremor. *Brain* (2014) 137:109–21. doi: 10.1093/brain/awt304
- Deppe M, Tabelow K, Kramer J, Tenberge JG, Schifferl P, Bittner S, et al. Evidence for early, non-lesional cerebellar damage in patients with multiple sclerosis: DTI measures correlate with disability, atrophy, and disease duration. *Mult Scler.* (2016) 22:73–84. doi: 10.1177/1352458515579439
- Muthuraman M, Raethjen J, Koirala N, Anwar A, Mideksa K, Elble R, et al. Cerebello-cortical network fingerprints differ among essential, Parkinson and mimicked tremors. *Brain* (2018) 141:1770–81. doi: 10.1093/brain/awy098
- Turrigiano G. Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu Rev Neurosci* (2011) 34:89–103. doi: 10.1146/annurev-neuro-060909-153238
- Harrison PJ, Weinberger DR. Schizophrenia genes, gene expression, and neuropathology: on the matter of their convergence. *Mol psychiatry* (2005) 10:40. doi: 10.1038/sj.mp.4001630
- Coyle JT. Glutamate and schizophrenia: beyond the dopamine hypothesis. *Cell Mol Neurobiol.* (2006) 26:365–84. doi: 10.1007/s10571-006-9062-8
- Belforte JE, Zsiros V, Sklar ER, Jiang Z, Yu G, Li Y, et al. Postnatal NMDA receptor ablation in corticolimbic interneurons confers schizophrenia-like phenotypes. *Nat Neurosci.* (2010) 13:76–83. doi: 10.1038/nn.2447
- Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, O'shea DJ, et al. Neocortical excitation/inhibition balance in information processing and social dysfunction. *Nature* (2011) 477:171–8. doi: 10.1038/nature10360
- Galvan CD, Hrachovy RA, Smith KL, Swann JW. Blockade of neuronal activity during hippocampal development produces a chronic focal epilepsy in the rat. *J Neurosci.* (2000) 20:2904–16. doi: 10.1523/JNEUROSCI.20-08-02904.2000
- Galvan CD, Wenzel JH, Dineley KT, Lam TT, Schwartzkroin PA, Sweatt JD, et al. Postsynaptic contributions to hippocampal network hyperexcitability induced by chronic activity blockade *in vivo*. *Eur J Neurosci.* (2003) 18:1861–72. doi: 10.1046/j.1460-9568.2003.02920.x
- Mizuno H, Hirano T, Tagawa Y. Pre-synaptic and post-synaptic neuronal activity supports the axon development of callosal projection neurons during different post-natal periods in the mouse cerebral cortex. *Eur J Neurosci.* (2010) 31:410–24. doi: 10.1111/j.1460-9568.2009.07070.x
- Ageta-Ishihara N, Takemoto-Kimura S, Nonaka M, Adachi-Morishima A, Suzuki K, Kamijo S, et al. Control of cortical axon elongation by a GABA-driven Ca2+/calmodulin-dependent protein kinase cascade. *J Neurosci.* (2009) 29:13720–9. doi: 10.1523/JNEUROSCI.3018-09.2009
- Mire E, Mezzerà C, Leyva-Diaz E, Paternain AV, Squarzonni P, Bluy L, et al. Spontaneous activity regulates Robo1 transcription to mediate a switch in thalamocortical axon growth. *Nat Neurosci.* (2012) 15:1134–43. doi: 10.1038/nn.3160
- Vogt J, Kirischuk S, Unichenko P, Schluter L, Pelosi A, Endle H, et al. Synaptic phospholipid signaling modulates axon outgrowth via glutamate-dependent Ca2+-mediated molecular pathways. *Cereb Cortex* (2017) 27:131–45. doi: 10.1093/cercor/bhw370
- Pinault D. N-methyl d-aspartate receptor antagonists ketamine and MK-801 induce wake-related aberrant gamma oscillations in the rat neocortex. *Biol Psychiatry* (2008) 63:730–5. doi: 10.1016/j.biopsych.2007.10.006
- Thalman C, Horta G, Qiao L, Endle H, Tegeder I, Cheng H, et al. Synaptic Phospholipids as a new target for cortical hyperexcitability and E/I-balance in psychiatric disorders. *Mol psychiatry* (In Press) 23:1699–710. doi: 10.1038/s41380-018-0053-1
- Hirano Y, Oribe N, Kanba S, Onitsuka T, Nestor PG, Spencer KM. Spontaneous gamma activity in schizophrenia. *JAMA Psychiatry* (2015) 72:813–21. doi: 10.1001/jamapsychiatry.2014.2642
- Brain Development Cooperative G. Total and regional brain volumes in a population-based normative sample from 4 to 18 years: the NIH MRI Study of Normal Brain Development. *Cereb Cortex* (2012) 22:1–12. doi: 10.1093/cercor/bhr018
- Rimol LM, Panizzon MS, Fennema-Notestine C, Eyler LT, Fischl B, Franz CE, et al. Cortical thickness is influenced by regionally specific genetic factors. *Biol Psychiatry* (2010) 67:493–9. doi: 10.1016/j.biopsych.2009.09.032
- Paus T, Keshavan M, Giedd JN. Why do many psychiatric disorders emerge during adolescence? *Nat Rev Neurosci.* (2008) 9:947–57. doi: 10.1038/nrn2513
- Paus T, Toro R, Leonard G, Lerner JV, Lerner RM, Perron M, et al. Morphological properties of the action-observation cortical network in adolescents with low and high resistance to peer influence. *Soc Neurosci.* (2008) 3:303–16. doi: 10.1080/17470910701563558
- Alexander-Bloch AF, Reiss PT, Rapoport J, McAdams H, Giedd JN, Bullmore ET, et al. Abnormal cortical growth in schizophrenia targets normative modules of synchronized development. *Biol Psychiatry* (2014) 76:438–46. doi: 10.1016/j.biopsych.2014.02.010

39. Seeley WW, Crawford RK, Zhou J, Miller BL, Greicius MD. Neurodegenerative diseases target large-scale human brain networks. *Neuron* (2009) 62:42–52. doi: 10.1016/j.neuron.2009.03.024
40. Zalesky A, Pantelis C, Cropley V, Fornito A, Cocchi L, McAdams H, et al. Delayed development of brain connectivity in adolescents with schizophrenia and their unaffected siblings. *JAMA psychiatry* (2015) 72:900–8. doi: 10.1001/jamapsychiatry.2015.0226
41. Gogtay N, Greenstein D, Lenane M, Clasen L, Sharp W, Gochman P, et al. Cortical brain development in nonpsychotic siblings of patients with childhood-onset schizophrenia. *Arch Gen Psychiatry* (2007) 64:772–80. doi: 10.1001/archpsyc.64.7.772
42. Greenstein D, Lerch J, Shaw P, Clasen L, Giedd J, Gochman P, et al. Childhood onset schizophrenia: cortical brain abnormalities as young adults. *J Child Psychol Psychiatry* (2006) 47:1003–12. doi: 10.1111/j.1469-7610.2006.01658.x
43. Galinowski A, Miranda R, Lemaitre H, Paillere Martinot ML, Artiges E, Vulser H, et al. Resilience and corpus callosum microstructure in adolescence. *Psychol Med.* (2015) 45:2285–94. doi: 10.1017/S0033291715000239
44. Dube CM, Molet J, Singh-Taylor A, Ivy A, Maras PM, Baram TZ. Hyperexcitability and epilepsy generated by chronic early-life stress. *Neurobiol Stress* (2015) 2:10–9. doi: 10.1016/j.yjnstr.2015.03.001
45. Selimbeyoglu A, Kim CK, Inoue M, Lee SY, Hong ASO, Kauvar I, et al. Modulation of prefrontal cortex excitation/inhibition balance rescues social behavior in CNTNAP2-deficient mice. *Sci Transl Med.* (2017) 9:eah6733. doi: 10.1126/scitranslmed.aah6733
46. Lam AD, Deck G, Goldman A, Eskandar EN, Noebels J, Cole AJ. Silent hippocampal seizures and spikes identified by foramen ovale electrodes in Alzheimer's disease. *Nat Med.* (2017) 23:678–80. doi: 10.1038/nm.4330
47. Cabeza R, Anderson ND, Locantore JK, McIntosh AR. Aging gracefully: compensatory brain activity in high-performing older adults. *Neuroimage* (2002) 17:1394–402. doi: 10.1006/nimg.2002.1280
48. Meunier D, Stamatakis EA, Tyler LK. Age-related functional reorganization, structural changes, and preserved cognition. *Neurobiol Aging* (2014) 35:42–54. doi: 10.1016/j.neurobiolaging.2013.07.003
49. Oh MM, Oliveira FA, Disterhoft JF. Learning and aging related changes in intrinsic neuronal excitability. *Front Aging Neurosci.* (2010) 2:2. doi: 10.3389/neuro.24.002.2010
50. Yeoman M, Scutt G, Faragher R. Insights into CNS ageing from animal models of senescence. *Nat Rev Neurosci.* (2012) 13:435–45. doi: 10.1038/nrn3230
51. Annunziato L, Pannaccione A, Cataldi M, Secondo A, Castaldo P, Di Renzo G, et al. Modulation of ion channels by reactive oxygen and nitrogen species: a pathophysiological role in brain aging? *Neurobiol Aging* (2002) 23:819–34. doi: 10.1016/S0197-4580(02)00069-6
52. Ota M, Yasuno F, Ito H, Seki C, Nozaki S, Asada T, et al. Age-related decline of dopamine synthesis in the living human brain measured by positron emission tomography with L-[beta-11C]DOPA. *Life Sci.* (2006) 79:730–6. doi: 10.1016/j.lfs.2006.02.017
53. Cardoso HD, Passos PP, Lagranha CJ, Ferraz AC, Santos Junior EF, Oliveira RS, et al. Differential vulnerability of substantia nigra and corpus striatum to oxidative insult induced by reduced dietary levels of essential fatty acids. *Front Hum Neurosci.* (2012) 6:249. doi: 10.3389/fnhum.2012.00249
54. Jensen N, Oliveira JR. Basal ganglia vulnerability to oxidative stress. *Front Neurosci.* (2014) 8:80. doi: 10.3389/fnins.2014.00080
55. Backman L, Lindenberger U, Li SC, Nyberg L. Linking cognitive aging to alterations in dopamine neurotransmitter functioning: recent data and future avenues. *Neurosci Biobehav Rev.* (2010) 34:670–7. doi: 10.1016/j.neubiorev.2009.12.008
56. Pakkenberg B, Gundersen HJ. Neocortical neuron number in humans: effect of sex and age. *J Comp Neurol.* (1997) 384:312–20.
57. Zappasodi F, Marzetti L, Olejarczyk E, Tecchio F, Pizzella V. Age-related changes in electroencephalographic signal complexity. *PLoS ONE* (2015) 10:e0141995. doi: 10.1371/journal.pone.0141995
58. Chiang AK, Rennie CJ, Robinson PA, Van Albada SJ, Kerr CC. Age trends and sex differences of alpha rhythms including split alpha peaks. *Clin Neurophysiol.* (2011) 122:1505–17. doi: 10.1016/j.clinph.2011.01.040
59. Finnigan S, Robertson IH. Resting EEG theta power correlates with cognitive performance in healthy older adults. *Psychophysiology* (2011) 48:1083–7. doi: 10.1111/j.1469-8986.2010.01173.x
60. Michels L, Muthuraman M, Luchinger R, Martin E, Anwar AR, Raethjen J, et al. Developmental changes of functional and directed resting-state connectivities associated with neuronal oscillations in EEG. *Neuroimage* (2013) 81:231–42. doi: 10.1016/j.neuroimage.2013.04.030
61. Vlahou EL, Thurm F, Kolassa IT, Schlee W. Resting-state slow wave power, healthy aging and cognitive performance. *Sci Rep.* (2014) 4:5101. doi: 10.1038/srep05101
62. Trammell JR, Macrae PG, Davis G, Bergstedt D, Anderson AE. The relationship of cognitive performance and the theta-alpha power ratio is age-dependent: an EEG study of short term memory and reasoning during task and resting-state in healthy young and old adults. *Front aging Neurosci.* (2017) 9:364. doi: 10.3389/fnagi.2017.00364
63. Berlinger M, Danelli L, Bottini G, Sberna M, Paulesu E. Reassessing the HAROLD model: is the hemispheric asymmetry reduction in older adults a special case of compensatory-related utilisation of neural circuits? *Exp Brain Res.* (2013) 224:393–410. doi: 10.1007/s00221-012-3319-x
64. Cabeza R. Hemispheric asymmetry reduction in older adults: the HAROLD model. *Psychol Aging* (2002) 17:85–100. doi: 10.1037/0882-7974.17.1.85
65. Fjell AM, Mcevoy L, Holland D, Dale AM, Walhovd KB, Initiative ASDN. What is normal in normal aging? Effects of aging, amyloid and Alzheimer's disease on the cerebral cortex and the hippocampus *Prog Neurobiol.* (2014) 117:20–40. doi: 10.1016/j.pneurobio.2014.02.004
66. Tamnes CK, Walhovd KB, Dale AM, Ostby Y, Grydeland H, Richardson G, et al. Brain development and aging: overlapping and unique patterns of change. *Neuroimage* (2013) 68:63–74. doi: 10.1016/j.neuroimage.2012.11.039
67. Raz N, Lindenberger U, Rodrigue KM, Kennedy KM, Head D, Williamson A, et al. Regional brain changes in aging healthy adults: general trends, individual differences and modifiers. *Cereb Cortex* (2005) 15:1676–89. doi: 10.1093/cercor/bhi044
68. Dennis NA, Cabeza R. Neuroimaging of healthy cognitive aging. In: Craik FIM, Salthouse TA, editors. *The Handbook of Aging and Cognition*. 3rd ed. New York, NY: Psychology Press (2008). p. 1–54.
69. Park DC, Reuter-Lorenz P. The adaptive brain: aging and neurocognitive scaffolding. *Annu Rev Psychol.* (2009) 60:173–96. doi: 10.1146/annurev.psych.59.103006.093656
70. Lockhart SN, Decarli C. Structural imaging measures of brain aging. *Neuropsychol Rev.* (2014) 24:271–89. doi: 10.1007/s11065-014-9268-3
71. Good CD, Johnsrude I, Ashburner J, Henson RN, Friston KJ, Frackowiak RS. Cerebral asymmetry and the effects of sex and handedness on brain structure: a voxel-based morphometric analysis of 465 normal adult human brains. *Neuroimage* (2001) 14:685–700. doi: 10.1006/nimg.2001.0857
72. Grieve SM, Clark CR, Williams LM, Peduto AJ, Gordon E. Preservation of limbic and paralimbic structures in aging. *Hum Brain Mapp.* (2005) 25:391–401. doi: 10.1002/hbm.20115
73. Long X, Zhang L, Liao W, Jiang C, Qiu B. Distinct laterality alterations distinguish mild cognitive impairment and Alzheimer's disease from healthy aging: statistical parametric mapping with high resolution MRI. *Human Brain Mapp.* (2013) 34:3400–10. doi: 10.1002/hbm.22157
74. Minkova L, Habich A, Peter J, Kaller CP, Eickhoff SB, Klöppel S. Gray matter asymmetries in aging and neurodegeneration: a review and meta-analysis. *Hum Brain Mapp.* (2017) 38:5890–904. doi: 10.1002/hbm.23772
75. Di X, Rypma B, Biswal BB. Correspondence of executive function related functional and anatomical alterations in aging brain. *Prog Neuropsychopharmacol Biol Psychiatry* (2014) 48:41–50. doi: 10.1016/j.pnpbp.2013.09.001
76. Sowell ER, Peterson BS, Thompson PM, Welcome SE, Henkenius AL, Toga AW. Mapping cortical change across the human life span. *Nat Neurosci.* (2003) 6:309–15. doi: 10.1038/nn1008
77. Raz N, Gunning FM, Head D, Dupuis JH, McQuain J, Briggs SD, et al. Selective aging of the human cerebral cortex observed *in vivo*: differential vulnerability of the prefrontal gray matter. *Cereb Cortex* (1997) 7:268–82. doi: 10.1093/cercor/7.3.268

78. Cherbuin N, Réglade-Meslin C, Kumar R, Sachdev P, Anstey KJ. Mild cognitive disorders are associated with different patterns of brain asymmetry than normal aging: the PATH through Life Study. *Front psychiatry* (2010) 1:11. doi: 10.3389/fpsy.2010.00011
79. Plessen KJ, Hugdahl K, Bansal R, Hao X, Peterson BS. Sex, age, and cognitive correlates of asymmetries in thickness of the cortical mantle across the life span. *J Neurosci* (2014) 34:6294–302. doi: 10.1523/JNEUROSCI.3692-13.2014
80. Nusbaum AO, Tang CY, Buchsbaum MS, Wei TC, Atlas SW. Regional and global changes in cerebral diffusion with normal aging. *AJNR Am J Neuroradiol.* (2001) 22:136–42. Available online at: <http://www.ajnr.org/content/22/1/136/tab-article-info>
81. O'sullivan M, Jones DK, Summers P, Morris R, Williams S, Markus H. Evidence for cortical “disconnection” as a mechanism of age-related cognitive decline. *Neurology* (2001) 57:632–38. doi: 10.1212/WNL.57.4.632
82. Salthouse TA. Neuroanatomical substrates of age-related cognitive decline. *Psychol Bull.* (2011) 137:753–84. doi: 10.1037/a0023262
83. Chételat G, Landeau B, Salmon E, Yakushev I, Bahri MA, Mézenge F, et al. Relationships between brain metabolism decrease in normal aging and changes in structural and functional connectivity. *Neuroimage* (2013) 76:167–77. doi: 10.1016/j.neuroimage.2013.03.009
84. Steiger TK, Weiskopf N, Bunzeck N. Iron level and myelin content in the ventral striatum predict memory performance in the aging brain. *J Neurosci.* (2016) 36:3552–8. doi: 10.1523/JNEUROSCI.3617-15.2016
85. Pajevic S, Basser PJ, Fields RD. Role of myelin plasticity in oscillations and synchrony of neuronal activity. *Neuroscience* (2014) 276:135–47. doi: 10.1016/j.neuroscience.2013.11.007
86. Peters A. The effects of normal aging on myelin and nerve fibers: a review. *J Neurocytol.* (2002) 31:581–93. doi: 10.1023/A:1025731309829
87. Callaghan MF, Freund P, Draganski B, Anderson E, Cappelletti M, Chowdhury R, et al. Widespread age-related differences in the human brain microstructure revealed by quantitative magnetic resonance imaging. *Neurobiol Aging* (2014) 35:1862–72. doi: 10.1016/j.neurobiolaging.2014.02.008
88. Berman S, West KL, Does MD, Yeatman JD, Mezer AA. Evaluating g-ratio weighted changes in the corpus callosum as a function of age and sex. *NeuroImage* (2017) 182:304–13. doi: 10.1016/j.neuroimage.2017.06.076
89. Keuken MC, Bazin PL, Backhouse K, Beekhuizen S, Himmer L, Kandola A, et al. Effects of aging on T(1), T(2)*, and QSM MRI values in the subcortex. *Brain Struct Funct.* (2017) 222:2487–505. doi: 10.1007/s00429-016-1352-4
90. Knight MJ, Mccann B, Tsivos D, Dillon S, Coulthard E, Kauppinen RA. Quantitative T2 mapping of white matter: applications for ageing and cognitive decline. *Phys Med Biol.* (2016) 61:5587–605. doi: 10.1088/0031-9155/61/15/5587
91. Longstreth WJ Jr, Manolio TA, Arnold A, Burke GL, Bryan N, Jungreis CA, et al. Clinical correlates of white matter findings on cranial magnetic resonance imaging of 3301 elderly people. *Stroke* (1996) 27:1274–82. doi: 10.1161/01.STR.27.8.1274
92. Rodrigue KM, Kennedy KM. The cognitive consequences of structural changes to the aging brain In: Schaie KW, Willis SL, editors. *Handbook of the Psychology of Aging*. 7th ed. New York, NY: Elsevier (2011). p. 73–91.
93. Sporns O, Tononi G, Edelman GM. Connectivity and complexity: the relationship between neuroanatomy and brain dynamics. *Neural Netw.* (2000) 13:909–22. doi: 10.1016/S0893-6080(00)00053-8
94. Friston KJ. Functional and effective connectivity: a review. *Brain Connect.* (2011) 1:13–36. doi: 10.1089/brain.2011.0008
95. Petti M, Toppi J, Babiloni F, Cincotti F, Mattia D, Astolfi L. EEG resting-state brain topological reorganization as a function of age. *Comput Intell Neurosci.* (2016) 2016:1–10. doi: 10.1155/2016/6243694
96. Krishnan V, Han MH, Graham DL, Berton O, Renthal W, Russo SJ, et al. Molecular adaptations underlying susceptibility and resistance to social defeat in brain reward regions. *Cell* (2007) 131:391–404. doi: 10.1016/j.cell.2007.09.018
97. Bradley RG, Binder EB, Epstein MP, Tang Y, Nair HP, Liu W, et al. Influence of child abuse on adult depression: moderation by the corticotropin-releasing hormone receptor gene. *Arch Gen Psychiatry* (2008) 65:190–200. doi: 10.1001/archgenpsychiatry.2007.26
98. Heim C, Newport DJ, Mletzko T, Miller AH, Nemeroff CB. The link between childhood trauma and depression: insights from HPA axis studies in humans. *Psychoneuroendocrinology* (2008) 33:693–710. doi: 10.1016/j.psyneuen.2008.03.008
99. Russo SJ, Murrrough JW, Han MH, Charney DS, Nestler EJ. Neurobiology of resilience. *Nat Neurosci* (2012) 15:1475–84. doi: 10.1038/nn.3234
100. McEwen BS. Protection and damage from acute and chronic stress: allostasis and allostatic overload and relevance to the pathophysiology of psychiatric disorders. *Ann N Y Acad Sci.* (2004) 1032:1–7. doi: 10.1196/annals.1314.001
101. Quirk GJ, Mueller D. Neural mechanisms of extinction learning and retrieval. *Neuropsychopharmacology* (2008) 33:56–72. doi: 10.1038/sj.npp.1301555
102. Kim MJ, Whalen PJ. The structural integrity of an amygdala-prefrontal pathway predicts trait anxiety. *J Neurosci.* (2009) 29:11614–8. doi: 10.1523/JNEUROSCI.2335-09.2009
103. Vytal KE, Overstreet C, Charney DR, Robinson OJ, Grillon C. Sustained anxiety increases amygdala-dorsomedial prefrontal coupling: a mechanism for maintaining an anxious state in healthy adults. *J Psychiatry Neurosci.* (2014) 39:321–9. doi: 10.1503/jpn.130145
104. Arnsten AF. Stress signalling pathways that impair prefrontal cortex structure and function. *Nature Reviews Neuroscience* (2009) 10:410. doi: 10.1038/nrn2648
105. Cabeza R, Dennis NA. Frontal lobes and aging: Deterioration and Compensation In: *Principles of frontal lobe function*. In: Stuss DT, Knight RT editors. (New York, NY: Oxford University Press) (2013). p. 628–52.
106. McEwen BS, Gianaros PJ. Stress- and allostasis-induced brain plasticity. *Annu Rev Med.* (2011) 62:431–45. doi: 10.1146/annurev-med-052209-100430
107. Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci.* (2009) 10:434–45. doi: 10.1038/nrn2639
108. Gotlib IH, Joormann J. Cognition and depression: current status and future directions. *Annu Rev Clin Psychol.* (2010) 6:285–312. doi: 10.1146/annurev.clinpsy.121208.131305
109. Sapolsky RM, Romero LM, Munck AU. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev.* (2000) 21:55–89. doi: 10.1210/er.21.1.55
110. McEwen BS. In pursuit of resilience: stress, epigenetics, and brain plasticity. *Ann N Y Acad Sci.* (2016) 1373:56–64. doi: 10.1111/nyas.13020
111. Videbech P, Ravnkilde B. Hippocampal volume and depression: a meta-analysis of MRI studies. *Am J Psychiatry* (2004) 161:1957–66. doi: 10.1176/appi.ajp.161.11.1957
112. Smith ME. Bilateral hippocampal volume reduction in adults with post-traumatic stress disorder: a meta-analysis of structural MRI studies. *Hippocampus* (2005) 15:798–807. doi: 10.1002/hipo.20102
113. Vythilingam M, Heim C, Newport J, Miller AH, Anderson E, Bronen R, et al. Childhood trauma associated with smaller hippocampal volume in women with major depression. *Am J Psychiatry* (2002) 159:2072–80. doi: 10.1176/appi.ajp.159.12.2072
114. Gilbertson MW, Shenton ME, Ciszewski A, Kasai K, Lasko NB, Orr SP, et al. Smaller hippocampal volume predicts pathologic vulnerability to psychological trauma. *Nat Neurosci.* (2002) 5:1242–7. doi: 10.1038/nn958
115. Charney DS, Manji HK. Life stress, genes, and depression: multiple pathways lead to increased risk and new opportunities for intervention. *Sci STKE.* (2004) 2004:re5. doi: 10.1126/stke.2252004re5
116. Sperling RA, Aisen PS, Beckett LA, Bennett DA, Craft S, Fagan AM, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement.* (2011) 7:280–92. doi: 10.1016/j.jalz.2011.03.003
117. Braak H, Braak E. Evolution of the neuropathology of Alzheimer's disease. *Acta Neurol Scand Suppl.* (1996) 165:3–12. doi: 10.1111/j.1600-0404.1996.tb05866.x
118. Albert MS, Dekosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Focus* (2013) 11:96–106. doi: 10.1176/appi.focus.11.1.96

119. Vlassenko AG, Benzinger TL, Morris JC. PET amyloid-beta imaging in preclinical Alzheimer's disease. *Biochim Biophys Acta* (2012) 1822:370–9. doi: 10.1016/j.bbadis.2011.11.005
120. Kashan N, Alon U. Spontaneous evolution of modularity and network motifs. *Proc Natl Acad Sci USA*. (2005) 102:13773–8. doi: 10.1073/pnas.0503610102
121. Gutchess AH, Welsh RC, Hedden T, Bangert A, Minear M, Liu LL, et al. Aging and the neural correlates of successful picture encoding: frontal activations compensate for decreased medial-temporal activity. *J Cogn Neurosci*. (2005) 17:84–96. doi: 10.1162/0898929052880048
122. Meunier D, Achard S, Morcom A, Bullmore E. Age-related changes in modular organization of human brain functional networks. *Neuroimage* (2009) 44:715–23. doi: 10.1016/j.neuroimage.2008.09.062
123. Franzmeier N, Gottler J, Grimmer T, Drzezga A, Araque-Caballero MA, Simon-Vermet L, et al. Resting-state connectivity of the left frontal cortex to the default mode and dorsal attention network supports reserve in mild cognitive impairment. *Front Aging Neurosci*. (2017) 9:264. doi: 10.3389/fnagi.2017.00264
124. Franzmeier N, Hartmann JC, Taylor ANW, Araque Caballero MA, Simon-Vermet L, Buerger K, et al. Left frontal hub connectivity during memory performance supports reserve in aging and mild cognitive impairment. *J Alzheimers Dis*. (2017) 59:1381–92. doi: 10.3233/JAD-170360
125. Chan MY, Park DC, Savalia NK, Petersen SE, Wig GS. Decreased segregation of brain systems across the healthy adult lifespan. *Proc Natl Acad Sci USA*. (2014) 111:E4997–5006. doi: 10.1073/pnas.1415122111
126. Zhao T, Cao M, Niu H, Zuo XN, Evans A, He Y, et al. Age-related changes in the topological organization of the white matter structural connectome across the human lifespan. *Hum Brain Mapp*. (2015) 36:3777–92. doi: 10.1002/hbm.22877
127. Gallen CL, Turner GR, Adnan A, D'Esposito M. Reconfiguration of brain network architecture to support executive control in aging. *Neurobiol Aging* (2016) 44:42–52. doi: 10.1016/j.neurobiolaging.2016.04.003
128. Pfefferbaum A, Adalsteinsson E, Sullivan EV. Frontal circuitry degradation marks healthy adult aging: evidence from diffusion tensor imaging. *Neuroimage* (2005) 26:891–9. doi: 10.1016/j.neuroimage.2005.02.034
129. Salat D, Tuch D, Greve D, Van Der Kouwe A, Hevelone N, Zaleta A, et al. Age-related alterations in white matter microstructure measured by diffusion tensor imaging. *Neurobiol Aging* (2005) 26:1215–27. doi: 10.1016/j.neurobiolaging.2004.09.017
130. Wu K, Taki Y, Sato K, Kinomura S, Goto R, Okada K, et al. Age-related changes in topological organization of structural brain networks in healthy individuals. *Hum Brain Mapp*. (2012) 33:552–68. doi: 10.1002/hbm.21232
131. Betzel RF, Byrge L, He Y, Goñi J, Zuo X-N, Sporns O. Changes in structural and functional connectivity among resting-state networks across the human lifespan. *Neuroimage* (2014) 102:345–57. doi: 10.1016/j.neuroimage.2014.07.067
132. Cao M, Huang H, Peng Y, Dong Q, He Y. Toward developmental connectomics of the human brain. *Front Neuroanat*. (2016) 10:25. doi: 10.3389/fnana.2016.00025
133. Vagberg M, Granasen G, Svenningsson A. Brain parenchymal fraction in healthy adults—a systematic review of the literature. *PLoS ONE* (2017) 12:e0170018. doi: 10.1371/journal.pone.0170018
134. Cai L, Chan JS, Yan JH, Peng K. Brain plasticity and motor practice in cognitive aging. *Front Aging Neurosci*. (2014) 6:31. doi: 10.3389/fnagi.2014.00031
135. Hamarat E, Thompson D, Steele D, Matheny K, Simons C. Age differences in coping resources and satisfaction with life among middle-aged, young-old, and oldest-old adults. *J Genet Psychol*. (2002) 163:360–7. doi: 10.1080/00221320209598689
136. Bowling A, Iliffe S. Psychological approach to successful ageing predicts future quality of life in older adults. *Health Qual Life Outcomes* (2011) 9:13. doi: 10.1186/1477-7525-9-13
137. Jeste DV, Savla GN, Thompson WK, Vahia IV, Glorioso DK, Martin AVS, et al. Association between older age and more successful aging: critical role of resilience and depression. *Am J Psychiatry* (2013) 170:188–96. doi: 10.1176/appi.ajp.2012.12030386
138. Buckner RL. Memory and executive function in aging and AD: multiple factors that cause decline and reserve factors that compensate. *Neuron* (2004) 44:195–208. doi: 10.1016/j.neuron.2004.09.006
139. Greenwood PM. Functional plasticity in cognitive aging: review and hypothesis. *Neuropsychology* (2007) 21:657–73. doi: 10.1037/0894-4105.21.6.657
140. Reuter-Lorenz PA, Park DC. Human neuroscience and the aging mind: a new look at old problems. *J Gerontol Series B* (2010) 65:405–15. doi: 10.1093/geronb/gbq035
141. Schneider-Garces NJ, Gordon BA, Brumback-Peltz CR, Shin E, Lee Y, Sutton BP, et al. Span, CRUNCH, and beyond: working memory capacity and the aging brain. *J Cogn Neurosci*. (2010) 22:655–69. doi: 10.1162/jocn.2009.21230
142. Park DC, Bischof, GN. Neuroplasticity, aging, and cognitive function, In: Schaie KW, Willis SL, editors. *Handbook of the Psychology of Aging*. San Diego, CA: Academic Press (2011). p. 109–117.
143. Rypma B, D'Esposito M. Age-related changes in brain-behaviour relationships: Evidence from event-related functional MRI studies. *Eur J Cogn Psychol*. (2001) 13:235–56. doi: 10.1080/0954144004200296
144. Vatansever D, Menon DK, Manktelow AE, Sahakian BJ, Stamatakis EA. Default mode dynamics for global functional integration. *J Neurosci*. (2015) 35:15254–62. doi: 10.1523/JNEUROSCI.2135-15.2015
145. Wen X, Zhang D, Liang B, Zhang R, Wang Z, Wang J, et al. Reconfiguration of the brain functional network associated with visual task demands. *PLoS ONE* (2015) 10:e0132518. doi: 10.1371/journal.pone.0132518
146. Liang X, Zou Q, He Y, Yang Y. topologically reorganized connectivity architecture of default-mode, executive-control, and salience networks across working memory task loads. *Cereb Cortex* (2016) 26:1501–11. doi: 10.1093/cercor/bhu316
147. Geerligns L, Maurits NM, Renken RJ, Lorist MM. Reduced specificity of functional connectivity in the aging brain during task performance. *Hum Brain Mapp*. (2014) 35:319–30. doi: 10.1002/hbm.22175
148. Geerligns L, Renken RJ, Saliassi E, Maurits NM, Lorist MM. A brain-wide study of age-related changes in functional connectivity. *Cereb Cortex* (2015) 25:1987–99. doi: 10.1093/cercor/bhu012
149. Honey CJ, Kötter R, Breakspear M, Sporns O. Network structure of cerebral cortex shapes functional connectivity on multiple time scales. *Proc Natl Acad Sci USA*. (2007) 104:10240–5. doi: 10.1073/pnas.0701519104
150. Hafkemeijer A, Altmann-Schneider I, Craen AJ, Slagboom PE, Grond J, Rombouts SA. Associations between age and gray matter volume in anatomical brain networks in middle-aged to older adults. *Aging Cell* (2014) 13:1068–74. doi: 10.1111/acel.12271
151. Liu K, Yao S, Chen K, Zhang J, Yao L, Li K, et al. Structural Brain Network Changes across the Adult Lifespan. *Front Aging Neurosci*. (2017) 9. doi: 10.3389/fnagi.2017.00275
152. Perry E, Blessed G, Tomlinson B, Perry R, Crow T, Cross A, et al. Neurochemical activities in human temporal lobe related to aging and Alzheimer-type changes. *Neurobiol Aging* (1981) 2:251–6. doi: 10.1016/0197-4580(81)90032-4
153. Salat DH, Kaye JA, Janowsky JS. Selective preservation and degeneration within the prefrontal cortex in aging and Alzheimer disease. *Arch Neurol*. (2001) 58:1403–8. doi: 10.1001/archneur.58.9.1403
154. De Haan W, Van Der Flier WM, Koene T, Smits LL, Scheltens P, Stam CJ. Disrupted modular brain dynamics reflect cognitive dysfunction in Alzheimer's disease. *Neuroimage* (2012) 59:3085–93. doi: 10.1016/j.neuroimage.2011.11.055
155. Van Den Heuvel MP, Fornito A. Brain networks in schizophrenia. *Neuropsychol Rev*. (2014) 24:32–48. doi: 10.1007/s11065-014-9248-7
156. Ganella EP, Seguin C, Bartholomeusz CF, Whittle S, Bousman C, Wannan CMJ, et al. Risk and resilience brain networks in treatment-resistant schizophrenia. *Schizophr Res* (2018) 193:284–92. doi: 10.1016/j.schres.2017.07.014
157. De Witte NAJ, Mueller SC. White matter integrity in brain networks relevant to anxiety and depression: evidence from the human

- connectome project dataset. *Brain Imaging Behav.* (2017) 11:1604–15. doi: 10.1007/s11682-016-9642-2
158. Kennis M, Van Rooij SJ, Van Den Heuvel MP, Kahn RS, Geuze E. Functional network topology associated with posttraumatic stress disorder in veterans. *Neuroimage Clin.* (2016) 10:302–9. doi: 10.1016/j.nicl.2015.12.008
159. Shim M, Im CH, Lee SH. Disrupted cortical brain network in post-traumatic stress disorder patients: a resting-state electroencephalographic study. *Transl Psychiatry* (2017) 7:e1231. doi: 10.1038/tp.2017.200
160. Fan Y, Shi F, Smith JK, Lin W, Gilmore JH, Shen D. Brain anatomical networks in early human brain development. *Neuroimage* (2011) 54:1862–71. doi: 10.1016/j.neuroimage.2010.07.025
161. Meunier D, Lambiotte R, Bullmore ET. Modular and hierarchically modular organization of brain networks. *Front Neurosci.* (2010) 4:200. doi: 10.3389/fnins.2010.00200
162. Meunier D, Foulupt P, Saive AL, Plailly J, Ravel N, Royet JP. Modular structure of functional networks in olfactory memory. *Neuroimage* (2014) 95:264–75. doi: 10.1016/j.neuroimage.2014.03.041

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Gonzalez-Escamilla, Muthuraman, Chirumamilla, Vogt and Groppa. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.