



Complex Role of Hypnotizability in the Cognitive Control of Pain

Enrica L. Santarcangelo* and Sybille Consoli

Department of Translational Research and New Technologies in Medicine and Surgery University of Pisa, Pisa, Italy

Keywords: hypnotizability, analgesia, BISBAS, cerebellum, expectation, insula, suggestions, pain

This opinion article deals with the role of hypnotizability in the efficacy of the suggestions of analgesia for the cognitive control of pain and of its physiological correlates.

Hypnotizability is a multidimensional trait including suggestibility (Raz, 2007) and other characteristics such as fantasy proneness, attitude to be deeply absorbed in specific tasks/mental images (Green and Lynn, 2011; Dasse et al., 2015), strong functional equivalence between imagery and perception (Papalia et al., 2014; Santarcangelo, 2014; Ibanez-Marcelo et al., 2018). It predicts the proneness to accept suggestions (Green et al., 2005; Elkins et al., 2015) and is measured by scales so that the general population is classified as high (*highs*, about 15%), medium (*mediums*, about 70%) and low (*lows*, about 15%) susceptible to hypnosis (De Pascalis et al., 2000). In the ordinary state of consciousness and in the absence of specific suggestions, different levels of hypnotizability are associated with different cerebral (Landry et al., 2017), cerebellar (Bocci et al., 2017; Picerni et al., 2018), sensorimotor and cardiovascular characteristics (Santarcangelo and Scattina, 2016).

The paper describes the interaction between explicit suggestions of analgesia and the expectation of pain relief (Huber et al., 2013; De Pascalis and Scacchia, 2016) on the basis of new findings regarding the hypnotisability-related polymorphism of opioid receptors $\mu 1$ (Presciuttini et al., 2018). In addition, the paper reports the observed joined influence of hypnotisability and cognitive-emotional traits (Madeo et al., 2015; Jensen et al., 2016) conceptualized as Behavioral Inhibition/Activation System (BIS/BAS) (Gray, 1990) and as Interoception/Interoceptive Awareness (Sebastiani et al., 2018; Varanini et al., 2018) on pain. Finally, the paper proposes that the observed morpho-functional peculiarities of the *highs*' salience network—insula, cingulate and prefrontal cortex (Landry et al., 2017)—and cerebellum (Bocci et al., 2017; Picerni et al., 2018) may play a role in the complex role of hypnotizability in pain modulation. In this respect, it should be noted that, although hypnotisability is an approximately stable individual trait, socio-cognitive factors such as relational disposition and the manipulation of expectation can modulate the efficacy of suggestions (Kirsch, 2018). The possible biological substrates for some of these factors, for instance the oxytocin release during hypnotic interventions, are now emerging (Kasos et al., 2018).

HYPNOTIZABILITY AND SUGGESTIONS FOR ANALGESIA

Suggestions are explicit instructions aimed at modifying perception, memory and behavior (Brafman and Kirsch, 1999). The suggestions for analgesia are widely employed owing to their efficacy in the cognitive control of pain, which can be predicted by the subjects' hypnotizability scores (Dillworth et al., 2012; Enea et al., 2014; Koban et al., 2017).

Hypnotisability scores predict the efficacy of the suggestions for analgesia in both the ordinary state of consciousness—that is in the absence of any procedure of hypnotic induction—and under hypnosis (Milling et al., 2005; Derbyshire et al., 2009; Meyer and Lynn, 2011). Explicit suggestions for analgesia can be associated or not with instructions for relaxation and pleasant imagery (Zachariae and Bjerring, 1994; Carlson et al., 2017; Hamlin and Robertson, 2017). Personalized suggestions (Berna et al., 2012; Koban et al., 2017) can be prepared for each patient according to her/his preference and directed to the sensory (Hofbauer et al., 2001) or cognitive-affective dimension of pain (Rainville et al., 1997) or both (Feldman, 2009).

OPEN ACCESS

Edited by:

Gianluca Castelnuovo,
Università Cattolica del Sacro Cuore,
Italy

Reviewed by:

Giorgia Varallo,
Istituto Auxologico Italiano (IRCCS),
Italy

Giuliano Carlo Geminiani,
Università degli Studi di Torino, Italy

*Correspondence:

Enrica L. Santarcangelo
enricals@dfb.unipi.it

Specialty section:

This article was submitted to
Clinical and Health Psychology,
a section of the journal
Frontiers in Psychology

Received: 01 February 2018

Accepted: 01 November 2018

Published: 20 November 2018

Citation:

Santarcangelo EL and Consoli S
(2018) Complex Role of Hypnotizability
in the Cognitive Control of Pain.
Front. Psychol. 9:2272.
doi: 10.3389/fpsyg.2018.02272

Neuroimaging studies (Del Casale et al., 2015) have shown that in *highs* the suggestions for analgesia modulate the functional connectivity among the regions of the pain matrix being able to modify pain perception, attention to pain, defensive responses and any other component of pain experience and behavior (Faymonville et al., 2003; Zeev-Wolf et al., 2016). EEG and EMG studies have also shown that in *highs* the suggestions for analgesia decrease pain and both cortical activity (De Pascalis et al., 1999, 2015; Valentini et al., 2013) and nociceptive reflexes (Kiernan et al., 1995; Danziger et al., 1998). Also *mediums* may respond to suggestions for analgesia, although to a lower extent (Fidanza et al., 2017). This enhances the number of subjects who may benefit from suggestion-induced analgesia from 15 to 85% of the general population (Montgomery et al., 2002a,b; Milling et al., 2006, 2007).

HYPNOTIZABILITY AND EXPECTATION OF PAIN RELIEF

Highs are more responsive than *lows* and *mediums* not only to the explicit suggestions of analgesia, but also to the conditioned analgesia, or Diffuse Noxious Inhibitory Control (Sandrini et al., 2000; Fidanza et al., 2017) which is mediated by endogenous opioids (Granot et al., 2008). This suggests that expectation-induced mechanisms, which are more effective in *highs* than in *lows*, are associated with those sustaining the conditioned analgesia. However, in *highs* the expectation of pain relief does not totally account for the suggestion induced analgesia (Gearan and Kirsch, 1993) and it is unlikely that the expectation of analgesia could be sustained by opioid mechanisms, in contrast to the general population (Amanzio and Benedetti, 1999; Benedetti et al., 1999; Petrovic et al., 2002; Zubieta et al., 2005; Scott et al., 2008; Babel et al., 2017). In fact, not only the effects of

suggestions is not abolished by naloxone (Moret et al., 1991) but, in addition, *highs* display the $\mu 1$ polymorphism (Presciuttini et al., 2018) which has been found associated with low sensitivity to opiates, low placebo response (Trescot and Faynboym, 2014; Bartošová et al., 2015; Peciña and Zubieta, 2015) and larger opiates consumption for post-surgery (Zhang et al., 2005; Boswell et al., 2013; Sia et al., 2013; Ren et al., 2015) and cancer pain (Gong et al., 2013; Wan et al., 2015; Yao et al., 2015).


In the general population, pain is associated with modulation of the activation and functional connectivity of the “pain matrix” that is the brain region sustaining the various dimensions of pain (Legrain et al., 2011). It includes the primary and secondary somatosensory areas, the insula and the anterior cingulate cortex. The emergence of pain depends on the flow and integration of information among these areas and is a function of individual characteristics and of the context (Iannetti and Mouraux, 2010).

Both explicit suggestions and placebo responses are due to top-down mechanisms (Zunhammer et al., 2018), but it has been shown that, in correspondence of similar subjective response to expectation-induced placebo, *highs* and *lows* exhibit opposite patterns of activity and functional connectivity (Huber et al., 2013). In fact, the former exhibit reduced functional connectivity between the right dorsolateral prefrontal cortex (rDLPFC) and the anterior midcingulate/medial prefrontal cortex, the left inferior frontal gyrus and the right cerebellum. In addition, placebo analgesia is associated with deactivation in the thalamus, basal ganglia, left precuneus and bilateral temporal gyrus only in *highs*. The observed differences are in line with earlier findings indicating that in the general population placebo analgesia is sustained by circuits involved in the regulation of emotional processes (Amanzio et al., 2011).

An observation relevant to clinical interventions, however, is that an experimental session including relaxation or distraction and suggestions for analgesia modulates pain experience also in chronic pain patients with low hypnotizability scores (Carli et al., 2008). This does not challenge the predictive role of hypnotizability as in *lows* analgesia is not time-locked with suggestions. This finding can be accounted for by a possible strong motivation to analgesia due to the presence of chronic pain, inducing expectation-induced placebo responses following suggestions (Hyland, 2011; Benedetti, 2013; Benedetti and Amanzio, 2013; Carlino et al., 2014) and making them indirectly effective also in *lows*. Thus, the suggestions for analgesia represent an easy and cheap tool for the cognitive control of pain in the large majority of acute (also procedure-related) and chronic pain patients (Elkins et al., 2007; Jensen et al., 2009; Stoelb et al., 2009; Didier et al., 2011; Jensen and Patterson, 2014; Mendoza et al., 2017a,b; Waisblat et al., 2017).

INTERACTION OF HYPNOTISABILITY WITH THE BEHAVIORAL INHIBITION/ACTIVATION SYSTEM AND INTEROCEPTION ABILITIES

Recent findings have challenged the established relation between the analgesic effects of suggestions and hypnotizability. In



fact, the interaction between hypnotizability and cognitive emotional traits such as those sustained by the Behavioral Inhibition/Activation System (BIS/BAS) (Gray, 1990) in pain imagery (Santarcangelo et al., 2013) and control (Jensen et al., 2016) and in its cortical correlates (Madeo et al., 2015) suggests hypnotizability may be just one of the factors involved in pain control by suggestions of analgesia.

BIS/BAS is based in limbic circuits (Gray, 1990; Angelides et al., 2017), concerns the proneness to approach or withdraw from possibly pleasant and unpleasant conditions, respectively, and is measured by scales (Carver, 2004). BIS is considered an attentional system sensitive to possible punishment, non-reward and novelty, while BAS reflects the motivation to follow one's goals and to approach fun and reward. High BIS is associated with enhanced attention, arousal and vigilance, high BAS with impulsivity, bipolar and attention deficit/ hyperactivity (De Pascalis et al., 2010). In particular, BIS/BAS modulates pain in patients with headache (Jensen et al., 2015) and musculoskeletal pain (Serrano-Ibáñez et al., 2018).

It has been shown that, even in the absence of significant differences between *highs*' and *lows*' scores, the BIS/BAS activity masks the hypnotizability-related differences in the vividness of pain imagery (Santarcangelo et al., 2013) and that the activity of BIS/BAS rather than hypnotizability itself is responsible for the hypnotizability-related EEG differences observed during tonic nociceptive stimulation associated and not associated with suggestions for analgesia in *highs*. On the other hand, in chronic pain patients the relation between BIS/BAS and hypnotizability is not linear (Jensen et al., 2016), which indicates a complex interaction.

Another trait potentially influencing the relation between hypnotizability and the effect of the suggestions for analgesia is the ability of interoception that is to detect and interpret bodily states and their changes pre-eminently related to the activity of the autonomic system. Interoceptive signals are monitored and processed at several levels of the central nervous system such as the insula, the orbitofrontal cortex and the cingulate cortex (Critchley and Harrison, 2013) and interoception has been found altered in mental disorders (Murphy et al., 2017; Khalsa et al., 2018) and chronic pain patients (Di Lernia et al., 2016). The role of interoception in pain has been found different in healthy *highs* and *lows*. In fact, a correlation between resting heart rate and pain threshold after suggestions of analgesia has been found in *highs* undergoing cold pressor test, but not in *lows*

(Varanini et al., 2018). In addition, preliminary findings indicate higher interoceptive awareness in *highs* than in *mediums* and *lows* (Sebastiani et al., 2018).

Morfo-functional differences between *highs* and *lows* have been observed in the insula and other limbic structures (Landry et al., 2017) and in the cerebellar cortex (Picerni et al., 2018). They consist of reduced gray matter volume (Landry et al., 2017; Picerni et al., 2018) and in a paradoxical increase in pain perception and amplitude of the cortically evoked observed after transcranial anodal stimulation of the cerebellum (Bocci et al., 2017). These morphofunctional differences could sustain the observed hypnotizability-related difference in the role of interoception and of the Behavioral Inhibition/Activation System in pain experience. In fact, interoception contributes to emotion (Critchley and Garfinkel, 2017), the insula and the cerebellum are involved in interoception /interpretation of bodily signals and autonomic monitoring and control, respectively (Di Lernia et al., 2016; Kuehn et al., 2016; Lu et al., 2016; Schulz, 2016; Adamaszek et al., 2017).

CONCLUSION

As summarized in **Figure 1**, (a) hypnotizability is just one of the individual traits involved in the ability to control pain through suggestions of analgesia; (b) in *highs* any method of cognitive control could be poorly sustained by opioid mechanisms; (c) hypnotizability-related morfo-functional characteristics of limbic circuits and of the cerebellum may sustain differences in cognitive-emotional traits contributing to peculiar pain processing; (d) the efficacy of the suggestions of analgesia in patients with low hypnotizability can be due to placebo responses elicited by suggestions.

The socio-cognitive views of hypnotizability and hypnosis (Lynn and Green, 2011) are the best reference frame to interpret the relation among hypnotizability and pain control. In fact, they allow to consider the joined role of a number of individual traits and of situational variables in pain perception and cognitive control.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

REFERENCES

- Adamaszek, M., D'Agata, F., Ferrucci, R., Habas, C., Keulen, S., Kirkby, K. C., et al. (2017). Consensus paper: cerebellum and emotion. *Cerebellum* 16, 552–576. doi: 10.1007/s12311-016-0815-8
- Amanzio, M., and Benedetti, F. (1999). Neuropharmacological dissection of placebo analgesia: expectation-activated opioid systems versus conditioning-activated specific subsystems. *J. Neurosci.* 19, 484–494. doi: 10.1523/JNEUROSCI.19-01-00484.1999
- Amanzio, M., Benedetti, F., Porro, C. A., Palermo, S., and Cauda, F. (2011). Activation likelihood estimation meta-analysis of brain correlates of placebo analgesia in human experimental pain. *Hum. Brain Mapp.* 34, 738–752. doi: 10.1002/hbm.21471
- Angelides, N. H., Gupta, J., and Vickery, T. J. (2017). Associating resting-state connectivity with trait impulsivity. *Soc. Cogn. Affect. Neurosci.* 12, 1001–1008. doi: 10.1093/scan/nsx031
- Babel, P., Bajcar, E. A., Adamczyk, W., Kicman, P., Lisinska, N., Swider, K., et al. (2017). Classical conditioning without verbal suggestions elicits placebo analgesia and nociceptive hyperalgesia. *PLoS ONE* 12:e0181856. doi: 10.1371/journal.pone.0181856
- Bartošová, O., Polanecky, O., Perlik, F., Adamek, S., and Slanar, O. (2015). OPRM1 and ABCB1 polymorphisms and their effect on postoperative pain relief with piritramide. *Physiol. Res.* 64(Suppl. 4), S521–S527.

- Benedetti, F. (2013). Placebo and the new physiology of the doctor-patient relationship. *Physiol. Rev.* 93, 1207–1246. doi: 10.1152/physrev.00043.2012
- Benedetti, F., and Amanzio, M. (2013). Mechanisms of the placebo response. *Pulm. Pharmacol. Ther.* 26, 520–523. doi: 10.1016/j.pupt.2013.01.006
- Benedetti, F., Arduino, C., and Amanzio, M. (1999). Somatotopic activation of opioid systems by target-directed expectations of analgesia. *J. Neurosci.* 19, 3639–3648. doi: 10.1523/JNEUROSCI.19-09-03639.1999
- Berna, C., Tracey, I., and Holmes, E. A. (2012). How a better understanding of spontaneous mental imagery linked to pain could enhance imagery-based therapy in chronic pain. *J. Exp. Psychopathol.* 3, 258–273. doi: 10.5127/jep.017911
- Bocci, T., Barloscio, D., Parenti, L., Sartucci, F., Carli, G., and Santarcangelo, E. L. (2017). High hypnotizability impairs the cerebellar control of pain. *Cerebellum* 16, 55–61. doi: 10.1007/s12311-016-0764-2
- Boswell, M. V., Stauble, M. E., Loyd, G. E., Langman, L., Ramey-Hartung, B., Baumgartner, R. N., et al. (2013). The role of hydromorphone and OPRM1 in postoperative pain relief with hydrocodone. *Pain Physic.* 16, e227–e235.
- Braffman, W., and Kirsch, I. (1999). Imaginative suggestibility and hypnotizability: an empirical analysis. *J. Pers. Soc. Psychol.* 77, 578–587. doi: 10.1037/0022-3514.77.3.578
- Carli, G., Suman, A. L., Biasi, G., Marcolongo, R., and Santarcangelo, E. L. (2008). Paradoxical experience of hypnotic analgesia in low hypnotizable fibromyalgic patients. *Arch. ItalBiol.* 146, 75–82.
- Carlino, E., Frisaldi, E., and Benedetti, F. (2014). Pain and the context. *Nat. Rev. Rheumatol.* 10, 348–355. doi: 10.1038/nrrheum.2014.17
- Carlson, L. E., Zelinski, E., Toivonen, K., Flynn, M., Qureshi, M., Piedalue, K. A., et al. (2017). Mind-body therapies in cancer: what is the latest evidence? *Curr. Oncol. Rep.* 19:67. doi: 10.1007/s11912-017-0626-1
- Carver, C. S. (2004). Negative affects deriving from the behavioral approach system. *Emotion* 4, 3–22. doi: 10.1037/1528-3542.4.1.3
- Critchley, H. D., and Garfinkel, S. N. (2017). Interoception and emotion. *Curr. Opin. Psychol.* 17, 7–14. doi: 10.1016/j.copsyc.2017.04.020
- Critchley, H. D., and Harrison, N. A. (2013). Visceral influences on brain and behavior. *Neuron* 77, 624–638. doi: 10.1016/j.neuron.2013.02.008
- Danziger, N., Fournier, E., Bouhassira, D., Michaud, D., De Broucker, T., Santarcangelo, E., et al. (1998). Different strategies of modulation can be operative during hypnotic analgesia: a neurophysiological study. *Pain* 75, 85–92. doi: 10.1016/S0304-3959(97)00208-X
- Dasse, M. N., Elkins, G. R., and Weaver, C. A. III. (2015). Correlates of the multidimensional construct of hypnotizability: paranormal belief, fantasy proneness, magical ideation, and dissociation. *Int. J. Clin. Exp. Hypn.* 63, 274–283. doi: 10.1080/00207144.2015.1031051
- De Pascalis, V., Bellusci, A., and Russo, P. M. (2000). Italian norms for the stanford hypnotic susceptibility scale, Form C1. *Int. J. Clin. Exp. Hypn.* 48, 315–323. doi: 10.1080/00207140008415249
- De Pascalis, V., Magurano, M. R., and Bellusci, A. (1999). Pain perception, somatosensory event-related potentials and skin conductance responses to painful stimuli in high, mid, and low hypnotizable subjects: effects of differential pain reduction strategies. *Pain* 83, 499–508. doi: 10.1016/S0304-3959(99)00157-8
- De Pascalis, V., and Scacchia, P. (2016). Hypnotizability 183 and placebo analgesia in waking and hypnosis as modulators of auditory startle responses in healthy women: an ERP study. *PLoS ONE* 11:e0159135. doi: 10.1371/journal.pone.0159135
- De Pascalis, V., Varriale, V., and Cacace, I. (2015). Pain modulation in waking and hypnosis in women: event-related potentials and sources of cortical activity. *PLoS ONE* 10:e0128474. doi: 10.1371/journal.pone.0128474
- De Pascalis, V., Varriale, V., and D'Antuono, L. (2010). Event-related components of the punishment and reward sensitivity. *Clin. Neurophysiol.* 121, 60–76. doi: 10.1016/j.clinph.2009.10.004
- Del Casale, A., Ferracuti, S., Rapinesi, C., Serata, D., Caltagirone, S. S., Savoja, V., et al. (2015). Pain perception and hypnosis: findings from recent functional neuroimaging studies. *Int. J. Clin. Exp. Hypn.* 63, 144–170. doi: 10.1080/00207144.2015.1002371
- Derbyshire, S. W., Whalley, M. G., and Oakley, D. A. (2009). Fibromyalgia pain and its modulation by hypnotic and non-hypnotic suggestion: an fMRI analysis. *Eur. J. Pain* 13, 542–550. doi: 10.1016/j.ejpain.2008.06.010
- Di Lernia, D., Serino, S., and Riva, G. (2016). Pain in the body. Altered interoception in chronic pain conditions: a systematic review. *Neurosci. Biobehav. Rev.* 71, 328–341. doi: 10.1016/j.neubiorev.2016.09.015
- Didier, H., Marchetti, C., Borromeo, G., Tullo, V., D'Amico, D., Bussone, G., et al. (2011). Chronic daily headache: suggestion for the neuromuscular oral therapy. *Neurol. Sci.* 32 (Suppl. 1), S161–S164. doi: 10.1007/s10072-011-0515-6
- Dillworth, T., Mendoza, M. E., and Jensen, M. P. (2012). Neurophysiology of pain and hypnosis for chronic pain. *Transl. Behav. Med.* 2, 65–72. doi: 10.1007/s13142-011-0084-5
- Elkins, G., Jensen, M. P., and Patterson, D. R. (2007). Hypnotherapy for the management of chronic pain. *Int. J. Clin. Exp. Hypn.* 55, 275–287. doi: 10.1080/00207140701338621
- Elkins, G. R., Barabasz, A. F., Council, J. R., and Spiegel, D. (2015). Advancing research and practice: the revised APA Division 30 definition of hypnosis. *Int. J. Clin. Exp. Hypn.* 63, 1–9. doi: 10.1080/00207144.2014.961870
- Enea, V., Dafinoiu, I., Opris, D., and David, D. (2014). Effects of hypnotic analgesia and virtual reality on the reduction of experimental pain among high and low hypnotizables. *Int. J. Clin. Exp. Hypn.* 62, 360–377. doi: 10.1080/00207144.2014.901087
- Faymonville, M. E., Roediger, L., Del Fiore, G., Delguedre, C., Phillips, C., Lamy, M., et al. (2003). Increased cerebral functional connectivity underlying the antinociceptive effects of hypnosis. *Brain Res. Cogn. Brain Res.* 17, 255–262. doi: 10.1016/S0926-6410(03)00113-7
- Feldman, J. B. (2009). Expanding hypnotic pain management to the affective dimension of pain. *Am. J. Clin. Hypn.* 51, 235–254. doi: 10.1080/00029157.2009.10401674
- Fidanza, F., Varanini, M., Ciaramella, A., Carli, G., and Santarcangelo, E. L. (2017). Pain modulation as a function of hypnotizability: diffuse noxious inhibitory control induced by cold pressor test vs explicit suggestions of analgesia. *Physiol. Behav.* 171, 135–141. doi: 10.1016/j.physbeh.2017.01.013
- Gearan, P., and Kirsch, I. (1993). Response expectancy as a mediator of hypnotizability modification: a brief communication. *Int. J. Clin. Exp. Hypn.* 41, 84–91. doi: 10.1080/00207149308414539
- Gong, X. D., Wang, J. Y., Liu, F., Yuan, H. H., Zhang, W. Y., Guo, Y. H., et al. (2013). Gene polymorphisms of OPRM1 A118G and ABCB1 C3435T may influence opioid requirements in Chinese patients with cancer pain. *Asian Pac. J. Cancer Prev.* 14, 2937–2943. doi: 10.7314/APJCP.2013.14.5.2937
- Granot, M., Weissman-Fogel, I., Crispel, Y., Pud, D., Granovsky, Y., Sprecher, E., et al. (2008). Determinants of endogenous analgesia magnitude in a diffuse noxious inhibitory control (DNIC) paradigm: do conditioning stimulus painfulness, gender and personality variables matter? *Pain* 136, 142–149. doi: 10.1016/j.pain.2007.06.029
- Gray, J. A. (1990). Brain systems that mediate both emotion and cognition. *Cogn. Emot.* 4, 269–288. doi: 10.1080/02699939008410799
- Green, J. P., Barabasz, A. F., Barrett, D., and Montgomery, G. H. (2005). Forging ahead: the 2003 APA Division 30 definition of hypnosis. *Int. J. Clin. Exp. Hypn.* 53, 259–264. doi: 10.1080/00207140590961321
- Green, J. P., and Lynn, S. J. (2011). Hypnotic responsiveness: expectancy, attitudes, fantasy proneness, absorption, and gender. *Int. J. Clin. Exp. Hypn.* 59, 103–121. doi: 10.1080/00207144.2011.522914
- Hamlin, A. S., and Robertson, T. M. (2017). Pain and complementary therapies. *Crit Care Nurs. ClinNorth Am.* 229, 449–460. doi: 10.1016/j.cnc.2017.08.005
- Hofbauer, R. K., Rainville, P., Duncan, G. H., and Bushnell, M. C. (2001). Cortical representation of the sensory dimension of pain. *J. Neurophysiol.* 86, 402–411. doi: 10.1152/jn.2001.86.1.402
- Huber, A., Lui, F., and Porro, C. A. (2013). Hypnotic susceptibility modulates brain activity related to experimental placebo analgesia. *Pain* 154, 1509–1518. doi: 10.1016/j.pain.2013.03.031
- Hyland, M. E. (2011). Motivation and placebos: do different mechanisms occur in different contexts? *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 366, 1828–1837. doi: 10.1098/rstb.2010.0391
- Iannetti, G. D., and Mouraux, A. (2010). From the neuromatrix to the pain matrix (and back). *Exp Brain Res.* 205, 1–12. doi: 10.1007/s00221-010-2340-1
- Ibanez-Marcelo, E., Campioni, L., Phinyomark, A., Petri, G., and Santarcangelo, E. L. (2018). Topology highlights mesoscopic functional equivalence between imagery and perception. *bioRxiv [Preprint]*. Available online at: <https://www.biorxiv.org/content/early/2018/02/20/268383>

- Jensen, M. P., Barber, J., Romano, J. M., Hanley, M. A., Raichle, K. A., Molton, I. R., et al. (2009). Effects of self-hypnosis training and EMG biofeedback relaxation training on chronic pain in persons with spinal-cord injury. *Int. J. Clin. Exp. Hypn.* 57, 239–268. doi: 10.1080/00207140902881007
- Jensen, M. P., Ehde, D., and Day, M. A. (2016). The behavioral activation and inhibition systems: implications for understanding and treating chronic pain. *J. Pain* 17, 529.e1–18. doi: 10.1016/j.jpain.2016.02.001
- Jensen, M. P., and Patterson, D. R. (2014). Hypnotic approaches for chronic pain management: clinical implications of recent research findings. *Am. Psychol.* 9, 167–177. doi: 10.1037/a0035644
- Jensen, M. P., Tan, G., and Chua, S. M. (2015). Pain intensity, headache frequency, and the behavioral activation and inhibition systems. *Clin. J. Pain.* 31, 1068–1074. doi: 10.1097/AJP.0000000000000215
- Kasos, E., Kasos, K., Pusztai, F., Polyák, Á., Kovács, K. J., and Varga, K., (2018). Changes in oxytocin and cortisol in active-alert hypnosis: hormonal changes benefiting low hypnotizable participants. *Int. J. Clin. Exp. Hypn.* 66, 404–427. doi: 10.1080/00207144.2018.1495009
- Khalsa, S. S., Adolphs, R., Cameron, O. G., Critchley, H. D., Davenport, P. W., Feinstein, J. et al. (2018). Interoception and mental health: a roadmap. *Biol. Psychiatry Cogn. Neurosci. Neuroimag.* 3, 501–513. doi: 10.1016/j.bpsc.2017.12.004
- Kiernan, B. D., Dane, J. R., Phillips, L. H., and Price, D. D. (1995). Hypnotic analgesia reduces R-III nociceptive reflex: further evidence concerning the multifactorial nature of hypnotic analgesia. *Pain* 60, 39–47. doi: 10.1016/0304-3959(94)00134-Z
- Kirsch, I. (2018). Response expectancy and the placebo effect. *Int. Rev. Neurobiol.* 138, 81–93. doi: 10.1016/bs.irn.2018.01.003
- Koban, L., Jepma, M., Geuter, S., and Wager, T. D. (2017). What's in a word? How instructions, suggestions, and social information change pain and emotion. *Neurosci. Biobehav. Rev.* 81, 29–42. doi: 10.1016/j.neubiorev.2017.02.014
- Kuehn, E., Mueller, K., Lohmann, G., and Schuetz-Bosbach, S. (2016). Interoceptive awareness changes the posterior insula functional connectivity profile. *Brain Struct. Funct.* 221, 1555–1571. doi: 10.1007/s00429-015-0989-8
- Landry, M., Lifshitz, M., and Raz, A. (2017). Brain correlates of hypnosis: a systematic review and meta-analytic exploration. *Neurosci. Biobehav. Rev.* 81(Pt A), 75–98. doi: 10.1016/j.neubiorev.2017.02.020
- Legrain, V., Iannetti, G. D., Plaghki, L., and Mouraux, A. (2011). The pain matrix Reloaded: a salience detection system for the body. *Prog. Neurobiol.* 93, 111–124. doi: 10.1016/j.pneurobio.2010.10.005
- Lu, C., Yang, T., Zhao, H., Zhang, M., Meng, F., Fu, H., et al. (2016). Insular cortex is critical for the perception, modulation, and chronicification of pain. *Neurosci Bull.* 32, 191–201. doi: 10.1007/s12264-016-0016-y
- Lynn, S. J., and Green, J. P. (2011). The sociocognitive and dissociation theories of hypnosis: toward a rapprochement. *Int. J. Clin. Exp. Hypn.* 59, 277–293. doi: 10.1080/00207144.2011.570652
- Madeo, D., Castellani, E., Mocenni, C., and Santarcangelo, E. L. (2015). Pain perception and EEG dynamics: does hypnotizability account for the efficacy of the suggestions of analgesia? *Physiol. Behav.* 145, 57–63. doi: 10.1016/j.physbeh.2015.03.040
- Mendoza, M. E., Capafons, A., Gralow, J. R., Syrjala, K. L., Suarez-Rodriguez, J. M., Fann, J. R., et al. (2017a). Randomized controlled trial of the Valencia model of waking hypnosis plus CBT for pain, fatigue, and sleep management in patients with cancer and cancer survivors. *Psychooncology* 26, 1832–1838. doi: 10.1002/pon.4232
- Mendoza, M. E., Capafons, A., and Jensen, M. P. (2017b). Hypnosis attitudes: treatment effects and associations with symptoms in individuals with cancer. *Am. J. Clin. Hypn.* 60, 50–67. doi: 10.1080/00029157.2017.1300570
- Meyer, E. C., and Lynn, S. J. (2011). Responding to hypnotic and nonhypnotic suggestions: performance standards, imaginative suggestibility, and response expectancies. *Int. J. Clin. Exp. Hypn.* 59, 327–349. doi: 10.1080/00207144.2011.570660
- Millington, L. S., Kirsch, I., Allen, G. J., and Reutenauer, E. L. (2005). The effects of hypnotic and nonhypnotic imaginative suggestion on pain. *Ann. Behav. Med.* 29, 116–127. doi: 10.1207/s15324796abm2902_6
- Millington, L. S., Reardon, J. M., and Carosella, G. M. (2006). Mediation and moderation of psychological pain treatments: response expectancies and hypnotic suggestibility. *J. Consult. Clin. Psychol.* 74, 253–262. doi: 10.1037/0022-006X.74.2.253
- Millington, L. S., Shores, J. S., Coursen, E. L., Menario, D. J., and Farris, C. D. (2007). Response expectancies, treatment credibility, and hypnotic suggestibility: mediator and moderator effects in hypnotic and cognitive-behavioral pain interventions. *Ann. Behav. Med.* 33, 167–178. doi: 10.1007/BF02879898
- Montgomery, G. H., David, D., Winkel, G., Silverstein, J. H., and Bovbjerg, D. H. (2002a). The effectiveness of adjunctive hypnosis with surgical patients: a meta-analysis. *Anesth Analg.* 94, 1639–1645.
- Montgomery, G. H., Wetz, C. R., Seltz, M., and Bovbjerg, D. H. (2002b). Brief presurgery hypnosis reduces distress and pain in excisional breast biopsy patients. *Int. J. Clin. Exp. Hypn.* 50, 17–32. doi: 10.1080/00207140208410088
- Moret, V., Forster, A., Laverriere, M. C., Lambert, H., Gaillard, R. C., Bourgeois, P., et al. (1991). Mechanism of analgesia induced by hypnosis and acupuncture: is there a difference? *Pain* 45, 135–140. doi: 10.1016/0304-3959(91)90178-Z
- Murphy, J., Brewer, R., Catmur, C., and Bird, G. (2017). Interoception and psychopathology: a developmental neuroscience perspective. *Dev. Cogn. Neurosci.* 23, 45–56. doi: 10.1016/j.dcn.2016.12.006
- Papalia, E., Manzoni, D., and Santarcangelo, E. L. (2014). Stabilizing posture through imagery. *Int. J. Lin. Exp. Hypn.* 62, 292–309. doi: 10.1080/00207144.2014.901080
- Peciña, M., and Zubieta, J. K. (2015). Molecular mechanisms of placebo responses in humans. *Mol. Psychiatry* 20, 416–423. doi: 10.1038/mp.2014.164
- Petrovic, P., Kalso, E., Petersson, K. M., and Ingvar, M. (2002). Placebo and opioid analgesia: imaging a shared neuronal network. *Science* 295, 1737–1740. doi: 10.1126/science.1067176
- Picerni, E., Santarcangelo, E. L., Laricchiuta, D., Cutuli, D., Petrosini, L., Spalletta, G., et al. (2018). Cerebellar structural variations in subjects with different hypnotizability. *Cerebellum*. doi: 10.1007/s12311-018-0965-y. [Epub ahead of print].
- Presciutti, S., Sciarrino, R., Curcio, M., Scatena, F., Jensen, M. P., and Santarcangelo, E. (2018). Polymorphism of opioids receptors $\mu 1$ in highly hypnotizable subjects. *Int. J. Clin. Exp. Hypn.* 66, 106–118. doi: 10.1080/00207144.2018.1396128
- Rainville, P., Duncan, G. H., Price, D. D., Carrier, B., and Bushnell, M. C. (1997). Pain affect encoded in human anterior cingulate but not somatosensory cortex. *Science* 277, 968–971.
- Raz, A. (2007). Suggestibility and hypnotizability: mind the gap. *Am. J. Clin. Hypn.* 49, 205–210. doi: 10.1080/00029157.2007.10401582
- Ren, Z. Y., Xu, X. Q., Bao, Y. P., He, J., Shi, L., Deng, J. H., et al. (2015). The impact of genetic variation on sensitivity to opioid analgesics in patients with postoperative pain: a systematic review and meta-analysis. *Pain Physician* 18, 131–152.
- Sandrini, G., Milanov, I., Malaguti, S., Nigrelli, M. P., Moglia, A., and Nappi, G. (2000). Effects of hypnosis on diffuse noxious inhibitory controls. *Physiol. Behav.* 69, 295–300. doi: 10.1016/S0031-9384(00)00210-9
- Santarcangelo, E. L. (2014). New views of hypnotizability. *Front. Behav. Neurosci.* 8:224. doi: 10.3389/fnbeh.2014.00224
- Santarcangelo, E. L., and Scattina, E. (2016). Complementing the latest APA definition of hypnosis: sensory-motor and vascular peculiarities involved in hypnotizability. *Int. J. Clin. Exp. Hypn.* 64, 318–330. doi: 10.1080/00207144.2016.1171093
- Santarcangelo, E. L., Varanini, M., Paoletti, G., Castellani, E., Palombo, C., and Carli, G. (2013). Pain-inducing imagery as a function of hypnotisability and of the activity of Gray's Behavioral Inhibition/Activation Systems. *Neurosci. Lett.* 557 Pt B, 184–187. doi: 10.1016/j.neulet.2013.06.049
- Schulz, S. M. (2016). Neural correlates of heart-focused interoception: a functional magnetic resonance imaging meta-analysis. *Phil. Trans. R. Soc. Lond. B* 371:20160018. doi: 10.1098/rstb.2016.0018
- Scott, D. J., Stohler, C. S., Egnatuk, C. M., Wang, H., Koeppe, R. A., and Zubieta, J. K. (2008). Placebo and nocebo effects are defined by opposite opioid and dopaminergic responses. *Arch. Gen. Psychiatry* 65, 220–231. doi: 10.1001/archgenpsychiatry.2007.34
- Sebastiani, L., Diolaiuti, F., Huber, A., Ciaramella, A., and Santarcangelo, E. L. (2018). Interoceptive awareness, conditioned emotion and hypnotisability. *Int. J. Psychophysiol.* 131S:S147.
- Serrano-Ibáñez, E. R., Ramírez-Maestre, C., López-Martínez, A. E., Esteve, R., Ruiz-Párraga, G. T., and Jensen, M. P. (2018). Behavioral inhibition and activation systems, and emotional regulation in individuals with chronic musculoskeletal pain. *Front. Psychiatry.* 9:394. doi: 10.3389/fpsy.2018.00394

- Sia, A. T., Lim, Y., Lim, E. C., Ocampo, C. E., Lim, W. Y., Cheong, P., et al. (2013). Influence of mu-opioid receptor variant on morphine use and self-rated pain following abdominal hysterectomy. *J. Pain* 14, 1045–1052. doi: 10.1016/j.jpain.2013.03.008
- Stoelb, B. L., Molton, I. R., Jensen, M. P., and Patterson, D. R. (2009). The efficacy of hypnotic analgesia in adults: a review of the literature. *Contemp. Hypn.* 26, 24–39. doi: 10.1002/ch.370
- Trescot, A. M., and Faynboym, S. (2014). A review of the role of genetic testing in pain medicine. *Pain Physician* 17, 425–445.
- Valentini, E., Betti, V., Hu, L., and Aglioti, S. M. (2013). Hypnotic modulation of pain perception and of brain activity triggered by nociceptive laser stimuli. *Cortex* 49, 446–462. doi: 10.1016/j.cortex.2012.02.005
- Varanini, M., Balocchi, R., Carli, G., Paoletti, G., and Santarcangelo, E. L. (2018). Hypnotizability and pain modulation: a body-mind perspective. *Int J. Clin. Exp. Hypn.* 66, 265–281. doi: 10.1080/00207144.2018.1460561
- Waisblat, V., Langholz, B., Bernard, F. J., Arnould, M., Benassi, A., Ginsbourger, F., et al. (2017). Impact of a hypnotically-based intervention on pain and fear in women undergoing labor. *Int. J. Clin. Exp. Hypn.* 65, 64–85. doi: 10.1080/00207144.2017.1246876
- Wan, X. S., Song, H. B., Chen, S., Zhang, W., Liu, J. Q., Huang, C., et al. (2015). Association of single nucleotide polymorphisms of ABCB1, OPRM1 and COMT with pain perception in cancer patients. *J. Huazhong Univ. Sci. Technol.* 35, 752–758. doi: 10.1007/s11596-015-1502-6
- Yao, P., Ding, Y. Y., Wang, Z. B., Ma, J. M., Hong, T., and Pan, S. N. (2015). Effect of gene polymorphism of COMT and OPRM1 on the preoperative pain sensitivity in patients with c cancer. *Int. J. Clin. Exp. Med.* 8, 10036–10039.
- Zachariae, R., and Bjerring, P. (1994). Laser-induced pain-related brain potentials and sensory pain ratings in high and low hypnotizable subjects during hypnotic suggestions of relaxation, dissociated imagery, focused analgesia, and placebo. *Int. J. Clin. Exp. Hypn.* 42, 56–80. doi: 10.1080/00207149408409341
- Zeev-Wolf, M., Goldstein, A., Bonne, O., and Abramowitz, E. G. (2016). Hypnotically induced somatosensory alterations: toward a neurophysiological understanding of hypnotic anaesthesia. *Neuropsychologia* 87, 182–191. doi: 10.1016/j.neuropsychologia.2016.05.020
- Zhang, Y., Wang, D., Johnson, A. D., Papp, A. C., and Sadee, W. (2005). Allelic expression imbalance of human mu opioid receptor (OPRM1) caused by variant A118G. *J. Biol. Chem.* 280, 32618–32624. doi: 10.1074/jbc.M504942200
- Zubieta, J. K., Bueller, J. A., Jackson, L. R., Scott, D. J., Xu, Y., Koeppe, R. A., et al. (2005). Placebo effects mediated by endogenous opioid activity on mu-opioid receptors. *J. Neurosci.* 25, 7754–7762. doi: 10.1523/JNEUROSCI.0439-05.2005
- Zunhammer, M., Bingel, U., Wager, T. D., and Placebo Imaging Consortium (2018). Placebo effects on the neurologic pain signature: a meta-analysis of individual participant functional magnetic resonance imaging data. *JAMA Neurol.* 75, 1321–1330. doi: 10.1001/jamaneurol.2018.2017

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Santarcangelo and Consoli. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.