



$\gamma\delta$ T Cells in Antimalarial Immunity: New Insights Into Their Diverse Functions in Protection and Tolerance

Kathleen W. Dantzler and Prasanna Jagannathan*

Department of Medicine, Stanford University, Stanford, CA, United States

Uniquely expressing diverse innate-like and adaptive-like functions, $\gamma\delta$ T cells exist as specialized subsets, but are also able to adapt in response to environmental cues. These cells have long been known to rapidly proliferate following primary malaria infection in humans and mice, but exciting new work is shedding light into their diverse functions in protection and following repeated malaria infection. In this review, we examine the current knowledge of functional specialization of $\gamma\delta$ T cells in malaria, and the mechanisms dictating recognition of malaria parasites and resulting proliferation. We discuss $\gamma\delta$ T cell plasticity, including changing interactions with other immune cells during recurrent infection and potential for immunological memory in response to repeated stimulation. Building on recent insights from human and murine experimental studies and vaccine trials, we propose areas for future research, as well as applications for therapeutic development.

Keywords: $\gamma\delta$ T cells, $V\gamma9V\delta2$ T cells, *Plasmodium*, cytotoxicity, cytokines, immunological memory

OPEN ACCESS

Edited by:

Kevin Couper,
University of Manchester,
United Kingdom

Reviewed by:

Bernd Lepenies,
University of Veterinary Medicine
Hannover, Germany
Nathan W. Schmidt,
University of Louisville, United States

*Correspondence:

Prasanna Jagannathan
prajs@stanford.edu

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Immunology

Received: 01 September 2018

Accepted: 03 October 2018

Published: 23 October 2018

Citation:

Dantzler KW and Jagannathan P
(2018) $\gamma\delta$ T Cells in Antimalarial
Immunity: New Insights Into Their
Diverse Functions in Protection and
Tolerance. *Front. Immunol.* 9:2445.
doi: 10.3389/fimmu.2018.02445

INTRODUCTION

$\gamma\delta$ T cells are unconventional T lymphocytes that are increasingly being appreciated for their unique role in integrating the innate and adaptive arms of the immune system. Comprising approximately 2–5% of peripheral blood T cells in healthy adults, they can uniquely recognize a broad range of antigens without the need for major histocompatibility complex (MHC) and can both establish and regulate the inflammatory response. Many of their individual functions—including production of pro-inflammatory cytokines, cytotoxic killing, antigen presentation, promotion of dendritic cell maturation, B cell help, recruitment of other immune cells, and secretion of growth factors—are shared with other immune cell types. However, a unique combination of antigen specificity, tissue distribution, kinetics and functional properties enable $\gamma\delta$ T cells to play an essential role in human immunity (1).

Long known to rapidly increase in number following systemic bacterial or parasitic infection (2–5), $\gamma\delta$ T cells may also be important in mediating protection against recrudescence and/or reinfection, particularly in the context of malaria (6–9). Recent investigations of $\gamma\delta$ T cell function during this disease caused by parasites in the *Plasmodium* genus are providing new insight into the processes underlying acute responses, as well as protection during chronic or recurrent infection. Despite progress in reducing worldwide incidence of malaria over the last decade, malaria remains a major global health problem, accounting for almost 500,000 deaths annually, predominantly in young children and pregnant women in sub-Saharan Africa (10). Improving our understanding of

the inflammatory and immunoregulatory roles of $\gamma\delta$ T cells during malaria infection may provide opportunities to manipulate this response therapeutically, potentially via combined targeting of $\gamma\delta$ T cells and B or T cell immunity as is currently being pursued for cancer. This review will integrate recent advances in understanding the diverse functions and plasticity of these fascinating cells in malaria. We discuss results from recent human and murine studies, including vaccine trials, and propose open areas for future research and development of novel antimalarial therapeutics targeting $\gamma\delta$ T cells.

THE UNIQUE FUNCTIONAL SPECIALIZATION OF $\gamma\delta$ T CELLS

Though $\gamma\delta$ T cells can carry out diverse innate- and adaptive-like functions, individual cell subsets have more restricted effector properties depending on expression of T cell receptor (TCR) V γ and V δ regions and associated tissue location (1). In humans, the V γ 9V δ 2 subset is the most abundant in adult human peripheral blood; approximately 50–90% of circulating $\gamma\delta$ T cells express this combination of chains, previously thought to be due to postnatal expansion. However, Dimova et al. recently demonstrated that V γ 9V δ 2 T cells with pre-programmed effector functions were the predominant $\gamma\delta$ T cell subset in fetal blood, suggesting that this subset of $\gamma\delta$ T cells may be prepared to respond before birth (11). The other major subset of $\gamma\delta$ T cells in humans, V δ 1+ $\gamma\delta$ T cells, are enriched in mucosal tissues where they sense host stress and stimulate leukocyte responses (12). In mice, $\gamma\delta$ T cells are most common in the skin and mucosal tissue (13) and act as the major initial IL-17 producers in various infectious and autoimmune models. Nearly all murine $\gamma\delta$ T cells in the epidermal layer of the skin, also known as dendritic epidermal T cells (DETC), express identical $\gamma\delta$ TCRs. In other animals like cattle, sheep, and chickens, $\gamma\delta$ T cells express highly diverse TCRs regardless of tissue localization (13). These differences between $\gamma\delta$ T cell subsets between species are essential to consider when interpreting conclusions from animal models. Subsets of $\gamma\delta$ T cells exhibiting different tissue tropism could have adapted to have differential potential for clonal expansion and therefore diverse roles in immunosurveillance.

Differential $\gamma\delta$ T cell subsets recognize different ligands; perhaps the best known interaction occurs between the stress-related phosphoantigens (PAgs) and the V γ 9V δ 2 subset (14). PAgs are intermediates of the eukaryotic mevalonate or the prokaryotic non-mevalonate pathway of isoprenoid synthesis; the former includes eukaryotic PAgs that are overproduced in tumor cells [e.g., isopentenyl pyrophosphate (IPP)] while the latter includes PAgs specifically produced by pathogens, such as (E)-4-hydroxy-3-methyl-but-2-enyl pyrophosphate (HMBPP). Importantly, recognition of these antigens is dependent on cell-cell contact involving the TCR but independent of antigen processing via MHC molecules. The potential for diversity in the $\gamma\delta$ TCR repertoire is currently under debate, but there is some evidence from deep sequencing of genomic DNA in a few individuals that though the majority of $\gamma\delta$ T cells in peripheral blood carry the same germline TCR γ

rearrangement, a substantial percentage (20%) have a more diverse TCR γ repertoire (15). Likely, this sequence diversity represents an evolutionary adaptation to bridge the innate and adaptive immune systems: universal sequences shared across individuals likely perform innate-like functions, while the diverse background repertoire plays a more adaptive role, as has been suggested in a recent study describing TCR repertoires within the V δ 2 compartment (16).

Regarding functional attributes of $\gamma\delta$ T cells, $\gamma\delta$ T cells can play numerous roles in response to infection, including direct anti-microbial roles, recruitment of innate immune cells (e.g., neutrophils, macrophages) and activation of the adaptive immune compartment (14). For example, $\gamma\delta$ T cells rapidly expand in response to mycobacterial phosphoantigen (17), mycobacterium infection (18), and vaccination with the tuberculosis vaccine Bacillus Calmette–Guerin (BCG) (18–20). V γ 9V δ 2 T cells can inhibit mycobacterial growth through soluble granzyme A (17, 21), and correlate with clearance of BCG bacteremia and immunity to fatal tuberculosis in BCG-vaccinated macaques (18). In contrast, human immunodeficiency virus (HIV) is associated with a loss of circulating V δ 2+ cells (22) and expansion of V δ 1+ cells that are able to proliferate and produce IFN γ and IL-17 in response to *Candida albicans* (23). Cytomegalovirus has also been shown to induce V δ 2^{neg} populations of $\gamma\delta$ T cells, including V δ 1+ cells, and these V δ 2^{neg} cells have antibody-dependent anti-cytomegalovirus activity through a process that is CD16 dependent (24). Finally, $\gamma\delta$ T cells display functional plasticity based on pre-programmed features (i.e., class of innate receptor, nature of inflammatory stimuli, and strength of the TCR signal) or more long-lasting effects induced by TCR signaling and environmental cues (1). For example, V γ 9V δ 2 cells responding to *E. coli* can adapt from a primarily cytokine-producing phenotype to a phenotype promoting phagocytosis (25). Clearly, $\gamma\delta$ T cells are capable of adopting a variety of anti-microbial functions, but the precise factors influencing this adaptability in specific infections or at different timepoints during infection require further characterization.

The functional attributes and plasticity of the $\gamma\delta$ T cell response to malaria resembles responses to other infections in some aspects, but also reflects the unique life cycle and epidemiology of *Plasmodium*. The epidemiology of malaria in many regions leading to recurrent infection over many years, as well as the diversity of mouse models for malaria, provide opportunities to gain insight into how $\gamma\delta$ T cells adapt their function following repeated stimulation.

IMPORTANCE OF $\gamma\delta$ T CELLS DURING MALARIA: INSIGHTS FROM NATURAL INFECTION AND EXPERIMENTAL MODELS IN HUMANS AND MICE

Whether in children, malaria-naïve adults, or malaria-experienced adults, it has long been known that $\gamma\delta$ T cells (in particular the phosphoantigen-responsive V γ 9V δ 2 subset) expand following infection with the most virulent human malaria

parasite, *Plasmodium falciparum* (*Pf*) (**Table 1**) (26, 29, 30, 47). Furthermore, higher frequencies and malaria-responsive cytokine production of V γ 9V δ 2 cells correlate with protection against subsequent infection in children living in endemic settings (31, 39). These associations, along with cytotoxic, anti-parasitic functions of V γ 9V δ 2 cells observed *in vitro* (48), suggest an important role for these cells in mediating protective immunity to malaria. Though most studies have focused on the V γ 9V δ 2 subset (**Figure 1**), earlier work suggested that V δ 1+, and not V δ 2+, T cells are expanded in the peripheral blood of individuals from endemic regions (28, 30), similar to what has been described in chronic HIV infection (22). Although this observation may be due to relative loss of circulating V δ 2+ T cells in repeatedly infected individuals rather than expansion of V δ 1+ T cells (34), a recent study demonstrated that non-V γ 9V δ 2, IL-10 producing $\gamma\delta$ T cells expand among individuals with uncomplicated malaria (38). Together this suggests that expansion of a non-V γ 9V δ 2 immunoregulatory population of $\gamma\delta$ T cells may also contribute to naturally acquired immunity.

In malaria-naïve individuals immunized with the attenuated *Pf* sporozoite (PfSPZ) vaccine, expansion and frequency of $\gamma\delta$ T cells (again, particularly the V δ 2 subset) was dose-dependent and a better correlate of protection compared to any other cellular immune responses (42, 43, 45). Numbers of memory V δ 2+ T cells similarly correlated with protection in the first PfSPZ trial in a malaria-endemic region; however, additional studies in the mouse led the authors to conclude that these cells were essential for induction of protective CD8+ T cell responses rather than directly exerting effector functions (46). Additional work is needed to further elucidate the mechanism of V δ 2+ $\gamma\delta$ T cell-induced protection, as well as to determine whether V δ 2 frequencies could be used as a biomarker for protection in PfSPZ vaccinations in malaria-endemic regions. Furthermore, in a trial immunizing malaria-naïve individuals with non-irradiated PfSPZ combined with chemoprophylaxis (PfSPZ-cVAC), V δ 2+ $\gamma\delta$ T cells, including cells expressing memory markers, also expanded in a dose-dependent manner and increased IFN γ expression (44). Together, these data suggest that $\gamma\delta$ T cells may be a correlate for both natural and vaccine-induced protection.

Studies in the mouse model have provided convincing evidence for a role for $\gamma\delta$ T cells in directly or indirectly mediating killing of blood-stage and/or liver-stage parasites and preventing parasite recrudescence. However, major differences in $\gamma\delta$ clones between mice and humans are an important caveat, as are differences between murine *Plasmodium* strains (**Table 2**). No subset corresponding to the human V γ 9V δ 2 subset exists in mice and most early studies examined all $\gamma\delta$ T cells without regard to antigen specificity. In mice infected with the *Plasmodium chabaudi* parasite, $\gamma\delta$ T cells expand by 10-fold (57, 58). Mice deficient in $\gamma\delta$ T cells experience higher parasitemia during acute *P. chabaudi* infection, as well as substantial parasitemic recrudescence (6–8). Depletion of $\gamma\delta$ T cells during chronic *P. chabaudi* infection in B cell-deficient mice also resulted in significantly worsened parasitemia (56). An exciting recent study demonstrated that $\gamma\delta$ T cells expand and become activated in later stages of *P. chabaudi* malaria (**Figure 2A**)—much later than CD4+ and CD8+ $\alpha\beta$ T cell activation—even when acute stages

are cleared early by drug treatment. This clonal expansion of $\gamma\delta$ T cell occurred primarily in murine blood, spleen, lung and liver, and effectively prevented late-stage parasite recurrence (9).

In the non-lethal *Plasmodium berghei* XAT model, control of parasitemia seems to be at least partially mediated by CD40 signaling and boosting of dendritic cell activation (53). Following vaccination with lethal *P. berghei* ANKA sporozoites, $\gamma\delta$ T cells contribute to pre-erythrocytic immunity by recruiting dendritic cells and CD8+ T cells during vaccination (46), but were not required to prevent infection upon blood-stage challenge. However, different results have been obtained utilizing the nonlethal *Plasmodium yoelii* model. In mice lacking $\alpha\beta$ T cells, $\gamma\delta$ T cells substantially influenced immunity to *P. yoelii* liver stages, but could not rescue immunity to blood stages (50), suggesting that at least in this parasite strain, $\gamma\delta$ T cells act as important effectors and their cytotoxicity may become more effective after interaction with CD4+ T cells. This same group showed that mice lacking $\gamma\delta$ T cells had significantly higher *P. yoelii* burden in the liver than similarly challenged immunocompetent mice, suggesting a potential role for $\gamma\delta$ T cells in the development of pre-erythrocytic immunity (51). These differences could potentially be explained solely by the different murine parasite strains used, as *P. yoelii* irradiated sporozoite vaccination does not induce sterile immunity while *P. berghei* vaccination does. Alternatively, the discrepant results could be explained by differing sporozoite preparations or the depleting monoclonal antibodies used (59, 60).

Finally, murine models have also suggested a role of $\gamma\delta$ T cell in disease pathogenesis. Mice depleted of $\gamma\delta$ T cells by monoclonal antibody were protected from cerebral malaria (the most severe form of malaria) due to *Plasmodium berghei*, but mice genetically depleted of $\gamma\delta$ T cells did not show this effect (52), implying that effective activation of $\gamma\delta$ T cells is extraordinarily time-sensitive. Likely, parasite species/mouse model and IFN γ levels at different infection timepoints strongly influence whether $\gamma\delta$ T cells contribute to protection or worsened pathogenesis.

RECOGNITION OF *PLASMODIUM* BY $\gamma\delta$ T CELLS

Numerous *in vitro* and *in vivo* studies have aimed to shed light on the mechanisms of $\gamma\delta$ T cell activation in response to malaria infection. *In vitro*, human $\gamma\delta$ (V γ 9V δ 2) T cells can proliferate in response to *Pf*-infected red blood cell (iRBC) lysates or schizont extract (34, 61–63), iRBC culture supernatants (62, 64, 65) and/or intact iRBCs (33, 48, 63, 66) (**Figure 1**). Presumably, these cells are responding to phosphoantigens present in *Pf* asexual blood stages (67), as earlier studies suggested that *Pf* sexual stages do not stimulate V γ 9V δ 2 T cells (68), though it was recently demonstrated that secreted phosphoantigens from iRBCs from all developmental stages (including gametocytes) are capable of stimulating V γ 9V δ 2 cells (69). Activation of V γ 9V δ 2 T cells in response to HMBPP has recently been shown to require butyrophilin 3A1 (BTN3A1, CD277), a type I glycoprotein in the B7 family (70). Though the precise molecular mechanisms

TABLE 1 | $\gamma\delta$ T cell responses to human malaria and associations with clinical outcomes.

Author, year	Country	Cohort	$\gamma\delta$ T cell subset	Impact of malaria exposure on $\gamma\delta$ T cell activation and function	Associations between $\gamma\delta$ T cell features and clinical outcomes
STUDIES OF MALARIA-NAÏVE TRAVELERS					
Roussilhon et al. (26)	France	Adults; acute <i>Pf</i>	All $\gamma\delta$	Expand after infection and remain elevated for months; subset respond <i>in vitro</i> to <i>Pf</i> schizont extract	
Howard et al. (27)	France	Adults; acute <i>Pf</i>	V γ 9V δ 2	<i>In vivo</i> exposure and <i>in vitro</i> stimulation associated with increased surface expression of APC-associated markers, induce naive $\alpha\beta$ T-cell responses, cross present soluble prototypical protein to antigen-specific CD8+ T cells	
STUDIES OF INDIVIDUALS IN MALARIA-ENDEMIC REGIONS					
Goodier et al. (28)	Benin	Adults and children	V γ 9+ and V δ 1+	Majority of $\gamma\delta$ T cells are V δ 1+; V γ 9+ cells not elevated compared to malaria-naïve controls but do proliferate after <i>in vitro</i> <i>Pf</i> stimulation	
Ho et al. (29)	Thailand	Age not reported; acute <i>Pf</i>	All $\gamma\delta$	Expand after acute infection and remain elevated for several weeks	
Hviid et al. (30)	Ghana	Children; acute <i>Pf</i>	V δ 1+	Increase after treatment and produce pro-inflammatory cytokines	
D'Ombrian et al. (31)	Papua New Guinea	Children	All $\gamma\delta$	Produce IFN γ following <i>in vitro</i> <i>Pf</i> stimulation	IFN γ from $\gamma\delta$ and $\alpha\beta$ T cells associated with immunity to symptomatic infection
Cairo et al. (32)	Cameroon	Neonates	V δ 2+	Placental malaria associated with increased proportions of central memory V γ 2V δ 2 cells in cord blood and altered V γ 2 chain repertoire <i>ex vivo</i> or after stimulation	
Stanisic et al. (33)	Papua New Guinea	Children	All $\gamma\delta$	Produce TNF, MIP-1 β , and MIP-1 α following <i>in vitro</i> <i>Pf</i> stimulation	Increased TNF from $\gamma\delta$ T cells and monocytes associated with severe malaria
Jagannathan et al. (34)	Uganda	Children	V δ 2+	Repeated infection associated with loss and dysfunction of V δ 2+ cells and increased V δ 2 expression of immunoregulatory genes including Tim3, CD57, CD16	Loss and dysfunction of V δ 2+ cells associated with clinical tolerance to infection
Farrington et al. (35)	Uganda	Children	V δ 2+	Frequencies and function lower and CD16 upregulated among children with high prior malaria exposure; antimalarial chemoprevention associated with enhanced V δ 2+ cytokine production	
Hsu et al. (36)	Malawi	Neonates	V δ 2+	Upregulate PD1 shortly after activation; after engagement of PD1 with PDL1, show dampened TNF α production and degranulation	
Schofield et al. (37)	Papua New Guinea	Children	All $\gamma\delta$	Elevated Tim-3+ $\gamma\delta$ T cells across whole cohort; IL-12 and IL-18 contribute to upregulation	Higher proportions of Tim-3+ $\gamma\delta$ T cells associated with asymptomatic malaria infection
Taniguchi et al. (38)	Laos	Adults and children; uncomplicated malaria	Non-V δ 2	Expand and produce IL-10 and IFN γ	
Jagannathan et al. (39)	Uganda	Children	V δ 2+	<i>In vivo</i> proliferative response attenuated with repeated exposure; repeated infection associated with loss and dysfunction of V δ 2+ cells	Higher pro-inflammatory cytokine production associate with protection from subsequent infection as well as increased odds of symptoms once infected

(Continued)

TABLE 1 | Continued

Author, year	Country	Cohort	$\gamma\delta$ T cell subset	Impact of malaria exposure on $\gamma\delta$ T cell activation and function	Associations between $\gamma\delta$ T cell features and clinical outcomes
VACCINATION STUDIES					
Teirlinck et al. (40); Roestenberg et al. (41)	The Netherlands	Malaria naïve adults; controlled-human malaria infection (CHMI) + chemoprophylaxis	All $\gamma\delta$	Produce IFN γ , even a year after infection	Long-term functional responses associated with protection against re-infection
Seder et al. (42); Ishizuka et al. (43)	USA	Malaria naïve adults; attenuated PfSPZ vaccination	V δ 2+	Expand after vaccination	Higher frequencies correlate with protection after CHMI
Mordmüller et al. (44)	Germany	Malaria naïve adults; non-irradiated PfSPZ vaccination + chemoprophylaxis	V γ 9V δ 2	Expand in dose-dependent manner and produce IFN γ	
Lyke et al. (45)	USA	Malaria naïve adults; attenuated PfSPZ vaccination	V δ 2+	Cell frequency increase after each vaccination and show activated phenotype	
Zaidi et al. (46)	Mali	Malaria-exposed adults; irradiated PfSPZ vaccination	V δ 2+		V δ 2+ T cells significantly elevated among vaccinees who remain uninfected during transmission season

underlying BTN3A1 essentiality remain controversial, there is now substantial evidence for a model in which the cytosolic B30.2 domain senses and binds to intracellular pyrophosphates and additional adaptor proteins. A resulting spatial redistribution or conformational change of the extracellular BTN3A1 domain is then recognized via an unknown mechanism by the V γ 9V δ 2 TCR (71–74). Other signaling pathways for human $\gamma\delta$ T cell activation involve the TCR interacting with ligands such as F1-ATPase or endothelial protein C receptor, or additional cell surface receptors such as natural killer group 2 member D (NKG2D) receptors or toll-like receptors (TLR). Some murine $\gamma\delta$ T cell subsets also appear to have a similar regulation pathway involving non-BTN3A1 butyrophilin-related molecules (75). It is unknown whether such a mechanism occurs during activation of $\gamma\delta$ T cells in malaria. In contrast to previous studies suggesting a requirement for cell-cell contact between $\gamma\delta$ T cells and parasites in initiating activation (48, 76), Guenot et al. demonstrated that at least for intact iRBCs, BTN3A is not present on the iRBC surface and cell-cell contact is not necessary for V γ 9V δ 2 activation (it may still be required for merozoites) (77). Rather, soluble molecules with characteristics of phosphoantigens seem to be released at the time of *Pf* egress from red blood cells (RBCs), leading the authors to hypothesize that V γ 9V δ 2 activation occurs via presentation by other $\gamma\delta$ T cells primarily in microvessels and in the red pulp of the spleen where later stage iRBCs sequester. It will be important to evaluate evidence for this theory, as well as to assess whether myeloid or other cells can present antigen to $\gamma\delta$ T cells.

PROLIFERATION, CYTOKINE, AND CYTOTOXIC RESPONSE OF $\gamma\delta$ T CELLS IN RESPONSE TO *PLASMODIUM*

The rapid proliferation of $\gamma\delta$ T cells in response to malaria infection likely depends on cytokine signaling and interaction with other immune cells (Figure 1). Proliferating $\gamma\delta$ T cells can then inhibit parasite growth *in vitro* (61, 78, 79). Intact iRBCs seem to generally be more effective than lysed iRBCs at inducing $\gamma\delta$ T cell expansion; however, lysed parasites can be made more effective with the addition of IL-2 (63). Similarly, activated CD4+ T cells may be required in the absence of IL-2 but are unnecessary when exogenous IL-2, IL-4 or IL-15 is added (64, 80). Proliferation of $\gamma\delta$ T cells in mouse spleen during chronic *P. chabaudi* infection appeared to depend on cytokines produced by CD4+ T cells (55). Additional molecules identified as required for IL-2-mediated survival and proliferation include the Ig superfamily receptor CD28 [demonstrated both with human samples and in the *Plasmodium berghei* mouse model (81)] and monocyte-derived cytokines, such as IL-10, IL-12, and IL-1 β (82), which also could increase cytokine production by $\gamma\delta$ T cells.

During malaria infection, $\gamma\delta$ T cells are a major source of IFN γ (Figures 1, 2), which is the cytokine most commonly associated with protection (83). Early IFN γ production has been associated with protection from clinical malaria in some cohorts (31, 84) but associated with worsened symptoms in others (33). There is some debate over whether $\gamma\delta$ T cells or NK cells are the predominant source of IFN γ ; several authors have suggested that




FIGURE 1 | Model for $V\gamma 9V\delta 2$ $\gamma\delta$ T cell response to *Plasmodium falciparum*. $V\gamma 9V\delta 2$ $\gamma\delta$ T cells recognize soluble phosphoantigens released from schizont stage parasites and potentially other iRBC stages. Though precise mechanisms of antigen presentation and recognition remain unclear, phosphoantigens are likely presented to $V\gamma 9V\delta 2$ TCR via BTN3A1. Other signals, such as CD28 and IL-2 from CD4+ cells and IL-15 from myeloid cells, contribute to $V\gamma 9V\delta 2$ proliferation and anti-parasitic activity, while $V\gamma 9V\delta 2$ anti-parasitic activity is dependent on granulysin production. Following activation, $V\gamma 9V\delta 2$ cells likely influence myeloid cell differentiation and activation through production of myeloid growth factors, and can themselves develop antigen-presenting cell (APC)-like functions such as activation of CD4+ T cells and cross-presentation of antigen to CD8+ T cells. After repeated parasite exposure, cells decrease production of pro-inflammatory cytokines and increase expression of CD16 and immunoregulatory molecules such as Tim-3. CD16 expression may enable alternative functions, such as antibody-dependent cellular cytotoxicity (ADCC). Immune cell images are adapted from the Reactome Icon Library (49).

$\gamma\delta$ T cells expressing NK cell receptors form the primary source of $IFN\gamma$ (33, 66, 85) whereas others have argued that $\gamma\delta$ T cells are important producers of $TNF\alpha$ but not $IFN\gamma$ (86). These differential results could be due to differences in timepoints, donors, parasite strains, surface markers used to differentiate cell populations, or immune cell activation conditions; further research, particularly in defining the impact of diversity in $\gamma\delta$ T cell numbers and TCR repertoires on heterogeneity of responses, is needed. It is also possible that $\gamma\delta$ T cells are required via an unknown mechanism for effective cytokine production by NK cells (86). Which cytokines are produced is likely determined by a combination of factors decided in cell development (e.g., expression of CD27) and epigenetic/transcriptional changes induced by environmental factors (87).

Several *in vitro* studies have provided further insight into the conditions required for $\gamma\delta$ T cell effector functions; while IL-2 induces $\gamma\delta$ T cell proliferation, for example, IL-15 is needed for anti-parasitic activity (48, 88, 89). Granulysin (but not perforin) released through cytotoxic granules also appears


to be required for anti-parasitic activity (48, 76), which is supported by the existence of granulysin-expressing $V\gamma 9V\delta 2$ cells in patients with malaria (48). Interestingly, Costa et al. showed that while both blood-stage parasites and extracellular merozoites activate $V\gamma 9V\delta 2$ cells and initiate degranulation, only merozoites trigger anti-parasitic activity by these cells (48). Though the RBC membrane has been suggested to be resistant to granulysin (90), it is currently unclear which parasite stages can be targeted by activated $\gamma\delta$ T cells and whether intracellular stages can be targeted via granulysin or other mechanisms.

$\gamma\delta$ T CELL ROLES IN STIMULATING OTHER INNATE AND ADAPTIVE IMMUNE RESPONSES IN RESPONSE TO *PLASMODIUM*

In addition to roles in cytokine production and cytotoxicity, there is increasing evidence that $\gamma\delta$ T cells can recruit and

TABLE 2 | Associations between $\gamma\delta$ T cells and protection in experimental *Plasmodium* infection models in mice.

Species	Strain	Characteristic	Finding	References
<i>Plasmodium yoelii</i>	Py17X (PyNL)	Non-lethal	<ul style="list-style-type: none"> $\gamma\delta$ T cells not essential for clearance of blood-stage parasites but do contribute to control of liver stages 	Tsuji et al. (50); McKenna et al. (51)
<i>Plasmodium berghei</i>	ANKA	Lethal	<ul style="list-style-type: none"> Mice with $\gamma\delta$ T cells depleted by monoclonal antibody are protected from cerebral malaria $\gamma\delta$ T cells affect CD8α+ dendritic cells in the liver, antigen-specific CD8+ T cell responses in the liver and spleen, and development of protective immunity 	Yañez et al. (52); Zaidi et al. (46)
<i>Plasmodium berghei</i>	XAT	Non-lethal	<ul style="list-style-type: none"> $\gamma\delta$ T cells essential for parasite clearance CD40 signaling between $\gamma\delta$ T cells and dendritic cells contributes to control of parasitemia $\gamma\delta$ T cells contribute to humoral immunity 	Inoue et al. (53); Inoue et al. (54)
<i>Plasmodium chabaudi</i>	AS	Non-lethal in C57BL/6 mice	<ul style="list-style-type: none"> $\gamma\delta$ T cells expand after infection and produce IFNγ Mice deficient in $\gamma\delta$ T cells have higher parasitemia $\gamma\delta$ T cells expanding later in infection protect against parasite recrudescence 	Van der Hyde et al. (55); Langhorne et al. (6); Seixas and Langhorne (7); Weidanz et al. (8); Weidanz et al. (56); Mamedov et al. (9)



stimulate other immune cells, and can adjust pro-inflammatory vs. regulatory effector functions depending on specific host or pathogen factors (for example, cytokines present in the microenvironment) (91, 92) (Figures 1, 2). As described above, non-V γ 9V δ 2 cells were found to proliferate among

individuals in a malaria-endemic region and produced the immunoregulatory cytokine IL-10, suggesting expansion of a non-cytotoxic, immunoregulatory population of $\gamma\delta$ T cells (38). Furthermore, though $\gamma\delta$ T cells have been known to function as antigen-presenting cells (APCs) in other contexts (93–96),

this phenomenon was recently demonstrated for the first time in malaria. $V\gamma9V\delta2$ T cells from malaria-infected individuals more highly expressed antigen-presenting and costimulatory molecules, such as HLA-DR, CD80, and CD86, compared with healthy and nonmalarial febrile control subjects (**Figure 1**) (27). These cells were found to have APC-like functions in response to *in vitro* iRBC stimulation, including activating naive $\alpha\beta$ T cell responses and cross-presenting protein to antigen-specific CD8+ T cells (27). Studies in humans have also shown that $V\gamma9V\delta2$ T cells express the myeloid growth factors GM-CSF and IL-3 following *in vitro* stimulation with iRBCs (34). In mice, Mamedov et al. found that a specific $\gamma\delta$ T cell clone (TRAVN-1+/V d6.3+) responsible for protection from *P. chabaudi* responded late in infection and prevented parasite recrudescence. Unlike IFN γ -producing $\gamma\delta$ T cells responding earlier in infection (9), this clone produced macrophage-colony stimulating factor (M-CSF) and accessory cytokines that influence the myeloid compartment (i.e., CCL5, CCL3) (**Figure 2A**). Precise differences between IFN γ - and M-CSF-producing $\gamma\delta$ T cells, including their direct vs. indirect roles in preventing parasite recrudescence, remain to be defined. Furthermore, whether malaria-induced expression of myeloid growth factors by $\gamma\delta$ T cells directly or indirectly influences macrophage function, including epigenetic reprogramming of the myeloid compartment [potentially inducing trained immunity (97)], remains to be determined.

Just as $\gamma\delta$ T cells could indirectly influence adaptive immunity through modulation of monocyte function, a parallel process could occur in dendritic cells (DCs). The previously mentioned PfSPZ vaccine trial and corresponding validation in the mouse model suggested that $V\delta2+$ $\gamma\delta$ T cells may be essential for induction of dendritic cell and protective CD8+ responses (46). The absence of $\gamma\delta$ T cells during murine vaccination led to dramatically reduced CD8 α + DCs in the liver, impaired antigen-specific CD8+ T cell responses in the liver and spleen, and resulting impaired development of protective immunity (46). These results highlight a possible role for $\gamma\delta$ T cells in promoting the migration and/or proliferation of CD8 α + DCs and/or a requirement for cross-talk between $\gamma\delta$ T cells and CD8 α + DCs in induction of downstream effector CD8+ T cell responses during PfSPZ vaccination (46). Interestingly, there was no impact of $\gamma\delta$ T cell absence on production of antibody targeting the circumsporozoite protein (CSP); however, this does not preclude an effect on antibodies targeting other parasite antigens. Interestingly, $\gamma\delta$ T cell expression of CD16, which is known to mediate antibody-dependent cellular cytotoxicity (ADCC) in CMV infection (24) is increased in children in malaria-endemic regions (34, 35), suggesting a potential role for $\gamma\delta$ T cells in inciting antibody-mediated parasite killing (**Figure 1**). Finally, paralleling previous observations during influenza infection (98), recent work in the *P. berghei* mouse model demonstrated a role for $\gamma\delta$ T cells in increasing levels of antigen-specific antibodies, T_H cells, and germinal center B cells via expression of IL-21 and IFN γ early in infection (54). Altogether, evidence suggests that malaria-responsive $\gamma\delta$ T cells are able to use diverse direct and indirect (via recruitment of monocytes, dendritic cells, and CD4+ cells) mechanisms to influence effector responses later in infection.

$\gamma\delta$ T CELL MODULATION DURING RECURRENT INFECTION

New insights into changing $\gamma\delta$ T cell functions during recurrent malaria infection bring up intriguing questions surrounding the relative importance of this modulation in natural immunity to malaria and whether this phenomenon represents a functional or dysfunctional response. Repeated *Pf* infection among Ugandan children was associated with reduced percentages of $V\delta2+$ $\gamma\delta$ T cells, decreased pro-inflammatory cytokine production in response to malaria antigens, and increased expression of CD16 and CD57 and immunoregulatory genes such as HAVCR2 (encoding the inhibitory receptor Tim-3) (34) (**Figures 1, 2B**). Importantly, though higher $V\delta2+$ pro-inflammatory cytokine production was associated with protection from subsequent infection, it was also associated with increased odds of having symptoms once infected (39). This suggests that $V\delta2+$ T cell dysfunction may represent a disease tolerance mechanism allowing for the development of “clinical immunity” to malaria—a decline in symptomatic infections and an increasing proportion of infections that are asymptomatic (34). Alternatively, it is also possible that with repeated exposure, $\gamma\delta$ T cell responses gain alternative functional capabilities (i.e., CD16-mediated processes like ADCC).

Regarding mechanisms driving this dysfunction, co-engagement of CD46 and $\gamma\delta$ TCR in cells stimulated by HMBPP has been shown to suppress production of IFN γ and TNF α , suggesting that CD46 could be involved in mediating $\gamma\delta$ T cell regulation (99). Alternatively, there is evidence for reduced $\gamma\delta$ T cell effector function in the setting of Tim-3 expression (37, 39), suggesting that repeated or chronic infection may induce Tim-3 mediated $\gamma\delta$ T cell exhaustion, similar to what has been described in Th1 T cells (100, 101) and other innate cells (102). Consistent with this hypothesis, a study found that Tim-3 blockade in murine malaria improved T cell-mediated immunity (103). Schofield et al. recently described that Tim-3+ $\gamma\delta$ T cells elevated in children living in malaria-endemic Papua New Guinea were independently associated with asymptomatic malaria infection, consistent with a role for Tim-3 mediated $\gamma\delta$ T cell immunoregulation in minimizing symptoms due to malaria (37). Tim-3 expression by $\gamma\delta$ T cells in this context was regulated by IL-12 and IL-18 (37). As IL-12/IL-18 can also induce IFN γ production, it will be important to assess the factors and timing differentiating this phenotype from Tim-3 expression, including the role of phosphoantigen.

Additional studies in neonates have highlighted that placental malaria affects the phenotype and repertoire of $V\delta2+$ lymphocytes in cord blood, potentially lowering the capacity for subsequent $V\delta2+$ responses to both malaria and other infectious diseases (32). Neonatal $V\delta2+$ T cells were recently shown to upregulate programmed death 1 (PD1), which when engaged by its ligand, PDL1, decreases TNF α production and degranulation by $V\delta2+$ cells (36). Intriguingly, PD1 expression by neonatal $V\delta2$ cells was inversely associated with promoter DNA methylation (36), suggesting a role for epigenetic programming in regulating inflammatory responses. Though much progress has been made in understanding the development of $\gamma\delta$ T regulation,

further work is needed to more precisely define the underlying mechanisms and to reconcile *in vivo* observations with *in vitro* results showing an increased responsiveness to lysed *Pf* iRBCs after priming with intact iRBCs (85).

POTENTIAL FOR IMMUNOLOGICAL MEMORY IN MALARIA-RESPONSIVE $\gamma\delta$ T CELLS

Altered $\gamma\delta$ T cell function and upregulated immunoregulatory markers following repeated infection lead to exciting questions concerning the capacity of $\gamma\delta$ T cells to develop immunological memory (104), whether similarly to canonical T cell memory or to innate memory. Though $\gamma\delta$ T cells have historically been thought of as primarily innate-like, quick-responding cells, there is increasing evidence that these cells are also important at later timepoints during infection and have important adaptive-like functions. In response to CMV, $\gamma\delta$ T cells developed a cytotoxic effector/memory phenotype, which in a secondary response led to a faster $\gamma\delta$ T cell expansion and a better resolution of infection than the primary response (105). Similarly, distinct primary and recall/memory responses were observed in response to mycobacterial infection in macaques, in which $\gamma\delta$ T cell ability to rapidly expand following BCG vaccination correlated with immunity to fatal tuberculosis (18). In controlled human malaria infections in malaria-naïve adults, $\gamma\delta$ T cells expand late after infection; elevated frequencies of cells expressing effector memory surface markers and enhanced responsiveness to *Pf* stimulation persisted for over 1 year (40). Furthermore, *in vitro* experiments demonstrated that the recall response of $\gamma\delta$ T cells post-challenge was independent of other PBMCs (40).

In contrast to adaptive T cell memory, innate-like memory in $\gamma\delta$ T cells is an intriguing alternative supported by evidence that monocytes can adapt secondary response to infection based on priming-induced epigenetic reprogramming (106, 107). A recent study has observed this phenomenon, termed “trained immunity”, in monocytes stimulated *in vitro* with *Pf*-iRBCs or from children living in malaria-endemic regions (108). Preliminary evidence showing epigenetic reprogramming at the PD1 locus in neonatal V γ 9V δ 2 cells (36) suggests that a similar process could be responsible for the immunoregulatory phenotype seemingly acquired among $\gamma\delta$ T cells after multiple infections (34, 39). Additional questions surrounding memory include the relative importance of adaptive-like vs. innate-like memory cells in protection from malaria, the localization of these cells, and the host and parasite factors influencing their development.

DIRECTIONS FOR FUTURE RESEARCH AND APPLICATION TO POTENTIAL THERAPEUTICS

Recent advances in our understanding of the role of $\gamma\delta$ T cells in recurrent malaria infection and vaccination highlight numerous open questions and areas for future research (Box 1). Whether $\gamma\delta$ T cell activation occurs via cell-cell contact, soluble

phosphoantigens released from rupturing schizonts (77) intact iRBCs (69), and/or other parasite stages, and how these cells are precisely able to target and kill *Pf*-iRBC, remains to be elucidated. Though V γ 9V δ 2 cells have been the most studied subset in human malaria, it will be important to determine the role of non-V γ 9V δ 2 cells in infection (38). In addition, evaluating the activation and role of $\gamma\delta$ T cells in other important human *Plasmodium* infections, including *Plasmodium vivax* (109), will benefit eradication programs in diverse settings. Furthermore, a better understanding of human donor variability (66, 77) and parallels and differences between human and murine $\gamma\delta$ T cell responses to malaria will enable better translation of observations between the experimental setting and the clinic.

Current work investigating plasticity of $\gamma\delta$ T cells in malaria have exciting implications for our understanding of $\gamma\delta$ T cell diversity and memory. For example, if V γ 9+ and V γ 9-V δ 2+ subsets serve distinct “innate-like” vs. “adaptive-like”

Box 1 | $\gamma\delta$ T cells in response to malaria: outstanding questions and areas for future research.

$\gamma\delta$ T cell activation and functional specialization.

- What are the precise mechanisms underlying V γ 9V δ 2 T cell activation in response to *Plasmodium falciparum*?
- What is the relative importance of intrinsic vs. extrinsic signals required for $\gamma\delta$ T cell activation, proliferation, and anti-parasitic activity?
- How do V γ 9V δ 2 T cells target and kill malaria-infected red blood cells?
- What is the functional role of non-V δ 2+ T cell subsets in human malaria? How are these cells activated?
- What is the role of $\gamma\delta$ T cell subsets in the response to non-*Plasmodium falciparum* strains? Given differences in mouse vs. human $\gamma\delta$ T cell subsets, how should differences in responses between mouse and human studies be interpreted?

$\gamma\delta$ T cell plasticity, memory, and altered function after repeated infection.

- How much diversity exists among malaria-responsive V δ 2+ cells (i.e., do separate “innate-like” and “adaptive-like” subsets coexist?)
- What malaria-induced cellular and microenvironmental cues drive $\gamma\delta$ T cell differentiation and plasticity?
- Does dysfunction of V γ 9V δ 2 T cells following repeated malaria represent exhaustion, anergy, or, alternatively, immunologic memory and/or gain of function (e.g., ADCC, antigen presentation)?
- What mechanisms drive V γ 9V δ 2 T cell loss and dysfunction following repeated malaria? (i.e., altered cellular metabolism, epigenetic modifications?)
- Can altered $\gamma\delta$ T cell function be reversed or made more functional?
- How do altered $\gamma\delta$ T cell functions impact interactions with other immune cells (monocytes, dendritic cells, $\alpha\beta$ T cells, B cells, T follicular helper cells)?
- What is the relative importance of innate-like “trained immunity” vs. canonical T cell memory in $\gamma\delta$ T cell immunological memory? Does this balance differ in natural infection vs. vaccine-induced exposure?

Applications to novel therapeutics and vaccines.

- Can therapeutic approaches that prevent the development of V γ 9V δ 2 T cell dysfunction enhance parasite clearance?
- Can adjuvant approaches targeting $\gamma\delta$ T cells (i.e., BCG, HMBPP, small molecules) influence the development of vaccine-induced malaria-specific humoral or $\alpha\beta$ T cell responses?

functions in malaria as has been recently shown in acute CMV (16), this would significantly impact our understanding of differentiation and clonal expansion across age and repeated malaria exposure. Differential cytokine expression in the tissue microenvironment in cancer (91) can induce functional plasticity and differentiation of diverse $\gamma\delta$ T cell subsets, but it remains to be determined whether cellular and/or environmental cues similarly drive differential cell fates in malaria. As repeated malaria has been shown to lead to dysfunction of V γ 9V δ 2 T cells, it will be important to assess whether this process represents exhaustion, anergy, or alternatively, immunologic memory and/or gain of function (e.g., ADCC). Mechanisms driving these alterations, including potential epigenetic and/or metabolic perturbations, should be examined (110). In addition, inflammation during recurrent malaria leads to suppression of functional memory B cells (MBC) (111) and expansion of “inferior” T follicular helper (Tfh) cells (112, 113) and “atypical” MBC exhibiting impaired proliferation, cytokine production and antibody secretion (114). Considering the overlapping timing of immunoregulatory phenotypes developing in $\gamma\delta$ T cells and adaptive cells, as well as evidence for $\gamma\delta$ T cells impacting germinal center reactions in murine malaria (54) and Tfh differentiation during influenza (98), future research should examine the impact of altered $\gamma\delta$ T cell function on expansion of MBC and Tfh subsets and interactions with other immune cells.

Novel tools that reduce the immense global burden of malaria by improving natural immunity are urgently needed. Potential therapies involving $\gamma\delta$ T cells could directly target $\gamma\delta$ T cell responses or could include $\gamma\delta$ T cell stimulation in an approach targeting antibody or T cell responses. The former approach could target activating or regulatory molecules or binding affinity/avidity to parasite antigens, induce intracellular accumulation of human V γ 9V δ 2 T cell agonists, and/or stimulate a particular $\gamma\delta$ T cell function (e.g., ADCC via increasing CD16 expression). Clinically available compounds performing such functions led to accelerated clearance of *Yersinia pestis* and repair of inflamed tissue in non-human primates (115), and co-administration of V γ 9V δ 2 T cell agonists and IL-2 in cancer patients induced efficient activation of $\gamma\delta$ T cells, and ultimately disease stabilization (116, 117). Interestingly, BCG has historically been used as treatment for bladder cancer, potentially by stimulating V γ 9V δ 2 T cells to more efficiently kill cancer cells (118, 119). Future studies investigating the role of BCG or other similar strategies may be useful to enhance responses to malaria and/or other infectious diseases. Therapeutic approaches to prevent $\gamma\delta$ T cell dysfunction should also be explored. For example, experimental malaria infection given under chloroquine prophylaxis leads to long-term

functional $\gamma\delta$ T cell responses associated with protection against re-infection (40, 41), and preventing blood stage infection with highly effective antimalarial chemoprevention was also recently shown to prevent $\gamma\delta$ T cell dysfunction in children (35). Drug development could also target inhibitory receptors involved in $\gamma\delta$ T cell dysfunction (e.g., Tim-3) or epigenetic pathways involved in modulating plasticity (and possibly trained immunity) of $\gamma\delta$ T cells (120).

Regarding approaches using adjuvant $\gamma\delta$ T cell stimulation to augment vaccine-induced antibody or T cell responses, one group immunized non-human primates with a subunit vaccine for tuberculosis combined with phosphoantigen. The authors identified a robust $\gamma\delta$ T cell response (including development of effector memory surface markers) following primary vaccination but anergy after subsequent boosts (121). In contrast, $\alpha\beta$ T cells proliferated after boost vaccinations. These promising results indicate a need for further work aiming to maximize protective responses, whether by preventing $\gamma\delta$ T cell anergy or optimizing timing of functional $\gamma\delta$ and $\alpha\beta$ T cell responses. As the balance between pro-inflammatory and anti-inflammatory responses changes drastically throughout malaria infection, the timing of any of these interventions would be essential to boosting responses without worsening pathology. Such interventions could also be useful for other infections that elicit chronic antigen exposure and/or an exhausted $\gamma\delta$ T cell phenotype.

Successful therapies targeting $\gamma\delta$ T cells for malaria will likely require a more thorough understanding of (1) functional differences between V γ 9V δ 2 T cell subpopulations, (2) migration and tissue infiltration of V γ 9V δ 2 T cells *in vivo*, (3) cellular interactions in the relevant microenvironment, (4) factors that influence $\gamma\delta$ T cell differentiation and exhaustion, (5) $\gamma\delta$ T cell detection of and response to metabolic changes in the host, and (6) factors that determine the balance between pro-inflammatory and immunoregulatory responses. The development of tools eliciting long-term, functional $\gamma\delta$ T cell responses will be a much-needed addition to the campaign to eliminate and eradicate malaria.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

This work was supported in part by U19AI057229 (pilot project grant to PJ). KD is supported by an Immunology training grant postdoctoral fellowship (5 T32 AI07290-33).

REFERENCES

- Bonneville M, O'Brien RL, Born WK. Gammadelta T cell effector functions: a blend of innate programming and acquired plasticity. *Nat Rev Immunol*. (2010) 10:467–78. doi: 10.1038/nri2781
- Galley HF, Lowes DA, Thompson K, Wilson ND, Wallace CA, Webster NR. Characterisation of gamma delta (gammadelta) T cell populations in patients with sepsis. *Cell Biol Int*. (2015) 39:210–6. doi: 10.1002/cbin.10361
- Hara T, Mizuno Y, Takaki K, Takada H, Akeda H, Aoki T, et al. Predominant activation and expansion of V gamma 9-bearing gamma delta T cells *in vivo* as well as *in vitro* in Salmonella infection. *J Clin Invest*. (1992) 90:204–10. doi: 10.1172/JCI115837
- Matsushima A, Ogura H, Fujita K, Koh T, Tanaka H, Sumi Y, et al. Early activation of gammadelta T lymphocytes in patients with severe systemic inflammatory response syndrome. *Shock* (2004) 22:11–5. doi: 10.1097/01.shk.0000129203.84330.b3

5. Schneider T, Jahn HU, Liesenfeld O, Steinhoff D, Riecken EO, Zeitz M, et al. The number and proportion of Vgamma9 Vdelta2 T cells rise significantly in the peripheral blood of patients after the onset of acute Coxiella burnetii infection. *Clin Infect Dis.* (1997) 24:261–4.
6. Langhorne J, Mombaerts P, Tonegawa S. Alpha beta and gamma delta T cells in the immune response to the erythrocytic stages of malaria in mice. *Int Immunol.* (1995) 7:1005–11.
7. Seixas EM, Langhorne J. Gammadelta T cells contribute to control of chronic parasitemia in *Plasmodium chabaudi* infections in mice. *J Immunol.* (1999) 162:2837–41.
8. Weidanz WP, Kemp JR, Batchelder JM, Cigel FK, Sandor M, Heyde HC. Plasticity of immune responses suppressing parasitemia during acute *Plasmodium chabaudi* malaria. *J Immunol.* (1999) 162:7383–8.
9. Mamedov MR, Scholzen A, Nair RV, Cumnock K, Kenkel JA, Oliveira JHM, et al. A Macrophage colony-stimulating factor-producing $\gamma\delta$ t cell subset prevents malarial parasitemic recurrence. *Immunity* (2018) 48:350–63. doi: 10.1016/j.immuni.2018.01.009
10. WHO. *WHO Malaria Report 2017* (2017).
11. Dimova T, Brouwer M, Gosselin F, Tassignon J, Leo O, Donner C, et al. Effector Vgamma9Vdelta2T cells dominate the human fetal gammadelta T-cell repertoire. *Proc Natl Acad Sci USA.* (2015) 112:E556–65. doi: 10.1073/pnas.1412058112
12. McCarthy NE, Eberl M. Human gammadelta T-cell control of mucosal immunity and inflammation. *Front Immunol.* (2018) 9:985. doi: 10.3389/fimmu.2018.00985
13. Chien YH, Meyer C, Bonneville M. Gammadelta T cells: first line of defense and beyond. *Annu Rev Immunol.* (2014) 32:121–55. doi: 10.1146/annurev-immunol-032713-120216
14. Lawand M, Dechanet-Merville J, Dieu-Nosjean MC. Key features of gamma-delta T-cell subsets in human diseases and their immunotherapeutic implications. *Front Immunol.* (2017) 8:761. doi: 10.3389/fimmu.2017.00761
15. Sherwood AM, Desmarais C, Livingston RJ, Andriessen J, Haussler M, Carlson CS, et al. Deep sequencing of the human TCRgamma and TCRbeta repertoires suggests that TCRbeta rearranges after alphabeta and gammadelta T cell commitment. *Sci Transl Med.* (2011) 3:90ra61. doi: 10.1126/scitranslmed.3002536
16. Davey MS, Willcox CR, Hunter S, Kasatskaya SA, Remmerswaal EBM, Salim M, et al. The human Vdelta2(+) T-cell compartment comprises distinct innate-like Vgamma9(+) and adaptive Vgamma9(-) subsets. *Nat Commun.* (2018) 9:1760. doi: 10.1038/s41467-018-04076-0
17. Lang F, Peyrat MA, Constant P, Davodeau F, David-Ameline J, Poquet Y, et al. Early activation of human V gamma 9V delta 2 T cell broad cytotoxicity and TNF production by nonpeptidic mycobacterial ligands. *J Immunol.* (1995) 154:5986–94.
18. Shen Y, Zhou D, Qiu L, Lai X, Simon M, Shen L, et al. Adaptive immune response of Vgamma2Vdelta2+ T cells during mycobacterial infections. *Science* (2002) 295:2255–8. doi: 10.1126/science.1068819
19. Mazzola TN, Da Silva MT, Moreno YM, Lima SC, Carniel EF, Morcillo AM, et al. Robust gammadelta+ T cell expansion in infants immunized at birth with BCG vaccine. *Vaccine* (2007) 25:6313–20. doi: 10.1016/j.vaccine.2007.06.039
20. Tastan Y, Arvas A, Demir G, Alikasifoglu M, Gur E, Kiray E. Influence of Bacillus Calmette-Guerin vaccination at birth and 2 months old age on the peripheral blood T-cell subpopulations [gamma/delta and alpha-beta T cell]. *Pediatr Allergy Immunol.* (2005) 16:624–9. doi: 10.1111/j.1399-3038.2005.00329.x
21. Spencer CT, Abate G, Sakala IG, Xia M, Truscott SM, Eickhoff CS, et al. Granzyme A produced by gamma(9)delta(2) T cells induces human macrophages to inhibit growth of an intracellular pathogen. *PLoS Pathog.* (2013) 9:e1003119. doi: 10.1371/journal.ppat.1003119
22. Pauza CD, Poonia B, Li H, Cairo C, Chaudhry S. Gammadelta T cells in HIV disease: past, present, and future. *Front Immunol.* (2014) 5:687. doi: 10.3389/fimmu.2014.00687
23. Fenoglio D, Poggi A, Catellani S, Battaglia F, Ferrera A, Setti M, et al. Vdelta2 T lymphocytes producing IFN-gamma and IL-17 are expanded in HIV-1-infected patients and respond to *Candida albicans*. *Blood* (2009) 113:6611–8. doi: 10.1182/blood-2009-01-198028
24. Couzi L, Pitard V, Sicard X, Garrigue I, Hawchar O, Merville P, et al. Antibody-dependent anti-cytomegalovirus activity of human gammadelta T cells expressing CD16 (FcgammaRIIIa). *Blood* (2012) 119:1418–27. doi: 10.1182/blood-2011-06-363655
25. Barisa M, Kramer AM, Majani Y, Moulding D, Saraiva L, Bajaj-Elliott M, et al. E. coli promotes human Vgamma9Vdelta2 T cell transition from cytokine-producing bactericidal effectors to professional phagocytic killers in a TCR-dependent manner. *Sci Rep.* (2017) 7:2805. doi: 10.1038/s41598-017-02886-8
26. Roussillon C, Agrapart M, Guglielmi P, Bensussan A, Brasseur P, Ballet JJ. Human TcR gamma delta+ lymphocyte response on primary exposure to *Plasmodium falciparum*. *Clin Exp Immunol.* (1994) 95:91–7.
27. Howard J, Loizon S, Tyler CJ, Duluc D, Moser B, Mechain M, et al. The antigen-presenting potential of Vgamma9Vdelta2 T cells during *Plasmodium falciparum* blood-stage infection. *J Infect Dis.* (2017) 215:1569–79. doi: 10.1093/infdis/jix149
28. Goodier M, Krause-Jauer M, Sanni A, Massougbdji A, Sadeler BC, Mitchell GH, et al. Gamma delta T cells in the peripheral blood of individuals from an area of holoendemic *Plasmodium falciparum* transmission. *Trans R Soc Trop Med Hyg.* (1993) 87:692–6.
29. Ho M, Tongtawe P, Kriangkum J, Wimonwattawatee T, Pattanapanyasat K, Bryant L, et al. Polyclonal expansion of peripheral gamma delta T cells in human *Plasmodium falciparum* malaria. *Infect Immun.* (1994) 62:855–62.
30. Hviid L, Kurtzhals JA, Adabayeri V, Loizon S, Kemp K, Goka BQ, et al. Perturbation and proinflammatory type activation of V delta 1(+) gamma delta T cells in African children with *Plasmodium falciparum* malaria. *Infect Immun.* (2001) 69:3190–6. doi: 10.1128/IAI.69.5.3190-3196.2001
31. D’Ombrain MC, Robinson LJ, Stanicic DI, Taraika J, Bernard N, Michon P, et al. Association of early interferon-gamma production with immunity to clinical malaria: a longitudinal study among Papua New Guinean children. *Clin Infect Dis.* (2008) 47:1380–7. doi: 10.1086/592971
32. Cairo C, Longinaker N, Cappelli G, Leke RG, Ondo MM, Djokam R, et al. Cord blood Vgamma2Vdelta2 T cells provide a molecular marker for the influence of pregnancy-associated malaria on neonatal immunity. *J Infect Dis.* (2014) 209:1653–62. doi: 10.1093/infdis/jit802
33. Stanicic DI, Cutts J, Eriksson E, Fowkes FJ, Rosanas-Urgell A, Siba P, et al. Gammadelta T cells and CD14+ monocytes are predominant cellular sources of cytokines and chemokines associated with severe malaria. *J Infect Dis.* (2014) 210:295–305. doi: 10.1093/infdis/jiu083
34. Jagannathan P, Kim CC, Greenhouse B, Nankya F, Bowen K, Eccles-James I, et al. Loss and dysfunction of Vdelta2(+) gammadelta T cells are associated with clinical tolerance to malaria. *Sci Transl Med.* (2014) 6:251ra117. doi: 10.1126/scitranslmed.3009793
35. Farrington LA, Jagannathan P, McIntyre TI, Vance HM, Bowen K, Boyle MJ, et al. Frequent malaria drives progressive Vdelta2 T-cell loss, dysfunction, and CD16 up-regulation during early childhood. *J Infect Dis.* (2016) 213:1483–90. doi: 10.1093/infdis/jiv600
36. Hsu H, Boudova S, Mvula G, Divala TH, Mungwira RG, Harman C, et al. Prolonged PD1 expression on neonatal Vdelta2 lymphocytes dampens proinflammatory responses: role of epigenetic regulation. *J Immunol.* (2016) 197:1884–92. doi: 10.4049/jimmunol.1600284
37. Schofield L, Ioannidis LJ, Karl S, Robinson LJ, Tan QY, Poole DP, et al. Synergistic effect of IL-12 and IL-18 induces TIM3 regulation of gammadelta T cell function and decreases the risk of clinical malaria in children living in Papua New Guinea. *BMC Med.* (2017) 15:114. doi: 10.1186/s12916-017-0883-8
38. Taniguchi T, Md Mannoor K, Nonaka D, Toma H, Li C, Narita M, et al. A unique subset of gammadelta T cells expands and produces IL-10 in patients with naturally acquired immunity against falciparum malaria. *Front Microbiol.* (2017) 8:1288. doi: 10.3389/fmicb.2017.01288
39. Jagannathan P, Lutwama F, Boyle MJ, Nankya F, Farrington LA, McIntyre TI, et al. Vdelta2+ T cell response to malaria correlates with protection from infection but is attenuated with repeated exposure. *Sci Rep.* (2017) 7:11487. doi: 10.1038/s41598-017-10624-3
40. Teirlinck AC, McCall MB, Roestenberg M, Scholzen A, Woestenenk R, de Mast Q, et al. Longevity and composition of cellular immune responses following experimental *Plasmodium falciparum* malaria infection in humans. *PLoS Pathog.* (2011) 7:e1002389. doi: 10.1371/journal.ppat.1002389

41. Roestenberg M, Teirlinck AC, McCall MB, Teelen K, Makamdop KN, Wiersma J, et al. Long-term protection against malaria after experimental sporozoite inoculation: an open-label follow-up study. *Lancet* (2011) 377:1770–6. doi: 10.1016/S0140-6736(11)60360-7
42. Seder RA, Chang LJ, Enama ME, Zephir KL, Sarwar UN, Gordon IJ, et al. Protection against malaria by intravenous immunization with a nonreplicating sporozoite vaccine. *Science* (2013) 341:1359–65. doi: 10.1126/science.1241800
43. Ishizuka AS, Lyke KE, DeZure A, Berry AA, Richie TL, Mendoza FH, et al. Protection against malaria at 1 year and immune correlates following PfSPZ vaccination. *Nat Med*. (2016) 22:614–23. doi: 10.1038/nm.4110
44. Mordmüller B, Surat G, Lagler H, Chakravarty S, Ishizuka AS, Lalremruata A, et al. Sterile protection against human malaria by chemoattenuated PfSPZ vaccine. *Nature* (2017) 542:445–9. doi: 10.1038/nature21060
45. Lyke KE, Ishizuka AS, Berry AA, Chakravarty S, DeZure A, Enama ME, et al. Attenuated PfSPZ Vaccine induces strain-transcending T cells and durable protection against heterologous controlled human malaria infection. *Proc Natl Acad Sci USA*. (2017) 114:2711–6. doi: 10.1073/pnas.1615324114
46. Zaidi I, Diallo H, Conteh S, Robbins Y, Kolasny J, Orr-Gonzalez S, et al. Gammadelta T cells are required for the induction of sterile immunity during irradiated sporozoite vaccinations. *J Immunol*. (2017) 199:3781–8. doi: 10.4049/jimmunol.1700314
47. Roussilhon C, Agrapart M, Ballet JJ, Bensussan A. T lymphocytes bearing the gamma delta T cell receptor in patients with acute *Plasmodium falciparum* malaria. *J Infect Dis*. (1990) 162:283–5. |
48. Costa G, Loizon S, Guenot M, Mocan I, Halary F, de Saint-Basile G, et al. Control of *Plasmodium falciparum* erythrocytic cycle: gammadelta T cells target the red blood cell-invasive merozoites. *Blood* (2011) 118:6952–62. doi: 10.1182/blood-2011-08-376111
49. Sidiropoulos K, Viteri G, Sevilla C, Juge S, Webber M, Orlic-Milacic M, et al. Reactome enhanced pathway visualization. *Bioinformatics* (2017) 33:3461–7. doi: 10.1093/bioinformatics/btx441
50. Tsuji M, Mombaerts P, Lefrançois L, Nussenzweig RS, Zavala F, Tonegawa S. Gamma delta T cells contribute to immunity against the liver stages of malaria in alpha beta T-cell-deficient mice. *Proc Natl Acad Sci USA*. (1994) 91:345–9.
51. McKenna KC, Tsuji M, Sarzotti M, Sacci JB Jr, Witney AA, Azad AF. $\gamma\delta$ T cells are a component of early immunity against preerythrocytic malaria parasites. *Infect Immun*. (2000) 68:2224–30. doi: 10.1128/IAI.68.4.2224-2230.2000
52. Yañez DM, Batchelder J, van der Heyde HC, Manning DD, Weidanz WP. Gamma delta T-cell function in pathogenesis of cerebral malaria in mice infected with *Plasmodium berghei* ANKA. *Infect Immun*. (1999) 67:446–8. doi: 10.1128/IAI.67.4.446-448.1999
53. Inoue S, Niikura M, Takeo S, Mineo S, Kawakami Y, Uchida A, et al. Enhancement of dendritic cell activation via CD40 ligand-expressing gammadelta T cells is responsible for protective immunity to *Plasmodium* parasites. *Proc Natl Acad Sci USA*. (2012) 109:12129–34. doi: 10.1073/pnas.1204480109
54. Inoue SI, Niikura M, Asahi H, Kawakami Y, Kobayashi F. $\gamma\delta$ T cells modulate humoral immunity against *Plasmodium berghei* infection. *Immunology* (2018). doi: 10.1111/imm.12997. [Epub ahead of print].
55. Van der Heyde HC, Manning DD, Weidanz WP. Role of CD4+ T cells in the expansion of the CD4-, CD8- gamma delta T cell subset in the spleens of mice during blood-stage malaria. *J Immunol*. (1993) 151:6311–7.
56. Weidanz WP, LaFleur G, Brown A, Burns JM JR, Gramaglia I, van der Heyde HC. Gammadelta T cells but not NK cells are essential for cell-mediated immunity against *Plasmodium chabaudi* malaria. *Infect Immun*. (2010) 78:4331–40. doi: 10.1128/IAI.00539-10
57. Langhorne J, Pells S, Eichmann K. Phenotypic characterization of splenic T cells from mice infected with *Plasmodium chabaudi*. *Scand J Immunol*. (1993) 38:521–8.
58. van der Heyde HC, Elloso MM, Roopenian DC, Manning DD, Weidanz WP. Expansion of the CD4-, CD8- gamma delta T cell subset in the spleens of mice during non-lethal blood-stage malaria. *Eur J Immunol*. (1993) 23:1846–50. doi: 10.1002/eji.1830230817
59. McKenna KC, Tsuji M. Comment on “gammadelta T cells are required for the induction of sterile immunity during irradiated sporozoite vaccinations.” *J Immunol*. (2018) 200:1533. doi: 10.4049/jimmunol.1701743
60. Zaidi I, Duffy PE. Response to comment on “gammadelta T cells are required for the induction of sterile immunity during irradiated sporozoite vaccinations.” *J Immunol*. (2018) 200:1533–4. doi: 10.4049/jimmunol.1701799
61. Behr C, Dubois P. Preferential expansion of V gamma 9V delta 2 T cells following stimulation of peripheral blood lymphocytes with extracts of *Plasmodium falciparum*. *Int Immunol*. (1992) 4:361–6.
62. Goodier MR, Lundqvist C, Hammarstrom ML, Troye-Blomberg M, Langhorne J. Cytokine profiles for human V gamma 9+ T cells stimulated by *Plasmodium falciparum*. *Parasite Immunol*. (1995) 17:413–23.
63. Waterfall M, Black A, Riley E. Gammadelta+ T cells preferentially respond to live rather than killed malaria parasites. *Infect Immun*. (1998) 66:2393–8.
64. Jones SM, Goodier MR, Langhorne J. The response of gamma delta T cells to *Plasmodium falciparum* is dependent on activated CD4+ T cells and the recognition of MHC class I molecules. *Immunology* (1996) 89:405–12.
65. Lindberg BG, Merritt EA, Rayl M, Liu C, Parmryd I, Olofsson B, et al. Immunogenic and antioxidant effects of a pathogen-associated prenyl pyrophosphate in *Anopheles gambiae*. *PLoS ONE* (2013) 8:e73868. doi: 10.1371/journal.pone.0073868
66. D’Ombrain MC, Hansen DS, Simpson KM, Schofield L. Gammadelta-T cells expressing NK receptors predominate over NK cells and conventional T cells in the innate IFN-gamma response to *Plasmodium falciparum* malaria. *Eur J Immunol*. (2007) 37:1864–73. doi: 10.1002/eji.200636889
67. Behr C, Poupot R, Peyrat MA, Poquet Y, Constant P, Dubois P, et al. *Plasmodium falciparum* stimuli for human gammadelta T cells are related to phosphorylated antigens of mycobacteria. *Infect Immun*. (1996) 64:2892–6.
68. Goodier MR, Targett GA. Polyclonal T-cell responses to *Plasmodium falciparum* gametocytes in malaria nonexposed donors. *Parasite Immunol*. (1997) 19:419–25.
69. Liu C, Emami SN, Pettersson J, Ranford-Cartwright L, Faye I, Parmryd I. V γ 9V δ 2 T cells proliferate in response to phosphoantigens released from erythrocytes infected with asexual and gametocyte stage *Plasmodium falciparum*. *Cell Immunol*. (in press). doi: 10.1016/j.cellimm.2018.08.012
70. Harly C, Guillaume Y, Nedellec S, Peigne CM, Monkkinen H, Monkkinen J, et al. Key implication of CD277/butyrophilin-3 (BTN3A) in cellular stress sensing by a major human gammadelta T-cell subset. *Blood* (2012) 120:2269–79. doi: 10.1182/blood-2012-05-430470
71. Boutin L, Scotet E. Towards deciphering the hidden mechanisms that contribute to the antigenic activation process of human Vgamma9Vdelta2 T cells. *Front Immunol*. (2018) 9:828. doi: 10.3389/fimmu.2018.00828
72. Rhodes DA, Chen HC, Price AJ, Keeble AH, Davey MS, James LC, et al. Activation of human gammadelta T cells by cytosolic interactions of BTN3A1 with soluble phosphoantigens and the cytoskeletal adaptor periplakin. *J Immunol*. (2015) 194:2390–8. doi: 10.4049/jimmunol.1401064
73. Sandstrom A, Peigne CM, Leger A, Crooks JE, Konczak F, Gesnel MC, et al. The intracellular B30.2 domain of butyrophilin 3A1 binds phosphoantigens to mediate activation of human Vgamma9Vdelta2 T cells. *Immunity* (2014) 40:490–500. doi: 10.1016/j.immuni.2014.03.003
74. Wang H, Morita CT. Sensor Function for Butyrophilin 3A1 in Prenyl Pyrophosphate Stimulation of Human Vgamma2Vdelta2 T Cells. *J Immunol*. (2015) 195:4583–94. doi: 10.4049/jimmunol.1500314
75. Di Marco Barros R, Roberts NA, Dart RJ, Vantourout P, Jandke A, Nussbaumer O, et al. Epithelia use butyrophilin-like molecules to shape organ-specific gammadelta T cell compartments. *Cell* (2016) 167:203–18 e17. doi: 10.1016/j.cell.2016.08.030
76. Farouk SE, Mincheva-Nilsson L, Krensky AM, Dieli F, Troye-Blomberg M. Gamma delta T cells inhibit *in vitro* growth of the asexual blood stages of *Plasmodium falciparum* by a granule exocytosis-dependent cytotoxic pathway that requires granulysin. *Eur J Immunol*. (2004) 34:2248–56. doi: 10.1002/eji.200424861
77. Guenot M, Loizon S, Howard J, Costa G, Baker DA, Mohabeer SY, et al. Phosphoantigen burst upon *Plasmodium falciparum* schizont rupture can instantly activate Vgamma9Vdelta2 T Cells. *Infect Immun*. (2015) 83:3816–24. doi: 10.1128/IAI.00446-15
78. Elloso MM, van der Heyde HC, vande Waa JA, Manning DD, Weidanz WP. Inhibition of *Plasmodium falciparum in vitro* by human gamma delta T cells. *J Immunol*. (1994) 153:1187–94.

79. Troye-Blomberg M, Worku S, Tangteerawatana P, Jamshaid R, Soderstrom K, Elghazali G, et al. Human gamma delta T cells that inhibit the *in vitro* growth of the asexual blood stages of the *Plasmodium falciparum* parasite express cytolytic and proinflammatory molecules. *Scand J Immunol.* (1999) 50:642–50.
80. Elloso MM, van der Heyde HC, Troutt A, Manning DD, Weidanz WP. Human gamma delta T cell subset-proliferative response to malarial antigen *in vitro* depends on CD4+ T cells or cytokines that signal through components of the IL-2R. *J Immunol.* (1996) 157:2096–102.
81. Ribot JC, Debarros A, Mancio-Silva L, Pamplona A, Silva-Santos B. B7-CD28 costimulatory signals control the survival and proliferation of murine and human gammadelta T cells via IL-2 production. *J Immunol.* (2012) 189:1202–8. doi: 10.4049/jimmunol.1200268
82. Pichyangkul S, Saengkrai P, Yongvanitchit K, Stewart A, Heppner DG. Activation of gammadelta T cells in malaria: interaction of cytokines and a schizont-associated *Plasmodium falciparum* antigen. *J Infect Dis.* (1997) 176:233–41.
83. McCall MB, Sauerwein RW. Interferon-gamma—central mediator of protective immune responses against the pre-erythrocytic and blood stage of malaria. *J Leukoc Biol.* (2010) 88:1131–43. doi: 10.1189/jlb.0310137
84. Robinson LJ, D'Ombrian MC, Stanicic DI, Taraika J, Bernard N, Richards JS, et al. Cellular tumor necrosis factor, gamma interferon, and interleukin-6 responses as correlates of immunity and risk of clinical *Plasmodium falciparum* malaria in children from Papua New Guinea. *Infect Immun.* (2009) 77:3033–43. doi: 10.1128/IAI.00211-09
85. Hensmann M, Kwiatkowski D. Cellular basis of early cytokine response to *Plasmodium falciparum*. *Infect Immun.* (2001) 69:2364–71. doi: 10.1128/IAI.69.4.2364-2371.2001
86. Horowitz A, Newman KC, Evans JH, Korbel DS, Davis DM, Riley EM. Cross-talk between T cells and NK cells generates rapid effector responses to *Plasmodium falciparum*-infected erythrocytes. *J Immunol.* (2010) 184:6043–52. doi: 10.4049/jimmunol.1000106
87. Schmolka N, Serre K, Grosso AR, Rei M, Pennington DJ, Gomes AQ, et al. Epigenetic and transcriptional signatures of stable versus plastic differentiation of proinflammatory gammadelta T cell subsets. *Nat Immunol.* (2013) 14:1093–100. doi: 10.1038/ni.2702
88. Caccamo N, Meraviglia S, Ferlazzo V, Angelini D, Borsellino G, Poccia F, et al. Differential requirements for antigen or homeostatic cytokines for proliferation and differentiation of human Vgamma9Vdelta2 naive, memory and effector T cell subsets. *Eur J Immunol.* (2005) 35:1764–72. doi: 10.1002/eji.200525983
89. Elloso MM, Wallace M, Manning DD, Weidanz WP. The effects of interleukin-15 on human gammadelta T cell responses to *Plasmodium falciparum* *in vitro*. *Immunol Lett.* (1998) 64:125–32.
90. Li Q, Dong C, Deng A, Katsumata M, Nakadai A, Kawada T, et al. Hemolysis of erythrocytes by granulysin-derived peptides but not by granulysin. *Antimicrob Agents Chemother.* (2005) 49:388–97. doi: 10.1128/AAC.49.1.388-397.2005
91. Lafont V, Sanchez F, Laprevotte E, Michaud HA, Gros L, Eliaou JF, et al. Plasticity of gammadelta T cells: impact on the anti-tumor response. *Front Immunol.* (2014) 5:622. doi: 10.3389/fimmu.2014.00622
92. Vermijlen D, Ellis P, Langford C, Klein A, Engel R, Willmann K, et al. Distinct cytokine-driven responses of activated blood $\gamma\delta$ T cells: insights into unconventional T cell pleiotropy. *J Immunol.* (2007) 178:4304–14. doi: 10.4049/jimmunol.178.7.4304
93. Himoudi N, Morgenstern DA, Yan M, Vernay B, Saraiva L, Wu Y, et al. Human gammadelta T lymphocytes are licensed for professional antigen presentation by interaction with opsonized target cells. *J Immunol.* (2012) 188:1708–16. doi: 10.4049/jimmunol.1102654
94. Zhu Y, Wang H, Xu Y, Hu Y, Chen H, Cui L, et al. Human gammadelta T cells augment antigen presentation in Listeria Monocytogenes infection. *Mol Med.* (2016) 22: 737–46. doi: 10.2119/molmed.2015.00214
95. Brandes M, Willmann K, Moser B. Professional antigen-presentation function by human gammadelta T Cells. *Science* (2005) 309:264–8. doi: 10.1126/science.1110267
96. Brandes M, Willmann K, Bioley G, Levy N, Eberl M, Luo M, et al. Cross-presenting human gammadelta T cells induce robust CD8+ alphabeta T cell responses. *Proc Natl Acad Sci USA.* (2009) 106:2307–12. doi: 10.1073/pnas.0810059106
97. Lewis MD, Langhorne J. A new link between gammadelta T cells and myeloid cells in malaria? *Immunity* (2018) 48:193–5. doi: 10.1016/j.immuni.2018.02.003
98. Chen Q, Wen K, Lv A, Liu M, Ni K, Xiang Z, et al. Human Vgamma9Vdelta2-T Cells synergize CD4(+) T follicular helper cells to produce influenza virus-specific antibody. *Front Immunol.* (2018) 9:599. doi: 10.3389/fimmu.2018.00599
99. Cardone J, Le Fric G, Vantourout P, Roberts A, Fuchs A, Jackson I, et al. Complement regulator CD46 temporally regulates cytokine production by conventional and unconventional T cells. *Nat Immunol.* (2010) 11:862–71. doi: 10.1038/ni.1917
100. Sabatos CA, Chakravarti S, Cha E, Schubart A, Sanchez-Fueyo A, Zheng XX, et al. Interaction of Tim-3 and Tim-3 ligand regulates T helper type 1 responses and induction of peripheral tolerance. *Nat Immunol.* (2003) 4:1102–10. doi: 10.1038/ni988
101. Sanchez-Fueyo A, Tian J, Picarella D, Domenig C, Zheng XX, Sabatos CA, et al. Tim-3 inhibits T helper type 1-mediated auto- and alloimmune responses and promotes immunological tolerance. *Nat Immunol.* (2003) 4:1093–101. doi: 10.1038/ni987
102. Finney CA, Ayi K, Wasmuth JD, Sheth PM, Kaul R, Loutfy M, et al. HIV infection deregulates Tim-3 expression on innate cells: combination antiretroviral therapy results in partial restoration. *J Acquir Immune Defic Syndr.* (2013) 63:161–7. doi: 10.1097/QAI.0b013e318285cf13
103. Hou N, Zou Y, Piao X, Liu S, Wang L, Li S, et al. T-cell immunoglobulin- and mucin-domain-containing molecule 3 signaling blockade improves cell-mediated immunity against malaria. *J Infect Dis.* (2016) 214:1547–56. doi: 10.1093/infdis/jiw428
104. Lalor SJ, McLoughlin RM. Memory gammadelta T Cells—newly appreciated protagonists in infection and immunity. *Trends Immunol.* (2016) 37:690–702. doi: 10.1016/j.it.2016.07.006
105. Pitard V, Roumanes D, Lafarge X, Couzi L, Garrigue I, Lafon ME, et al. Long-term expansion of effector/memory Vdelta2-gammadelta T cells is a specific blood signature of CMV infection. *Blood* (2008) 112:1317–24. doi: 10.1182/blood-2008-01-136713
106. Ifrim DC, Quintin J, Joosten LA, Jacobs C, Jansen T, Jacobs L, et al. Trained immunity or tolerance: opposing functional programs induced in human monocytes after engagement of various pattern recognition receptors. *Clin Vaccine Immunol.* (2014) 21:534–45. doi: 10.1128/CVI.00688-13
107. Saeed S, Quintin J, Kerstens HH, Rao NA, Aghajani-farah A, Matarese F, et al. Epigenetic programming of monocyte-to-macrophage differentiation and trained innate immunity. *Science* (2014) 345:1251086. doi: 10.1126/science.1251086
108. Schrum JE, Crabtree JN, Dobbs KR, Kiritsy MC, Reed GW, Gazzinelli RT, et al. Cutting edge: *Plasmodium falciparum* induces trained innate immunity. *J Immunol.* (2018) 200:1243–8. doi: 10.4049/jimmunol.1701010
109. Perera MK, Carter R, Goonewardene R, Mendis KN. Transient increase in circulating gamma/delta T cells during *Plasmodium vivax* malarial paroxysms. *J Exp Med.* (1994) 179:311–5.
110. Arts RJW, Carvalho A, La Rocca C, Rodrigues F, Silvestre R, et al. Immunometabolic pathways in BCG-induced trained immunity. *Cell Rep.* (2016) 17:2562–71. doi: 10.1016/j.celrep.2016.11.011
111. Keitany GJ, Kim KS, Krishnamurthy AT, Hondowicz BD, Hahn WO, Dambrauskas N, et al. Blood Stage malaria disrupts humoral immunity to the pre-erythrocytic stage circumsporozoite protein. *Cell Rep.* (2016) 17:3193–205. doi: 10.1016/j.celrep.2016.11.060
112. Obeng-Adjey N, Portugal S, Tran TM, Yazew TB, Skinner J, Li S, et al. Circulating Th1-cell-type Tfh cells that exhibit impaired b cell help are preferentially activated during acute malaria in children. *Cell Rep.* (2015) 13:425–39. doi: 10.1016/j.celrep.2015.09.004
113. Ryg-Cornejo V, Ioannidis LJ, Ly A, Chiu CY, Tellier J, Hill DL, et al. Severe malaria infections impair germinal center responses by inhibiting t follicular helper cell differentiation. *Cell Rep.* (2016) 14:68–81. doi: 10.1016/j.celrep.2015.12.006

114. Portugal S, Moebius J, Skinner J, Doumbo S, Doumtable D, Kone Y, et al. Exposure-dependent control of malaria-induced inflammation in children. *PLoS Pathog.* (2014) 10:e1004079. doi: 10.1371/journal.ppat.1004079
115. Huang D, Chen CY, Ali Z, Shao L, Shen L, Lockman HA, et al. Antigen-specific Vgamma2Vdelta2 T effector cells confer homeostatic protection against pneumonic plaque lesions. *Proc Natl Acad Sci USA.* (2009) 106:7553–8. doi: 10.1073/pnas.0811250106
116. Dieli F, Caccamo N, Meraviglia S. Advances in immunotherapy of castration-resistant prostate cancer: bisphosphonates, phosphoantigens and more. *Curr Opin Investig Drugs* (2008) 9:1089–94.
117. Wilhelm M, Kunzmann V, Eckstein S, Reimer P, Weissinger F, Ruediger T, et al. Gammadelta T cells for immune therapy of patients with lymphoid malignancies. *Blood* (2003) 102:200–6. doi: 10.1182/blood-2002-12-3665
118. Alexandroff A, Black J, Bollina P, James K. Differential production of gamma delta T cells in the urine of bladder cancer patients receiving Bacillus Calmette Guerin immunotherapy. *Int J Oncol.* (1997) 10:387–93.
119. Tanaka Y, Iwasaki M, Murata-Hirai K, Matsumoto K, Hayashi K, Okamura H, et al. Anti-tumor activity and immunotherapeutic potential of a bisphosphonate prodrug. *Sci Rep.* (2017) 7:5987. doi: 10.1038/s41598-017-05553-0
120. Bhat J, Kouakanou L, Peters C, Yin Z, Kabelitz D. Immunotherapy With Human Gamma Delta T Cells-Synergistic Potential of Epigenetic Drugs? *Front Immunol* (2018) 9:512. doi: 10.3389/fimmu.2018.00512
121. Cendron D, Ingoure S, Martino A, Casetti R, Horand F, Romagne F, et al. A tuberculosis vaccine based on phosphoantigens and fusion proteins induces distinct gammadelta and alphabeta T cell responses in primates. *Eur J Immunol.* (2007) 37:549–65. doi: 10.1002/eji.200636343

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Dantzler and Jagannathan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.