

# Conservation priorities for terrestrial mammals in Dobrogea Region, Romania

Iulia V. Miu<sup>1</sup>, Gabriel B. Chisamera<sup>2</sup>, Viorel D. Popescu<sup>1,3</sup>, Ruben Iosif<sup>4</sup>,  
Andreea Nita<sup>1</sup>, Steluta Manolache<sup>1</sup>, Viorel D. Gavril<sup>5,6</sup>,  
Ioana Cobzaru<sup>6</sup>, Laurentiu Rozyłowicz<sup>1</sup>

**1** University of Bucharest, Center for Environmental Research and Impact Studies, 1 N. Balcescu, 010041, Bucharest, Romania **2** National Museum of Natural History Grigore Antipa, 1 Kiseleff Blvd., 011341, Bucharest, Romania **3** Department of Biological Sciences, Ohio University, Athens, OH, USA **4** University Ovidius Constanța, Faculty of Natural Sciences and Agricultural Sciences, 1 Al. Universității, corp B, 900470, Constanța, Romania **5** University of Bucharest, Faculty of Biology, 91–95 Splaiul Independenței, 050095, Bucharest, Romania **6** Romanian Academy, Institute of Biology, 296 Splaiul Independenței, 060031 Bucharest, Romania

Corresponding author: Iulia V. Miu (iulia.miu@drd.unibuc.ro)

---

Academic editor: J. Maldonado | Received 27 March 2018 | Accepted 30 August 2018 | Published 23 October 2018

---

<http://zoobank.org/8F7EBA17-46D6-4236-B91B-A3B2DE8CC950>

---

**Citation:** Miu IV, Chisamera GB, Popescu VD, Iosif R, Nita A, Manolache S, Gavril VD, Cobzaru I, Rozyłowicz L (2018) Conservation priorities for terrestrial mammals in Dobrogea Region, Romania. ZooKeys 792: 133–158. <https://doi.org/10.3897/zookeys.792.25314>

---

## Abstract

Based on species occurrence records of museum collections, published literature, and unpublished records shared by mammalian experts, we compiled a distribution database for 59 terrestrial mammals populating the extensively protected Dobrogea Region of Romania. The spatial patterns of mammal distribution and diversity was evaluated and systematic conservation planning applied to identify priority areas for their conservation. The spatial analyses revealed that intensive sampling was not directly correlated to mammal diversity but rather to accessibility for inventory. The spatial prioritisation analysis indicated a relatively aggregated pattern of areas with a high or low conservation value with virtually no connecting corridors between them. The significant overlap between Natura 2000 sites and national protected areas induced an over-optimistic vision of the effectiveness and representativeness of existing Natura 2000 network for species found in Annexes II and IV of the Habitats Directive. These results represent a key step in identifying core areas for the protection of mammal diversity and dispersal corridors for improved connectivity, and to guide future conservation efforts in increasing the effectiveness of the existing protected areas in the context of environmental changes.

**Keywords**

Dobrogea, Natura 2000, species distribution, species richness, systematic conservation planning, terrestrial mammals

**Introduction**

Terrestrial mammals are well-studied taxa, yet their distribution and conservation status are not fully understood (Crooks et al. 2011). Mammalian population decline accelerates the loss of ecosystem services and poses a substantial threat to species diversity at the community level (Ceballos 2002, Rodrigues et al. 2004). Since mammals display diverse traits and can exploit a wide range of ecological niches, they are also effective focal species for conservation, and their population status might be a proxy for both fragmentation and connectivity across landscapes (Crooks et al. 2011).

A common conservation strategy to prevent the loss of biodiversity is the creation of protected areas (Margules and Pressey 2000, Williams et al. 2002). Protected areas must ensure the long-term persistence and viability of species and should ideally support many rare, threatened, or endemic taxa, particularly those with low mobility and high sensitivity to environmental alterations (Rodrigues et al. 2004, Possingham et al. 2006). However, typically, the effectiveness of protected areas is undermined by poor governance (Eklund et al. 2011, Manolache et al. 2018, Nita et al. 2018) and lack of funding and relevant resources (Sánchez-Fernández et al. 2017).

The Natura 2000 network of protected areas of European importance represents one of the most extensive networks of conservation areas worldwide (Nita et al. 2017). Scientists and policymakers often question the effectiveness of this network due to the Member States allocating fewer funds than needed to implement conservation programs (Nita et al. 2017, Sánchez-Fernández et al. 2017). Natura 2000 is more effective in protecting species listed in Birds Directive because of a better overlap between ancillary conservation investments such as Common Agricultural Policy and biodiversity value (Lung et al. 2014, Maiorano et al. 2015), and because birds are more intensely studied than other vertebrate groups. To be more effective, Natura 2000 network must incorporate potential changes in species distributions (Popescu et al. 2013, Kukkala et al. 2016). Failure to acknowledge changes in species ranges may lead to gaps in protecting species that are sensitive to climate change and other anthropogenic pressures (Araújo et al. 2011).

One of Romania's legal obligations since joining the European Union in 2007 was to designate Natura 2000 sites in a short time (Ioja et al. 2010). Due to the lack of adequate species and habitat distribution data, regions that already benefited from protection under national laws were preferred for the first phase of the designation process. Consequently, the EU conservation goals were not met, which resulted in the designation of additional protected areas (Ioja et al. 2010, Popescu et al. 2013) and a disproportionate increase of land protected in some regions such as it is the case of Dobrogea (now 63% under protection, 9700 km<sup>2</sup>). The Natura 2000 network from Dobrogea includes 67 sites (35 Sites of Community Importance - SCI and 32 Special

Protection Areas – SPA, most of the SCIs and SPAs spatially overlap). Within Dobrogea, highlands and floodplains gained extensive protection while lowlands occupied by arable lands remained largely unprotected. However, the latter areas are inhabited by endangered species such as the marbled polecat (*Vormela peregusna*) and the steppe polecat (*Mustela eversmanii*) (Murariu et al. 2009, 2010).

Due to the diverse landforms, climatic influences, and habitats, Dobrogea harbours a large number of mammal species (Murariu 1996, Murariu et al. 2010). To date, 59 mammal species have been documented in this region, three of which reach the outer limit of their geographic range (the marbled polecat *Vormela peregusna*, the stoat *Mustela erminea*, and the common hamster *Cricetus cricetus*), and two other species have their core range in Dobrogea (the Romanian hamster *Mesocricetus newtoni* and the Southern birch mouse *Sicista nordmanni*) (Bunescu 1959, 1961, Popescu and Murariu 2001, Murariu and Munteanu 2005). Of the 59 mammal species, 14 are protected by Habitats Directive. Despite the focus of many Natura 2000 sites within Dobrogea on protecting mammal species, limited and outdated distributional databases are available for individual species, e.g., the Romanian hamster *Mesocricetus newtoni* (Hamar and Schutowa 1966), the Eurasian beaver *Castor fiber* (Kiss et al. 2012, Kiss et al. 2014), the European mink *Mustela lutreola* (Cuzic and Marinov 2004), and the Southern birch mouse *Sicista nordmanni* (Ausländer and Hellwing 1957). Moreover, with few exceptions, (e.g., Murariu 1996, 2006, Murariu et al. 2009, Murariu et al. 2010) the Dobrogea Region lacks actual regional species distribution data.

One tool supporting management decisions and for investigating species population coverage within protected areas is spatial conservation prioritisation (Pouzols et al. 2014). As part of systematic conservation planning (Margules and Pressey 2000) and accounting for complementarity, spatial prioritisation can be an efficient instrument in identifying spatial priorities and in achieving conservation goals (Pressey et al. 2007) even in broadly protected and underfunded regions such as Dobrogea (Rozyłowicz et al. 2017). In this study, we evaluate priority areas for mammal conservation in Dobrogea, Romania and assess the spatial patterns of distribution and diversity of terrestrial mammals by: (1) compiling mammal distribution records from published papers, museum records, and unpublished data, (2) analyzing spatial patterns of distribution data, and (3) using systematic conservation planning in identifying high priority areas for conservation of terrestrial mammal listed in Annexes II and IV of Habitats Directive within the regional Natura 2000 network.

## Materials and methods

### Mammal species occurrences


To map the distribution of mammals in Dobrogea, we extracted species occurrence records from three primary sources: museum collections, published data, and unpublished field data. Occurrences that could not be georeferenced to a location (e.g., as-

signed to a large watershed or geographical province), or associated with unspecified taxa within genera, were not included in this geodatabase. The species taxonomy considered in this paper is based on Wilson and Reeder (2005) and Arslan et al. (2016). Sibling species which are difficult to discriminate in the field, such as the yellow-necked mouse *Apodemus flavicollis*, the wood mouse *Apodemus sylvaticus* (Bartolommei et al. 2016), the common vole *Microtus arvalis*, and the East European vole *Microtus levis* (Jaarola et al. 2004), were included as individual species, as their occurrences were acquired through museum collections and published data. Red list status was based on Temple and Terry (2009).

The dataset used to map the species distribution includes 6724 occurrence records for 59 mammal species. For spatial pattern analyses, we excluded species found exclusively in fenced areas (the European mouflon *Ovis aries musimon*), the vagrant species (elk *Alces alces*), and synanthropic species (the rats *Rattus rattus*, *Rattus norvegicus* and the house mouse *Mus musculus* (Table 2)), resulting in 5593 occurrence records for 54 species. For creating species distribution maps, we aggregated the occurrence records at a Universal Traverse Mercator spatial resolution of 25 km<sup>2</sup> (UTM 5 × 5 km). Following Cogalniceanu et al. (2013), the occurrence records were classified based on the year of observation into *old records*, if recorded before 1990, and *recent records*, if recorded after 1990. For spatial pattern analyses, we increased the cell size to UTM 10 × 10 km, allowing us to highlight regional patterns in richness, rarity, and dissimilarity, and to reduce the potential bias in sampling (Graham and Hijmans 2006). For spatial prioritisation of mammal conservation within Natura 2000 sites, we used the UTM 5 × 5 km occurrences maps of 14 native species listed in Annexes II and IV of Habitats Directive (Figure 1).

### Spatial bias in species occurrence

Potential bias at the scale of the study area was assessed using the overall spatial autocorrelation in mammal records per 5 × 5 km grid cell. We used Global Moran's I test (Fortin and Dale 2005) to evaluate spatial pattern of sampling per grid cell being significantly clustered ( $Z > 0$ ) or dispersed ( $Z < 0$ ) across Dobrogea. To assess the local patterns of sampling bias we used the Getis Ord  $G_i^*$  spatial statistic. This analysis identifies clusters of records with values numerically higher than expected by random chance within a specified searching distance (Ord and Getis 1995). The distance threshold for the aggregation patterns was set up to 7100 m to include the neighbouring eight grid cells for each UTM grid of interest. The Getis Ord  $G_i^*$  test returns a Z-score for every cell, which, depending on the level of aggregation describes spatial clusters of high or low sampling effort. We identified clusters of UTM 5 × 5 km cells where the sampling effort was significantly higher (hotspots of occurrence,  $G_iZScore > 1.87$ ) or lower (cold spots of occurrence,  $G_iZScore < 1.87$ ). All spatial analyses were performed using ARCMAP 10.3 (ESRI, CA) (Figure 1).



**Figure 1.** Flowchart of diversity analyses and spatial prioritisation of conservation of terrestrial mammals within Dobrogea Region, Romania.

**Estimating species richness, rarity, and dissimilarity**

To emphasise regional patterns of richness, rarity, and dissimilarity of mammals of Dobrogea, we aggregated the occurrence records at 5 × 5 km and 10 × 10 km and imported them into BIODIVERSE software (v. 1.1) (Laffan et al. 2010), a tool for spatial analysis of biodiversity (Figure 1).

Richness index was measured as the number of species in each grid cell. Species rarity was assessed by dividing the corrected weighted rarity (CWE) by the total number of species in the respective cell, where CWE is (Equation 1).

$$CWE = WE / Richness \tag{1}$$

Weighted rarity (WE) of a species represents the occurrence records of sample counts of the respective species divided by the number of occurrence records of all species in the dataset (Equation 2).

$$WE = \sum_{t \in T} \frac{S_t}{S_t} \tag{2}$$

where  $t$  is a taxon in the set of taxa  $T$  across neighbourhood set 1,  $s_t$  is the sum of the sample counts for  $t$  across the elements in neighbouring sets 1 and 2, and  $S_t$  represents the total number of samples across the data set for  $t$  (Laffan et al. 2010). In our case, only one neighbouring set is specified.

To calculate the differences in species composition across Dobrogea, we used the turnover index ( $S_2$ ), which refers to changes in species composition from one community to another along a gradient and across different sites (Whittaker 1972).  $S_2$  calculates the dissimilarity between two sets of species. We compared a focal quadrat with one of its eight neighbours (Equation 3).

where  $a$  is the total number of species found in both neighbour sets,  $b$  is the number of species unique to the neighbour set 1, and  $c$  is the number of species unique to the neighbour set 2 (Laffan et al. 2010).

$$S_2 = 1 - \frac{a}{a + \min(b, c)} \quad (3)$$

Selecting the smallest values of  $b$  or  $c$  in the  $S_2$  equation denominator reduces the impact of imbalances of species richness on neighbour dissimilarity. The highest value that  $S_2$  can result is the value of one (1), which indicates the focal quadrat has no species in common with any neighbour and the lowest possible value is zero (0), indicating that all quadrats have an identical set of species (Lennon et al. 2001).

### Identifying high-priority areas for Natura 2000 mammal species conservation

To identify high-priority areas for mammal species conservation across Natura 2000 sites within Dobrogea Region, we used systematic conservation planning software ZONATION v4 (Lehtomäki and Moilanen 2013, Moilanen et al. 2014). This software uses a complementarity-based algorithm including connectivity, with the result that landscapes can be zoned according to their conservation potential. Using a deterministic iterative process, ZONATION creates a hierarchical ranking of the landscape from the highest to the lowest conservation value (Moilanen et al. 2014).

For priority analysis, we used  $5 \times 5$  km raster layers for presence/absence data for 14 mammal species listed in Annexes II and IV of the Habitats Directive and a hierarchic mask of the Natura 2000 Sites of Community Importance within Dobrogea Region (Figure 1). A hierarchic mask represents a mask layer specifying priority land uses, in our case the Natura 2000 network. This planning design forces the prioritisation algorithm to undertake ranking cells outside the Natura 2000 network, followed by ranking those in the Natura 2000 network, allowing the application to analyse an optimal conservation area network. We sequenced the prioritisation model using an additive benefit function with exponent  $z = 0.25$ , which is a default value representing the exponent of the species-area curve (Moilanen et al. 2014). In this prioritisation model, the function sums the loss across features, converted

via feature-specific benefit functions, giving high importance to the cells containing many species (Arponen et al. 2005).

The outputs of the analysis are conservation priority ranking of the landscape, derived from the order of iterative cell ranking whereby each grid cell has a value between 0 and 1, indicating that ranking close to 0 are removed first (low priority), while ranking close to 1 are retained until the end of the iteration. The outputs show the most important areas for mammal species conservation across Natura 2000 sites and a set of curves describing the absolute performance levels of species conservation. We considered as high-priority areas for conservation, all grid cells falling in the top 20% of the predicted priority ranks, a proportion that maximises mammal species representation at the regional level (Arponen et al. 2005). Suppl. material 3 presents the methodology used to identify high-priority Natura 2000 sites with Zonation v4.

The data underpinning the analysis reported in this paper are deposited at GBIF, the Global Biodiversity Information Facility, <http://ipt.pensoft.net/resource?r=mammalsdobrogea>.

## Results

### Mammal species occurrences in Dobrogea

We collected 4451 records from published museum collections data (66%), 1326 personal records shared by experts (20%), and 947 records from other papers reporting the results of fauna inventories (14%). Of all the accessible papers (published museum collections and fauna inventories) 67% were published before the year 1990 and 33% after 1990 (Suppl. material 1). Over 54% of all the records were reported before 1990, and 46% are records collected after 1990. Occurrences maps for 59 mammal species aggregated at 5 × 5 km resolution are presented in Suppl. material 2.

The rate of accumulation of mammal occurrences increased in 1956 by 688 records, due to the rediscovery of the Southern birch mouse (*Sicista nordmanni*) at Valu lui Traian in 1955. That report attracted additional fieldwork by mammalogists the following year, consequently, an increase in the number of records for other rodent species. After 1990, and up to 2017, the peak number of records per year took place in 2007 with 456 new records (Figure 2, Table 1).

### Spatial patterns in mammal species occurrences in Dobrogea

Of 757 UTM 5 × 5 km grid cells encompassing the Dobrogea Region, only 335 grid cells (i.e., 44%) include reported mammal sightings (Figure 3). At the regional scale, Global Moran's I test indicated a random pattern in the number of mammal occurrences per UTM 5 × 5 grid cell ( $Z = 1.87$ ,  $p = 0.06$ ). However, the local Getis Ord  $G_i^*$

**Table I.** Checklist of mammals of Dobrogea Region, Romania.

Order	Family	Species	Total number of records	New records (after 1990)	Total number of UTM 5 × 5 occupied cells	Habitats Directive Annexes	European Red List status	
Rodentia	Sciuridae	<i>Sciurus vulgaris</i> (Linnaeus, 1758)	6	3	4	–	Least concern	
		<i>Spermophilus citellus</i> (Linnaeus, 1766)	214	92	95	II/IV	Vulnerable	
	Gliridae	<i>Dryomys nitedula</i> (Pallas, 1778)	35	22	20	IV	Least concern	
		<i>Muscardinus avellanarius</i> (Linnaeus, 1758)	1	1	1	–	Least concern	
		<i>Glis glis</i> (Linnaeus, 1766)	3	3	3	–	Least concern	
	Castoridae	<i>Castor fiber</i> (Linnaeus, 1758)	12	12	5	II/IV	Least concern	
	Dipodidae	<i>Sicista nordmanni</i> (Keyserling & Blasius, 1840)	76	0	2	II/IV	Vulnerable	
	Spalacidae	<i>Nannospalax leucodon</i> (Nordmann, 1840)	163	82	57	–	Least concern	
	Cricetidae	<i>Cricetus cricetus</i> (Linnaeus, 1758)	2	0	1	IV	Least concern	
		<i>Mesocricetus newtoni</i> (Nehring, 1898)	98	13	31	II/IV	Near threatened	
		<i>Ondatra zibethicus</i> (Linnaeus, 1766)	87	37	57	–	Invasive	
		<i>Arvicola amphibius</i> (Linnaeus, 1758)	29	12	22	–	Least concern	
		<i>Microtus agrestis</i> (Linnaeus, 1761)	28	11	18	–	Least concern	
		<i>Microtus arvalis</i> (Pallas, 1779)	187	40	44	–	Least concern	
		<i>Microtus levis</i> (Miller, 1908)	29	9	13	–	Least concern	
		<i>Microtus subterraneus</i> (Sclys-Longchamps, 1836)	16	7	9	–	Least concern	
		<i>Myodes glareolus</i> (Schreber, 1780)	1	0	1	–	Least concern	
		Muridae	<i>Micromys minutus</i> (Pallas, 1771)	36	15	24	–	Least concern
			<i>Apodemus agrarius</i> (Pallas, 1771)	451	96	50	–	Least concern
			<i>Apodemus flavicollis</i> (Melchior, 1834)	134	80	34	–	Least concern
			<i>Apodemus sylvaticus</i> (Linnaeus, 1758)	1327	330	65	–	Least concern
	<i>Apodemus uralensis</i> (Pallas, 1811)		16	6	8	–	Least concern	
	<i>Mus spicilegus</i> (Petényi, 1882)		20	20	19	–	Least concern	
Myocastoridae	<i>Myocastor coypus</i> (Molina, 1782)	5	2	5	–	Invasive		
Lagomorpha	Leporidae	<i>Lepus europaeus</i> (Pallas, 1778)	262	255	102	–	Least concern	
Erinaceomorpha	Erinaceidae	<i>Erinaceus roumanicus</i> (Barrett-Hamilton, 1900)	52	40	39	–	Least concern	




Order	Family	Species	Total number of records	New records (after 1990)	Total number of UTM 5 × 5 occupied cells	Habitats Directive Annexes	European Red List status
Soricomorpha	Soricidae	<i>Crocidura leucodon</i> (Hermann, 1780)	85	14	18	–	Least concern
		<i>Crocidura suaveolens</i> (Pallas, 1811)	131	40	36	–	Least concern
		<i>Neomys anomalus</i> (Cabrer, 1907)	12	4	8	–	Least concern
		<i>Neomys fodiens</i> (Pennant, 1771)	5	1	4	–	Least concern
		<i>Sorex araneus</i> (Linnaeus, 1758)	63	14	25	–	Least concern
		<i>Sorex minutus</i> (Linnaeus, 1766)	15	9	10	–	Least concern
	Talpidae	<i>Talpa europaea</i> (Linnaeus, 1758)	65	53	55	–	Least concern
	Carnivora	Felidae	<i>Felis silvestris</i> (Schreber, 1777)	101	94	52	IV
Felidae		<i>Lynx lynx</i> (Linnaeus, 1758)	2	1	2	II/IV	Least concern
Canidae		<i>Canis aureus</i> (Linnaeus, 1758)	214	198	94	–	Least concern
		<i>Canis lupus</i> (Linnaeus, 1758)	27	22	14	II/IV	Least concern
		<i>Nyctereutes procyonoides</i> (Gray, 1834)	87	35	41	–	Invasive
		<i>Vulpes vulpes</i> (Linnaeus, 1758)	230	223	122	–	Least concern
Mustelidae		<i>Mustela erminea</i> (Linnaeus, 1758)	25	7	23	IV	Vulnerable
		<i>Mustela eversmannii</i> (Lesson, 1827)	31	24	25	II/IV	Vulnerable
		<i>Mustela lutreola</i> (Linnaeus, 1761)	119	109	50	II/IV	Endangered
		<i>Mustela nivalis</i> (Linnaeus, 1766)	67	54	50	–	Least concern
		<i>Mustela putorius</i> (Linnaeus, 1758)	89	74	61	–	Least concern
		<i>Vormela peregusna</i> (Güldenstädt, 1770)	70	16	39	II/IV	Vulnerable
		<i>Martes foina</i> (Erxleben, 1777)	98	97	58	–	Least concern
		<i>Martes martes</i> (Linnaeus, 1758)	36	36	20	–	Least concern
		<i>Meles meles</i> (Linnaeus, 1758)	102	92	60	–	Least concern
	<i>Neovison vison</i> (Schreber, 1777)	2	2	1	–	Invasive	
	<i>Lutra lutra</i> (Linnaeus, 1758)	55	49	35	II/IV	Near threatened	
Artiodactyla	Suidae	<i>Sus scrofa</i> (Linnaeus, 1758)	221	204	105	–	Least concern
	Cervidae	<i>Dama dama</i> (Linnaeus, 1758)	46	29	21	–	Least concern
		<i>Cervus elaphus</i> (Linnaeus, 1758)	38	32	25	–	Least concern
		<i>Capreolus capreolus</i> (Linnaeus, 1758)	262	190	119	–	Least concern

**Table 2.** Checklist of synanthropic and vagrant mammals of Dobrogea Region, Romania.

Order	Family	Species	Total number of records	New records (after 1990)	Total number of UTM 5 × 5 occupied cells
Rodentia	Muridae	<i>Rattus norvegicus</i> (Berkenhout, 1769)	114	49	64
Rodentia	Muridae	<i>Rattus rattus</i> (Linnaeus, 1758)	3	2	3
Rodentia	Muridae	<i>Mus musculus</i> (Linnaeus, 1758)	1001	139	78
Artiodactyla	Cervidae	<i>Alces alces</i> (Linnaeus, 1758)	4	0	3
Artiodactyla	Bovidae	<i>Ovis aries musimon</i> (Pallas, 1881)	9	5	4


spatial statistic indicates 3 hotspots for mammal sightings: Valu lui Traian Biological Research Station (mean  $Z = 7.73$ ), North Dobrogea Plateau Natura 2000 site (mean  $Z = 3.26$ ), and Letea Forest, a natural reserve within Danube Delta (mean  $Z = 2.75$ ). Additionally, there are few moderately sampled regions such as Măcin Mountains National Park in the northwest, Dumbrăveni-Urului Valley-Vederoasa Lake Natura 2000 site and Canaraua-Fetii Iortmac Natura 2000 site in the southwest, and Hagieni – Cotul Văii Forest Natura 2000 site in the southeast (Figure 4).

The mammal occurrences at 5 × 5 km resolution ranged between 1 and 35 reported species per quadrat (Figure 5). The map highlights a lower sampling effort in southern and central Dobrogea, areas with intensive agriculture, and the highest diversity in the northern and southwest parts of Dobrogea, comprising mostly forested habitat.


**Figure 2.** Accumulation of mammals' occurrence records (blue) and the number of records per year (red) within Dobrogea Region, Romania.

### Species richness, rarity, and dissimilarity

When aggregating species records at  $10 \times 10$  km, the number of reported species ranged from 2 to 45 per cell grid, with the highest species diversity located in the northern part of Dobrogea Region overlapping the following Natura 2000 sites: North Dobrogea Plateau with a maximum richness of 45 species, western part of Danube Delta with 39 species and Agighiolului Hills with 38 species. Most of the grid cells with species richness are concentrated in the northern region reflecting an optimal sampling of mammal species (number of species from 29 to 37) (Figure 6), while grid



**Figure 3.** Mammals reported occurrences in Dobrogea Region, Romania at  $5 \times 5$  km resolution. Grids with reported occurrences before 1990 were plotted as old records whereas those with reported occurrences after 1990 were considered new records (reports of synanthropic and vagrant mammals were excluded).




**Figure 4.** Hotspots of sampling efforts within Dobrogea. The numbered statistically significant hot-spots are **1** Valu lui Traian Biological Research Station and Fântânița-Murfatlar **2** North Dobrogea Plateau **3** Letea Forest Natural Reserve in the Danube Delta.

cells with the lowest richness values are distributed in the southern and central part of Dobrogea Region.

Corrected weighted rarity (CWE) varied across Dobrogea from 0.0087 for cell grids with widespread species to 0.62 grid cells with species of restricted distribution. The highest value of corrected weighted rarity can be found in the Danube Delta, specifically in the levee complex of Puiu – Roșu – Lumina, with a value up to 0.62 (Figure 7).

The values of dissimilarity index S2 ranged from 0 to 1 with the highest turnover quadrats in the southern area of Dobrogea where there are low richness zones. The




**Figure 5.** The mammal species richness at  $5 \times 5$  km grid resolution within Dobrogea.

value of 1 implies that the quadrat has no species in common with any neighbour (Figure 8). We found that areas with the higher richness of species have more species in common with their neighbours.


### High-priority areas for conservation within Natura 2000 sites

Based on the Zonation analysis results, the top spatial conservation priorities overlap Danube Delta, North Dobrogea Plateau, and the Măcin Mountains in the northern part of Dobrogea region, where a relatively aggregated pattern of top conservation value areas appear due to their extensive wetland area and forested



**Figure 6.** Mammal species richness of Dobrogea at 10 × 10 km. Grids with high richness partially overlap. **1** North Dobrogea Plateau **2** Danube Delta, and **3** Agighiolului Hills.


habitats. Isolated hotspots are represented by Dumbrăveni-Urliua Valley-Vederoasa Lake in the southwest, Hagieni – Cotul Văii Forest in the southeast, and Cheia Jurassic Reefs in Central Dobrogea. Grid cells with the lowest ranking are located in the central and southern part of Dobrogea Region, where the majority of the regions' agricultural lands are clustered (Figure 9). Nevertheless, the Natura 2000 network encompasses 45% of mammal species distribution listed in Annexes II and IV of the Habitats Directive when top 20% of the landscape is protected by Natura 2000 sites (Figure 10).



**Figure 7.** Corrected weighted rarity map of Dobrogea mammal species.

## Discussion

By using an updated distribution of terrestrial mammals, we identified high priority areas for protecting mammal diversity to guide future conservation efforts in an extensively protected Romanian region. In the broader context of systematic conservation planning, the prioritisation analysis is a useful tool to identify key areas for biodiversity conservation, e.g., where species are more likely to survive (Ferrier and Wintle 2009, Wilson et al. 2009, Kukkala and Moilanen 2013).




**Figure 8.** Dissimilarity map of Dobrogea mammal species, Romania (S2 index).

The number of reported occurrences in Dobrogea varied among species. The largest number of records (20%) are for the wood mouse (*Apodemus sylvaticus*), mostly because they are widespread within the region, have a higher population abundance, and are evidently. The wood mouse may be easily misidentified as a yellow-necked mouse (*Apodemus flavicollis*) (Bartolommei et al. 2016), but it still retains the first rank because of their higher population in Dobrogea (Popescu and Murariu 2001).


The lowest number of records in Dobrogea is recorded for the hazel dormouse (*Muscardinus avellanarius*) and the bank vole (*Myodes glareolus*), with only one record per each species. Two other widespread species, but with an uncharacteristically low number of reported presences are the red fox (*Vulpes vulpes*) and the roe deer (*Capreolus capreolus*),





**Figure 9.** Priority conservation areas for mammal species listed in Annexes II and IV of the Habitats Directive within Natura 2000 sites of Dobrogea. Areas have been graded according to their priority rank, with highest priorities (top 20%) shown in red.

possibly because they are common species, with minor interest for biologists. The research effort for species sampling was focused on selected areas: Măcin Mountains National Park, Danube Delta Biosphere Reserve, and the North Dobrogea Plateau. Here, we recorded a higher than expected number of species occurrences per grid cell, mostly because the long-term protection status attracted faunistic inventory projects over time. Typically, the sampling bias is higher in protected areas because they attract more conservation funds leading to greater efforts for biodiversity research (e.g., Botts et al. 2011). This process describes most of the biodiversity spatial databases (Lobo et al. 2007). A



**Figure 10.** Zonation performance curves quantifying the proportion of remaining species occurrences covered by Natura 2000 sites in Dobrogea. When considering 20% of landscape within Natura 2000 sites as protected (e.g., conservation measures are enforced), 45% of Natura 2000 mammal occurrences are protected.

higher than expected sampling effort also was evident near cities, major roads, and research facilities, which are easily accessible to researchers (e.g., Valu lui Traian Biological Research Station - see Figure 4). We noted a lack of research interest in central and southern Dobrogea, where most of the agricultural landscape is located, and only small patches of natural habitats remain as wildlife refuges (Rey et al. 2007). However, agricultural landscapes are essential for many species protected by Habitats Directive, such as *Vormela peregusna*, *Mustela eversmanii*, *Spermophilus citellus*, *Mesocricetus newtoni*, and *Cricetus cricetus* (Popescu and Murariu 2001, Murariu and Munteanu 2005, Murariu et al. 2009, 2010). Those identified species have restrictive ecological requirements, and hence, the researchers should focus on increasing sampling effort in these neglected areas to draft appropriate conservation plans.

We found that high species richness did not match all the hotspots of sampling efforts, such as in Valu lui Traian Biological Research Station and Fântânița-Murfatlar (location 1 in Figure 4) and Letea Forest Natural Reserve in the Danube Delta (location 3 in Figure 4). This validates the finding that intensive sampling was not directly correlated with mammal diversity, but rather ease of access to the regions (Santos et al. 2017). To better understand the patterns of species richness at the regional level, we expanded the resolution to 10 × 10 km, but the spatial pattern did not change between the two mapping resolutions. In both cases, the highest diversity (35 species at 5 × 5 km resolution, see Figure 5); 45 mammal species at 10 × 10 km resolution, see Figure 6), is found in the northern part of Dobrogea, overlapping North Dobrogea Plateau Natura 2000 site, where habitat heterogeneity is high (Rey et al. 2007).

The spatial turnover index ( $S_2$ ) suggests that areas with lower species richness are dissimilar compared to their neighbours. Notably, we observed some affinities of par-

ticular species towards low species richness areas (Lennon et al. 2001), e.g., species dependent on steppe or agricultural landscape (Popescu and Murariu 2001, Murariu and Munteanu 2005). Typically, the spatial turnover tends to be correlated with species richness (Gaston et al. 2007), but in our study, the variation in turnover is determined by the rarity of the species which then tend to have narrower habitat niches and drive turnover patterns more than widespread species. By analysing species richness and turnover index maps, we found low congruency between the Natura 2000 sites and areas with high species richness and areas occupied by species with a narrower range. Species with narrow ranges (e.g. *Lutra lutra*, Memedemin et al. 2017) are often underrepresented in protected areas, potentially resulting in suboptimal effectiveness of the Natura 2000 network in protecting such species in Dobrogea, despite the large area protected under conservation.

Our results regarding the identification of high-priority areas in Dobrogea for mammal species listed in Annexes II and IV of the Habitats Directive highlight a relatively aggregated pattern of the grid cells with high conservation value in the north-eastern and northern Dobrogea Region, where there are extensive wetlands and forests (i.e., Danube Delta, Măcin Mountains National Park, and North Dobrogea Plateau) (Rey et al. 2007). Additionally, we identified small isolated high-priority areas in agricultural landscapes of southern and central Dobrogea, where biodiversity-friendly agricultural practices should be considered as a conservation method. Distribution of top spatial conservation priorities demonstrated a lack of connectivity between Natura 2000 sites with high conservation values from the northern part of Dobrogea, isolated priority areas in the center of the region, as well as those in the southern part of the region. The distribution of high-priority areas for conservation suggests the necessity of addressing the lack of connectivity, as non-priority areas are essential for the dispersal of species (Christie and Knowles 2015).

The significant overlap between Natura 2000 sites and the other protected areas statutes leads to misunderstandings in law enforcement and an over-optimistic vision of their effectiveness (Ioja et al. 2010). As an example, species whose distributions are limited to the EU Steppic Biogeographic Region or reach the boundaries of their geographic range in Dobrogea tend to be under-represented (Popescu and Murariu 2001, Murariu and Munteanu 2005) as in the case of reptiles and amphibians (Popescu et al. 2013). In Dobrogea, isolation of protected areas leads to low connectivity between habitat patches, which then need to be addressed in future conservation planning and protected area management plans. The lack of research in agricultural landscapes may potentially lead to the populational decline of certain species by not being aware of their distribution and by using flawed species range data (Grant et al. 2007).

The absence of buffer zones and corridors between Natura 2000 sites and small isolated protected areas (the area of the smallest Natura 2000 site in Dobrogea is 0.11 km<sup>2</sup>), are not beneficial in maintaining viable populations, causing the isolation of species with low mobility and specific habitat requirements (Christie and Knowles 2015). Establishing corridors between Natura 2000 sites, especially in the central and southern part of Dobrogea increases connectivity and promotes species dispersal.

Our study is limited by the lack of viable and current distribution data. Most records do not identify geographical coordinates, but localities or toponymies. This makes the niche modelling at a fine scale a challenge. Furthermore, elusive species such as *Mesocricetus newtoni*, *Sicista nordmanni*, and *Vormela peregusna*, are data deficient, and the lack of records (false absences) may influence the results of the analysis. Similarly, misidentification of sibling species may lead to over- or under- estimation of their range. Notably, a study analysing the distribution of amphibians in Dobrogea (Székely et al. 2009) indicates similar issues regarding biased and incomplete distribution data due to the lack of comprehensive surveys of areas with difficult accesses. Another similarity is that some amphibians (e.g., *Bombina bombina*, *Bufo viridis*, *Hyla arborea*), as well as some mammals (e.g., *Capreolus capreolus*, *Apodemus sylvaticus*) are considered widespread and highly detectable, while amphibians such as *Pelobates fuscus* and *Pelobates syriacus*, are cryptic and elusive species and therefore, have low detectability and incomplete distributions (Székely et al. 2013), and that includes species such as *Vormela peregusna*, *Sicista nordmanni* or *Mesocricetus newtoni*. However, biased data lead to more priority areas to protect fewer species (Grant et al. 2007), which is not a shortcoming. Furthermore, Rodrigues et al. (2011) concluded that decision-based on incomplete taxonomic and/or phylogenetic data (such as misidentified sibling species) are robust, and the researcher can safely make use of the best available systematic data.

Future research may focus on identifying buffer zones around Natura 2000 sites to minimise potential negative impacts, particularly in Natura 2000 sites that are adjacent to agricultural areas. From this assessment, we envisage further mapping of corridor networks between small isolated protected areas in southern and central Dobrogea. New research should focus on systematic surveys of agricultural landscapes in central and southern Dobrogea, where vegetation patches remain as refuges for some species listed in Annexes II and IV of the Habitats Directive (*Vormela peregusna*, *Mustela eversmannii*, *Spermophilus citellus*, *Mesocricetus newtoni*, *Cricetus cricetus*, and *Sicista nordmanni*).

## Acknowledgments

IVM, VDP, AN, SM, and LR were supported by a grant from the Romanian National Authority for Scientific Research, PN-III-P4-ID-PCE-2016-0483. Editing and proof-reading by Edward F. Rozyłowicz streamlined the flow of the manuscript. We thank Maria Patroescu, Tiberiu Sahlean, and an anonymous reviewer for their comments that improved the manuscript. We are grateful to Dumitru Murariu, Nastase Radulet, Bogdan Bejenariu, Catalin Stanciu, Dragos Mantoiu, Ionut Stefan Iorgu, Daniyar Memedemin, Razvan Popescu Mirceni, Silviu Petrovan, Marius Skolka, Razvan Zaharia, Alexandra Telea, Dragos Balasoiu, Mihaela Stanescu, and Elena Buhaciuc-Ionita for sharing their data on species occurrences.

## References

- Araújo MB, Alagador D, Cabeza M, Nogués-Bravo D, Thuiller W (2011) Climate change threatens European conservation areas. *Ecology Letters* 14: 484–492. <https://doi.org/10.1111/j.1461-0248.2011.01610.x>
- Arponen A, Heikkinen RK, Thomas CD, Moilanen A (2005) The value of biodiversity in reserve selection: Representation, species weighting, and benefit functions. *Conservation Biology* 19: 2009–2014. <https://doi.org/10.1111/j.1523-1739.2005.00218.x>
- Arslan A, Kryštufek B, Matur F, Zima J (2016) Review of chromosome races in blind mole rats (*Spalax* and *Nannospalax*). *Folia Zoologica* 65: 249–301. <https://doi.org/10.25225/fozo.v65.i4.a1.2016>
- Ausländer D, Hellwing S (1957) Beiträge zur Variabilität und Biologie der Streifenmaus (*Sicista subtilis nordmanni* Keys. et Blas, 1840). *Travaux du Muséum d'Histoire naturelle Grigore Antipa* 1: 255–274.
- Bartolommei P, Sozio G, Bencini C, Cinque C, Gasperini S, Manzo E, Petre S, Solano E, Cozzolino R, Mortelletti A (2016) Field identification of *Apodemus flavicollis* and *Apodemus sylvaticus*: a quantitative comparison of different biometric measurements. *Mammalia* 80: 541–547. <https://doi.org/10.1515/mammalia-2014-0051>
- Botts EA, Erasmus BFN, Alexander GJ (2011) Geographic sampling bias in the South African Frog Atlas Project: Implications for conservation planning. *Biodiversity and Conservation* 20: 119–139. <https://doi.org/10.1007/s10531-010-9950-6>
- Bunescu A (1959) Contribution à l'étude de la répartition géographique de quelques mammifères méditerranéens en Roumanie. *Säugerierkundliche Mitteilungen* 1: 1–4.
- Bunescu A (1961) Contribuții la studiul răspândirii geografice a unor animale mediteraneene din R. P. Română, Nota II – Vertebrate. *Probleme de Geografie* 8: 123–144.
- Christie MR, Knowles LL (2015) Habitat corridors facilitate genetic resilience irrespective of species dispersal abilities or population sizes. *Evolutionary Applications* 8: 454–463. <https://doi.org/10.1111/eva.12255>
- Cogalniceanu D, Rozyłowicz L, Székely P, Samoilă C, Stanescu F, Tudor M, Székely D, Iosif R (2013) Diversity and distribution of reptiles in Romania. *ZooKeys* 341: 49–76. <https://doi.org/10.3897/zookeys.341.5502>
- Crooks KR, Burdett CL, Theobald DM, Rondinini C, Boitani L (2011) Global patterns of fragmentation and connectivity of mammalian carnivore habitat. *Philosophical transactions of the Royal Society of London: Series B, Biological sciences* 366: 2642–2651. <https://doi.org/10.1098/rstb.2011.0120>
- Cuzic M, Marinov M (2004) Date privind situația populației de nură europeană (*Mustela lutreola* L, 1761) (Mammalia, Carnivora, Mustelidae) în câteva zone din Rezervația Biosferei Delta Dunării. *Muzeul Brukenthal. Studii și Comunicări Științele Naturii* 29: 231–239.
- Eklund J, Arponen A, Visconti P, Cabeza M (2011) Governance factors in the identification of global conservation priorities for mammals. *Philosophical transactions of the Royal Society of London: Series B, Biological sciences* 366: 2661–2669. <https://doi.org/10.1098/rstb.2011.0114>

- Ferrier S, Wintle BA (2009) Quantitative approaches to spatial conservation prioritization: matching the solution to the need. In: Moilanen KA, Wilson HPP (Eds) Spatial Conservation Prioritization: Quantitative Methods & Computational Tools. Oxford University Press, Oxford, 1–13.
- Fortin M, Dale M (2005) Spatial Analysis – A Guide for Ecologists. Cambridge University Press, Cambridge, 365 pp.
- Gaston KJ, Davies RG, Orme CDL, Olson VA, Thomas GH, Ding TS, Rasmussen PC, Lennon JJ, Bennett PM, Owens IP, Blackburn TM (2007) Spatial turnover in the global avifauna. *Proceedings of the Royal Society B: Biological Sciences* 274: 1567–1574. <https://doi.org/10.1098/rspb.2007.0236>
- Grand J, Cummings MP, Rebelo TG, Ricketts TH, Neel MC (2007) Biased data reduce efficiency and effectiveness of conservation reserve networks. *Ecology Letters* 10: 364–374. <https://doi.org/10.1111/j.1461-0248.2007.01025.x>
- Hamar M, Schutowa M (1966) Neue daten über die geographische veränderlichkeit und die entwicklung der gattung *Mesocricetus* Nehring, 1898 (Gires, Mammalia). *Zeitschrift-Saeugetierkunde* 31: 237–251.
- Ioja CI, Patroescu M, Rozyłowicz L, Popescu VD, Vergheș M, Zotta M, Felciuc M (2010) The efficacy of Romania's protected areas network in conserving biodiversity. *Biological Conservation* 143: 2468–2476. <https://doi.org/10.1016/j.biocon.2010.06.013>
- Jaarola M, Martínková N, Gündüz İ, Brunhoff C, Zima J, Nadachowski A, Amori G, Bulatova NS, Chondropoulos B, Fragedakis-Tsolis S, González-Esteban J, José López-Fuster M, Kandaurov AS, Kefelioglu H, da Luz Mathias M, Villate I, Searle JB (2004) Molecular phylogeny of the speciose vole genus *Microtus* (Arvicolinae, Rodentia) inferred from mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 33: 647–663. <https://doi.org/10.1016/j.ympev.2004.07.015>
- Kiss JB, Doroșencu A, Marinov ME, Alexe V, Bozagiievici R (2012) Considerations regarding the occurrence of the Eurasian Beaver (*Castor fiber* Linnaeus 1758) in the Danube Delta (Romania). *Scientific Annals of the Danube Delta Institute* 18: 49–56. <https://doi.org/10.7427/DDI.18.05>
- Kiss JB, Marinov M, Alexe V, Doroșenco A (2014) Eurasian Beaver (*Castor fiber* L. 1758), Pine Marten (*Martes martes* L. 1758) and Stone Marten (*Martes foina* Erxleben, 1777) in the Danube Delta (Romania). *Beiträge zur Jagd-und Wildforschung* 39: 347–355.
- Kiss JB, Marinov M, Alexe V, Doroșenco A, Sandor A (2013) Pătrunderea jderului de copac (*Martes martes*, Linnaeus 1758) în Delta Dunării (România) și considerații privind urmările ecologice scontate al acestui fenomen. *Revista Pădurilor* 128(3): 38–47.
- Kiss JB, Marinov M, Alexe V, Sandor A (2012) First record of the Pine Marten (*Martes martes*) in the Danube Delta. *North-Western Journal of Zoology* 8: 195–197.
- Kukkala AS, Moilanen A (2013) Core concepts of spatial prioritisation in systematic conservation planning. *Biological Reviews* 88: 443–464. <https://doi.org/10.1111/brv.12008>
- Kukkala AS, Santangeli A, Butchart SHM, Maiorano L, Ramirez I, Burfield IJ, Moilanen A (2016) Coverage of vertebrate species distributions by Important Bird and Biodiversity Areas and Special Protection Areas in the European Union. *Biological Conservation* 202: 1–9. <https://doi.org/10.1016/j.biocon.2016.08.010>

- Laffan SW, Lubarsky E, Rosauer DF (2010) Biodiverse, a tool for the spatial analysis of biological and related diversity. *Ecography* 33: 643–647. <https://doi.org/10.1111/j.1600-0587.2010.06237.x>
- Lehtomäki J, Moilanen A (2013) Methods and workflow for spatial conservation prioritization using Zonation. *Environmental Modelling and Software* 47: 128–137. <https://doi.org/10.1016/j.envsoft.2013.05.001>
- Lennon JJ, Koleff P, Greenwood JJD, Gaston KJ (2001) The geographical structure of British bird distributions: Diversity, spatial turnover and scale. *Journal of Animal Ecology* 70: 966–979. <https://doi.org/10.1046/j.0021-8790.2001.00563.x>
- Lung T, Meller L, van Teeffelen AJA, Thuiller W, Cabeza M (2014) Biodiversity Funds and Conservation Needs in the EU Under Climate Change. *Conservation Letters* 7: 390–400. <https://doi.org/10.1111/conl.12096>
- Maiorano L, Amori G, Montemaggiori A, Rondinini C, Santini L, Saura S, Boitani L (2015) On how much biodiversity is covered in Europe by national protected areas and by the Natura 2000 network: Insights from terrestrial vertebrates. *Conservation Biology* 29: 986–995. <https://doi.org/10.1111/cobi.12535>
- Manolache S, Nita A, Ciocanea CM, Popescu VD, Rozyłowicz L (2018) Power, influence and structure in Natura 2000 governance networks. A comparative analysis of two protected areas in Romania. *Journal of Environmental Management* 212: 54–64. <https://doi.org/10.1016/j.jenvman.2018.01.076>
- Margules CR, Pressey RL (2000) Systematic conservation planning. *Nature* 405: 243–253. <https://doi.org/10.1038/35012251>
- Memedemin D, Tudor M, Cogalniceanu D, Skolka M, Banica G, Rozyłowicz L (2017) Update on the Geographic Distribution of *Lutra lutra* at the Romanian Black Sea Coast. *Travaux du Muséum National d'Histoire Naturelle Grigore Antipa* 60: 413–417. <https://doi.org/10.1515/travmu-2017-0004>
- Moilanen A, Veach V, Meller L, Arponen A, Kujala H (2014) Spatial conservation planning methods and software. Version 4. User Manual. University of Helsinki, Helsinki, Finland, 290 pp. <https://doi.org/10.1017/CBO9781107415324.004>
- Murariu D (1996) Mammals of the Danube Delta (Romania). *Travaux du Muséum National d'Histoire Naturelle* 36: 361–371
- Murariu D (2006) Mammal ecology and distribution from North Dobrogea (Romania). *Travaux du Muséum national d'Histoire naturelle* 49: 387–399.
- Murariu D, Atanasova I, Raykov I, Chișamera G (2009) Results on mammal (Mammalia) survey from Bulgarian and Romanian Dobrogea. *Travaux du Muséum national d'Histoire naturelle* 52: 371–386.
- Murariu D, Chișamera G, Petrescu A, Atanasova I, Raykov I (2010) Terrestrial vertebrates of Dobrogea – Romania and Bulgaria. *Travaux du Muséum national d'Histoire naturelle Grigore Antipa* 53: 357–375. <https://doi.org/10.2478/v10191-010-0027-2>
- Murariu D, Munteanu D (2005) Fauna României – Mammalia – Carnivora. Editura Academiei Române, Bucharest, 224 pp

- Nita A, Rozyłowicz L, Manolache S, Ciocanea CM, Miu IV, Popescu VD (2016) Collaboration Networks in Applied Conservation Projects across Europe. PLoS ONE 11(10): e0164503. <https://doi.org/10.1371/journal.pone.0164503>
- Nita A, Ciocanea CM, Manolache S, Rozyłowicz L (2018) A network approach for understanding opportunities and barriers to effective public participation in the management of protected areas. Social Network Analysis and Mining 8. <https://doi.org/10.1007/s13278-018-0509-y>
- Ord JK, Getis A (1995) Local spatial autocorrelation statistics: distributional issues and an application. Geographical Analysis 27: 286–306. <https://doi.org/10.1111/j.1538-4632.1995.tb00912.x>
- Popescu A, Murariu D (2001) Fauna României. Mammalia. Rodentia. Editura Academiei Române, Bucharest, 214 pp.
- Popescu VD, Rozyłowicz L, Cogalniceanu D, Niculae IM, Cucu AL (2013) Moving into protected areas? Setting conservation priorities for Romanian reptiles and amphibians at risk from climate change. PLoS ONE 8: e79330. <https://doi.org/10.1371/journal.pone.0079330>
- Possingham HP, Wilson KA, Andelman SJ, Vynne CH (2006) Protected areas: Goals, limitations, and design. In: Groom MJ, Meffe GK, Carroll CR (Eds) Principles of Conservation Biology. Sinauer Associates, Sunderland, 509–533.
- Pouzols F, Toivonen T, Di Minin E, Kukkala AS, Kullberg P, Kuusterä J, Lehtomäki J, Tenkanen H, Verburg PH, Moilanen A (2014) Global protected area expansion is compromised by projected land-use and parochialism. Nature 516: 383–386. <https://doi.org/10.1038/nature14032>
- Pressey RL, Cabeza M, Watts ME, Cowling RM, Wilson K (2007) Conservation planning in a changing world. Trends in Ecology & Evolution 22: 583–92. <https://doi.org/10.1016/j.tree.2007.10.001>
- Rey V, Ianos I, Groza O, Patroescu M (2007) Atlas de la Roumanie Nouvelle edition. Reclus, Montpellier, 208 pp.
- Santos BPA, Bitencourt C, Rapini A (2017) Distribution patterns of *Kielmeyera* (Calophyllaceae): the Rio Doce basin emerges as a confluent area between the northern and southern Atlantic Forest. Neotropical Biodiversity 3: 1–9. <https://doi.org/10.1080/23766808.2016.1266730>
- Rodrigues ASL, Andelman SJ, Bakarr MI, Boitani L, Brooks TM, Cowling RM, Fishpool LDC, Da Fonseca GAB, Gaston KJ, Hoffmann M, Long JS, Marquet PA, Pilgrim JD, Pressey RL, Schipper J, Sechrest W, Stuart SN, Underhill LG, Waller RW, Watts MEJ, Yan X (2004) Effectiveness of the global protected area network in representing species diversity. Nature 428: 640–3. <https://doi.org/10.1038/nature02422>
- Rodrigues ASL, Grenyer R, Baillie JEM, Bininda-Emonds ORP, Gittlemann JL, Hoffmann M, Safi K, Schipper J, Stuart SN, Brooks T (2011) Complete, accurate, mammalian phylogenies aid conservation planning, but not much. Philosophical Transactions of the Royal Society B: Biological Sciences 366: 2652–2660. <https://doi.org/10.1098/rstb.2011.0104>



- Rozyłowicz L, Nita A, Manolache S, Ciocanea CM, Popescu VD (2017) Recipe for success: A network perspective of partnership in nature conservation. *Journal for Nature Conservation* 38: 21–29. <https://doi.org/10.1016/j.jnc.2017.05.005>
- Sánchez-Fernández D, Abellán P, Aragón P, Varela S, Cabeza M (2017) Matches and mismatches between conservation investments and biodiversity values in the European Union. *Conservation Biology* 32: 109–115. <https://doi.org/10.1111/cobi.12977>
- Székely P, Iosif R, Székely D, Stanescu F, Cogalniceanu D (2013) Range extension for the Eastern spadefoot toad *Pelobates syriacus* (Boettger, 1889) (Anura: Pelobatidae). *Herpetology Notes* 6: 481–484.
- Székely P, Plaiasu R, Tudor M, Cogalniceanu D (2009) The distribution and conservation status of amphibians in Dobrudja (Romania). *Turkish Journal of Zoology* 33: 147–156. <https://doi.org/10.3906/zoo-0805-12>
- Temple HJ, Terry A (2009) European mammals: Red List status, trends, and conservation priorities. *Folia zoologica* 58: 248
- Whittaker RH (1972) Evolution and measurement of species diversity. *Taxon* 21: 213–251. <https://doi.org/10.2307/1218190>
- Williams PH, Margules CR, Hilbert DW (2002) Data requirements and data sources for biodiversity priority area selection. *Journal of Biosciences* 27: 327–338. <https://doi.org/10.1007/BF02704963>
- Wilson DE, Reeder DM (2005) *Mammal Species of the World. A Taxonomic and Geographic Reference* (3<sup>rd</sup> edn). Johns Hopkins University Press, 2142 pp.
- Wilson KA, Cabeza M, Klein CJ (2009) Fundamental concepts of spatial conservation prioritization. In: Moilanen A, Wilson KA, Possingham HP (Eds) *Spatial Conservation Prioritization: Quantitative Methods and Computational Tools*. Oxford University Press, New York, 16–27.

## Supplementary material I

### Publications used to compile distribution of terrestrial mammal species from Dobrogea Region, Romania (field reports and data from museum collections)

Authors: Iulia V. Miu, Gabriel B. Chisamera, Viorel D. Popescu, Ruben Iosif, Andreea Nita, Steluta Manolache, Viorel D. Gavril, Ioana Cobzaru, Laurentiu Rozyłowicz

Data type: reference

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.792.25314.suppl1>

## **Supplementary material 2**

### **Occurrences maps for 59 mammal species**

Authors: Iulia V. Miu, Gabriel B. Chisamera, Viorel D. Popescu, Ruben Iosif, Andreea Nita, Steluta Manolache, Viorel D. Gavril, Ioana Cobzaru, Laurentiu Rozyłowicz

Data type: occurrence

Explanation note: The mammal species old (before 1990) and new (after 1990) occurrence records at a 5 × 5 km grid resolution within Dobrogea, Romania.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.792.25314.suppl2>

## **Supplementary material 3**

### **Methodology used to identify high-priority Natura 2000 sites with Zonation v4**

Authors: Iulia V. Miu, Gabriel B. Chisamera, Viorel D. Popescu, Ruben Iosif, Andreea Nita, Steluta Manolache, Viorel D. Gavril, Ioana Cobzaru, Laurentiu Rozyłowicz

Data type: methodology

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.792.25314.suppl3>