



Endogenous Galectin-1 in T Lymphocytes Regulates Anti-prostate Cancer Immunity

Enrique Corapi^{1,2}, Gustavo Carrizo^{1,2}, Daniel Compagno^{1,2} and Diego Laderach^{1,2,3*}

¹ Laboratorio de Glico-Oncología Molecular y Funcional, Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina, ² CONICET-Instituto de Química Biológica de la Facultad de Ciencias Exactas y Naturales, Buenos Aires, Argentina, ³ Departamento de Ciencias Básicas, Universidad Nacional de Luján, Buenos Aires, Argentina

OPEN ACCESS

Edited by:

Jin S. Im,
University of Texas MD Anderson
Cancer Center, United States

Reviewed by:

Charles J. Dimitroff,
Harvard Medical School,
United States
Francisco Sanchez-Madrid,
Universidad Autonoma de Madrid,
Spain

*Correspondence:

Diego Laderach
diegoladerach@qb.fcen.uba.ar

Specialty section:

This article was submitted to
Cancer Immunity and Immunotherapy,
a section of the journal
Frontiers in Immunology

Received: 09 June 2018

Accepted: 04 September 2018

Published: 26 September 2018

Citation:

Corapi E, Carrizo G, Compagno D and
Laderach D (2018) Endogenous
Galectin-1 in T Lymphocytes
Regulates Anti-prostate Cancer
Immunity. *Front. Immunol.* 9:2190.
doi: 10.3389/fimmu.2018.02190

The identification of effective new therapies for prostate cancer (PCa) requires a better understanding of the multiple molecular interactions between tumor cells and their associated microenvironment. In this context, galectin-1 (Gal-1) is a key molecule in the determination of the prostatic carcinoma microenvironment; therefore, it is essential to understand all the molecular processes in which this protein is involved. Most of the previous studies found in the literature have focused on the microenvironment remodeling properties of tumor-secreted Gal-1, through its interactions with the glyco-receptors at the cell membrane and the extracellular matrix. This report shows original aspects of the lectin by focusing on the role of lymphocyte endogenous Gal-1 in controlling anti-prostate tumor immunity. Using a murine preclinical model of prostate cancer, our results demonstrate that endogenous Gal-1 in lymphocytes modulates their proliferative rate and cytotoxic function in conditions of high extracellular Gal-1 concentration, mainly derived from tumor cells. In such conditions, the absence of Gal-1 in T lymphocytes potentiates anti-tumor immune responses. Further studies demonstrated that endogenous Gal-1 in CD4+, but mainly in CD8+T cells, acts as a negative regulator of anti-tumor immunity. In conclusion, prostate tumors require Gal-1 in lymphocytes to evade immune responses. This report lays the foundation for an original immunotherapy strategy for prostate cancer.

Keywords: prostate cancer, endogenous galectin-1 in lymphocytes, cancer immunotherapy, tumor microenvironment, tumor immune escape

INTRODUCTION

Prostate cancer (PCa) is a major challenge for public health worldwide; epidemiological studies place it as the second-screening and the fifth-leading cause of cancer death in adult men (1). Suitable treatments currently exist for this type of cancer if detected early; in contrast, new treatment options are required for advanced stages of the disease (2). Immunotherapy is a validated alternative for the advanced stages of PCa, e.g., Sipuleucel-T (Provenge[®]) is the first immune-based therapy approved by the US Food and Drug Administration for this type of cancer (3). In addition, several immunotherapy strategies are currently being evaluated for this and other types of cancers involving active immunization against tumor-associated antigens and blocking immune checkpoint molecules (4, 5). In this context, Galectin-1 (Gal-1) is a clue molecule secreted by different types of cancers to promote a tolerogenic microenvironment (6). These tolerogenic

effects occur locally at the primary tumor site but also at a distance since Gal-1 is found soluble in plasma and is transported inside microvesicles, such as exosomes (7). Indeed, most of the immunological studies about Gal-1 are focused on its lectin properties resulting from its interaction with β -galactosides and promotion of lattice formation with membrane receptors and extracellular matrix proteins [reviewed in (8)]. This kind of interactions induces lymphocyte apoptosis after activation (9–11), T cell exhaustion (7) and inhibits T-cell migration across endothelial cells (12); actions that favor the tumor-inefficient immune responses.

However, Gal-1 is also expressed intracellularly in several cell types; it has been reported on transformed tumor cells (13–16), regulatory T cells (17, 18) and antigen-presenting cells (19, 20), where the endogenous Gal-1 plays a role as a regulator of the anti-tumor immune responses. Effector T lymphocytes also express endogenous Gal-1; the absence of Gal-1 in lymphocytes promotes a Th1 response during parasitic diseases (21, 22). Few data are available about the role of the endogenous Gal-1 in the control of cytotoxic CD8+ T cell responses. On one hand, Gal-1 negatively regulates CD8+ T cell expansion, survival upon activation, and function fate (23). On the other hand, another study has demonstrated that Gal-1-deficient mice have impaired *in vivo* anti-viral CD8+ T cell-mediated immune responses (24). There are no data about the role of endogenous Gal-1 expressed by CD8+ T lymphocytes in the control of the anti-tumor properties. However, translating the second concept, in which the absence of Gal-1 impairs immune cytotoxicity, seems controversial, due to the compelling evidence in the literature. In fact, *Lgals1*^{-/-} mice have Foxp3+ Treg cell suppressive dysfunction (17) and are prone to autoimmunity (10, 19, 25–27). More interestingly, the injection of tumorigenic cell lines into *Lgals1*^{-/-} mice demonstrated reduction of tumorigenesis (28, 29), proving the relative importance of the stroma (including immune cells) in determining tumorigenesis potential. Given this scenario, this research aimed to challenge the role of Gal-1 as an endogenous modulator of T cell properties in PCa. Although expressed at low levels in lymphocytes, endogenous Gal-1 plays a major role in the control of the lymphocyte anti-tumor functions in a PCa context. The results allow us to propose the modulation of the endogenous load of Gal-1 in T cells as a novel immunotherapeutic strategy for PCa.

MATERIALS AND METHODS

Animals

Animal procedures complied with the Guidelines for the Welfare of Animals in Experimental Neoplasia (UK) and approved by the University of Buenos Aires's Institutional Animal Care and Use Committee (IACUC; FCEN protocol #2014-038). Six to eight-week-old male C57BL/6 mice were housed in the animal facility

Abbreviations: PCa, prostate cancer; Gal-1, Galectin-1; TRAMP-C1, prostate tumor cell line used in this preclinical model; WT, wild type mice; Gal-1 KO (*Lgals1*^{-/-}), Galectin-1 deficient mice; AOD, average stage of division; CFSE, Carboxyfluorescein succinimidyl ester used to trace cell division by flow cytometry; Th, T helper profile of the immune response; DC, dendritic cells; Treg, regulatory T cells; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; BAIM: brachial, axillary, inguinal and mesenteric lymph nodes.

of the School of Sciences, University of Buenos Aires (Argentina). *Lgals1*^{-/-} mice (originally produced by Dr. F. Poirier, France), were provided by Dr. Gabriel A. Rabinovich (Argentina). Ly5.1 C57BL/6 (CD45.1) mice were provided by Dr. Sophie Ezine (CNRS CDTA; Orleans, France). Athymic nude *Foxn1*^{tmu} mice were acquired from the animal facility of the Veterinary School, National University of La Plata (Argentina).

Cell Culture

Murine PCa cell line TRAMP-C1 (obtained from ATCC) was cultured in Dulbecco's Modified Eagle Medium (DMEM; Invitrogen), 10% heat-inactivated FBS (Gibco), antibiotics (1 U/ml penicillin, 1 μ g/ml streptomycin, 2.5 ng/ml amphotericin) and 0.25 U/ml Insulin (Humulin N; Eli Lilly and Co). Cell morphology was routinely evaluated, and cells were periodically examined for androgen sensitivity (MTT assay) and mycoplasma contamination (PCR). Lymph node cell primary cultures were carried out in proliferation medium: RPMI1640 (Invitrogen) containing 10% heat-inactivated FBS (PAA), 1 unit/mL penicillin, 1 μ g/mL streptomycin, 2.5 ng/mL amphotericin B, 2 mM L-glutamine and 2×10^{-5} M β -mercaptoethanol.

Stable Gal-1 downregulated TRAMP-C1 cell line was produced by transduction with a shRNA lentivirus previously reported (30). Briefly, after 1 week, transduced (GFP+) cells were purified by cell sorting using a FACSAria II cytometer (BD Bioscience). Purification of the transduced cells was carried out if GFP+ cells did not exceed 20%, in order to minimize the number of viral integrations and thus guarantee a minimum perturbation of the genome. As previously reported, this shRNA sequence produced 85% of Gal-1 down-regulation at the protein level.

Murine Samples

Lymph node (brachial, axillary, inguinal and mesenteric; BAIM) samples were harvested and single-cell suspensions were obtained by mechanical disruption.

Cell Purification and Immunoblotting

CD3+T lymphocytes were purified (>98% purity) by cell sorting (FACSAria; BD Biosciences) using PE labeled anti-CD3 mAb (1452C11, BD Pharmingen). Antigen presenting cells (APC) were purified by adherence to plastic (>90% CD14+ purity). TRAMP-C1 cells were obtained from an exponentially growing culture. Cells were lysed in radioimmunoprecipitation assay (RIPA) buffer [50 mmol/L Tris-HCl pH 8, 150 mmol/L NaCl, 1% IGEPAL, 0.5% sodium deoxycholate, 0.1% SDS, 10 mmol/L EDTA, 1 mmol/L sodium vanadate, and Protease Inhibitor Cocktail Set III (Calbiochem)]. Equal amounts of protein (20 μ g) were resolved by 15% SDS-PAGE, blotted onto polyvinylidene difluoride (PVDF) membranes (GE Healthcare), blocked with 5% bovine serum albumin (BSA, Sigma-Aldrich), and probed with anti-galectin-1 (H-45 1:500, Santa Cruz) or anti- β -actin (H196 1:1,000, Santa Cruz) antibodies. Specificity of the used antibodies had been previously evaluated by immunoblotting (30, 31). Bound antibodies were detected with peroxidase-labeled anti-rabbit total immunoglobulins (1:3,000; Sigma-Aldrich). Peroxidase activity was detected using a luminol-based method and chemiluminescence was determined using a Fuji Photo Film

Analyzer. Images were analyzed using ImageJ software (NIH, USA).

Lymphocyte Proliferation, Apoptosis and Degranulation Assays

For T-cell proliferation assays, 5×10^5 CFSE-stained murine lymph node cells ($2.5 \mu\text{M}$, 5 min; Sigma-Aldrich) were seeded in a 96-well U-bottomed plate in proliferation medium. To obtain optimal co-stimulation, 10% APC (adherent splenocytes, 5×10^4) was added to each well. 1×10^4 mitomycin-arrested TRAMP-C1 cells (3-h treatment, $10 \mu\text{g/ml}$; Sigma-Aldrich) were added into cultures to mimic a tumor microenvironment. For T-cell proliferation, cells were stimulated for 72 h with coated anti-CD3 antibody ($1 \mu\text{g/ml}$; 145-2C11 purified mAb) and proliferation assessed by CFSE dilution. In the case of experiences with purified cells, CD4+ and CD8+ T cells were purified to homogeneity (>95% purity) by cell sorting (FACSARIA, BD Biosciences) using fluorescent-conjugated mAbs (anti-CD8, clone 53-6.7, anti-CD4 clone GK1.5; anti-CD3 clone 1452C11; all from BD Pharmingen), mixed at a 2:3 CD8:CD4 ratio and cultured in presence of APC, tumor and anti-CD3 stimuli as before. For transwell assays, 250,000 lymph node cells from Lgals1-/- and WT mice were co-cultivated separated by a $0.4 \mu\text{m}$ pore size transwell system (Falcon). The low chamber was loaded with 250,000 KO lymphocytes, 2,500 TC-1 and 25,000 spleen adherent cells (APC) whereas the top chamber of the transwell was loaded with 250,000 WT (CD45.1); 2,500 TC-1 and 25,000 APC cells. Lymphocytes were activated by $0.25 \mu\text{g/ml}$ anti-CD3 mAb (145-2C11, used soluble to activate T cells in both chambers). After 72 h of co-culture the membrane were removed and the lymphocyte proliferation were assess by CFSE dilution in the CD8+ CD45.2+ gate. In all cases, average stage of division (AOD) was calculated as before (23). Briefly, percentage of cells in each CFSE division was calculated on live CD8+ T cells and inserted into the following equation:

$$\text{AOD} = \{[(\% \text{ undivided}) + (2 \times \% \text{ division 1}) + (3 \times \% \text{ division 2}) + \dots] / 100\} - 1.$$

Apoptosis was determined by Annexin V (BD Pharmingen) staining, performed as before (32). For T-cell degranulation, cells were stimulated for 48 h with coated anti-CD3 antibody ($1 \mu\text{g/ml}$; 145-2C11) and anti-CD107a was added (1 h, 37°C ; 1D4B, BD Pharmingen) followed by an additional 4 h-culture with monensin ($3 \mu\text{M}$, Sigma-Aldrich). In all the cases, cells were then stained for CD8 (53-6.7, BD Pharmingen) in staining buffer (PBS 1% FBS, 0.01% sodium azide) for 30 min on ice. Flow cytometric analysis was performed in a FACSARIA (BD Biosciences) and analyzed using the FlowJo software.

Generation of Experienced Anti-tumor Lymphocytes

Dendritic cells (DC) were prepared from C57BL/6 bone marrows as previously described (33). Briefly, bone marrow cells were flushed with ice-cold PBS and erythrocytes lysed using ACK buffer ($8.7 \text{ g/L NH}_4\text{Cl}$, 1 g/L KHCO_3 , 0.05 mM EDTA , pH 7.3). Cells were cultured in DMEM 20% heat-inactivated FBS (Gibco), 1 mM sodium pyruvate, 2 mM glutamine and antibiotics (1 U/ml penicillin, $1 \mu\text{g/ml}$ streptomycin) (Invitrogen), in presence of 20 ng/ml mrGM-CSF and 10 ng/ml mrIL-4

(ImmunoTools). Cells were cultured for 5 days, with addition of GM+IL-4 every 2 days. Immature DCs were pulsed during 5 h with TRAMP-C1 lysates (4 freeze/thawing steps), and then matured with 2 ng/ml CpG-1826 (phosphorothioate 5'-TCCATGACGTTCCCTGACGTT-3', IDT) and $20 \mu\text{g/ml}$ Poly-U (Sigma-Aldrich) during an overnight culture. To obtain experienced anti-tumor lymphocytes, wild type or Lgals1-/- mice were *s.c.* immunized with tumor lysate-pulsed dendritic cells. At day 7 post-immunization, lymphocytes obtained from axillary, brachial and inguinal lymph nodes were used for immune reconstitution of nude mice.

Tumor Model in Nude Mice Reconstituted With Wt or Lgals1-/- Experienced Lymphocytes

Six to eight-week-old male nude mice were *i.v.* reconstituted with immune cells from one experienced wild type or Lgals1-/- mice. The day after reconstitution, 1×10^6 TRAMP-C1 cells (log phase of growth) were *s.c.* injected in mice in Matrigel ($4\text{--}5 \text{ mg/ml}$; Corning). No changes in mice weights were detected during the protocol. Euthanasia was practiced when tumor volume reached 2500 mm^3 . Tumor size was measured every 2 days with caliper and calculated as $W^2 \times L/2$, where W = width and L = length.

Statistical Analysis


At least three biological replicates (i.e., independent experiments) were carried out in each case, unless stated otherwise. Data represent mean \pm SD. Prism software (GraphPad) was used. Two groups were compared with the Student *t*-test for unpaired data (two-sided tests). $P < 0.05$ was considered statistically significant.

RESULTS

Endogenous Gal-1 in Lymphocytes Negatively Regulates Their Functional Properties in a Tumor Context

A vast literature in different animal models and experimental design are found describing the tumor cells as the main producer of Gal-1 (13, 15, 34–44). Regarding PCa, we previously described that Gal-1 is expressed at high levels in several human PCa cell lines and, more importantly, in patient prostatectomies, being upregulated during disease progression (30, 45). Gal-1 is also highly expressed in murine TRAMP-C1 cells (Figures 1A,B) and the downregulation of Gal-1 in these cells by RNAi reduces their immunosuppressive potential. In fact, the *in-vitro* polyclonal activation of lymphocytes in the presence of Gal-1 downregulated TRAMP-C1 led to higher proliferation of CD8+T cells (Figure 1C). Consequently, Gal-1 is an important immune escape mechanism activated by the tumors.

Besides tumor cells themselves, other cells in the tumor associated microenvironment play fundamental roles in the determination of the tumor evolution. Understanding how endogenous Gal-1 in each of these cells affects their own properties seems, therefore, essential to have a complete functional picture of the PCa microenvironment. In particular,



little is known about the endogenous role of Gal-1 in immune cells. Therefore, we first checked whether Gal-1 is expressed in immune cells and compared its expression levels with those observed in tumor cells. Interestingly, the expression levels of Gal-1 in immune cells [T lymphocytes and antigen presenting cells (APC)] were considerably lower (10–20 times less) than those of tumor cells (**Figures 1A,B**). Low levels of Gal-1 expression in immune cells do not indicate *a priori* that lymphocyte endogenous Gal-1 does not play any role as an intrinsic modulator of their properties. Thus, we decided

to design an *in vitro* experimental model where the function of *Lgals1*^{-/-} lymphocytes would be compared with the wild-type (WT) ones in the presence or absence of wild type tumor (responsible of secreting high levels of “exogenous Gal-1”). As readout, we focused on the function of CD8+ T lymphocytes, as these cells are critical in the anti-tumor cytotoxic response (46). Our results demonstrate that without endogenous Gal-1, CD8+ T cells showed higher proliferative rates (**Figure 2**). A detailed analysis of these results shows that, in the absence of tumor, there was a trend toward greater proliferation of Gal-1-deficient CD8+ T lymphocytes mainly at low conditions of stimulation (**Figures 2A,B**, 1 μ g/mL anti-CD3). However, the results were considerably more significant in the presence of an immunosuppressive microenvironment generated by the tumor cells (**Figures 2C,D**); the absence of endogenous Gal-1 led to higher CD8+ T cell proliferative capacity even in such conditions. We highlight that, the major source of extracellular Gal-1 in these experimental settings are tumor cells that almost saturate the extracellular medium by the secretion of high quantities of Gal-1 [dosed by ELISA at levels of 20 ng/mL, supported also by the high cellular levels of Gal-1 shown in **Figure 1**, and evaluated in patients in our previous study (30) and reported at high levels in the serum of cancer patients (47–50)]. The most potent abrogation of tumor-immunosuppression was achieved by the absence of endogenous Gal-1 in lymphocytes combined with the down-regulation of Gal-1 in tumor cells (**Figures 1C,D**). While the immune effects of Gal-1 in tumor cells have been extensively evaluated (13–15, 51), the endogenous effect of Gal-1 effect in lymphocytes is original and requires more understanding.

Since a widely described action of exogenous Gal-1 is the induction of cell death on activated T lymphocytes (9), we evaluated if lymphocytes with no endogenous Gal-1 were differentially induced into apoptosis in our tumor experimental model. **Figure 2E** shows the percentage of apoptotic cells determined by Annexin V staining inside the CD8+ T cell subpopulation, and **Figure 2F** shows the same staining inside each peak of division of CD8+ T cells. Altogether, these results indicate that the absence of endogenous Gal-1 in CD8+ T lymphocytes protects these cells from apoptosis in the presence of tumor cells. As in proliferative studies, the major source of extracellular Gal-1 was the tumor.

Degranulation is a prerequisite to perforin-granzyme-mediated cytotoxic function by responding to antigen-specific CD8+ T cells. The lytic granules are surrounded by a lipid bilayer containing numerous lysosomal-associated membrane proteins (LAMPs, CD107a and b); thus, labeling responding cells with anti-CD107 antibodies and measuring their expression by flow cytometry can directly identify degranulating CD8+ T cells. The results shown in **Figure 3** demonstrate that the absence of endogenous Gal-1 in CD8+ T lymphocytes increases their ability to degranulate. This was also observed in conditions without tumor cells and at different percentages of TRAMP-C1 (**Figure 3** and data not shown). Interestingly, endogenous Gal-1 in lymphocytes affects not only their percentages of degranulating cells (**Figure 3A**), but also





FIGURE 2 | Role of endogenous Galectin-1 in the proliferation and activation-induced cell death of CD8+ T cells. Proliferation rates of wild-type and Lgals1^{-/-} CD8+ T lymphocytes are compared upon activation with different anti-CD3 concentrations in a 72 h proliferative assay. **(A,B)** In the absence of TRAMP-C1 tumor cells. **(A)** Representative CFSE dilution proliferative experience. **(B)** Graph bars indicating the average stage of division obtained from three independent experiences; White bars: wild-type cells; Black bars: Lgals1^{-/-} cells (* $p < 0.05$, t -test). **(C,D)** In the presence of 1% TRAMP-C1 tumor cells. **(C)** Representative CFSE dilution proliferative experience. **(D)** Graph bars indicating the average stage of division obtained from three independent experiences; White bars: wild-type cells; Black bars: Lgals1^{-/-} cells (* $p < 0.05$, t -test). **(E)** Role of endogenous Galectin-1 in the apoptosis of CD8+ T cells upon polyclonal activation in the presence of 1% TRAMP-C1 cells. Apoptosis rates of wild-type ($n = 3$) and Lgals1^{-/-} ($n = 3$) lymphocytes are comparatively analyzed by Annexin V staining inside total CD8+ T cell gate **(E)** or inside each peak of division (CFSE) in the CD8+ T cell gate **(F)**. White bars: wild-type cells; black bars: Lgals1^{-/-} cells (* $p < 0.05$, t -test).

the CD107a mean fluorescence intensity (MFI). This last result indicates that the intrinsic expression/exposition of CD107a is regulated by this lectin on a *per cell* basis (Figure 3B). Altogether, these results imply that endogenous Gal-1 in CD8+T cells an important role in the control of their function, particularly in the presence of tumor cells.

Lymphocyte Endogenous Gal-1 Modulates Their Function Independently of Exogenous Factors

To evaluate if the immunosuppressive role of lymphocyte endogenous Gal-1 involves cell-to-cell contacts or the production of soluble factors, we performed transwell experiments. Figure 4 shows that Gal-1 deficient lymphocytes proliferate more



extensively than wild-type ones when activated in the presence of tumor (as in **Figure 2**). We next co-cultured wild-type and Gal-1 deficient lymphocytes (they differed in the expression of CD45 isoform; both were activated and proliferation was evaluated in CD45.2+ CFSE labeled). The presence of wild-type cells reduced the proliferation of Gal-1 deficient CD8+T cells. However, when these two lymphocyte sources were separated through a transwell membrane, such a reduction in proliferation was no longer observed (**Figure 4**). This experiment clearly demonstrates the fundamental role of endogenous lymphocyte Gal-1 in controlling their function upon activation, a phenomena that is independent of the tumor secretion of Gal-1 and can be controlled through lymphocyte-lymphocyte interactions. This introduces new variants for the current understanding of the role of Gal-1 in defining a tumor-associated tolerogenic context.

Gal-1 Negatively Regulates *in vivo* Immunity Against PCa

Previously, it has been reported that several tumor cells lose their tumorigenicity when injected into Lgals1^{-/-} mice, compared with their injection into wild-type mice (28, 29). We obtained the same behavior in the case of the TRAMP-C1 prostate tumor model. In fact, while 92% of wild-type mice injected *s.c.* with TRAMP-C1 cells developed tumors (lag time: 47 ± 5 ; duplication time: 7 ± 2 days, $n = 5$), no Lgals1^{-/-} mice (0/4) developed tumors after up to 80 days. However, the conclusions obtained from this type of experimental model are complex, and the real impact of immune-mediated effects cannot be established since Gal-1 is absent in all cells of the body, including endothelial cells in which it was previously demonstrated Gal-1 plays a major regulatory role. To evaluate more precisely whether endogenous Gal-1 in lymphocytes affects the capacity of mice to control tumor growth under *in vivo* settings, we developed the following experimental procedure: anti-tumor lymphocytes were induced by the immunization of Lgals1^{-/-} and wild-type mice with

dendritic cells pulsed with TRAMP-C1 lysates. At day 7 post-immunization, immune cells were obtained from the BAIM lymph nodes; no significant phenotype changes were detected at this stage (total absolute cell numbers, %CD4+, %CD8+, %CD4+CD25+Foxp3+Tregs; data not shown). Lymphocytes obtained from the lymph nodes of immunized mice (wild type or Lgals1^{-/-}) were then transferred into an immunodeficient nude mouse (1-to-1 mice ratio). Under such reconstitution settings, all non-immune host compartments (e.g., stroma and particularly endothelial cells) expressed Gal-1 at normal levels, with the exception of T lymphocyte (transferred) cells. This system allowed us to compare the ability of Lgals1^{-/-} and wild-type lymphocytes to control tumor growth in a WT micro- and macro-environment. An in-depth comparison analysis demonstrated that mice reconstituted with Lgals1^{-/-} lymphocytes showed a marked increase in tumor latency and a slight decrease in tumor duplication time (**Figure 5** and **Table 1**). In contrast, tumorigenesis, defined as the percentage of mice developing tumors, was 100% in both experimental groups, implying that endogenous Gal-1 on lymphocytes plays an important role in controlling tumor growth in initial phases, but later other molecular pathways can circumvent its action.

Cellular Mechanisms Involved in Lymphocyte Modulation by Endogenous Gal-1

Considering the previous *in-vitro* and *in-vivo* results, we analyzed the impact of the endogenous Gal-1 absence in the different subpopulations of cells involved in the immune response. For this purpose, we purified APC, CD4+, and CD8+ T cells to homogeneity and performed *in-vitro* proliferation assays by mixing the subpopulations of different genotypes in physiological proportions. The results shown in **Figure 6** demonstrate that the absence of Gal-1 in CD4+ and CD8+ T cells results in higher proliferative rates of CD8+ T lymphocytes. Higher effect was observed when Gal-1 was absent in the responding CD8+ T cells compared to CD4+ T cells or all wild-type immune cells (AOD = 1.7 vs. 1.1 or 0.8, respectively); however, maximal CD8+ T cell proliferation occurs when Gal-1 is absent in both lymphoid subpopulations (AOD = 2.8). The absence of Gal-1 in APC did not produce any change in the proliferative potential of CD8+ T lymphocytes (data not shown). Altogether, these results clearly demonstrate that endogenous Gal-1 regulates the functional properties of CD4+ and CD8+ T cells.

DISCUSSION

The pleiotropic functions of Gal-1 are partly due to the complex cellular distribution of this lectin. At the extracellular level, Gal-1 is able to interact with glycosylated receptors (through their N-acetylglucosamine glycans), thereby forming complex membrane lattices [reviewed in (52)]. Through this outside-in process, Gal-1 regulates several cellular properties, acting as a negative regulator of immune responses [reviewed in (53, 54)]. Consistently, most of the current Gal-1-based immune strategies proposed for autoimmunity and cancer are focused at the extracellular level.






TABLE 1 | Different Tumor Growth on Nude Mice Reconstituted with Gal-1-deficient or competent lymphocytes.

	Latency (days)	Duplication time (days)
WT lymphocytes	32 ± 2	6.0 ± 1.2
Lgals1 ^{-/-} lymphocytes	54 ± 4	8.2 ± 0.4

Additionally, the intracellular regulation of Gal-1 is expected to perform extra regulatory functions. At this subcellular localization, Gal-1 regulates several cellular processes, such as

growth, survival, and differentiation; several of these effects are carbohydrate-independent [reviewed in (52)]. It has been demonstrated that the downregulation of the endogenous Gal-1 expressed by tumor cells has a direct effect on the immune response [our results and (13–15, 51)]. As expected, these strategies also modify the extracellular levels of Gal-1 (38), as the transformed cells are the main Gal-1 producers. Interestingly, Gal-1 is also expressed by other cells of the tumor microenvironment; the endogenous function of Gal-1 in those cells may affect tumor growth. Indeed, several reports have demonstrated the decreased tumorigenic



potential of various tumor cell lines when injected into Gal-1-deficient mice (28, 29). The absence of Gal-1 on endothelial cells of *Lgals1*^{-/-} hosts is responsible for the loss of tumorigenicity; angiogenesis defaults prevent an adequate supply of nutrients, oxygen, and hormones, factors that are essential for optimal tumor growth (28). While this report does not exclude the importance of this scenario, it proposes a complementary alternative in which tumors require Gal-1 to be endogenously expressed in T lymphocytes to escape from immune attack.

In fact, our results demonstrate that lymphocytes isolated from Gal-1-deficient mice have higher CD8⁺ T cell proliferative rates and effector functions, even when activated in the presence of high levels of extracellular Gal-1 secreted by tumor cells. This biological observation could be due to a direct endogenous effect of Gal-1 on CD8⁺ T cells or through indirect effects on other cell types; this report dissected the functional role of endogenous Gal-1 in immune cells. Our results show that the expression of Gal-1 on antigen-presenting cells is indifferent at potentiating CD8⁺ T cell functions. However, the absence of Gal-1 in CD4⁺T cells induced higher levels of CD8⁺ T proliferation. Gal-1 plays a fundamental role in the control of differentiation and function of conventional CD4⁺CD25⁺Foxp3⁺ Tregs (17). While we did not detect changes in the frequency of Tregs in our experimental settings (data not shown), such Tregs are functionally deficient (17). Consequently, CD4⁺T cell subpopulations from *Lgals1*^{-/-}

and wild-type mice differ at least at this functional level. However, other endogenous modifications are also plausible in the CD4⁺T cell compartment, as was previously demonstrated in parasitic models of infection (21, 22). More interestingly, the absence of endogenous Gal-1 induces higher functional properties of CD8⁺ T cells. These results agree with a previous report that demonstrated that Gal-1 has a major impact over proliferation burst, lymphocyte survival upon activation, and effector fate (23).

Mechanistically, our transwell experiments demonstrated that the immunoregulation mediated by endogenous Gal-1 in immune cells involves cell-to-cell interactions. In fact, the co-culture of Gal-1 deficient together with wild-type lymphocytes abolished the immunopotentiality observed in *Lgals1*^{-/-} cells alone. In addition, our results demonstrate that soluble factors are not involved in this type of regulation. Altogether, these results led us to hypothesize that Gal-1 deficient lymphocytes lack immune-inhibitory membrane receptors (or, alternatively, acquire immune-activating receptors) that have a direct effect on lymphocyte function. It has been reported that intracellular Gal-1 could affect cell behavior through regulation of the transcriptional machinery (Gal-1 forms part of the spliceosome), miRNA, and signaling pathways (52). Effects coming from intracellular Gal-1 interactors in lymphocytes have not yet been investigated.

Interestingly, the relevant roles of lymphocyte endogenous Gal-1 have recently been proposed in active systemic lupus erythematosus. In fact, patient lymphocytes demonstrate lower levels of endogenous Gal-1 upon activation, a mechanism that may make them prone to autoimmunity (55). In cancer, our results support a scenario where the high levels of exogenous Gal-1 produced by tumor cells are not enough to evade immune attack; tumors require additional licensing by lymphocyte endogenous Gal-1 to be efficiently immunosuppressive. These results define a new and potent target for immunotherapy in PCa. From a clinical point of view, it is interesting to note that the immunopotentiality resulting from the downregulation of Gal-1 in lymphocytes is more significant in the presence of a tumor; consequently, an anti-PCa therapeutic strategy based on these results should have few side effects.

AUTHOR CONTRIBUTIONS

EC and GC performed and analyzed the work. DC data interpretation, critical revision. DL conceived the idea, analyzed data, supervised the project and wrote the paper. All authors discussed the results and contributed to the final manuscript.

FUNDING

Research at the Molecular and Functional Glico-Oncology Laboratory is supported by ANPCyT PICT 2012-1533; ANPCyT PICT 2014-0983; Fundación Florencio Fiorini, Instituto Nacional del Cáncer (all from Argentina) and Worldwide Cancer Research (UK).

ACKNOWLEDGMENTS

We thank Lic. Guillermo Plaza at the flow cytometry facility and staff at the FCEN-Central animal facility for animal husbandry.

We acknowledge Laura Giribaldi for initial studies, Dr. Gabriel A. Rabinovich (IBYME, Argentina) for providing Lgals1-/- mice and Dr. Sophie Ezine (Institut Necker Enfants Malades, France) for providing C57BL/6 Ly5.1 mice.

REFERENCES

- International Agency for Research on Cancer, WHO. *World Cancer Report 2014*. Stewart BW, Wild CP, editors. (2014).
- Evans AJ. Treatment effects in prostate cancer. *Mod Pathol.* (2018) 31:S110–21. doi: 10.1038/modpathol.2017.158
- Sipuleucel-T. APC 8015, APC-8015, prostate cancer vaccine–Dendreon. *Drugs R D* (2006) 7:197–201.
- Coulie PG, Van den Eynde BJ, van der Bruggen P, Boon T. Tumour antigens recognized by T lymphocytes: at the core of cancer immunotherapy. *Nat Rev Cancer* (2014) 14:135–46. doi: 10.1038/nrc3670
- Nishino M, Ramaia NH, Hatabu H, Hodi F. S. Monitoring immune-checkpoint blockade: response evaluation and biomarker development. *Nat Rev Clin Oncol.* (2017) 14:655–68. doi: 10.1038/nrclinonc.2017.88
- Wu AA, Drake V, Huang HS, Chiu S, Zheng L. Reprogramming the tumor microenvironment: tumor-induced immunosuppressive factors paralyze T cells. *Oncoimmunology* (2015) 4:e1016700. doi: 10.1080/2162402X.2015.1016700
- Maybruck BT, Pfannenstiel LW, Diaz-Montero M, Gastman BR. Tumor-derived exosomes induce CD8(+) T cell suppressors. *J Immunother Cancer* (2017) 5:65. doi: 10.1186/s40425-017-0269-7
- Elola MT, Blidner AG, Ferragut F, Bracalente C, Rabinovich GA. Assembly, organization and regulation of cell-surface receptors by lectin-glycan complexes. *Biochem J.* (2015) 469:1–16. doi: 10.1042/BJ20150461
- Perillo NL, Pace KE, Seilhamer JJ, Baum LG. Apoptosis of T cells mediated by galectin-1. *Nature* (1995) 378:736–9. doi: 10.1038/378736a0
- Toscano MA, Bianco GA, Ilarregui JM, Croci DO, Correale J, Hernandez JD, et al. Differential glycosylation of TH1, TH2 and TH-17 effector cells selectively regulates susceptibility to cell death. *Nat Immunol.* (2007) 8:825–34. doi: 10.1038/ni1482
- Motran CC, Molinder KM, Liu SD, Poirier F, Miceli MC. Galectin-1 functions as a Th2 cytokine that selectively induces Th1 apoptosis and promotes Th2 function. *Eur J Immunol.* (2008) 38:3015–27. doi: 10.1002/eji.200838295
- He J, Baum LG. Endothelial cell expression of galectin-1 induced by prostate cancer cells inhibits T-cell transendothelial migration. *Lab Invest.* (2006) 86:578–90. doi: 10.1038/labinvest.3700420
- Rubinstein N, Alvarez M, Zwirner NW, Toscano MA, Ilarregui JM, Bravo A, et al. Targeted inhibition of galectin-1 gene expression in tumor cells results in heightened T cell-mediated rejection; A potential mechanism of tumor-immune privilege. *Cancer Cell* (2004) 5:241–51. doi: 10.1016/S1535-6108(04)00024-8
- Banh A, Zhang J, Cao H, Bouley DM, Kwok S, Kong C, et al. Tumor galectin-1 mediates tumor growth and metastasis through regulation of T-cell apoptosis. *Cancer Res.* (2011) 71:4423–31. doi: 10.1158/0008-5472.CAN-10-4157
- Verschuere T, Toelen J, Maes W, Poirier F, Boon L, Tousseyn T, et al. Glioma-derived galectin-1 regulates innate and adaptive antitumor immunity. *Int J Cancer* (2014) 134:873–84. doi: 10.1002/ijc.28426
- Mathieu V, de Lassalle EM, Toelen J, Mohr T, Bellahcene A, Van Goietsenoven G, et al. Galectin-1 in melanoma biology and related neo-angiogenesis processes. *J Invest Dermatol.* (2012) 132:2245–54. doi: 10.1038/jid.2012.142
- Garin MI, Chu CC, Golshayan D, Cernuda-Morollon E, Wait R, Lechler RI. Galectin-1: a key effector of regulation mediated by CD4+CD25+ T cells. *Blood* (2007) 109:2058–65. doi: 10.1182/blood-2006-04-016451
- Wang J, Lu ZH, Gabius HJ, Rohowsky-Kochan C, Ledeen RW, Wu G. Cross-linking of GM1 ganglioside by galectin-1 mediates regulatory T cell activity involving TRPC5 channel activation: possible role in suppressing experimental autoimmune encephalomyelitis. *J Immunol.* (2009) 182:4036–45. doi: 10.4049/jimmunol.0802981
- Ilarregui JM, Croci DO, Bianco GA, Toscano MA, Salatino M, Vermeulen ME, et al. Tolerogenic signals delivered by dendritic cells to T cells through a galectin-1-driven immunoregulatory circuit involving interleukin 27 and interleukin 10. *Nat Immunol.* (2009) 10, 981–91. doi: 10.1038/ni1772
- Ortner D, Grabher D, Hermann M, Kremmer E, Hofer S, Heufler C. The adaptor protein Bam32 in human dendritic cells participates in the regulation of MHC class I-induced CD8+ T cell activation. *J Immunol.* (2011) 187:3972–8. doi: 10.4049/jimmunol.1003072
- Bunn PT, Montes de Oca M, Rivera FL, Kumar R, Edwards CL, Faleiro RJ, et al. Galectin-1 impairs the generation of anti-parasitic Th1 cell responses in the liver during experimental visceral leishmaniasis. *Front Immunol.* (2017) 8:1307. doi: 10.3389/fimmu.2017.01307
- Lonnberg T, Svensson V, James KR, Fernandez-Ruiz D, Sebina I, Montandon R, et al. Single-cell RNA-seq and computational analysis using temporal mixture modelling resolves Th1/Th fate bifurcation in malaria. *Sci Immunol.* (2017) 2:eaa12192. doi: 10.1126/sciimmunol.aal2192
- Liu SD, Tomassian T, Bruhn KW, Miller JF, Poirier F, Miceli MC. Galectin-1 tunes TCR binding and signal transduction to regulate CD8 burst size. *J Immunol.* (2009) 182:5283–95. doi: 10.4049/jimmunol.0803811
- Clemente T, Vieira NJ, Cerliani JP, Adrain C, Luthi A, Dominguez MR, et al. Proteomic and functional analysis identifies galectin-1 as a novel regulatory component of the cytotoxic granule machinery. *Cell Death Dis.* (2017) 8:e3176. doi: 10.1038/cddis.2017.506
- Perez CV, Gomez LG, Gualdoni GS, Lustig L, Rabinovich GA, Guazzone VA. Dual roles of endogenous and exogenous galectin-1 in the control of testicular immunopathology. *Sci Rep.* (2015) 5:12259. doi: 10.1038/srep12259
- Iqbal AJ, Cooper D, Vulger A, Gittens BR, Moore A, Perretti M. Endogenous galectin-1 exerts tonic inhibition on experimental arthritis. *J Immunol.* (2013) 191:171–7. doi: 10.4049/jimmunol.1203291
- Blois SM, Ilarregui JM, Tometten M, Garcia M, Orsal AS, Cordo-Russo R, et al. A pivotal role for galectin-1 in fetomaternal tolerance. *Nat Med.* (2007) 13, 1450–7. doi: 10.1038/nm1680
- Thijssen VL, Barkan B, Shoji H, Aries IM, Mathieu V, Deltour L, et al. Tumor cells secrete galectin-1 to enhance endothelial cell activity. *Cancer Res.* (2010) 70:6216–24. doi: 10.1158/0008-5472.CAN-09-4150
- Thijssen VL, Postel R, Brandwijk RJ, Dings RP, Nesmelova I, Satijn S, et al. Galectin-1 is essential in tumor angiogenesis and is a target for antiangiogenesis therapy. *Proc Natl Acad Sci USA.* (2006) 103:15975–80. doi: 10.1073/pnas.0603883103
- Laderach DJ, Gentilini LD, Giribaldi L, Delgado VC, Nugnes L, Croci DO, et al. A Unique galectin signature in human prostate cancer progression suggests Galectin-1 as a key target for treatment of advanced disease. *Cancer Res.* (2013) 73:86–96. doi: 10.1158/0008-5472.CAN-12-1260
- Gentilini LD, Jaworski FM, Tiraboschi C, Perez IG, Kotler ML, Chauchereau A, et al. Stable and high expression of Galectin-8 tightly controls metastatic progression of prostate cancer. *Oncotarget* (2017) 8:44654–68. doi: 10.18632/oncotarget.17963
- Jaworski FM, Gentilini LD, Gueron G, Meiss RP, Ortiz EG, Berguer PM, et al. *In vivo* hemin conditioning targets the vascular and immunologic compartments and restrains prostate tumor development. *Clin Cancer Res.* (2017) 23:5135–48. doi: 10.1158/1078-0432.CCR-17-0112
- Tavare R, Escuin-Ordinas H, Mok S, McCracken MN, Zettlitz KA, Salazar FB, et al. An effective immuno-PET imaging method to monitor CD8-dependent responses to immunotherapy. *Cancer Res.* (2016) 76:73–82. doi: 10.1158/0008-5472.CAN-15-1707
- Croci DO, Salatino M, Rubinstein N, Cerliani JP, Cavallin LE, Leung HJ, et al. Disrupting galectin-1 interactions with N-glycans suppresses hypoxia-driven angiogenesis and tumorigenesis in Kaposi's sarcoma. *J Exp Med.* (2012) 209:1985–2000. doi: 10.1084/jem.20111665
- Saussez S, Decaestecker C, Lorfèvre F, Cucu DR, Mortuaire G, Chevalier D, et al. High level of galectin-1 expression is a negative prognostic predictor

- of recurrence in laryngeal squamous cell carcinomas. *Int J Oncol.* (2007) 30:1109–17. doi: 10.3892/ijo.30.5.1109
36. Su Y, Wang W, Xu Y, Liangjun W, Wang Y, Li C, et al. Clinicopathological significance of galectin-1 expression and percentage of galectin-1-expressing T cells in clear-cell renal cell carcinoma. *Can Urol Assoc J.* (2018) 12:E243–9. doi: 10.5489/auaj.4573
 37. Juszczynski P, Ouyang J, Monti S, Rodig SJ, Takeyama K, Abramson J, et al. The AP1-dependent secretion of galectin-1 by Reed Sternberg cells fosters immune privilege in classical Hodgkin lymphoma. *Proc Natl Acad Sci USA.* (2007) 104:13134–9. doi: 10.1073/pnas.0706017104
 38. Dalotto-Moreno T, Croci DO, Cerliani JP, Martinez-Allo VC, Dergan-Dylon S, Mendez Huelgo SP, et al. Targeting galectin-1 overcomes breast cancer associated immunosuppression and prevents metastatic disease. *Cancer Res.* (2013) 73:1107–17. doi: 10.1158/0008-5472.CAN-12-2418
 39. Barrow H, Rhodes JM, Yu LG. The role of galectins in colorectal cancer progression. *Int J Cancer.* (2011) 129:1–8. doi: 10.1002/ijc.25945
 40. Szoke T, Kayser K, Baumhake JD, Trojan I, Furak J, Tizlavicz L, et al. Prognostic significance of endogenous adhesion/growth-regulatory lectins in lung cancer. *Oncology* (2005) 69 167–74. doi: 10.1159/000087841
 41. Chow SN, Chen RJ, Chen CH, Chang TC, Chen LC, Lee WJ, et al. Analysis of protein profiles in human epithelial ovarian cancer tissues by proteomic technology. *Eur J Gynaecol Oncol.* (2010) 31:55–62.
 42. Rorive S, Belot N, Decaestecker C, Lefranc F, Gordower L, Micik S, et al. Galectin-1 is highly expressed in human gliomas with relevance for modulation of invasion of tumor astrocytes into the brain parenchyma. *Glia* (2001) 33:241–55. doi: 10.1002/1098-1136(200103)33:3<241::AID-GLIA1023>3.0.CO;2-1
 43. D'Haene N, Maris C, Sandras F, Dehou MF, Rummelink M, Decaestecker C, et al. The differential expression of Galectin-1 and Galectin-3 in normal lymphoid tissue and non-Hodgkin's and Hodgkin's lymphomas. *Int J Immunopathol Pharmacol.* (2005) 18:431–43. doi: 10.1177/039463200501800304
 44. Koopmans SM, Bot FJ, Schouten HC, Janssen J, van Marion AM. The involvement of Galectins in the modulation of the JAK/STAT pathway in myeloproliferative neoplasia. *Am J Blood Res.* (2012) 2:119–27.
 45. Compagno D, Laderach DJ, Gentilini L, Jaworski FM, Rabinovich GA. Delineating the “galectin signature” of the tumor microenvironment. *Oncoimmunology* (2013) 2:e23565. doi: 10.4161/onci.23565
 46. Fridman WH, Zitvogel L, Sautes-Fridman C, Kroemer G. The immune contexture in cancer prognosis and treatment. *Nat Rev Clin Oncol.* (2017) 14:717–34. doi: 10.1038/nrclinonc.2017.101
 47. Verschuere T, Van Woensel M, Fieuws S, Lefranc F, Mathieu V, Kiss R, et al. Altered galectin-1 serum levels in patients diagnosed with high-grade glioma. *J Neurooncol.* (2013) 115, 9–17. doi: 10.1007/s11060-013-1201-8
 48. Ouyang J, Plutschow A, Pogge von Strandmann E, Reiners KS, Ponader S, Rabinovich GA, et al. Galectin-1 serum levels reflect tumor burden and adverse clinical features in classical Hodgkin lymphoma. *Blood* (2013) 121:3431–3. doi: 10.1182/blood-2012-12-474569
 49. Chen L, Yao Y, Sun L, Zhou J, Liu J, Wang J, et al. Clinical implication of the serum galectin-1 expression in epithelial ovarian cancer patients. *J Ovarian Res.* (2015) 8:78. doi: 10.1186/s13048-015-0206-7
 50. Saussez S, Glinier D, Chantrain G, Pattou F, Carnaille B, Andre S, et al. Serum galectin-1 and galectin-3 levels in benign and malignant nodular thyroid disease. *Thyroid* (2008) 18:705–12. doi: 10.1089/thy.2007.0361
 51. Buchel G, Schulte JH, Harrison L, Batzke K, Schuller U, Hansen W, et al. Immune response modulation by Galectin-1 in a transgenic model of neuroblastoma. *Oncoimmunology* (2017) 5:e1131378. doi: 10.1080/2162402X.2015.1131378
 52. Compagno D, Jaworski FM, Gentilini LD, Contrufo G, Gonzalez Perez I, Pregi N, et al. Galectins: Signaling modulators inside and outside the cell. *Curr. Mol. Med.* (2014) 1–22. doi: 10.2174/1566524014666140603101953
 53. Sundblad V, Morosi LG, Geffner JR, Rabinovich GA. Galectin-1: a jack-of-all-trades in the resolution of acute and chronic inflammation. *J Immunol.* (2017) 199:3721–30. doi: 10.4049/jimmunol.1701172
 54. Ito K, Stannard K, Gabutero E, Clark AM, Neo SY, Onturk S, et al. Galectin-1 as a potent target for cancer therapy: role in the tumor microenvironment. *Cancer Metastasis Rev.* (2012) 31:763–78. doi: 10.1007/s10555-012-9388-2
 55. Deak M, Hornung A, Novak J, Demydenko D, Szabo E, Czibula A, et al. Novel role for galectin-1 in T-cells under physiological and pathological conditions. *Immunobiology* (2015) 220:483–9. doi: 10.1016/j.imbio.2014.10.023

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Corapi, Carrizo, Compagno and Laderach. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.