

Flavonoids Regulate Lipid Droplets Biogenesis in *Drosophila melanogaster*

Natural Product Communications
 May 2019: 1–8
 © The Author(s) 2019
 Article reuse guidelines:
sagepub.com/journals-permissions
 DOI: 10.1177/1934578X19852430
journals.sagepub.com/home/npx



Marianna Fantin^{1*}, Francesca Garelli^{2*}, Barbara Napoli¹,
 Alessia Forgiarini², Sentiljana Gumeni³, Sara De Martin², Monica Montopoli²,
 Chiara Vantaggiato¹, and Genny Orso²

Abstract

Lipid droplets (LDs), cytosolic fat storage organelles, are emerging as major regulators of lipid metabolism, trafficking, and signaling in various cells and tissues. LDs are altered in cardiovascular and neuronal disorders, inflammation, obesity, and cancer. Flavonoids comprise different classes of molecules, characterized by a well-known antioxidant activity and a beneficial effect in several diseases. However, the cellular mechanism by which different classes of flavonoids improve health is poorly understood, in particular as far as LDs biogenesis is concerned. Here we used *Drosophila melanogaster* as a model system to investigate the effects of a selected group of flavonoids on larval tissues by examining LDs biogenesis. In our study, fruit flies were grown in xanthohumol-, isoquercetin-, and genistein-enriched food and larval tissues were analyzed using a LD marker. Total mRNA expression of two main enzymes (minotaur and midway) responsible for triacylglycerides synthesis was evaluated after treatments. Among the flavonoids analyzed, xanthohumol and isoquercetin resulted to be potent regulators of LDs biogenesis in a tissue-specific manner, inducing fat storage decrease in fat bodies and accumulation of LDs in nerves. Since LDs have been suggested to play a protective role against intracellular stress in nonadipocyte cells, our data support the hypothesis that some phytochemicals could act as strong modulators of LDs biogenesis in vivo. The knowledge of how different flavonoids act on lipid metabolism in different tissues can help to manage the use of phytochemicals with the aim of selectively ameliorating specific neuronal and metabolic diseases' manifestations.

Keywords

lipid droplets biogenesis, flavonoids, *Drosophila melanogaster*

Received: January 15th, 2019; Accepted: March 7th, 2019.

Flavonoids are the largest group of phenolic compounds in plants and are synthesized through phenyl propanoid pathway against microbial infection.¹ Numerous studies have shown that food and beverage containing flavonoids, such as fruits, vegetables, cereals, red wine, and green tea, can improve health and participate in the prevention of numerous diseases.^{2,3} Moreover, emerging evidences suggest that flavonoids may have beneficial effects on cardiovascular diseases, neuronal disorders, inflammation, and obesity in animals as well as in humans.^{4–8} This protective mechanism seems to be related to their free radicals scavenging activity, inhibition of enzymes activity (such as aldose reductase, xanthine oxidase, lipoxygenase, and cyclic nucleotide phosphodiesterases [PDEs]), and modulation of protein kinase (e.g., AMP kinase), lipid kinase signaling, and peroxisome proliferator-activated receptor pathways.^{9,10} Several in vivo studies had revealed that flavonoids help to reduce body weight gain, food consumption, and fat accumulation.^{11,12} In mammals, fat is stored in the adipose tissue, which is made

principally of white and brown adipocytes. It is well known that selected flavonoids can induce lipolysis in adipose tissue, likely through inhibition of PDEs and antagonism of cyclic adenosine monophosphate degradation.¹⁰

¹ Scientific Institute, IRCCS E. Medea, Laboratory of Molecular Biology, Bosisio Parini, Lecco, Italy

² Department of Pharmaceutical and Pharmacological Sciences, University of Padova, Italy

³ Department of Cell Biology and Biophysics, Faculty of Biology, National and Kapodistrian University of Athens, Greece

*The authors Marianna Fantin and Francesca Garelli contributed equally to the work.

Corresponding Author:

Genny Orso, Department of Pharmaceutical and Pharmacological Sciences, University of Padova, Padova, Italy.
 Email: genny.orso@unipd.it



The lipids are stored as single or multiple droplets called lipid droplets (LDs). LDs are cellular specialized organelles that store neutral lipids in all living organisms composed by a core, containing mainly triacylglycerides (TAG) and sterol esters, enclosed by a phospholipid monolayer.¹³ LDs are particularly important in tissues specialized for energy storage or lipid turnover, such as the adipose tissue, the liver, and the intestine, and accumulate in skeletal muscles and nervous system.¹⁴⁻¹⁷ LDs not only provide substrates for energy metabolism and building blocks for membranes, but also play a pivotal role in various cellular pathways, such as protein trafficking, protein degradation, and modulation of nuclear receptors.¹⁴ Moreover, LDs exhibit a protective function against oxidative damages caused by different stimuli leading to endoplasmic reticulum (ER) stress.¹⁸⁻²² Simple animal models as *Caenorhabditis elegans* and *Drosophila melanogaster* have been used to study the effects of flavonoids-rich diet on longevity, locomotors activity, fertility, and lethality. However, only few studies were conducted to investigate the relationship between the antioxidant effect of flavonoids and LDs biogenesis at cellular and tissue levels.²³⁻²⁵ How different classes of flavonoids modulate lipid metabolism and intracellular LDs in various tissues is poorly understood. On the basis of this considerations, the aim of this study was to evaluate the effects of flavonoids on cytosolic LDs content in *Drosophila* fat bodies and oenocytes (adipose- and liver-like larval tissues), and as well in tissues with high energy demands such as the central nervous system and muscles.

The effects of genistein (an isoflavone), xanthohumol (a prenylated flavonol), and isoquercetin (a flavonol) on LDs mobilization in fruit fly fat body were evaluated. *Drosophila* fat body is considered the analogous of white adipose tissue in humans. It is known that some flavonoids exhibit lipid-lowering activity and decrease LDs in adipocytes through lipolysis by different mechanisms reducing LDs' number.²⁶⁻²⁸

Fat tissues were dissected from control larvae and stained with a lipophilic dye (BODIPY 493/503) to detect neutral lipids (Figure 1a). In the fat body cells, all the flavonoids induced a decrease of the percentage of cytoplasm area occupied by LDs, from 35.15% of the control to 30.17%, 32.56%, 26.82%, and 29.04% in the genistein, xanthohumol, and isoquercetin exposure, respectively (Figure 1b). Genistein induced similar changes, but only the increase of the smaller LDs' size (of about 6.4 μm) resulted statistically different (Figure 1c and d).

The *Drosophila* specialized cells called oenocytes are considered the analogous of mammal hepatocytes and could be used as a model for lipid metabolism studies.²⁹ When LDs from fat bodies are mobilized in order to furnish energy, the LDs content in oenocytes increased. In addition, under starved conditions, stimulated lipolysis results in high levels of circulating lipids and subsequently causes strong LDs




Figure 1. (a) Representative confocal images of larval fat bodies in control and treated animals. The compounds utilized were genistein (10 μM), xanthohumol (10 μM), and isoquercetin (10 μM). The LDs were visualized by using BODIPY 493/503. Scale bar = 50 μm . Percentage of fat body cell cytoplasm occupied by LDs was calculated (b). Quantification of LDs number (c) and diameter (d). (e) The results of the LDs' size were plotted as box-and-whisker plot, showing the interquartile range, the median as horizontal bar, and the whiskers are the minimum and maximum values. Values are mean \pm SD from at least 3 independent experiments. Significance was calculated using unpaired *t*-test (two-tailed) for LDs number and one-way analysis of variance with Tukey's post hoc test for LDs size. $P < 0.05^*$ and $P < 0.01^{**}$ vs control. LD: lipid droplet.

staining in oenocytes. To correlate the effect of lipid mobilization in the fat body after chronic flavonoids treatment, we analyzed LDs distribution changes in oenocytes (Figure 2a). As expected, LDs in control larvae were barely present, while their number increased upon treatments with almost all the flavonoids used, except genistein (Figure 2b). Indeed, xanthohumol and isoquercetin led to increase of LDs content and induced the formation of smaller LDs (Figure 2c and d). The enhancement of oenocyte LDs amount by xanthohumol (1.24 LDs number/ μm^2) and isoquercetin (0.44 LDs number/ μm^2) treatment suggests an active lipolysis and breakdown of LDs in the fat body, whereas genistein, at least at the concentrations used in this study, has no clear lipolytic effect.

In *Drosophila*, LDs are produced in glial cells and are demonstrated to be important modulators of neuronal function and maintenance.³⁰ In *Drosophila* wild-type larvae, all the treatments led to a statistically significant increase of LDs number (Figure 3a and b) without affecting (Figure 3c and d). We found that in control larvae were present 0.044 LDs/ μm^2 , while in the larvae treated with genistein were observed 0.05 LDs/ μm^2 . Moreover, the number of LDs upon xanthohumol and isoquercetin treatment was increased to 0.06 LDs/ μm^2 . Our data show that glial cells are sensitive to flavonoids supplementation.

Muscle fibers are full of LD in insect and mammals and are localized in close proximity to mitochondria in the subsarcolemmal region near the plasma membrane or in-between the




Figure 2. (a) Representative confocal images of larval oenocytes in control and treated animals. The compounds utilized were genistein (10 μM), xanthohumol (10 μM), and isoquercetin (10 μM). The LDs were visualized by using BODIPY 493/503. Scale bar = 20 μm . Both number (b) and diameter (c) of LDs were measured. (d) The results of the LDs size were plotted as box-and-whisker plot, showing the interquartile range, the median as horizontal bar, and the whiskers are the minimum and maximum values. Significance was calculated using unpaired *t*-test (two-tailed) for LDs number and one-way analysis of variance with Tukey's post hoc test for LDs size. $P < 0.05^*$ and $P < 0.01^{**}$ vs control. LD: lipid droplet.

myofibrils. The effects of flavonoids on LDs in the muscle tissues were analyzed (Figure 4a). Xanthohumol and isoquercetin led to a significant increase of LDs number: 0.114 LDs/ μm^2 and 0.165 LDs/ μm^2 , respectively, compared to 0.091 LDs/ μm^2 of the control (Figure 4b). Intriguingly, the treatment with genistein was not able to modify the number or the mean size of muscle LDs, compared to control; however, it changed the LDs' profile size, similar to the xanthohumol administration. In control larvae, we observed that LDs diameters range was between 0.4 and 1 μm , while in muscle larval tissues exposed to genistein and xanthohumol, the LDs diameter ranged from 0.6 to 0.8 μm (Figure 4d). Taken together, our results show that the flavonoids tested have also an undeniable effect on LDs biogenesis in muscles.

To assess the effect of different classes of flavonoids on LDs biogenesis, we evaluated the expression of the 2 key enzymes of *Drosophila* fatty acid metabolism, using a real-time reverse transcriptase polymerase chain reaction (RT-PCR) approach. The human glycerol-3-phosphate acyltransferase and diacylglycerol *O*-acyltransferase 1 (DGAT1) are encoded by *Drosophila minotaur (mino)* and *midway*




Figure 3. (a) Maximum-intensity projections of proximal ventral ganglion nerves from control or treated *Drosophila* third instar larvae. The compounds utilized were genistein (10 μM), xanthohumol (10 μM), and isoquercetin (10 μM). Nerves were labeled with anti-horseradish peroxidase antibody (red) to visualize neuronal membranes and BODIPY 493/503 dye (green) to determine the number and size of LDs. Scale bar = 10 μm . Both number (b) and size (c) of LDs were measured. (d) The results of the LDs size were plotted as box-and-whisker plot, showing the interquartile range, the median as horizontal bar, and the whiskers are the minimum and maximum values. Values are mean \pm SD from at least 3 independent experiments. Significance was calculated using unpaired *t*-test (two-tailed) for LDs number and one-way analysis of variance with Tukey's post hoc test for LDs size. $P < 0.05^*$ and $P < 0.01^{**}$ vs control. LD: lipid droplet.

(*mdy*) genes, respectively³⁰ (Figure 5a). We, thus, investigated whether the protective effects of flavonoids were linked to lipid droplets metabolism. Total RNA from larvae grown in standard and flavonoids-enriched food were extracted and subjected to quantitative real-time PCR procedure. As shown in Figure 5(b), isoquercetin and xanthohumol strongly enhanced both the transcripts expression of *mino* and *mdy*, whereas genistein administration had the minor effect.

This is the first in vivo study conducted in *Drosophila* reporting the modulatory effects on LDs in different tissues due to phytochemical consumption. Our data show that different classes of flavonoids, at the concentrations here reported, can regulate lipid droplets biogenesis in *Drosophila* larvae and their effects are tissue specific. Almost all the flavonoids tested decreased LDs accumulation in fat bodies and increased LDs number and/or size in oenocytes and muscles,




Figure 4. (a) Representative images of *Drosophila* larval muscles labeled with BODIPY 493/503. The compounds utilized were genistein (10 μ M), xanthohumol (10 μ M), and isoquercetin (10 μ M). Scale bar = 20 μ m. Quantification of LDs number (b) and diameter (c) of LDs. (d) The results of the LDs size were plotted as box-and-whisker plot, showing the interquartile range, the median as horizontal bar, and the whiskers are the minimum and maximum values. Values are mean \pm SD from at least 3 independent experiments. Significance was calculated using unpaired *t*-test (two-tailed) for LDs number and one-way analysis of variance with Tukey's post hoc test for LDs size. $P < 0.05^*$ and $P < 0.01^{**}$ vs control. LD: lipid droplet.

with the exceptions of genistein, which displayed a marginal effect on LDs in all the tissues analyzed. Furthermore, the effects of flavonoids on LDs seem to be correlated to the transcriptional levels of the metabolic enzymes *mino* (GPAT) and *mdy* (DGAT1), both involved in LDs biosynthesis. Accordingly, the transcriptional level of *mino* and *mdy* were barely upregulated after genistein administration. To our knowledge, the effects of flavonoids on LDs are not well studied and the research of LDs regulation and homeostasis has been principally focused on adipocyte cells. Flavonoids are considered antiobesity and lipid-lowering agents but they are also shown to display neuroprotective activity and prevent muscle atrophy.³¹⁻³⁴ Adipocyte specialized cells and their LDs have distinct properties compared to nonadipocyte cells. Increased LDs storage in adipocytes is considered a pathologic condition, while an increase of LDs in nervous system and muscles is considered a protective mechanism.^{16,30,35-38}

Our results confirm that flavonoids administrated to *Drosophila*, similar to mammalian systems, decreased fat storage in adipocyte-like tissues. LDs number, size, and total area occupied by LDs are decreased after flavonoids




Figure 5. (a) TAG de novo synthesis in mammals and *Drosophila*. (b) Relative mRNA levels of minotaur and midway after treatments. The compounds utilized were genistein (10 μ M), xanthohumol (10 μ M), and isoquercetin (10 μ M). Values are mean \pm SD from at least 3 different experiments. Significance was calculated using unpaired *t*-test (two-tailed). $P < 0.05^*$ vs control. GPAT: glycerol-3-phosphate acyltransferase; LPA: lysophosphatidic acid; AGPAT: 1-acylglycerol-3-phosphate acyltransferase; PA: phosphatidic acid; DAG: diacylglycerol; DGAT: diacylglycerol acyltransferase; TAG: triacylglycerol.

administration. This finding is in accordance with a previous study reporting that polyphenols as theaflavin, epigallocatechin gallin, and epigallocatechin suppressed LDs accumulation in larval fat body of a *Drosophila* model of obesity. The authors also reported that the 3 polyphenols prevented the changes of expression in lipid metabolism-related genes, induced by high-fat diet and thereby improved lipid metabolism.³⁹ Antidiabetic and dyslipidemia-correcting effects of polyphenols may be explained by different mechanisms that modulate the adipocyte number causing apoptosis and decreasing adipogenesis, or by modulating the lipid accumulation through activation of lipolysis and oxidation of fatty acids, or by decreasing lipid synthesis.^{12,40,41} In our study, the increase of LDs in oenocytes suggests that fat body lipolysis is increased after flavonoids administration.

On the other hand, treatment with isoquercetin, genistein, and xanthohumol in *Drosophila* greatly modified LDs in nerves and muscles. Changes in the number of LDs have been associated to neurological disorders^{42,43} and mutations of several LDs proteins are linked to motor neuron diseases.⁴⁴ Diets rich in polyphenolic compounds, like flavonoids, were associated with better cognitive functions and reduced risk of dementia and age-associated degenerative diseases.^{8,45} In *Drosophila*, oxidative stress and hypoxia stimulate LDs biosynthesis in order to protect both neuroblasts and glia from peroxidation. LDs accumulation limits the reactive oxygen species (ROS) levels and inhibits the increase of polyunsaturated fatty acids.⁴⁶ Furthermore, studies performed in mice and *Drosophila* showed that LDs accumulation occurs in glial cells in a mitochondria dysfunction model as a protective mechanism.⁴⁷ This protection is attributed to the metabolic exchange between glia and neurons: glial lactate can supply neuronal lipogenesis in response to ROS production, and neuronal lipids are transported and stored in glia as lipid droplets.¹⁰ Flavonoids can protect against cognitive deficits, neuronal injury, and oxidative stress induced in a rat model of Alzheimer's disease, whereas genistein administration had the minor effect.⁴⁸ Similarly, isoquercetin and quercetin protect neurons from different cellular stresses.³³ It is also demonstrated that certain flavonoids protect muscle functionality,^{34,49} and proved that LDs accumulation in muscles is a protective mechanism against the toxicity of lipid intermediates accumulation, such as lipid-derived DAGs and ceramides.⁵⁰

In this scenario, the isolation of certain lipids in specific cell compartments, such as LDs, is conceived as a lifeguard compensatory strategy. Considering that several studies demonstrate that some phytochemicals exert their beneficial effects through the activation of the adaptive stress response-signaling pathway, the effects of the flavonoids used could be due to the hormetic mechanism. The main pathways implicated in adaptive response include the activation of Nrf2/Keap1 and its interaction with other signaling such as Sirtuin-FOXO and NF- κ B signaling.⁵¹ More precisely, several studies suggest that phytochemicals act as a low-dose stressor activating adaptive response to prepare cell against stress condition. The accumulation of LDs could be a cell-type response to protect nonadipocytes from ER stress and lipotoxicity. Moreover, we found that LD changes in *Drosophila* tissues were associated with increased *mdy* and *mino* expression levels, suggesting the potential ability of certain flavonoids to promote LDs biogenesis. DGAT1 upregulation, which we observed after flavonoids treatment, has been demonstrated to protect cells from lipotoxicity and ER stress in both *Drosophila* and mammalian in vitro and in vivo models. DGAT1, in fact, functions in fatty acids re-esterification and formation of new small LDs after lipolysis to protect ER from lipotoxic stress and adipose tissue inflammation.²² Moreover, DGAT1 protects mammalian cells during starvation-induced autophagy and increases LDs by

activating the biogenesis to sequester FAs in TAG-rich LDs, preventing acylcarnitine accumulation and subsequent mitochondrial dysfunction.²¹ In a *Drosophila* model of Huntington's disease, the flies carrying a mutated Huntington protein (Htt) showed an alteration in the size and distribution of LDs.⁴³ In this light, the study of LDs biogenesis and turnover in cellular and animal models of neurological and muscle genetic disorders such as amyotrophic lateral sclerosis, myoclonic epilepsy, ataxia, and muscular dystrophy,⁵²⁻⁵⁶ for which pharmacological treatments are still not available, could help to better clarify the biological role and importance of LDs in healthy and diseased states, as well as their pharmacological impact in dysfunctional conditions. In conclusion, the pharmacological modulation of lipid metabolism could be a potential novel molecular-based approach for treating diseases due to dysfunctions of lipid storage. A major comprehension of phytochemicals mechanism of action and their biological targets can promote a more selective use of phytochemicals to repair pathological conditions. The knowledge of how different flavonoids act on lipid metabolism in different tissues can help to manage the use of phytochemicals with the aim of selectively ameliorating specific neuronal and metabolic diseases manifestations.

Experimental

Drosophila melanogaster Strain and Maintenance Conditions

A white mutant strain w1118 was obtained from the Blomington stock center and used in all experiments. The flies were reared on a standard corn flour medium (4.4% w/v yeast granules, 4.4% w/v sucrose, 1.1% agar, and 2.6% w/v nipagin dissolved in ethanol 96%) at constant temperature of $25 \pm 1^\circ\text{C}$ and under a 12-hour dark/12-hour light cycle.

Antibodies, Chemicals, and Reagents

Antibodies and probe used for immunofluorescence analysis were anti-horseradish peroxidase antibody Cy3 or Cy5 conjugated (Jackson ImmunoResearch, Inc., West Grove, PA, USA), to bind *Drosophila* neuronal membranes, and BODIPY 493/503 dye (Thermo Fisher Scientific Inc., Waltham, MA, USA), to visualize and determine the LDs number and size. Trizol reagent was obtained from Thermo Fisher Scientific Inc (Waltham, MA, USA), and Direct-Zol TM RNA MiniPrep kit was purchased from Zymo Research (Irvine, CA, USA). Xanthohumol was purchased from HWI ANALYTIK GMBH (Rülzheim, Germania). All other chemicals used were purchased from Sigma-Aldrich (St Louis, MO, USA).

Flavonoids Treatments

Three different compounds (xanthohumol, isoquercetin, and genistein) were added to regular *Drosophila* food, all at the

final physiological concentration of 10 μM as recommended in Kanazawa⁵⁷. Flavonoids were dissolved in dimethyl sulfoxide. For control flies, the same amount of vehicle without chemicals was added to standard food. Briefly, adult flies from white1118 strain were allowed to lay eggs in flavonoids-enriched food, and in standard medium. Embryos were reared in these conditions until third instar wandering larval stage was concluded. The transcription levels of the main enzymes involved in LDs biogenesis and changes in LDs number and size in fat bodies, oenocytes, muscles, and nerves were then evaluated.

Immunohistochemistry and LDs Quantification

Third instar larvae were raised on regular and flavonoids-enriched food. After harvesting larvae, they were dissected in phosphate-buffered saline (PBS) and fixed in 4% paraformaldehyde in PBS for 10 minutes and washed in PBT 0.3% Triton X-00 (PBTX) as in Antonioli et al.⁵⁸. To detect LDs in muscles, fat bodies, and oenocytes, the permeabilized larvae were incubated with BODIPY 493/503 (1:100 dilution in PBT 0.3%) for 1 hour at room temperature and mounted on glass slides with Mowiol 4-88. For LDs detection in proximal nerves the permeabilized larvae were stained with BODIPY 493/503 (1:100 dilution in PBT 0.3%) and with anti-horseradish peroxidase conjugated to Cy3 (1:500 dilution with PBT 0.3) for 1 hour at room temperature to mark neuronal membranes and mounted on glass slides with Mowiol 4-88. Images were acquired with a Nikon EZ-C1 confocal microscope (Tokyo, Japan) equipped with a Nikon Plan APO 60.0 \times /1.40 oil immersion objective as in Papadopoulos et al.⁵⁹. Sixty Z-stacks with a step size of 1 μm were taken using identical settings. Each stack consisted of 10 to 12 plane images. Ten animals per treatment were analyzed. The area and number of LDs were calculated using Fiji (NIH) particle analyzer tool.

RNA Extraction and Real-Time Quantitative PCR

Total RNA was isolated from 5 third instar larvae per group and was extracted by Trizol reagent and purified using Direct-Zol TM RNA MiniPrep kit according to the manufacturer's instructions. The RNA concentration was determined using a NanoDrop 2000c spectrophotometer (Thermo Fisher Scientific), and the purity of RNA was determined using the 260/280 nm absorbance ratio. A real-time quantitative PCR was performed on Eco Real-Time PCR System (Illumina Inc., San Diego, CA, USA), using One Step SYBR[®] Prime Script TM RT-PCR Kit II (Takara-Clontech, Kusatsu, Japan) as previously described.⁶⁰ The real-time PCR cycling conditions were reverse transcription 50°C for 15 minutes, polymerase activation 95°C for 2 minutes, followed by 40 cycles of 95°C for 15 seconds, 60°C for 1 minute; melting curve 95°C for 15 seconds, 55°C for 15 seconds, and 95°C for 15 seconds for all target genes. Primers used are listed below.

Relative mRNA expression levels were adjusted with the housekeeping gene *rp49* and were calculated by the threshold cycle (Cq) value of each PCR product and normalized using comparative $2^{-\Delta\Delta\text{Ct}}$ method, which normalizes the expression of the target genes relative to a single reference gene and expressed relative to a reference sample. All experiments were performed in triplicate.

midway 5'-TGG TTA TGG CCT TGT TTC AA-3'
 5'-CAC GTC CAT ATT GGA GAA CG-3'
 minotaur 5'-CCC GAG AAG ATG GAA ACA TT-3'
 5'-GGA GGA GGT GGA CTG ATT GT-3'
 rp49 5'-AGG CCC AAG ATC GTG AAG AA-3'
 5'-TCG ATA CCC TTG GGC TTG C-3'

Statistical Analysis

Statistics was performed with Graph Pad Prism 7.0 software. Significance was calculated using unpaired *t*-test (two-tailed) to assess the differences in the number and area of LDs, and one-way analysis of variance followed by post hoc Tukey's honestly significant difference test to assess the differences in the size distribution of LDs. Differences were considered statistically significant at $P < 0.05$. For real-time PCR data were analyzed using unpaired *t*-test (two-tailed). Data are presented as mean \pm SEM. At least 5 images of each sample were acquired and a minimum of 2 random regions of interest (ROI) of 35.5 μm^2 were created from each image and the area of the LDs contained in the ROI was retrieved and analyzed.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported in part by the Ricerca Corrente of Ministry of Italian Health; Fondazione Cariparo, Young Investigator Grant on Pediatric Research; PRID-J SID18-01 from Univeristy of Padova.

References

1. Falcone Ferreyra ML, Rius SP, Casati P. Flavonoids: biosynthesis, biological functions, and biotechnological applications. *Front Plant Sci.* 2012;3:1-15.
2. Pandey KB, Rizvi SI. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid Med Cell Longev.* 2009;2(5):270-278.
3. Scalbert A, Manach C, Morand C, Rémésy C, Jiménez L. Dietary polyphenols and the prevention of diseases. *Crit Rev Food Sci Nutr.* 2005;45(4):287-306.
4. Bakhtiari M, Panahi Y, Ameli J, Darvishi B. Protective effects of flavonoids against Alzheimer's disease-related neural dysfunctions. *Biomed Pharmacoth.* 2017;93:218-229.

5. Kawser Hossain M, Abdal Dayem A, Han J, Hossain MK, Dayem AA, et al. Molecular mechanisms of the anti-obesity and anti-diabetic properties of flavonoids. *Int J Mol Sci*. 2016;17(4):569.
6. Mozaffarian D, Wu JHY. Flavonoids, dairy foods, and cardiovascular and metabolic health. *Circ Res*. 2018;122(2):369-384.
7. Gabbia D, Dall'Acqua S, Di Gangi I, et al. The Phytocomplex from *Fucus vesiculosus* and *Ascophyllum nodosum* controls postprandial plasma glucose levels: an in vitro and in vivo study in a mouse model of NASH. *Marine Drugs*. 2017;15(2):41-53.
8. Joseph JA, Shukitt-Hale B, Lau FC. Fruit polyphenols and their effects on neuronal signaling and behavior in senescence. *Ann N Y Acad Sci*. 2007;1100:470-485.
9. Panche AN, Diwan AD, Chandra SR. Flavonoids: an overview. *J Nutr Sci*. 2016;5:1-15.
10. Janda E, Lascala A, Martino C, et al. Molecular mechanisms of lipid- and glucose-lowering activities of bergamot flavonoids. *Pharma Nut*. 2016;4:S8-S18.
11. Liu C, Ma J, Sun J, et al. Flavonoid-rich extract of *Paulownia fortunei* flowers attenuates diet-induced hyperlipidemia, hepatic steatosis and insulin resistance in obesity mice by AMPK pathway. *Nutrients*. 2017;9(9):959-974.
12. Assini JM, Mulvihill EE, Huff MW. Citrus flavonoids and lipid metabolism. *Curr Opin Lipidol*. 2013;24(1):34-40.
13. Kraemer N, Farese RV, Walther TC. Balancing the fat: lipid droplets and human disease. *EMBO Mol Med*. 2013;5(7):973-983.
14. Onal G, Kutlu O, Gozuacik D, Dokmeci Emre S. Lipid droplets in health and disease. *Lipids Health Dis*. 2017;16(1):128-138.
15. Fujimoto T, Ohsaki Y, Cheng J, Suzuki M, Shinohara Y. Lipid droplets: a classic organelle with new outfits. *Histochem Cell Biol*. 2008;130(2):263-279.
16. Morales PE, Bucarey JL, Espinosa A. Muscle lipid metabolism: role of lipid droplets and perilipins. *J Diabetes Res*. 2017;2017(2 part 1):1-10.
17. Walther TC, Farese RV. Lipid droplets and cellular lipid metabolism. *Annu Rev Biochem*. 2012;81:687-714.
18. Ito J, Ishii N, Akihara R, et al. A high-fat diet temporarily renders SOD1-deficient mice resistant to an oxidative insult. *J Nutr Biochem*. 2017;40:44-52.
19. Lee J, Homma T, Fujii J. Mice in the early stage of liver steatosis caused by a high fat diet are resistant to thioacetamide-induced hepatotoxicity and oxidative stress. *Toxicol Lett*. 2017;277:92-103.
20. Zheng P, Xie Z, Yuan Y, et al. PLIN5 alleviates myocardial ischaemia/reperfusion injury by reducing oxidative stress through inhibiting the lipolysis of lipid droplets. *Sci Rep*. 2017;7(1):1-10.
21. Nguyen TB, Louie SM, Daniele JR, et al. DGAT1-Dependent lipid droplet biogenesis protects mitochondrial function during starvation-induced autophagy. *Dev Cell*. 2017;42(1):9-21.
22. Chitraju C, Mejhert N, Haas JT, et al. Triglyceride synthesis by DGAT1 protects adipocytes from lipid-induced ER stress during lipolysis. *Cell Metab*. 2017;26(2):407-418.
23. Proshkina E, Lashmanova E, Dobrovolskaya E, et al. Geroprotective and radioprotective activity of quercetin, (-)-epicatechin, and ibuprofen in *Drosophila melanogaster*. *Front Pharmacol*. 2016;7:505-521.
24. Saul N, Pietsch K, Menzel R, Stürzenbaum SR, Steinberg CEW. Catechin induced longevity in *C. elegans*: from key regulator genes to disposable soma. *Mech Ageing Dev*. 2009;130(8):477-486.
25. Morselli E, Maiuri MC, Markaki M, et al. Caloric restriction and resveratrol promote longevity through the Sirtuin-1-dependent induction of autophagy. *Cell Death Dis*. 2010;1(1):e10-10.
26. Ke JY, Kliewer KL, Hamad EM, et al. The flavonoid, naringenin, decreases adipose tissue mass and attenuates ovariectomy-associated metabolic disturbances in mice. *Nutr Metab*. 2015;12(1):1-10.
27. Lim S, Soh K, Kuppusamy U. Effects of naringenin on lipogenesis, lipolysis and glucose uptake in rat adipocyte primary culture: a natural antidiabetic agent. *Int J Altern Med*. 2007;5(2):1-5.
28. Ma L, Li K, Wei D, Xiao H, Niu H, Huang W. High Anti-oxidative and Lipid-lowering Activities of Flavonoid Glycosides-rich Extract from the Leaves of *Zanthoxylum bungeanum* in Multi-system. *J Food Nutr Res*. 2015;3(1):62-68.
29. Gutierrez E, Wiggins D, Fielding B, Gould AP. Specialized hepatocyte-like cells regulate *Drosophila* lipid metabolism. *Nature*. 2007;445(7125):275-280.
30. Cabriol-Pol M-J, Khalil B, Rival T, Faivre-Sarrailh C, Besson MT. Glial lipid droplets and neurodegeneration in a *Drosophila* model of complex I deficiency. *Glia*. 2018;66(4):874-888.
31. Eid HM, Nachar A, Thong F, Sweeney G, Haddad PS. The molecular basis of the antidiabetic action of quercetin in cultured skeletal muscle cells and hepatocytes. *Pharmacogn Mag*. 2015;11(41):74-81.
32. Grosso C, Valentão P, Ferreres F, Andrade PB. The use of flavonoids in central nervous system disorders. *Curr Med Chem*. 2013;20(37):4694-4719.
33. Magalingam KB, Radhakrishnan A, Haleagrahara N. Protective effects of quercetin glycosides, rutin, and isoquercitrin against 6-hydroxydopamine (6-OHDA)-induced neurotoxicity in rat pheochromocytoma (PC-12) cells. *Int J Immunopathol Pharmacol*. 2016;29(1):30-39.
34. Mantle D, Falkous G, Perry EK. Effect of flavonoids on protease activities in human skeletal muscle tissue in vitro. *Clin Chim Acta*. 1999;285(1-2):13-20.
35. Liu Z, Huang X. Lipid metabolism in *Drosophila*: development and disease. *Acta Biochim Biophys Sin*. 2013;45(1):44-50.
36. Konige M, Wang H, Sztalryd C. Role of adipose specific lipid droplet proteins in maintaining whole body energy homeostasis. *Biochim Biophys Acta (BBA) - Mol Basis Dis*. 2014;1842(3):393-401.
37. Velázquez AP, Tatsuta T, Ghillebert R, Drescher I, Graef M. Lipid droplet-mediated ER homeostasis regulates autophagy and cell survival during starvation. *J Cell Biol*. 2016;212(6):621-631.
38. Nguyen TB, Olzmann JA. Lipid droplets and lipotoxicity during autophagy. *Autophagy*. 2017;13(11):2002-2003.

39. Kayashima Y, Murata S, Sato M, et al. Tea polyphenols ameliorate fat storage induced by high-fat diet in *Drosophila melanogaster*. *Biochem Biophys Rep.* 2015;4:417-424.
40. Hajiaghaalipour F, Khalilpourfarshbafi M, Arya A. Modulation of glucose transporter protein by dietary flavonoids in type 2 diabetes mellitus. *Int J Biol Sci.* 2015;11(5):508-524.
41. Rigano D, Sirignano C, Tagliatela-Scafati O. The potential of natural products for targeting PPAR α . *Acta Pharm Sin B.* 2017;7(4):427-438.
42. Marshall LL, Stimpson SE, Hyland R, Coorsen JR, Myers SJ. Increased lipid droplet accumulation associated with a peripheral sensory neuropathy. *J Chem Biol.* 2014;7(2):67-76.
43. Aditi K, Shakarad MN, Agrawal N. Altered lipid metabolism in *Drosophila* model of Huntington's disease. *Sci Rep.* 2016;6(1):1-12.
44. Welte MA. Expanding roles for lipid droplets. *Curr Biol.* 2015;25(11):R470-R481.
45. Letenneur L, Proust-Lima C, Le Gouge A, Dartigues JF, Barberger-Gateau P. Flavonoid intake and cognitive decline over a 10-year period. *Am J Epidemiol.* 2007;165(12):1364-1371.
46. Bailey AP, Koster G, Guillemier C, et al. Antioxidant role for lipid droplets in a stem cell niche of *Drosophila*. *Cell.* 2015;163(2):340-353.
47. Liu L, Zhang K, Sandoval H, et al. Glial lipid droplets and ROS induced by mitochondrial defects promote neurodegeneration. *Cell.* 2015;160(1-2):177-190.
48. Khan MB, Khan MM, Khan A, et al. Naringenin ameliorates Alzheimer's disease (AD)-type neurodegeneration with cognitive impairment (AD-TNDCI) caused by the intracerebroventricular-streptozotocin in rat model. *Neurochem Int.* 2012;61(7):1081-1093.
49. Mukai R, Terao J. Role of dietary flavonoids in oxidative stress and prevention of muscle atrophy. *J Phys Fit Sports Med.* 2013;2(4):385-392.
50. Richter EA, Hargreaves M. Exercise, GLUT4, and skeletal muscle glucose uptake. *Physiol Rev.* 2013;93(3):993-1017.
51. Speciale A, Chirafisi J, Saija A, Cimino F. Nutritional antioxidants and adaptive cell responses: an update. *Curr Mol Med.* 2011;11(9):770-789.
52. Rossetto MG, Zanarella E, Orso G, et al. Defhc1.1, a homologue of the juvenile myoclonic gene EFHC1, modulates architecture and basal activity of the neuromuscular junction in *Drosophila*. *Hum Mol Genet.* 2011;20(21):4248-4257.
53. Crimella C, Cantoni O, Guidarelli A, et al. A novel nonsense mutation in the APTX gene associated with delayed DNA single-strand break removal fails to enhance sensitivity to different genotoxic agents. *Hum Mutat.* 2011;32(4):E2118-E2133.
54. Miglietta D, De Palma C, Sciorati C, et al. Naproxinod shows significant advantages over naproxen in the mdx model of Duchenne muscular dystrophy. *Orphanet J Rare Dis.* 2015;10:101-114.
55. Mushtaq Z, Choudhury SD, Gangwar SK, Orso G, Kumar V. Human Senataxin modulates structural plasticity of the neuromuscular junction in *Drosophila* through a neuronally conserved TGF β signalling pathway. *Neurodegener Dis.* 2016;16(5-6):324-336.
56. Fischer J, Lefèvre C, Morava E, et al. The gene encoding adipose triglyceride lipase (PNPLA2) is mutated in neutral lipid storage disease with myopathy. *Nat Genet.* 2007;39(1):28-30.
57. Kanazawa K. Bioavailability of non-nutrients for preventing lifestyle-related diseases. *Trends Food Sci Technol.* 2011;22(12):655-659.
58. Antonioli L, Pellegrini C, Fornai M, et al. Colonic motor dysfunctions in a mouse model of high-fat diet-induced obesity: an involvement of A2B adenosine receptors. *Purinergic Signal.* 2017;13(4):497-510.
59. Papadopoulos C, Orso G, Mancuso G, et al. Spastin binds to lipid droplets and affects lipid metabolism. *PLoS Genet.* 2015;11(4):e1005149-27.
60. De Martin S, Gabbia D, Albertin G, et al. Differential effect of liver cirrhosis on the pregnane X receptor-mediated induction of CYP3A1 and 3A2 in the rat. *Drug Metab Dispos.* 2014;42(10):1617-1626.