



REVIEW

Recent advances in understanding corticotroph pituitary tumor initiation and progression [version 1; peer review: 2 approved]

Ulrich Renner , Denis Ciato , Günter K. Stalla

Max Planck Institute of Psychiatry, Clinical Neuroendocrinology Group, Munich, Germany


V1 First published: 29 Aug 2018, 7(F1000 Faculty Rev):1354 (<https://doi.org/10.12688/f1000research.14789.1>)Latest published: 29 Aug 2018, 7(F1000 Faculty Rev):1354 (<https://doi.org/10.12688/f1000research.14789.1>)**Abstract**

Cushing's disease is the most frequent form of hypercortisolism and is caused by hypophyseal corticotroph adenomas secreting excessive amounts of adrenocorticotrophic hormone. Most of the tumors develop sporadically and only a limited number of corticotroph adenomas have been found to be associated with different neuroendocrine syndromes or with familial isolated pituitary adenomas. The pathogenic mechanisms of corticotroph adenomas are largely unknown, but the discovered aberrant chaperoning activity of heat shock protein 90 on the one hand and the presence of ubiquitin-specific protease 8 mutations on the other hand partially explained the causes of their development. Corticotroph tumors arise initially as benign microadenomas but with time form invasively growing aggressive macroadenomas which can switch to corticotroph carcinomas in extremely rare cases. The mechanisms through which corticotroph tumors escape from glucocorticoid negative feedback are still poorly understood, as are the processes that trigger the progression of benign corticotroph adenomas toward aggressive and malignant phenotypes. This review summarizes recent findings regarding initiation and progression of corticotroph pituitary tumors.

Keywords

corticotroph tumor, USP8 mutation, progression, corticotroph carcinoma

Open Peer ReviewReviewer Status  

	Invited Reviewers	
	1	2
version 1 published 29 Aug 2018		

F1000 Faculty Reviews are written by members of the prestigious **F1000 Faculty**. They are commissioned and are peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

- Francesca Pecori Girdali**, Istituto Auxologico Italiano, Milan, Italy
- Xavier Bertagna**, Universite Paris, Paris, France

Any comments on the article can be found at the end of the article.

Corresponding author: Ulrich Renner (renner@mpipsykl.mpg.de)**Author roles:** Renner U: Conceptualization, Writing – Original Draft Preparation; Ciato D: Writing – Review & Editing; Stalla GK: Writing – Review & Editing**Competing interests:** No competing interests were disclosed.**Grant information:** This work was supported by German Research Council (DFG).*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.***Copyright:** © 2018 Renner U *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**How to cite this article:** Renner U, Ciato D and Stalla GK. **Recent advances in understanding corticotroph pituitary tumor initiation and progression [version 1; peer review: 2 approved]** F1000Research 2018, 7(F1000 Faculty Rev):1354 (<https://doi.org/10.12688/f1000research.14789.1>)**First published:** 29 Aug 2018, 7(F1000 Faculty Rev):1354 (<https://doi.org/10.12688/f1000research.14789.1>)

Introduction

Corticotroph adenomas (CAs) derive from normal corticotroph cells of the anterior pituitary and account for about 4% to 8% of all clinically hormone-active anterior pituitary tumor types¹. Including clinically inactive silent corticotroph tumors, CAs encompass up to 15% of all pituitary tumors². The prevalence for hormone-active CAs is about 40 cases per million and the incidence ranges from 1.2 to 2.4 per million per year. In adults, CAs are diagnosed mainly in the fourth to sixth decade of life and are about three times more prevalent in women¹. No gender preference has been observed in CAs in children, in whom these tumors are rare (<10% of all CAs)³.

CAs have rarely been described with hereditary background, and in the vast majority of sporadic CAs, the genetic background is still largely unknown⁴. Whereas silent CAs are immunopositive for adrenocorticotrophic hormone (ACTH), but do not manifest biochemical or clinical hypercortisolism⁵, the majority of clinically active CAs secrete excessive amounts of ACTH, causing chronic hypercortisolism due to lack of hypothalamus–pituitary–adrenal (HPA) axis feedback regulation at the level of the CA, leading to manifestations of the clinical features of Cushing’s disease¹. Normally, there is a delay of several years between the onset of the disease and its diagnosis because early symptoms of Cushing’s disease are rather unspecific. At the time of diagnosis, about 90% of the patients present with corticotroph microadenomas less than 1 cm in diameter; in some cases, these tumors are hardly detectable with magnetic resonance imaging¹. Transsphenoidal surgery is generally the initial treatment of choice but is not always successful. In the case of incomplete resection (for example, if the tumor is critically located in proximity to the optical nerves), the tumor might expand and become more aggressive over time, requiring repeated surgery, radiotherapy, pharmaceutical therapy, or adrenalectomy¹. In rare cases, CAs finally transform to metastasizing corticotroph carcinomas for which temozolomide treatment has recently been established with some success in single cases^{6,7}.

Genetics of hereditary corticotroph adenomas

Less than 5% of all CAs are familial adenomas representing rare tumor manifestations in various hereditary endocrine syndromes—multiple endocrine neoplasia 1 (MEN1), MEN4, Carney complex (CNC), and DICER1 syndrome—or in familial isolated pituitary adenomas (FIPAs)^{4,8}. Patients with MEN1, which is caused by a mutation of the *MEN1* gene encoding the tumor suppressor menin, mainly manifest prolactinomas and somatotropinomas and only about 5% of their totality is represented by corticotroph tumors^{8,9}. So far, a single case of CA has been described in MEN4, in which the cell cycle regulator p27^{Kip1} is impaired by inactivating mutations of the *CDKN1B* gene¹⁰. In CNC, caused by inactivating germline mutations of the *PRKARIA* gene which encodes the type 1 α regulatory subunit of protein kinase A, one case of CA has been reported¹¹. In DICER1 syndrome, which is associated with the development of different types of blastomas during embryogenesis, 2% of the affected patients develop corticotroph blastomas as the only pituitary blastoma manifestation^{12,13}. Patients with DICER1 syndrome bear an autosomal dominant mutation of the *DICER1*

gene which encodes the DICER protein, a microRNA (miR) processing endoribonuclease. This suggests that a disturbed production of miRs and thus an altered regulation of their target mRNAs may be responsible for the development of the corticotroph blastomas. However, it is not yet known which miRs are affected and how they induce corticotroph tumor development or why only a small subset of affected patients develops corticotroph blastomas^{12,13}.

Hereditary disorders, in which the patients manifest only pituitary tumors, are designated FIPAs. About 20% of all patients with FIPA bear mutations of the aryl hydrocarbon receptor-interacting protein (AIP), whereas in the majority of FIPAs, the underlying mutations are not known. CAs account for about 5% of all pituitary tumors in FIPAs¹⁴.

In several studies with big cohorts of patients with sporadic pituitary adenoma, only three cases of pathogenic AIP mutations and three cases of menin mutations have been found in sporadic corticotropinomas, suggesting that mutations of genes causing hereditary corticotroph tumors play only a minor role in sporadic CA development^{4,15,16}.

Genetic and epigenetic modifications in sporadic corticotroph adenomas

Most of the CAs (>95%) are sporadic tumors of monoclonal origin^{17,18}. Loss of heterozygosity in combination with different mutations leading to loss of tumor suppressor genes or activation of oncogenes (or both) may play a role in the initiation of sporadic CAs¹⁹.

Despite considerable progress during the last decade, sufficient knowledge about the genetic background of CAs is still missing. Important progress has been made with the recent detection of recurrent gain-of-function mutations in the gene encoding a protein called ubiquitin-specific protease 8 (USP8) in about 20% to 60% of CAs in adult patients^{20–22}. As the mutation was not found in any other type of pituitary tumors, it is evident that the mutation is specific for CAs²³. USP8 is involved in the ubiquitination/deubiquitination process, which is a modality of post-translational protein modification^{24,25}. In this complex process, specific proteins are ubiquitinated and then degraded by the lysosome or are rescued from degradation by deubiquitination. USP8 is an enzyme that mediates the deubiquitination of target proteins, and described mutations in CAs lead to an increased activity of the protein. CAs with mutated USP8 exhibit more stable or higher levels of proteins whose activities are regulated by ubiquitination/deubiquitination^{20,21}.

Multiple proteins are known to be regulated at least in part by ubiquitination/deubiquitination, and in principal all USP8 regulated proteins, if expressed in CAs, could be affected by USP8 mutations^{24,25}. In this context, two candidate proteins that might play a pathogenic role and be a pharmacological target in CAs with mutated USP8 have already been identified: the epidermal growth factor receptor (EGFR)^{20,21} and the somatostatin receptor type 5 (SSTR5)²⁶. The tyrosine kinase receptor EGFR is expressed in corticotroph cells and mediates the ACTH-stimulating

effect of the epidermal growth factor (EGF) and transforming growth factor- α (TGF- α)^{27–29}. In CAs with USP8 mutations and thus enhanced deubiquitination, EGFR is stabilized and its expression is increased. Therefore, the ACTH-stimulating activity of EGF/TGF- α is elevated and may contribute to the excessive ACTH production. Indeed, comparative studies between CAs with and without mutated USP8 have already shown that the expression and activity of the EGFR are enhanced in CAs with USP8 mutations,^{20,21} although this could not be confirmed in another study²⁶. In the case of SSTR5, USP8 mutations would lead to an increased expression and activity of this receptor and suggest that CAs with mutated USP8 may respond better to the treatment with pasireotide, a somatostatin analogue that suppresses CA growth and ACTH secretion through SSTR5^{26,30}. In addition, a Cushing's disease-relevant protein, whose expression or function or both are regulated at least in part by ubiquitination/deubiquitination (and thus may be affected by USP8 mutations), is the glucocorticoid receptor (GR)³¹. The latter regulates the responsiveness to glucocorticoids, resistance to which is a typical feature of Cushing's disease. Whether a reduced sensitivity to glucocorticoids by CAs is related to the mutational status of USP8 still needs to be clarified. Interestingly, a recent article has shown evidence that in normal rat corticotroph cells the proopiomelanocortin (POMC) peptide is directly degraded by ubiquitination and therefore that the treatment of rat corticotropes with the polyubiquitination inhibitor K48R enhanced ACTH production³². Whether this is also relevant in human corticotroph tumor cells has yet to be proven. Comparative investigations have demonstrated that CAs with mutated USP8 are smaller and that the prevalence for USP8 mutations is higher in women^{20,21,33}. In a recent study, a lower incidence of USP8 mutations was found in children with Cushing's disease compared with adults^{33,34}. Whereas the prevalence of USP8 mutations was reduced in Crooke's cell adenomas²⁶, the proportion of mutant USP8 in so-called Nelson's tumors (see the "Corticotroph pituitary tumor types" section) was identical to that in corticotropinomas in general³⁵. Patients with USP8 mutations were diagnosed significantly earlier and had higher preoperative 24-hour urinary-free cortisol levels³⁶. After surgery, recurrences were more abundant and appeared significantly earlier in patients with USP8 mutant corticotropinomas³⁶. In contrast to ACTH-secreting CAs, no USP8 mutations have been detected in a small number (n = 13) of silent CAs studied³³ but this has to be confirmed in a bigger cohort of this tumor type. Interestingly, no mutant USP8 has been found in tumors from a big cohort of dogs with Cushing's disease, suggesting that the mutation is primate-specific³⁷.

So far, apart from mutated USP8, no other recurrent mutations have been found in CAs^{4,23,38}. Thus, more than half of the CA cases may have a variable genetic background and may be associated with mutations of different single genes or may be caused by epigenetic mechanisms (see below). So far, several mutations of different genes have been described in CAs, which mainly affected cell cycle-regulating proteins, signal transduction proteins, oncogenes, and tumor suppressors^{4,23,38,39}. Very recently, loss-of-function mutations of the Cdk5 and ABL enzyme substrate 1 (*CABLES1*) gene were found in four patients from a

cohort of 146 pediatric and 35 adult patients with Cushing's disease: two in children and two in young adults⁴⁰. The mutated *CABLES1* protein lost its inhibitory action on corticotropinoma cell growth, which may explain why all affected patients had corticotroph macroadenomas⁴⁰.

Given that mutations play a tumorigenic role in only a part of the corticotropinomas, it is thought that epigenetic changes such as DNA methylation, histone methylation/acetylation, and miR are of considerable impact for corticotropinoma formation^{41,42}. Most studies on this subject have been performed in unselected cohorts of different types of pituitary adenomas; among them, corticotropinomas and epigenetic changes of different factors have been detected in subsets of the adenomas. Corticotropinoma-specific studies on histone deacetylases have demonstrated that a loss of HDAC2 might play a role in inducing glucocorticoid resistance⁴³, and treatment with histone deacetylase inhibitors reduced survival and ACTH secretion in corticotroph tumor cells⁴⁴. Many of the epigenetic changes caused by DNA methylation or histone methylation/acetylation have direct or indirect effects on the expression of the tumor suppressors p53 and retinoblastoma (Rb) protein, suggesting that these factors are prominent targets of epigenetic changes^{41,42}. A case report of loss of Rb expression in a corticotroph carcinoma⁴⁵ supports the concept that this tumor suppressor plays a role in corticotropinoma development but apparently it is involved in tumor progression rather than in adenoma initiation.

In regard to miRs, which bind to specific messenger RNAs and block their translation into corresponding proteins, several entities have been reported to be either over- or under-expressed in pituitary adenomas in general^{46–48} or specifically in CAs^{49–51}. For instance, invasive characteristics of pituitary tumors were found to be associated with the over-expression of the miR-106b~25 cluster consisting of miR-25, miR-93, and miR-106b⁵⁰. Moreover, aggressive corticotroph Crooke's cell adenomas had a higher expression of miR-106b~25 than other corticotropinomas, suggesting that this miR cluster is also associated with corticotroph tumor aggressiveness⁵⁰. In another study about miR expression specifically in corticotropinomas, patients presenting reduced miR-141 levels showed a better chance of remission, whereas the reduced expression of other miRs (miR-15a, miR-16, miR-21, miR-141, miR-143, miR-145, miR-150, and let-7a) had no association with corticotropinoma phenotype or clinical parameters of the affected patients⁴⁹. The targets and pathological consequences of miRs in CAs have been identified in only a few cases^{48,51}, and much more work has to be done to identify the pathological mechanisms of aberrant miR expression.

Glucocorticoid resistance in corticotroph adenomas

In normal corticotroph cells, ACTH production and release are under negative feedback control of glucocorticoids and are well balanced. If needed, the ACTH and thus glucocorticoid production can be transiently adapted to specific physiological or pathophysiological conditions (stress, infections, inflammation, and so on)⁵². Corticotroph tumor cells produce excessive amounts of ACTH despite strongly elevated glucocorticoid serum levels, indicating that the negative glucocorticoid

feedback is impaired. It is still not clear why corticotroph tumor cells are resistant to glucocorticoids. Inactivating mutations of the GR encoding nuclear receptor subfamily 3 group C member 1 (*NR3C1*) gene are rare, and as corticotroph tumor cells show mostly enhanced GR levels, the missing response to glucocorticoids cannot be explained by GR downregulation^{23,39,53,54}. Thus, an alternate impairment of the function of GR must be responsible for the glucocorticoid resistance, and two candidates—the testicular orphan nuclear receptor 4 (TR4) and heat shock protein 90 (HSP90)—are discussed to play important roles in GR dysfunction.

TR4 is a nuclear receptor which can both activate and repress transcription in different types of organs. TR4 is overexpressed in CAs and corticotroph tumor cell lines and is able to activate the ACTH encoding *POMC* gene by binding to its promoter^{55,56}. The activation is further enhanced if TR4 is phosphorylated through the mitogen-activated protein kinase/extracellular signal-regulated kinase (MAPK/ERK) pathway. In nude mice harboring corticotroph tumors, TR4 overexpression stimulated ACTH secretion and tumor cell growth⁵⁵. It was shown that TR4 interacts with the GR and could overcome the negative regulation of GR on *POMC* transcription and ACTH secretion, suggesting that TR4 might promote resistance to negative glucocorticoid feedback^{55,56}. Suppression of ERK-mediated TR4 phosphorylation by the ERK inhibitor MEK-162 in a murine model of corticotroph tumors reduced ACTH production and inhibited tumor growth, indicating that targeting TR4 action or suppression (or both) could be a novel therapy for Cushing's disease⁵⁷.

HSP90 is a chaperone protein that stabilizes and regulates different proteins by inducing conformational changes. HSP90 interferes with the GR to facilitate ligand binding and receptor translocation to the nucleus and therefore plays an important role in the function of the GR⁵⁸. However, under certain conditions, HSP90 can impair GR function^{59,60}. HSP90 expression is strongly enhanced in the anterior pituitary in patients with CA in comparison with the normal anterior pituitary, and it was shown that HSP90 overexpression restrains the release of mature GR from the chaperone, leading to partial glucocorticoid resistance⁶¹. C-terminal inhibitors of HSP90 such as novobiocin and the herbal compound silibinin could induce the release of mature GR in corticotroph tumor cells *in vitro*, and *in vivo* silibinin was able to revert glucocorticoid resistance in a corticotroph tumor allograft mouse model, leading to a partial restoration of symptoms of the disease⁶¹. This suggests that silibinin, a safe drug already used in humans for the treatment of liver disease and amanitin intoxication⁶², may represent a novel therapeutic option for the treatment of Cushing's disease and corresponding clinical studies are currently in preparation.

Progression of corticotroph adenomas

Corticotroph pituitary tumor types

According to the 2017 World Health Organization (WHO) classification of pituitary tumors, CAs are characterized by the expression of Tpit, a transcription factor specifically expressed in the ACTH-producing pituitary cell lineage. Histologically, hormone-secreting CAs are further subclassified into densely and

sparsely granulated CAs and Crooke's cell adenomas². Not classified as a specific entity in the 2017 WHO classification are Nelson's tumors, the original designation of aggressively progressing corticotroph macroadenomas developing in 8% to 29% of patients, in which the chronic hypercortisolism had been treated by bilateral adrenalectomy^{63–65}. It was thought that the removal of cortisol excess could play a causative role in the development of Nelson's tumors. However, as CAs in adrenalectomized patients mostly remain stable or progress slowly and as the patients receive adequate cortisol replacement therapy, it is unlikely that the adrenalectomy-associated removal of excessive cortisol alone is responsible for the formation of Nelson's tumors^{63–65}. Nevertheless, owing to elevated proliferative index and enhanced invasive growth, Nelson's tumors can evolve to carcinomas in rare cases⁶⁶. There is no evidence for a common specific genetic background leading to the formation of Nelson's tumors, and in a recent study the proportion of USP8 mutations (45%) was found to be similar to that of corticotroph tumors in general³⁵. Thus, the mechanisms triggering the development of Nelson's tumors are still unknown^{64,65}.

Crooke's cell adenomas are characterized by the replacement of cytoplasmic granules by hyaline material², a process that has been designated Crooke's change. In a recent study, it was shown that depending on the degree of hypercortisolism, densely/sparsely corticotroph tumors show focal Crooke's changes⁶⁷. This suggests that Crooke's cell adenomas have no specific genetic/epigenetic background but derive from densely/sparsely CAs at a high hypercortisolemic state. This is supported by the observation that there is little difference in the proportion of USP8-mutated and non-mutated tumors between densely/sparsely granulated CAs and Crooke's cell adenomas²⁶. At present, the underlying mechanisms triggering the development of Crooke's cell adenomas are not known. In the 2017 WHO high-risk pituitary adenoma subclassification, Crooke's cell adenomas are listed as an aggressive tumor type as these tumors have an elevated proliferative index and are already invasively growing at a microadenoma state². As Crooke's cell adenomas are at risk to transform to corticotroph carcinomas, chemotherapeutic treatment of these adenomas is recommended to prevent the transformation⁶⁸. Silent CAs also belong to the aggressive pituitary tumor types in the new WHO classification. This tumor entity is immunopositive for Tpit and expresses different POMC precursor proteins but does not secrete ACTH; moreover, genes related to ACTH synthesis are differently expressed in comparison with ACTH-secreting CAs^{2,69}. Thus, silent CAs clinically represent a member of the group of non-functioning pituitary adenomas (NFPAs), and according to recent studies, up to 20% of all NFPAs are silent CAs⁵. There are several hypotheses and speculations about silent CAs and their inability to secrete ACTH. Silent CAs may not derive from normal anterior pituitary corticotroph cells but may originate from POMC-expressing cells of the pars intermedia and thus may have characteristics different from those of "normal" CAs⁶⁹. There is also electron microscope-based evidence that silent CAs have elevated lysosome numbers and show fusions of these lysosomes with secretory granules and this has led to speculation that ACTH is intracellularly destroyed before it can be secreted⁷⁰. Alternatively, the

function or expression of POMC product-processing enzymes may be disturbed in silent CAs and indeed abnormalities of these prohormone convertases have been described⁷¹. However, the processes responsible for silent CA development still need to be clarified in future studies. The observation that silent corticotropinomas probably have no mutated USP8 may indicate that silent CAs have a different genetic background compared with classic CAs³³. However, in some cases, silent CAs have spontaneously changed to secretory CAs and vice versa, suggesting that at least in some cases silent and secretory CAs may have the same origin⁷². Like most NFPAs, silent CAs are diagnosed quite late when they have already formed invasively growing macroadenomas. At this state, complete surgical resection is not possible and further clinical management of these tumors is quite challenging; in rare cases, silent CAs finally transform to carcinomas^{73,74}.

Microadenomas, macroadenomas, and aggressive corticotroph tumors

At diagnosis, most of the ACTH-secreting CAs are microadenomas and in some cases are hardly visible by routine imaging techniques¹. When CAs have reached a critical size of more than 2 mm in diameter, oxygen transport to the tumor cells per diffusion will no longer be sufficient and will lead to tumor neovascularization through angiogenesis forced by intratumoral hypoxia^{75,76}. The most important regulator of angiogenesis is the hypoxia-inducible factor-1 (HIF-1), which is inactive under normoxic conditions but active under hypoxia⁷⁷. Activated HIF-1 induces the production of multiple angiogenic factors—for example, vascular endothelial growth factor-A (VEGF-A), basic fibroblast growth factor, and platelet-derived growth factor—and suppresses the formation of anti-angiogenic components. This finally stimulates the proliferation of endothelial cells and pericytes, promotes the formation of tubular structures, and induces the targeted sprouting of the newly formed vessels into the hypoxic zones of the growing tumor⁷⁷. HIF-1 is a heterodimeric protein composed of two subunits: oxygen-regulated HIF-1 α and constitutively expressed HIF-1 β ⁷⁸. Under normoxic conditions, HIF-1 α has a very short half-life due to permanent degradation by ubiquitination and therefore little or no active HIF-1 is present. During hypoxia, the ubiquitination of HIF-1 α is blocked and active HIF-1 is formed to induce the angiogenic processes⁷⁸. As the previously mentioned USP8 protein interferes with HIF-1 α deubiquitination processes^{25,79}, it would be interesting to study whether HIF-1 levels are different in corticotropinomas with and without mutated USP8 and whether corticotropinoma vascularization might be affected as in the case of the product of the *RWWD3* gene, RWD-domain-containing sumoylation enhancer (RSUME)⁸⁰. The latter is upregulated under hypoxia and overexpressed in pituitary tumors, including corticotropinomas⁸¹. It has been shown that RSUME could stabilize HIF-1 and thus enhance VEGF-A secretion. Therefore, the knock-down of RSUME was associated with reduced production of HIF-1 and VEGF-A in pituitary tumors *in vitro* and *in vivo*⁸¹. Whether USP8 has similar HIF-1-stabilizing properties and thus influences angiogenic growth factor production in corticotroph tumors needs to be clarified.

The intratumoral formation of new vessels requires the disruption of cell–cell contacts between tumor cells. Thus, during neovascularization, not only factors stimulating the growth of vessels but also factors degrading the extracellular matrix of tumor cells are produced⁷⁶. In these processes, enzymes of the matrix metalloproteinase (MMP) family and their negative regulators, the tissue inhibitors of MMPs (TIMPs), play important roles^{82,83}, but numerous other factors are also involved⁷⁶. These soluble factors not only may locally act to pave the way for invading vessels but may diffuse into the tumor cell matrix and into the anatomical structures surrounding the adenomas. Therefore, the process of neovascularization bears the risk that matrix degrading components support the migration of tumor cells to invade surrounding tissues at the edge of the tumors^{84,85}. The link between angiogenesis and invasiveness is supported by the observation that RSUME not only has angiogenic properties in corticotroph tumors but also stimulates migration and invasive growth of corticotroph tumor cells⁸⁶. Normally, corticotroph macroadenomas show invasive growth characteristics, but in Crooke's cell adenomas, for instance, invasive growth has already been observed in microadenomas².

Comparative studies between the normal anterior pituitary and CAs have identified hundreds of genes, miRs, proteins, and peptides that are up- or down-regulated in the adenomas⁸⁷. This is not surprising as, in the adult anterior pituitary, cells normally do not grow and neither angiogenic nor invasive processes take place. Thus, all factors related to these events are somehow differentially expressed in the corticotropinomas, but it should be emphasized that these aberrant expression patterns are mainly a consequence, not a cause, of CAs. Correspondingly, there are also differences between non-invasive and invasive growing corticotroph tumors^{88,89}, such as differential expression of the miR106b~25 cluster, the cyclin D2 (*CCND2*), zinc finger protein 676 (*ZNF676*), death-associated protein kinase 1 (*DAPK1*), and *TIMP2* genes as well as genes associated with TGF- β and G protein signaling pathways, DNA damage response pathways, focal adhesion-associated pathways, and others^{48,50,88–92}. Although it has been speculated that all of the aberrantly expressed factors play a role in corticotropinoma expansion and thus represent putative targets for pharmacological treatment, this has been proven *in vivo* in mouse allograft corticotroph tumor models in only a few cases but has not led to corresponding clinical studies.

Corticotroph carcinomas

According to the 2017 WHO classification, pituitary carcinomas, which represent less than 0.1% of all pituitary tumors, are characterized by their ability to form metastases and both hereditary and sporadic pituitary tumors can transform into carcinomas². Although the proportion of CAs within all pituitary tumor types is relatively small, they represent about 35% to 48% of all pituitary carcinomas^{6,7,93}. This may be a consequence of the relative high proportion of aggressive tumor types (Crooke's cell adenomas, Nelson's tumors, and silent CAs) among the corticotroph tumors². In most reports, several common features of corticotroph carcinomas are evident, such as repeated recurrence of the adenomas after incomplete surgical resection and

transient response but finally resistance to different pharmacological therapies or to radiotherapy^{6,7,66,68,73,74,94}. Usually, decades separate the first diagnosis of the CA and the full expression of the carcinoma phenotype. At present, there is no evidence of a specific genetic or epigenetic background or a common factor that is responsible for the development of pituitary carcinomas⁹⁵. In a recent meta-analysis, in which pituitary carcinomas were compared with aggressive adenomas, a number of factors related to the stimulation of cell growth, angiogenesis, and invasiveness (cyclin D1, VEGF, MMP9, miR-122, and miR-493) were found to be up-regulated, whereas several factors, including growth inhibitors and apoptosis inducers (p16Ink4A, p27Kip1, MT3, BCL-2, Bax, Bcl-X, and O⁶-methylguanine DNA methyltransferase [MGMT]), were down-regulated^{95,96}. However, these factors are also differently expressed in invasive versus non-invasive adenomas, in micro- versus macro-adenomas, or in poorly versus densely vascularised pituitary tumors^{76,84,87}. These data suggest that an accumulation of changes associated with increased pituitary tumor aggressiveness rather than a carcinoma-specific mutation or epigenetic change (or both) will finally lead to development of metastasizing pituitary carcinomas^{93,95}.

Temozolomide, an alkylating chemotherapeutic drug, has been identified as an effective compound in the treatment of pituitary carcinomas^{7,94}. The effect of temozolomide is reduced in pituitary tumors with high expression of MGMT. Crooke's cell adenomas, Nelson's tumors, invasively growing corticotroph macroadenomas, and corticotroph carcinomas mostly have low MGMT expression^{2,97-100}. This may explain the relatively good response of corticotroph carcinomas to temozolomide therapy in several cases^{7,94}. However, this also suggests that advanced aggressive corticotropinoma types may already be treated with temozolomide to prevent transformation into corticotroph carcinomas¹⁰¹.

Conclusions

CAs are mostly sporadic tumors; only a small proportion of CAs have a hereditary background and are associated with neuroendocrine syndromes or are FIPAs. In sporadic CAs, a recurrent activating mutation of the *USP8* gene, which is present in about 40% of ACTH-secreting CAs, has recently been found. The *USP8* mutation participates in CA pathophysiology by stimulating EGFR expression and thus enhances EGF-induced ACTH production. As mutated *USP8* also increases *SSTR5* expression, CAs with *USP8* mutation respond better to treatment with the somatostatin analogue pasireotide which acts through *SSTR5*. No other recurrent mutations have been identified in CAs,

suggesting that single different mutations or epigenetic changes (or both) may play a role in CA initiation. The ACTH secretion by corticotroph tumor cells, in contrast to normal corticotroph cells, is resistant to inhibitory glucocorticoid feedback control, and HSP90 overexpression has recently been shown to be involved in Cushing's disease. Thus, HSP90 inhibitors like silibinin may be promising drugs to restore glucocorticoid responsiveness of CAs and to reverse the pathophysiological changes of Cushing's disease. During the progression of CAs from microadenomas to aggressive macroadenomas and finally rare cases of corticotroph carcinomas, multiple factors (genes, miRs, proteins, and peptides) are up- or down-regulated and thus may provide putative targets for new pharmacological treatment strategies. However, despite remarkable progress in understanding corticotroph tumor initiation and progression, a lot of work is still needed to explore the underlying mechanisms, and the translation of new findings into clinical applications for diagnosis and therapy of corticotropinomas is still unsatisfactory.

Abbreviations

ACTH, adrenocorticotrophic hormone; AIP, aryl hydrocarbon receptor-interacting protein; CA, corticotroph adenoma; CABLES1, Cdk5 and ABL enzyme substrate 1; CNC, Carney complex; EGF, epidermal growth factor; EGFR, epidermal growth factor receptor; ERK, extracellular signal-regulated kinase; FIPA, familial isolated pituitary adenomas; GR, glucocorticoid receptor; HIF-1, hypoxia-inducible factor 1; HSP90, heat shock protein 90; MAPK, mitogen-activated protein kinase; MEN, multiple endocrine neoplasia; MGMT, O⁶-methylguanine DNA methyltransferase; miR, microRNA; MMP, matrix metalloproteinase; NFPA, non-functioning pituitary adenoma; POMC, proopiomelanocortin; Rb, retinoblastoma; RSUME, RWD-domain-containing sumoylation enhancer; *SSTR5*, somatostatin receptor type 5; TGF, transforming growth factor; TIMP, tissue inhibitor of metalloproteinase; TR4, testicular orphan nuclear receptor 4; USP8, ubiquitin-specific protease 8; VEGF-A, vascular endothelial growth factor-A; WHO, World Health Organization.

Grant information

This work was funded by a grant from the German Research Council (DFG) to GKS (STA 285/20-1).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Pivonello R, De Leo M, Cozzolino A, *et al.*: **The Treatment of Cushing's Disease.** *Endocr Rev.* 2015; **36**(4): 385–486.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mete O, Lopes MB: **Overview of the 2017 WHO Classification of Pituitary Tumors.** *Endocr Pathol.* 2017; **28**(3): 228–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lodish M: **Cushing's syndrome in childhood: update on genetics, treatment, and outcomes.** *Curr Opin Endocrinol Diabetes Obes.* 2015; **22**(1): 48–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Xiong Q, Ge W: **Gene mutations in Cushing's disease.** *Biomed Rep.* 2016; **5**(3): 277–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)



5. Ben-Shlomo A, Cooper O: **Silent corticotroph adenomas.** *Pituitary.* 2018; **21**(2): 183–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Yoo F, Kuan EC, Heaney AP, et al.: **Corticotrophic pituitary carcinoma with cervical metastases: case series and literature review.** *Pituitary.* 2018; **21**(3): 290–301.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. McCormack A, Dekkers OM, Petersenn S, et al.: **Treatment of aggressive pituitary tumours and carcinomas: results of a European Society of Endocrinology (ESE) survey 2016.** *Eur J Endocrinol.* 2018; **178**(3): 265–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
8. Scherthaner-Reiter MH, Trivellin G, Stratakis CA: **MEN1, MEN4, and Carney Complex: Pathology and Molecular Genetics.** *Neuroendocrinology.* 2016; **103**(1): 18–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Vannucci L, Marini F, Giusti F, et al.: **MEN1 in children and adolescents: Data from patients of a regional referral center for hereditary endocrine tumors.** *Endocrine.* 2018; **59**(2): 438–48.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Georgitsi M, Raitila A, Karhu A, et al.: **Germline *CDKN1B/p27^{Kip1}* mutation in multiple endocrine neoplasia.** *J Clin Endocrinol Metab.* 2007; **92**(8): 3321–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Kiefer FW, Winhofer Y, Iacovazzo D, et al.: ***PRKAR1A* mutation causing pituitary-dependent Cushing disease in a patient with Carney complex.** *Eur J Endocrinol.* 2017; **177**(2): K7–K12.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. de Kock L, Sabbaghian N, Plourde F, et al.: **Pituitary blastoma: a pathognomonic feature of germ-line *DICER1* mutations.** *Acta Neuropathol.* 2014; **128**(1): 111–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Sahakitrungruang T, Srichomthong C, Pornkunwilai S, et al.: **Germline and somatic *DICER1* mutations in a pituitary blastoma causing infantile-onset Cushing's disease.** *J Clin Endocrinol Metab.* 2014; **99**(8): E1487–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Hernández-Ramírez LC, Gabrovská P, Dénes J, et al.: **Landscape of Familial Isolated and Young-Onset Pituitary Adenomas: Prospective Diagnosis in AIP Mutation Carriers.** *J Clin Endocrinol Metab.* 2015; **100**(9): E1242–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Lecoq AL, Bouligand J, Hage M, et al.: **Very low frequency of germline *GPR101* genetic variation and no biallelic defects with AIP in a large cohort of patients with sporadic pituitary adenomas.** *Eur J Endocrinol.* 2016; **174**(4): 523–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Cazabat L, Bouligand J, Salenave S, et al.: **Germline AIP mutations in apparently sporadic pituitary adenomas: prevalence in a prospective single-center cohort of 443 patients.** *J Clin Endocrinol Metab.* 2012; **97**(4): E663–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Herman V, Fagin J, Gonsky R, et al.: **Clonal origin of pituitary adenomas.** *J Clin Endocrinol Metab.* 1990; **71**(6): 1427–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Gicquel C, Le Bouc Y, Luton JP, et al.: **Monoclonality of corticotroph macroadenomas in Cushing's disease.** *J Clin Endocrinol Metab.* 1992; **75**(2): 472–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Biller BM, Alexander JM, Zervas NT, et al.: **Clonal origins of adrenocorticotrophin-secreting pituitary tissue in Cushing's disease.** *J Clin Endocrinol Metab.* 1992; **75**(5): 1303–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Reincke M, Sbierra S, Hayakawa A, et al.: **Mutations in the deubiquitinase gene *USP8* cause Cushing's disease.** *Nat Genet.* 2015; **47**(1): 31–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
21. Ma ZY, Song ZJ, Chen JH, et al.: **Recurrent gain-of-function *USP8* mutations in Cushing's disease.** *Cell Res.* 2015; **25**(3): 306–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
22. Ballmann C, Thiel A, Korah HE, et al.: ***USP8* Mutations in Pituitary Cushing Adenomas-Targeted Analysis by Next-Generation Sequencing.** *J Endocr Soc.* 2018; **2**(3): 266–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
23. Song ZJ, Reitman ZJ, Ma ZY, et al.: **The genome-wide mutational landscape of pituitary adenomas.** *Cell Res.* 2016; **26**(11): 1255–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
24. He M, Zhou Z, Shah AA, et al.: **The emerging role of deubiquitinating enzymes in genomic integrity, diseases, and therapeutics.** *Cell Biosci.* 2016; **6**: 62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Lei H, Shan H, Wu Y: **Targeting deubiquitinating enzymes in cancer stem cells.** *Cancer Cell Int.* 2017; **17**: 101.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Hayashi K, Inoshita N, Kawaguchi K, et al.: **The *USP8* mutational status may predict drug susceptibility in corticotroph adenomas of Cushing's disease.** *Eur J Endocrinol.* 2016; **174**(2): 213–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
27. Renner U, Paez-Pereda M, Arzt E, et al.: **Growth factors and cytokines: function and molecular regulation in pituitary adenomas.** *Front Horm Res.* 2004; **32**: 96–109.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Theodoropoulou M, Arzberger T, Gruebler Y, et al.: **Expression of epidermal growth factor receptor in neoplastic pituitary cells: evidence for a role in corticotropinoma cells.** *J Endocrinol.* 2004; **183**(2): 385–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Cooper O, Vlotides G, Fukuoka H, et al.: **Expression and function of ErbB receptors and ligands in the pituitary.** *Endocr Relat Cancer.* 2011; **18**(6): R197–211.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Pivonello R, Petersenn S, Newell-Price J, et al.: **Pasireotide treatment significantly improves clinical signs and symptoms in patients with Cushing's disease: results from a Phase III study.** *Clin Endocrinol (Oxf).* 2014; **81**(3): 408–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Weikum ER, Knuesel MT, Ortlund EA, et al.: **Glucocorticoid receptor control of transcription: precision and plasticity via allostery.** *Nat Rev Mol Cell Biol.* 2017; **18**(3): 159–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Sesta A, Cassarino MF, Cavagnini F, et al.: **Role of the ubiquitin/proteasome system on ACTH turnover in rat corticotropes.** *Endocrine.* 2018; **61**(3): 511–517.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
33. Perez-Rivas LG, Theodoropoulou M, Ferràù F, et al.: **The Gene of the Ubiquitin-Specific Protease 8 Is Frequently Mutated in Adenomas Causing Cushing's Disease.** *J Clin Endocrinol Metab.* 2015; **100**(7): E997–1004.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
34. Faucz FR, Tirosh A, Tatsi C, et al.: **Somatic *USP8* Gene Mutations Are a Common Cause of Pediatric Cushing Disease.** *J Clin Endocrinol Metab.* 2017; **102**(8): 2836–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. Pérez-Rivas LG, Theodoropoulou M, Puar TH, et al.: **Somatic *USP8* mutations are frequent events in corticotroph tumor progression causing Nelson's tumor.** *Eur J Endocrinol.* 2018; **178**(1): 59–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Albani A, Pérez-Rivas LG, Dimopoulou C, et al.: **The *USP8* mutational status may predict long-term remission in patients with Cushing's disease.** *Clin Endocrinol (Oxf).* 2018.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Sbierra S, Tryfonidou MA, Weigand I, et al.: **Lack of Ubiquitin Specific Protease 8 (*USP8*) Mutations in Canine Corticotroph Pituitary Adenomas.** *PLoS One.* 2016; **11**(2): e0169009.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
38. Uzilov AV, Cheesman KC, Fink MY, et al.: **Identification of a novel *RASD1* somatic mutation in a *USP8*-mutated corticotroph adenoma.** *Cold Spring Harb Mol Case Stud.* 2017; **3**(3): a001602.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
39. Karl M, Von Wichert G, Kempter E, et al.: **Nelson's syndrome associated with a somatic frame shift mutation in the glucocorticoid receptor gene.** *J Clin Endocrinol Metab.* 1996; **81**(1): 124–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Hernández-Ramírez LC, Gam R, Valdés N, et al.: **Loss-of-function mutations in the *CABLES1* gene are a novel cause of Cushing's disease.** *Endocr Relat Cancer.* 2017; **24**(8): 379–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
41. Zhou Y, Zhang X, Klibanski A: **Genetic and epigenetic mutations of tumor suppressive genes in sporadic pituitary adenoma.** *Mol Cell Endocrinol.* 2014; **386**(1–2): 16–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Ezzat S, Cheng S, Asa SL: **Epigenetics of pituitary tumors: Pathogenetic and therapeutic implications.** *Mol Cell Endocrinol.* 2018; **469**: 70–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
43. Bilodeau S, Vallette-Kasic S, Gauthier Y, et al.: **Role of Brg1 and HDAC2 in GR trans-repression of the pituitary *POMC* gene and misexpression in Cushing disease.** *Genes Dev.* 2006; **20**(20): 2871–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
44. Lu J, Chatain GP, Bugarini A, et al.: **Histone Deacetylase Inhibitor SAHA Is a Promising Treatment of Cushing Disease.** *J Clin Endocrinol Metab.* 2017; **102**(8): 2825–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
45. Hinton DR, Hahn JA, Weiss MH, et al.: **Loss of Rb expression in an ACTH-secreting pituitary carcinoma.** *Cancer Lett.* 1998; **126**(2): 209–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Wierinckx A, Roche M, Legras-Lachuer C, et al.: **MicroRNAs in pituitary tumors.** *Mol Cell Endocrinol.* 2017; **456**: 51–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Zhang T, Yang Z, Gao H: **Advancements in the study of miRNA regulation during the cell cycle in human pituitary adenomas.** *J Neurooncol.* 2017; **134**(2): 253–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

48. **F** Wang Y, Yin X, Zhao L, *et al.*: **MicroRNA-200b inhibits pituitary tumor cell proliferation and invasion by targeting PKC α .** *Exp Ther Med.* 2017; **14**(2): 1706–14. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
49. Amaral FC, Torres N, Saggioro F, *et al.*: **MicroRNAs differentially expressed in ACTH-secreting pituitary tumors.** *J Clin Endocrinol Metab.* 2009; **94**(1): 320–3. [PubMed Abstract](#) | [Publisher Full Text](#)
50. **F** Garbicz F, Mehlich D, Rak B, *et al.*: **Increased expression of the microRNA 106b-25 cluster and its host gene MCM7 in corticotroph pituitary adenomas is associated with tumor invasion and Crooke's cell morphology.** *Pituitary.* 2017; **20**(4): 450–63. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
51. Gentilin E, Tagliati F, Filieri C, *et al.*: **miR-26a plays an important role in cell cycle regulation in ACTH-secreting pituitary adenomas by modulating protein kinase C δ .** *Endocrinology.* 2013; **154**(5): 1690–700. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Smith SM, Vale WW: **The role of the hypothalamic-pituitary-adrenal axis in neuroendocrine responses to stress.** *Dialogues Clin Neurosci.* 2006; **8**(4): 383–95. [PubMed Abstract](#) | [Free Full Text](#)
53. Karl M, Lamberts SW, Koper JW, *et al.*: **Cushing's disease preceded by generalized glucocorticoid resistance: clinical consequences of a novel, dominant-negative glucocorticoid receptor mutation.** *Proc Assoc Am Physicians.* 1996; **108**(4): 296–307. [PubMed Abstract](#)
54. Dahia PL, Honegger J, Reincke M, *et al.*: **Expression of glucocorticoid receptor gene isoforms in corticotropin-secreting tumors.** *J Clin Endocrinol Metab.* 1997; **82**(4): 1088–93. [PubMed Abstract](#) | [Publisher Full Text](#)
55. **F** Du L, Bergsneider M, Mirsadraei L, *et al.*: **Evidence for orphan nuclear receptor TR4 in the etiology of Cushing disease.** *Proc Natl Acad Sci U S A.* 2013; **110**(21): 8555–60. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
56. **F** Zhang D, Du L, Heaney AP: **Testicular Receptor-4: Novel Regulator of Glucocorticoid Resistance.** *J Clin Endocrinol Metab.* 2016; **101**(8): 3123–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
57. **F** Zhang D, Bergsneider M, Wang MB, *et al.*: **Targeting the ERK pathway for the treatment of Cushing's disease.** *Oncotarget.* 2016; **7**(43): 69149–58. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
58. **F** Kirschke E, Goswami D, Southworth D, *et al.*: **Glucocorticoid receptor function regulated by coordinated action of the Hsp90 and Hsp70 chaperone cycles.** *Cell.* 2014; **157**(7): 1685–97. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
59. Kang KI, Meng X, Devin-Leclerc J, *et al.*: **The molecular chaperone Hsp90 can negatively regulate the activity of a glucocorticosteroid-dependent promoter.** *Proc Natl Acad Sci U S A.* 1999; **96**(4): 1439–44. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. **F** Freeman BC, Yamamoto KR: **Disassembly of transcriptional regulatory complexes by molecular chaperones.** *Science.* 2002; **296**(5576): 2232–5. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
61. **F** Riebold M, Kozany C, Freiburger L, *et al.*: **A C-terminal HSP90 inhibitor restores glucocorticoid sensitivity and relieves a mouse allograft model of Cushing disease.** *Nat Med.* 2015; **21**(3): 276–80. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
62. Saller R, Meier R, Brignoli R: **The use of silymarin in the treatment of liver diseases.** *Drugs.* 2001; **61**(4): 2035–63. [PubMed Abstract](#) | [Publisher Full Text](#)
63. Assié G, Bahurel H, Bertherat J, *et al.*: **The Nelson's syndrome... revisited.** *Pituitary.* 2004; **7**(4): 209–15. [PubMed Abstract](#) | [Publisher Full Text](#)
64. Patel J, Eloy JA, Liu JK: **Nelson's syndrome: a review of the clinical manifestations, pathophysiology, and treatment strategies.** *Neurosurg Focus.* 2015; **38**(2): E14. [PubMed Abstract](#) | [Publisher Full Text](#)
65. **F** Assié G, Bahurel H, Coste J, *et al.*: **Corticotroph tumor progression after adrenalectomy in Cushing's Disease: A reappraisal of Nelson's Syndrome.** *J Clin Endocrinol Metab.* 2007; **92**(1): 172–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
66. Kemink SA, Wesseling P, Pieters GF, *et al.*: **Progression of a Nelson's adenoma to pituitary carcinoma; a case report and review of the literature.** *J Endocrinol Invest.* 1999; **22**(1): 70–5. [PubMed Abstract](#) | [Publisher Full Text](#)
67. Oldfield EH, Vance ML, Louis RG, *et al.*: **Crooke's Changes In Cushing's Syndrome Depends on Degree of Hypercortisolism and Individual Susceptibility.** *J Clin Endocrinol Metab.* 2015; **100**(8): 3165–71. [PubMed Abstract](#) | [Publisher Full Text](#)
68. **F** Kurowska M, Tarach JS, Malicka J, *et al.*: **Long-term complete remission of Crooke's corticotropinoma after temozolomide treatment.** *Endokrynol Pol.* 2016; **67**(5): 526–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
69. Tateno T, Izumiyama H, Doi M, *et al.*: **Differential gene expression in ACTH-secreting and non-functioning pituitary tumors.** *Eur J Endocrinol.* 2007; **157**(6): 717–24. [PubMed Abstract](#) | [Publisher Full Text](#)
70. Kovacs K, Horvath E, Bayley TA, *et al.*: **Silent corticotroph cell adenoma with lysosomal accumulation and crinophagy. A distinct clinicopathologic entity.** *Am J Med.* 1978; **64**(3): 492–9. [PubMed Abstract](#) | [Publisher Full Text](#)
71. Tateno T, Izumiyama H, Doi M, *et al.*: **Defective expression of prohormone convertase 1/3 in silent corticotroph adenoma.** *Endocr J.* 2007; **54**(5): 777–82. [PubMed Abstract](#) | [Publisher Full Text](#)
72. **F** Righi A, Faustini-Fustini M, Morandi L, *et al.*: **The changing faces of corticotroph cell adenomas: the role of prohormone convertase 1/3.** *Endocrine.* 2017; **56**(2): 286–97. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
73. Roncaroli F, Scheithauer BW, Young WF, *et al.*: **Silent corticotroph carcinoma of the adenohypophysis: a report of five cases.** *Am J Surg Pathol.* 2003; **27**(4): 477–86. [PubMed Abstract](#) | [Publisher Full Text](#)
74. **F** He L, Forbes JA, Carr K, *et al.*: **Response of silent corticotroph pituitary carcinoma to chemotherapy: case report.** *J Neurosurg Sci.* 2016; **60**(2): 272–80. [PubMed Abstract](#) | [F1000 Recommendation](#)
75. Cohen AB, Lessell S: **Angiogenesis and pituitary tumors.** *Semin Ophthalmol.* 2009; **24**(3): 185–9. [PubMed Abstract](#) | [Publisher Full Text](#)
76. Lloyd RV, Vidal S, Horvath E, *et al.*: **Angiogenesis in normal and neoplastic pituitary tissues.** *Microsc Res Tech.* 2003; **60**(2): 244–50. [PubMed Abstract](#) | [Publisher Full Text](#)
77. Schito L, Semenza GL: **Hypoxia-Inducible Factors: Master Regulators of Cancer Progression.** *Trends Cancer.* 2016; **2**(12): 758–70. [PubMed Abstract](#) | [Publisher Full Text](#)
78. Koyasu S, Kobayashi M, Goto Y, *et al.*: **Regulatory mechanisms of hypoxia-inducible factor 1 activity: Two decades of knowledge.** *Cancer Sci.* 2018; **109**(3): 560–71. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. Troilo A, Alexander I, Muehl S, *et al.*: **HIF1 α deubiquitination by USP8 is essential for ciliogenesis in normoxia.** *EMBO Rep.* 2014; **15**(1): 77–85. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. Carbia-Nagashima A, Gerez J, Perez-Castro C, *et al.*: **RSUME, a small RWD-containing protein, enhances SUMO conjugation and stabilizes HIF-1 α during hypoxia.** *Cell.* 2007; **131**(2): 309–23. [PubMed Abstract](#) | [Publisher Full Text](#)
81. Shan B, Gerez J, Haedo M, *et al.*: **RSUME is implicated in HIF-1-induced VEGF-A production in pituitary tumour cells.** *Endocr Relat Cancer.* 2012; **19**(1): 13–27. [PubMed Abstract](#) | [Publisher Full Text](#)
82. **F** Liu HY, Gu WJ, Wang CZ, *et al.*: **Matrix metalloproteinase-9 and -2 and tissue inhibitor of matrix metalloproteinase-2 in invasive pituitary adenomas: A systematic review and meta-analysis of case-control trials.** *Medicine (Baltimore).* 2016; **95**(23): e3904. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
83. **F** Liu X, Feng M, Zhang Y, *et al.*: **Expression of Matrix Metalloproteinase-9, Pituitary Tumor Transforming Gene, High Mobility Group A 2, and Ki-67 in Adrenocorticotrophic Hormone-Secreting Pituitary Tumors and Their Association with Tumor Recurrence.** *World Neurosurg.* 2018; **113**: e213–e221. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
84. Mete O, Ezzat S, Asa SL: **Biomarkers of aggressive pituitary adenomas.** *J Mol Endocrinol.* 2012; **49**(2): R69–78. [PubMed Abstract](#) | [Publisher Full Text](#)
85. Hui P, Xu X, Xu L, *et al.*: **Expression of MMP14 in invasive pituitary adenomas: relationship to invasion and angiogenesis.** *Int J Clin Exp Pathol.* 2015; **8**(4): 3556–67. [PubMed Abstract](#) | [Free Full Text](#)
86. **F** He W, Huang L, Shen X, *et al.*: **Relationship between RSUME and HIF-1 α VEGF-A with invasion of pituitary adenoma.** *Gene.* 2017; **603**: 54–60. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
87. Seltzer J, Ashton CE, Scotton TC, *et al.*: **Gene and protein expression in pituitary corticotroph adenomas: a systematic review of the literature.** *Neurosurg Focus.* 2015; **38**(2): E17. [PubMed Abstract](#) | [Publisher Full Text](#)
88. **F** Lan X, Gao H, Wang F, *et al.*: **Whole-exome sequencing identifies variants in invasive pituitary adenomas.** *Oncol Lett.* 2016; **12**(4): 2319–28. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
89. **F** de Araújo LJ, Lerario AM, de Castro M, *et al.*: **Transcriptome Analysis Showed a Differential Signature between Invasive and Non-invasive Corticotrophinomas.** *Front Endocrinol (Lausanne).* 2017; **8**: 55. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
90. **F** Liu B, Pang B, Wang Q, *et al.*: **EZH2 upregulation correlates with tumor invasiveness, proliferation, and angiogenesis in human pituitary adenomas.** *Hum Pathol.* 2017; **66**: 101–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
91. Renjie W, Haiqian L: **MIR-132, miR-15a and miR-16 synergistically inhibit pituitary tumor cell proliferation, invasion and migration by targeting Sox5.**

Cancer Lett. 2015; **356**(2 Pt B): 568–78.

[PubMed Abstract](#) | [Publisher Full Text](#)

92. **F** Zheng Z, Zhang Y, Zhang Z, *et al.*: **Effect of miR-106b on Invasiveness of Pituitary Adenoma via PTEN-PI3K/AKT.** *Med Sci Monit.* 2017; **23**: 1277–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
93. **F** Trouillas J, Burman P, McCormack A, *et al.*: **Aggressive pituitary tumours and carcinomas: Two sides of the same coin?** *Eur J Endocrinol.* 2018; **178**(6): C7–C9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
94. **F** Lasolle H, Cortet C, Castinetti F, *et al.*: **Temozolomide treatment can improve overall survival in aggressive pituitary tumors and pituitary carcinomas.** *Eur J Endocrinol.* 2017; **176**(6): 769–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
95. **F** Yang Z, Zhang T, Gao H: **Genetic aspects of pituitary carcinoma: A systematic review.** *Medicine (Baltimore).* 2016; **95**(47): e5268.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
96. Stilling G, Sun Z, Zhang S, *et al.*: **MicroRNA expression in ACTH-producing pituitary tumors: Up-regulation of microRNA-122 and -493 in pituitary carcinomas.** *Endocrine.* 2010; **38**(1): 67–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
97. **F** Dai C, Sun B, Liu X, *et al.*: **O-6-Methylguanine-DNA methyltransferase expression is associated with pituitary adenoma tumor recurrence: a systematic meta-analysis.** *Oncotarget.* 2017; **8**(12): 19674–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
98. Takeshita A, Inoshita N, Taguchi M, *et al.*: **High incidence of low O⁶-methylguanine DNA methyltransferase expression in invasive macroadenomas of Cushing's disease.** *Eur J Endocrinol.* 2009; **161**(4): 553–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
99. Salehi F, Scheithauer BW, Moyes VJ, *et al.*: **Low immunohistochemical expression of MGMT in ACTH secreting pituitary tumors of patients with Nelson syndrome.** *Endocr Pathol.* 2010; **21**(4): 227–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
100. Salehi F, Scheithauer BW, Kovacs K, *et al.*: **O-6-methylguanine-DNA methyltransferase (MGMT) immunohistochemical expression in pituitary corticotroph adenomas.** *Neurosurgery.* 2012; **70**(2): 491–6; discussion 496.
[PubMed Abstract](#) | [Publisher Full Text](#)
101. **F** Raverot G, Burman P, McCormack A, *et al.*: **European Society of Endocrinology Clinical Practice Guidelines for the management of aggressive pituitary tumours and carcinomas.** *Eur J Endocrinol.* 2018; **178**(1): G1–G24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:  

Editorial Note on the Review Process

F1000 Faculty Reviews are written by members of the prestigious F1000 Faculty. They are commissioned and are peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

- Xavier Bertagna**
Faculte de Medicine Paris-Descartes, Universite Paris, Paris, France
Competing Interests: No competing interests were disclosed.
- Francesca Pecori Giraldi**
University of Milan, Istituto Auxologico Italiano, Milan, Italy
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research