



A bioinformatics approach to identifying *Wolbachia* infections in arthropods

Jane Pascar^{1,2} and Christopher H. Chandler¹

¹Department of Biological Sciences, State University of New York at Oswego, Oswego, NY, United States of America

²Department of Biology, Syracuse University, Syracuse, NY, United States of America

ABSTRACT

Wolbachia is the most widespread endosymbiont, infecting >20% of arthropod species, and capable of drastically manipulating the host's reproductive mechanisms. Conventionally, diagnosis has relied on PCR amplification; however, PCR is not always a reliable diagnostic technique due to primer specificity, strain diversity, degree of infection and/or tissue sampled. Here, we look for evidence of *Wolbachia* infection across a wide array of arthropod species using a bioinformatic approach to detect the *Wolbachia* genes *ftsZ*, *wsp*, and the *groE* operon in next-generation sequencing samples available through the NCBI Sequence Read Archive. For samples showing signs of infection, we attempted to assemble entire *Wolbachia* genomes, and in order to better understand the relationships between hosts and symbionts, phylogenies were constructed using the assembled gene sequences. Out of the 34 species with positively identified infections, eight species of arthropod had not previously been recorded to harbor *Wolbachia* infection. All putative infections cluster with known representative strains belonging to supergroup A or B, which are known to only infect arthropods. This study presents an efficient bioinformatic approach for post-sequencing diagnosis and analysis of *Wolbachia* infection in arthropods.

Subjects Bioinformatics, Entomology, Evolutionary Studies, Genomics, Microbiology

Keywords *Wolbachia*, Insects, Bioinformatics, NCBI SRA, Anopheles

Submitted 14 September 2017

Accepted 30 July 2018

Published 3 September 2018

Corresponding author

Jane Pascar, jpascar@oswego.edu

Academic editor

Joseph Gillespie

Additional Information and
Declarations can be found on
page 23

DOI 10.7717/peerj.5486

© Copyright

2018 Pascar and Chandler

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

INTRODUCTION

Symbiotic relationships are ubiquitous in nature and can vary between parasitic, commensal, and mutualistic. *Wolbachia* is a diverse and widespread α -proteobacterium and obligatory endosymbiont ([Werren, 1997a](#); [Werren, 1997b](#); [Saridaki & Bourtzis, 2010](#)). *Wolbachia* was first described in *Culex pipiens* ([Hertig, 1936](#)) and has since been identified in various clades of arthropods including Chelicerata ([Werren & Windsor, 2000](#)), Myriapoda ([Mock et al., 2016](#)), Crustacea ([Bouchon, Rigaud & Juchault, 1998](#); [Cordaux, Michel-Salvat & Bouchon, 2001](#); [Cordaux et al., 2012](#)), and Hexapoda ([Werren & Windsor, 2000](#); [Clark et al., 2001](#); [Augustinos et al., 2011](#); [Bing et al., 2014](#)). Conservative estimates suggest that the frequency of *Wolbachia* infection in arthropods is at least 20% ([Werren, 1997b](#); [Werren & Windsor, 2000](#)), while one study suggests a prevalence as high as 76% of arthropod species ([Jeyaprakash & Hoy, 2000](#)). Meta-analysis indicates that the infection distribution in the total number of species may be closer to 66% ([Hilgenboecker et al., 2008](#)).

Wolbachia is normally transmitted vertically, from mother to offspring, and can manipulate the host's reproduction through five mechanisms: cytoplasmic incompatibility, parthenogenesis, male killing, feminization (Cordaux, Bouchon & Grève, 2011) and meiotic drive (Kageyama et al., 2017). However, there is evidence that *Wolbachia* can be horizontally transmitted (Vavre et al., 1999; Cordaux, Michel-Salzat & Bouchon, 2001; Raychoudhury et al., 2009; Kraaijeveld et al., 2011). Recently, discrete reciprocal benefits provided by infection have been observed including a positive impact on host immunity (Teixeira, Ferreira & Ashburner, 2008; Osborne et al., 2009), immunocompetence (Braquart-Varnier et al., 2008), fecundity (Weeks et al., 2007), and metabolic activity (Darby et al., 2012).

Currently, all *Wolbachia* strains are classified as a single species, with further classification into at least sixteen supergroups, A–Q (Lo et al., 2007; Lindsey et al., 2016a). The four most well studied clades are supergroups A–D. Supergroups A and B are monophyletic and are the most common supergroups known to infect arthropods, while supergroups C and D infect filarial nematodes (Gerth et al., 2014). Supergroup G was discovered to be a recombinant between supergroups A and B; thus it is no longer considered a distinct lineage (Baldo & Werren, 2007). Supergroups E–Q infect a variety of hosts including nematodes, springtails, termites, fleas, aphids, and mites (Lo et al., 2002; Casiraghi et al., 2005; Ros et al., 2009; Haegeman et al., 2009; Augustinos et al., 2011; Bing et al., 2014; Glowska et al., 2015).

Wolbachia has a relatively small genome at about 0.9–1.5 Mbp. Historically, *Wolbachia* infection was diagnosed using 16S rRNA sequences; however, strains range in divergence from 0.2% to 2.6%, and when used independently, 16S provides limited information for inferring phylogenetic relationships (O'Neill et al., 1992). *wsp*, *ftsZ* and the *groE* operon are all protein-encoding genes used for the detection and phylogenetic analysis of *Wolbachia* (Van Borm et al., 2003). The *ftsZ* gene is involved in cell division and is highly conserved in uncultivable bacteria species (Holden, Brookfield & Jones, 1993), but regions that are relatively higher in divergence make it a candidate for better phylogenetic resolution allowing the distinction between supergroups A and B to become apparent (Werren, 1997a). The *wsp* gene, which codes for the surface protein WSP in *Wolbachia*, shows an even higher variability and faster evolutionary rate than 16S or *ftsZ* and can be used in identifying groups and strains of *Wolbachia* (Zhou, Rousset & O'Neil, 1998; Braig et al., 1998), but also displays recombination, which can be misleading when used in phylogenetic analyses (Baldo & Werren, 2007). The *groE*- homologous operon has been noted as another candidate for resolving strain taxonomy (Masui, Sasaki & Ishikawa, 1997). Only a single copy of the operon exists in the genome and it includes the genes that encode the heat shock proteins GroES and GroEL, which are separated by a non-coding intergenic region that is thought to be faster evolving than either of the coding regions (Masui, Sasaki & Ishikawa, 1997).

With the use of antibiotics, *Wolbachia* infections in some species have been cured and the phenotypic changes that are induced by infection are consequently reversed (Stouthamer, Luck & Hamilton, 1990; Bourtzis et al., 1994; Giordano, Jackson & Robertson, 1997). More recently, *Wolbachia* has been proposed as a natural solution to controlling the spread of vector-borne diseases like malaria, yellow fever, and dengue (Hoffmann et al., 2011; Walker & Moreira, 2011; Baldini et al., 2014). Arthropods are present in nearly every habitat on

Earth and they play important ecological roles in a variety of niches. With an estimated 2.4–10.2 million species of arthropods ([Ødegaard, 2000](#)) it is important to quantify the prevalence and distribution of *Wolbachia* infection.

Wolbachia infections are typically diagnosed via polymerase chain reaction (PCR), using *Wolbachia*- specific primers. However, PCR-based tests may produce false positives or false negatives, depending on the strain of *Wolbachia* and the presence of other related bacterial symbionts ([Simões et al., 2011](#)). A metagenomics-based approach can also be useful for characterizing microbiomes, including looking for *Wolbachia* and other symbionts (e.g. [Dittmer & Bouchon, 2018](#)), and can even provide whole-genome sequence information for the symbiont (e.g., [Salzberg et al., 2005](#); [Richardson et al., 2012](#); [Saha et al., 2012](#); [Campana, Robles García & Tuross, 2015](#); [Derks et al., 2015](#); [Wang & Chandler, 2016](#); [Lindsey et al., 2016b](#); [Gerth & Hurst, 2017](#)). While performing a high-throughput sequencing-based screen for *Wolbachia* involving hundreds of different species would require a huge sampling effort and could be cost-prohibitive, screening existing sequence datasets generated for other projects offers a powerful opportunity to diagnose novel infections and better characterize variation in symbionts.

Here, using publicly accessible next-generation sequencing data available in the NCBI Sequence Read Archive (SRA), we looked for evidence of *Wolbachia* infection in a diverse assemblage of arthropod species. We present methods for bioinformatically identifying *Wolbachia* infections in genomic samples. We then used these sequence data to assemble a draft genome sequence for each *Wolbachia* isolate and reconstruct the phylogenetic relationships among the identified *Wolbachia* strains. Using this approach, we uncover novel *Wolbachia* infections, as well as find possible evidence for horizontal transfer of *Wolbachia* between hosts and parasites. These results illustrate how existing genetic databases can provide a wealth of information on symbiotic microbes as a byproduct of host sequencing.

MATERIALS & METHODS

Retrieving data

All samples tested are available through the NCBI Sequence Read Archive (SRA) ([Table S1](#)). To identify samples for testing, all accession numbers that matched the criteria of Arthropoda genomic DNA were sent to the NCBI Run Selector (as of January 2017). In the Run Selector samples were selected based on the criteria that they were run on an Illumina platform, have a genomic library source, a random library selection and the library layout is paired. Transcriptome samples were excluded because of the possibility that some RNA preparation methods may select against bacterial RNA (e.g., poly-A enrichment [Westermann, Gorski & Vogel, 2012](#)) thus increasing the likelihood of false negatives and because assembling *Wolbachia* genomes would be impossible with these data. Similarly, targeted sequencing (e.g., RAD-seq) samples were excluded due to the possibility that the sequences used for detecting *Wolbachia* infections might be excluded during the library preparation process. Only paired-end datasets were considered in order to facilitate whole-genome assembly for positive samples, but there were relatively few species (only

22) in the database with single-end datasets that otherwise met our criteria. Every species that had a sample that met our criteria was chosen for sampling. Some species were over-represented in the number of runs that are available in the SRA; depending on the number of samples available in the SRA, an appropriate amount to include in our dataset was determined on a case-by-case basis (Table S2). Fastq-dump v. 2.8.0 from the SRA Toolkit (NCBI SRA) was used to download, at most, 5×10^7 reads from each accession.

Diagnosing *Wolbachia* Infection

Magic-Blast v1.1.0 (NCBI) was used to compare the SRA reads to selected reference *wsp*, *ftsZ*, and *groE* operon sequences isolated from *Wolbachia* samples that are representative of supergroups A–D (Table 1). A custom R script identified SRA samples where there were matches at least 98 bp in length, $\geq 95\%$ identity to one or more of the reference genes, and with three or more matching sequence reads. All samples that met these criteria were called *Wolbachia* positive samples.

To look for previous reports of *Wolbachia* infection in the species that tested positive, first Google Scholar was used. [species] + *Wolbachia* was used for the search terms. If no published results were found, next we used NCBI GenBank with the same search parameters to look for deposited sequences that may be unpublished that would indicate that *Wolbachia* had been found in the host species previously.

Assembling the *Wolbachia* gene and genome sequences

From all the samples that tested positive (Table S3) if there were more than 3 samples from one species a maximum of 3 samples were chosen for downstream analysis (Table S4).

Velvet v1.2.10 (Zerbino & Birney, 2008) was used to separately assemble the *wsp*, *ftsZ*, and *groE* sequences for each biological sample using the sequence reads that aligned to each gene in the previous step. It was run for kmer values of 21, 31, 41, and 51, using the automatic coverage cutoff flag. To select the optimal assembly of each gene for each sample, we performed BLASTn v2.28 (Altschul et al., 1990), which searched against a database made of each respective reference gene (Table 1). BEDTools v2.25.0 (Quinlan & Hall, 2010) and a custom script was used to parse the BLASTn results for the single longest contig matching each gene from each sample.

To assemble draft genomes for each *Wolbachia* isolate we identified, an iterative bait-and-assemble approach was used. Independent SRA experiments or runs from the same BioSample were first combined into a single dataset. For each sample, the mirabait tool from MIRA v4.0.2 (Chevreux, Wetter & Suhai, 1999) was then used to extract all reads from the full dataset that shared at least one kmer with at least one of seven reference *Wolbachia* genomes representing *Wolbachia* isolates from insects and nematodes (wPip, GCF_000073005.1; wMel, GCF_000008025.1; wNo, GCF_000376585.1; wRi, GCA_000022285.1; wVol, GCF_000530755.1; wCle, GCF_000829315.1; wTpre, GCF_001439985.1), using $k = 31$. These reads, and their corresponding paired-end partners, were assembled using SPAdes 3.11.1 (Bankevich et al., 2012). All resulting contigs were then aligned to the reference *Wolbachia* genomes using dc-megablast 2.7.0+ (Camacho et al., 2009), and any contig that matched any of the reference genomes with an e-value of

Table 1 Reference Wolbachia genes. Gene sequences from *Wolbachia*-infected hosts used to create the reference database for Magic-BLAST searches of SRA accessions to diagnose novel *Wolbachia* infections.

Host Classification								
Phylum	Class	Order	Species	Supergroup	Strain	Accession number	Gene	Citation
Arachnida	Trombidiformes		<i>Bryobia praetiosa</i>	B	–	JN572870.1	wsp	Ros et al. (2012)
			<i>Diaeа circumlita</i>	A	wDiacir3	AY486091.1	wsp	Rowley, Raven & McGraw (2004)
	Coleoptera		<i>Tribolium confusum</i>	–	NFR114	AB469356.1	wsp	D Kageyama, S Narita, T Immura and A Miyanoshita (2008, unpublished data)
			<i>Tribolium confusum</i>	–	–	DQ842337.1	ftsZ	Baldo et al. (2006)
			<i>Dicladispa armigera</i>	A	wDic	DQ243935.1	groE	Wiwatana ratanabutr et al. (2009)
			<i>Culex pipiens</i>	B	–	DQ900650.1	wsp	ND Djadid, N Daneshinia, S Gholizadeh and S Zakeri (2006, unpublished data)
			<i>Culex quinquefasciatus</i>	B	–	AY462861.1	wsp	Tsai et al. (2004)
Diptera			<i>Drosophila melanogaster</i>	A	wMel	FJ403332.1	wsp	YF Wang and Y Zheng (2008, unpublished data)
			<i>Drosophila simulans</i>	A	wMel	DQ412101.1	wsp	Mateos et al. (2006)
			<i>Protocalliphora sialia</i>	B	wProtPA	AF448376.1	wsp	Werren & Bartos (2001)
			<i>Culex quinquefasciatus</i>	B	22	GU901159.1	ftsZ	JC Rondan-Duenas, A Blanco, and CN Gardenal (2010, unpublished data)
			<i>Drosophila melanogaster</i>	A	Canton-S	X71906.1	ftsZ	Holden, Brookfield & Jones (1993)
			<i>Drosophila recens</i>	A	–	U28174.1	ftsZ	Werren, Zhang & Guo (1995)
			<i>Drosophila simulans</i>	A	wHa	AY508998.1	ftsZ	JWO Ballard (2003, unpublished data)
			<i>Drosophila simulans</i>	A	wMa(Ma)	AY508999.1	ftsZ	JWO Ballard (2003, unpublished data)
			<i>Aedes albopictus</i>	A	wAlbA	DQ243927.1	groE	Wiwatana ratanabutr et al. (2009)
			<i>Culex fusciceps</i>	B	wFusc	AJ511284.1	groE	S Wiwatana ratanabutr and P Kittayapong (2002, unpublished data)
			<i>Drosophila simulans</i>	A	–	AB002287.1	groE	Masui, Sasaki & Ishikawa (1997)
			<i>Drosophila simulans</i>	A	–	AB002288.1	groE	Masui, Sasaki & Ishikawa (1997)
			<i>Drosophila tristis</i>	A	–	AY563553.1	groE	Haine, Pickup & Cook (2005)

(continued on next page)

Table 1 (continued)

Host Classification								
Phylum	Class	Order	Species	Supergroup	Strain	Accession number	Gene	Citation
Arthropoda	Insecta		<i>Encarsia formosa</i>	B	–	KC161951.1	wsp	F Lu and MX Jiang (2012, unpublished data)
			<i>Muscidifurax uniraptor</i>	A	–	DQ380857.1	wsp	Kyei-Poku <i>et al.</i> (2006)
			<i>Nasonia giraulti</i>	–	wNGirVA	AF448381.1	wsp	Werren & Bartos (2001)
			<i>Nasonia vitripennis</i>	–	wNvi-2	KC161919.1	wsp	F Lu and MX Jiang (2012, unpublished data)
	Hymenoptera		<i>Trichogramma cordubensis</i>	B	Sib	AF245164.1	wsp	Pintureau <i>et al.</i> (2000)
			<i>Diplolepis rosae</i>	B	Type I	U83887.1	ftsZ	Schilthuizen & Stouthamer, 1998
			<i>Habrocytus bedeguaris</i>	B	Type I	U83886.1	ftsZ	Schilthuizen & Stouthamer, 1998
			<i>Trichogramma</i> n. spec. (nr. deion)	B	–	U59696.1	ftsZ	Schilthuizen & Stouthamer, 1998
			<i>Asobara tabida</i>	–	–	AJ634749.1	groE	Haine, Pickup & Cook (2005)
			<i>Bombyx mandarina</i>	–	–	KJ659909.1	wsp	Zha <i>et al.</i> (2014)
Lepidoptera	Acraea		<i>Acraea encedon</i>	B	–	AJ130892.1	ftsZ	Hurst & Jiggins (1999)
			<i>Ephestia kuehniella</i>	A	Type II	U62126.1	ftsZ	Schilthuizen, Honda & Stouthamer (1998)
			<i>Acraea pharsalus</i>	B	–	AJ318481.1	groE	Jiggins <i>et al.</i> (2002)
	Ephestia		<i>Ephestia cautella</i>	A	–	AB002289.1	groE	Masui, Sasaki & Ishikawa (1997)
			<i>Ephestia cautella</i>	B	–	AB002290.1	groE	Masui, Sasaki & Ishikawa (1997)
Malacostraca	Orthoptera		<i>Gryllus pennsylvanicus</i>	B	–	U28195.1	ftsZ	Werren, Windsor & Guo (1995)
	Isopoda		<i>Chaetophiloscia elongata</i>	–	–	AM087239.1	groE	Wiwatwanaratana butr <i>et al.</i> (2009)

(continued on next page)

Table 1 (continued)

Host Classification				Supergroup	Strain	Accession number	Gene	Citation
Phylum	Class	Order	Species					
Nematoda	Secementea	Spirurida	<i>Dirofilaria repens</i>	C	–	AJ252176.1	wsp	Bazzocchi et al. (2000)
			<i>Litomosoides sigmodontis</i>	D	–	AJ252177.1	wsp	Bazzocchi et al. (2000)
			<i>Onchocerca gibsoni</i>	C	–	AJ252178.1	wsp	Bazzocchi et al. (2000)
			<i>Brugia malayi</i>	D	–	AJ252061.1	wsp	Bazzocchi et al. (2000)
			<i>Brugia pahangi</i>	D	–	AJ252175.1	wsp	Bazzocchi et al. (2000)
			<i>Dirofilaria immitis</i>	C	–	AJ252062.1	wsp	Bazzocchi et al. (2000)
			<i>Wuchereria bancrofti</i>	D	–	AF285273.1	groE	S Salahuddeen and TB Nutman (2000, unpublished data)

Table 2 *Wolbachia* sequences of known origin for phylogenetic analysis. *Wolbachia* genes used as controls and the species name from which they were isolated. The supergroup of the *Wolbachia* strain is listed and these genes served as a control during the creation of the phylogeny.

Host Classification			Gene & Accession Number		Supergroup	Citation
Phylum	Order	Species	<i>ftsZ</i>	<i>groEL/groES</i>		
Arthropoda	Isopoda	<i>Armidillidium vulgare</i>	DQ778101	DQ778104	B	Verne et al. (2007)
	Hemiptera	<i>Bemisia afer</i>	KF452573	KF452533	B	Bing et al. (2014)
	Hemiptera	<i>Bemisia tabaci</i>	KF452577	KF452536	B	Bing et al. (2014)
	Diptera	<i>Drosophila ambigua</i>	AY563550	AY563552	A	Haine, Pickup & Cook (2005)
	Diptera	<i>Drosophila melanogaster</i>	DQ235339	DQ235379	A	Paraskevopoulos et al. (2006)
	Diptera	<i>Drosophila tristis</i>	AY563551	AY563553	A	Haine, Pickup & Cook (2005)

10^{-10} or better, alignment length of at least 100 bp, and percent identity of at least 70%, was retained. This process was then repeated for a total of five iterations, in each cycle using mirabait to identify reads sharing one or more kmers with the last set of assembled contigs, re-assembling these putatively *Wolbachia*-derived reads, and retaining any of the newly assembled contigs that show similarity to a *Wolbachia* reference genome in BLAST searches. The quality of each final assembly was evaluated using QUAST v4.4 ([Gurevich et al., 2013](#)) and BUSCO v3.0.2b ([Simão et al., 2015](#)) with the Bacteria *odb9* reference gene set. Finally, we mapped all sequencing reads from each associated BioSample (not just those used for the assembly process) to the corresponding assembly using bwa mem v.0.7.17 ([Li, 2013](#)), and then used the sambamba depth command ([Tarasov et al., 2015](#)) to extract coverage information for each assembled contig over 400 bp in length, excluding 150 bp from the ends of the contigs (where coverage tends to drop off because reads extending beyond the contig may fail to map successfully).

Phylogenetic analysis

We first constructed phylogenies using the assembled *ftsZ* and *groE* sequences, as well as from a concatenated dataset of both genes; *wsp* was excluded from phylogenetic analyses because of its high frequency of recombination ([Baldo & Werren, 2007](#)). *Wolbachia* gene sequences representing *ftsZ* and the *groE* operon from other studies where the supergroup classification was determined were used as control samples; in this analysis, we included only reference sequences where both genes had been sequenced from the same biological sample (Table 2). MAFFT v7.310 ([Katoh & Standley, 2013](#)) was used to align the sequences for each respective gene. Samples that lacked sufficient length of matching base pairs (at least 800 bp in total across both genes) were discarded from downstream analysis. GBLOCKS v0.91b ([Castresana, 2000](#)) removed the poorly aligned portions of the sequences from each gene alignment using the default parameters. MEGA v7.0 ([Kumar, Stecher & Tamura, 2016](#)) was used to construct phylogenies using maximum likelihood. The model for which the phylogenies were constructed was chosen according to MEGA's suggestion for best fit based on the lowest Bayesian information criterion (BIC) (Table S5). Node support was assessed by bootstrapping with 1,000 replicates.

We also constructed phylogenies based on whole-genome data from a subset of the assemblies which appeared the most complete based on the BUSCO assessment. For

these phylogenies, we used REALPHY ([Bertels et al., 2014](#)), to align genome sequences and identify loci for inclusion in the phylogenetic analysis, using as a reference the seven *Wolbachia* genomes used in the assembly process and merging the reference alignments with the default parameters. We then performed phylogenetic analysis by maximum likelihood in RAxML v8.2.11 ([Stamatakis, 2014](#)) using the TVM+I+G model as selected by ModelTest-NG v0.1.2 (the successor to jModelTest); ([Darriba et al., 2012](#)) using AIC. The RAxML analysis included 100 independent replicate searches for the best-scoring tree and 200 bootstrap replicates to assess node support.

RESULTS

Diagnosing *Wolbachia* infections from publicly available sequence data

A total of 2,545 individual 'runs' from the SRA, representing 288 species and subspecies were tested for *Wolbachia* ([Table S1](#)). Of those, 173 runs from 34 unique species tested positive for the selected reference *Wolbachia* genes ([Table S3](#)). That is, 11.8% of species tested positive for *Wolbachia* in at least one sample and only 6.8% of all SRA runs tested positive. All samples that tested positive were from samples that are in the class Insecta and representative of five orders: Coleoptera, Diptera, Hymenoptera, Hemiptera, and Lepidoptera. According to our literature search eight of these species have not previously been confirmed to have *Wolbachia* infections—*Bembidion lapponicum*, *Ceratina calcarata*, *Delias oraia*, *Diachasma alloeum*, *Diploeciton nevermanni*, *Ectophyla simulans*, *Gerris buenoi* and *Isocolus centaureae* ([Table 3](#)).

Assembling *Wolbachia* genomes

In total, we assembled draft genomes for 51 *Wolbachia* isolates ([Table 4](#)), including at least one for each of the 34 unique host species. There were only two cases in which the assembly was substantially smaller than the expected genome size. In one of those (*Biorhiza pallida* 3), infection was confirmed in independent biological samples, and in the other (*Mycopsylla proxima*) the small assembly probably resulted from the small size of the input dataset. The rest of these assemblies appeared nearly complete, with total assembly sizes of at least 1 Mb and high numbers of BUSCO reference genes represented by a single gene in the assembly. All assemblies were missing at least 13 of the BUSCO reference genes.

We also sought to determine whether each sample that tested positive was likely to represent an actual *Wolbachia* infection, or the result of *Wolbachia* sequences horizontally transferred into the host genome. If the sequencing depth of the *Wolbachia*-like contigs in the assembly differs substantially from the sequencing depth of the host genome, then horizontal transfer can be ruled out. However, performing whole-genome assembly with every sequence dataset to estimate the sequencing depth of the host genome was computationally time-consuming, and our attempts to estimate sequencing depth more rapidly by counting k -mers in the raw data were unsuccessful in most cases because of low sequencing depth. Therefore, we obtained estimates of the genome size of host species from other sources, such as draft assemblies available at NCBI ([Table 4](#)), when available; although draft assemblies can differ substantially in size from actual genome sizes, for

Table 3 Species showing evidence of *Wolbachia* infection. List of unique species (class; order) that tested positive for the presence of *Wolbachia* genes.

Phylum	Class	Order	Species	Supergroup (this study)	Supergroup (previous study)	Citation	GenBank Accession numbers
Coleoptera			* <i>Bembidion lapponicum</i>	B	—	—	—
			<i>Callosobruchus chinensis</i>	A (but possible double infection)	A/B	<i>Kondo, Shimada & Fukatsu (1999), Kondo et al. (2002)</i> , D Kageyama, S Narita, T Ima-mura and A Miyanoshita (2008, unpublished data)	AB025965; AB080664, AB080665, and AB081842; AB469358
			<i>Diabrotica virgifera virgifera</i>	A	A	<i>Giordano, Jackson & Robertson (1997); R Giordano, L Clark, R Alvarez-Zagoya and JF Perez-Dominguez (2005, unpublished data)</i>	U83098, AF011270–AF011271; DQ091306, DQ091307, DQ091308, DQ091309
			* <i>Diploeciton nevermanni</i>	B	—	—	—
Diptera			* <i>Ectophyia simulans</i>	B	—	—	—
			<i>Anopheles gambiae</i>	B	A	<i>Baldini et al. (2014)</i>	KJ728739–KJ728755
			<i>Drosophila melanogaster</i>	A	A	<i>Bourtzis et al. (1994), Woolfit et al. (2013)</i>	Z28981 Z28982 Z28983; KI440871–KI440895
			<i>Drosophila simulans</i>	A/B	A/B	<i>Riegler et al. (2004); Ellegaard et al. (2013)</i>	AY227739, AY227742; CP003883, CP003884
			<i>Drosophila triauraria</i>	A	A	<i>Cordaux et al. (2008)</i>	EU714523
			<i>Drosophila yakuba</i>	A	A	<i>Charlat, Ballard & Mercot (2004), Zabalou et al. (2004), Ioannidis et al. (2007), Cordaux et al. (2008)</i>	AY291346, AY291348; AJ620679; DQ498875; EU714519
			<i>Rhagoletis pomonella</i>	A	A	<i>Schuler et al. (2011)</i>	HQ333145, HQ333146, HQ333147, HQ333148, HQ333149, HQ333150, HQ333151, HQ333152, HQ333153, HQ333154, HQ333155, HQ333156, HQ333157, HQ333158, HQ333159
			<i>Rhagoletis zephyria</i>	A	A	<i>Schuler et al. (2011)</i>	—

(continued on next page)

Table 3 (continued)

Phylum	Class	Order	Species	Supergroup (this study)	Supergroup (previous study)	Citation	GenBank Accession numbers
Arthropoda	Insecta		<i>Dactylopius coccus</i>	B	A/B	<i>Ramirez-Puebla et al. (2016)</i>	LSYX00000000, LSYY00000000
			<i>Diaphorina citri</i>	B	B	<i>Subandiyah et al. (2000), Lindsey et al. (2016b)</i>	AB038366–AB038370
			* <i>Gerris buenoi</i>	B	–	–	–
			<i>Homalodisca vitripennis</i>	B	B	<i>Rogers & Backus (2014)</i>	KF636751
			<i>Maconellicoccus hirsutus</i>	B	B	<i>Husnik & McCutcheon (2016)</i>	PRJEB12066 (European Nucleotide Archive)
	Hemiptera		<i>Megacopta cribraria</i>	A	A	<i>Kikuchi & Fukatsu (2003), TM Jenkins, TD Eaton and C Krauss (2011, unpublished data)</i>	AB109601, AB109602; JQ266093
			<i>Mycopsylla fici</i>	–	–	<i>C Fromont, M Riegler and JM Cook (2015, unpublished data)</i>	KT273254, KT273255, KT273261, KT273277
			<i>Mycopsylla proxima</i>	–	–	<i>C Fromont, M Riegler and JM Cook (2015, unpublished data)</i>	KT273257, KT273259, KT273260, KT273278
	Hymenoptera		<i>Acromyrmex echinatior</i>	A	A	<i>Frost et al. (2010)</i>	HM211007–HM211071
			<i>Biorhiza pallida</i>	A	–	<i>Rokas et al. (2001)</i>	AF339629
			* <i>Ceratina calcarata</i>	–	–	–	–
			<i>Cynipini</i> sp.	A	A	<i>Abe & Miura (2002)</i>	AB052667
			* <i>Diachasma alloeum</i>	A	–	–	–
			<i>Diplolepis spinosa</i>	B	A	<i>Plantard et al. (1999)</i>	AF034987
			* <i>Isocolus centaureae</i>	B	–	–	–
			<i>Pediaspis aceris</i>	A	A	<i>Rokas et al. (2002)</i>	–
			<i>Pseudomyrmex</i> sp. PSW-54	A	–	<i>Kautz, Rubin & Moreau (2013)</i>	KF015789
			<i>Trichogramma pretiosum</i>	B	B	<i>Lindsey et al. (2016a); Lindsey et al. (2016b)</i>	LKEQ00000000
Lepidoptera			* <i>Delias oraia</i>	B	–	–	–
			<i>Operophtera brumata</i>	B	B	<i>Derks et al. (2015)</i>	JYPC00000000
			<i>Pararge aegeria</i>	B	–	<i>Russell et al. (2012)</i>	KC137224
			<i>Polygonia c-album</i>	B	B	<i>Kodandaramaiah et al. (2011)</i>	JN093149, JN093150, JN093151, JN093152, JN093153

Notes.

*Species indicated with a * are species that have not previously been identified, according to our literature search, to harbor *Wolbachia* strains. The supergroup classification of the *Wolbachia* strain according to this study and previously studies is listed if known.

Table 4 *Wolbachia* genome assemblies Information on *Wolbachia* draft genome assemblies. Expected host coverage is calculated as (total sequence data/host genome size). “Evidence of multiple infections” indicates whether or not the assembly contains signs pointing to multiple, distinct *Wolbachia* strains within the same biological host sample used for generating the sequence data (though some of these consisted of pooled individuals). BUSCO comp., BUSCO dup., BUSCO frag., and BUSCO missing refer to the number of BUSCO orthologs that were found to be complete and single copy, duplicated, fragmented, and missing from the *Wolbachia* assembly, out of 148 BUSCOs present in the Bacteria *odb9* reference gene set. Grey rows at the bottom of the table were omitted from the whole-genome phylogenetic analysis because the assemblies appeared less complete (as indicated by missing BUSCO genes) or showed evidence of being chimeric or a mixture of two independent strains.

Host species/ ID	Description/ common name	BioSample accession number	SRA accession numbers	Total seq. data (Gb)	Host genome size (ref.)	Expected host coverage (x)	<i>Wolbachia</i> median cov. (x)	<i>Wolbachia</i> assembly size (Mbp)	<i>Wolbachia</i> assembly N50 (kb)	BUSCO comp.	BUSCO dup.	BUSCO frag.	BUSCO missing	Evidence of multiple infections?	Sample notes
<i>Anopheles gambiae</i>	Mosquito	SAMEA3911293	ERR1554906 ERR1554870 ERR1554834	9.1	280 Mb <i>Holt (2002)</i>	32	9.0	1,212	23.12	125	0	2	21	No	
<i>Biorhiza pallida</i> 1	Wasp	SAMEA2053316	ERR233308	8.7			16	1,249	9.5	125	0	5	18	No	
<i>Biorhiza pallida</i> 2	Wasp	SAMEA2053315	ERR233309	8.3			16	1,246	10.29	128	0	3	17	No	
<i>Delias oraea</i>	Butterfly	SAMN05712507	SRR4341246	13.2			26	1,207	13.35	124	0	4	20	No	Wild caught whole insect
<i>Diabrotica virgifera</i> 1	Western corn rootworm	SAMN02373824	SRR1106898 SRR1106897 SRR1106544	98.9	2.4 Gb (GCA_003013835.1)	41	900	1,505	31	128	0	3	17	No	5 animals
<i>Diabrotica virgifera</i> 2	Western corn rootworm	SAMN02373827	SRR1106912 SRR1106546	95.1	2.4 Gb (GCA_003013835.1)	40	750	1,487	35.01	128	0	3	17	No	5 animals
<i>Diabrotica virgifera</i> 3	Western corn rootworm	SAMN02373842	SRR1107707 SRR1107708 SRR1107710 SRR1107712	88.9	2.4 Gb (GCA_003013835.1)	37	660	1,376	32.16	128	0	3	17	No	5 animals
<i>Diachasma alloeum</i>	Wasp	SAMN03701895	SRR2042503 SRR2046752	56.8	390 Mb (GCA_001412515.1)	150	830	1,377	21.39	127	1	2	18	No	Adults collected from <i>Rhagoletis</i> pupae, so there may be some contamination with <i>Rhagoletis</i> DNA
<i>Diaphorina citri</i> 1	Asian citrus psyllid	SAMN00100712	SRR189238 SRR183690	25	490 Mb (GCA_000475195.1)	51	180	1,379	25.7	128	0	2	18	No	

(continued on next page)

Table 4 (continued)

Host species/ID	Description/ common name	BioSample accession number	SRA accession numbers	Total seq. data (Gb)	Host genome size (ref.)	Expected host coverage (x)	Wolbachia median cov. (x)	Wolbachia assembly size (Mbp)	Wolbachia assembly N50 (kb)	BUSCO comp.	BUSCO dup.	BUSCO frag.	BUSCO missing	Evidence of multiple infections?	Sample notes
<i>Diaphorina citri</i> 2	Asian citrus psyllid	SAMN01886038	SRR649417 SRR649429 SRR649431 SRR649432 SRR649434	27.1	490 Mb (GCA_000475195.1)	55	250	1,425	25.7	128	0	2	18	No	
<i>Diploeciton nevermanni</i>	Beetle	SAMN05860871	SRR4342174	24.8		63	1,698	10.59	124	0	5	19	No		
<i>Dipolepis spinosa</i> 1	Gall wasp	SAMEA3930570	ERR1359308	6.8		57	1,398	12.93	127	0	3	18	No		
<i>Dipolepis spinosa</i> 2	Gall wasp	SAMEA3930574	ERR1359312	7.1		53	1,382	12.13	121	0	3	24	No		
<i>Drosophila melanogaster</i> 1	Fruit fly	SAMEA3634594	ERR1092813 ERR1092814 ERR1092815 ERR1092816 ERR1092817 ERR1092818	18.1	~175 Mb	100	1600	1,208	19.19	127	0	3	18	No	FM7a-23229-hemizygous
<i>Drosophila melanogaster</i> 2	Fruit fly	SAMN04017483	SRR2347338	3.5	~175 Mb	20	21	1,198	13.83	125	0	4	19	No	Haploid embryos; natural population
<i>Drosophila simulans</i> 1	Fruit fly	SAMEA4395362	ERR1597896	25.3	~150 Mb	170	390	1,265	14.67	130	0	2	16	No	
<i>Drosophila simulans</i> 2	Fruit fly	SAMEA4394322	ERR1597899	23.4	~150 Mb	160	1100	1,294	15.57	130	0	2	16	No	
<i>Drosophila simulans</i> 3	Fruit fly	SAMEA4394323	ERR1597900	37	~150 Mb	250	1600	1,313	15.72	130	0	2	16	No	
<i>Drosophila triauraria</i> 1	Fruit fly	SAMD00051863	DRR061000	22.6	~170 for other <i>Drosophila</i>	130	150	1,306	22.94	129	0	3	16	No	Female
<i>Drosophila yakuba</i> 1	Fruit fly	SAMN04044077	SRR2318687	8	~170 Mb	47	42	1,254	14.75	128	0	2	18	No	Pooled isofemale lines (11 lines, 4 females per line)

(continued on next page)

Table 4 (continued)

Host species/ ID	Description/ common name	BioSample accession number	SRA accession numbers	Total seq. data (Gb)	Host genome size (ref.)	Expected host coverage (x)	Wolbachia median cov. (x)	Wolbachia assembly size (Mbp)	Wolbachia assembly N50 (kb)	BUSCO comp.	BUSCO dup.	BUSCO frag.	BUSCO missing	Evidence of multiple infections?	Sample notes
<i>Drosophila yakuba</i> 2	Fruit fly	SAMN04044078	SRR2318706	4.4	~170 Mb	26	35	1.26	10.24	127	0	2	19	No	Pooled isofemale lines (11 lines, 4 females per line)
<i>Ectophya simulans</i>	Rove beetle	SAMN05833357	SRR4301374	22.7			170	1.437	42.92	127	1	3	17	No	Whole insect
<i>Gerris buenoi</i> 1	Water striders	SAMN02439785	SRR1197265	27.4	990 Mb (GCA_001010745.2)	28	69	1.538	13.14	127	0	3	18	No	Female, whole individuals, adults
<i>Gerris buenoi</i> 2	Water striders	SAMN02439786	SRR1197267	27.8	990 Mb (GCA_001010745.2)	28	39	1.537	13.14	127	0	3	18	No	Male, whole individuals, adults
<i>Homalodisca vitripennis</i> 1	Glassy-winged sharpshooter (leafhopper)	SAMN02209956 SRR941996 SRR941997	SRR941995	107.7	1.45 Gb (GCA_000696855.2)	74	260	1.675	14.19	121	9	3	15	Maybe: BUSCO duplications	Lab reared Florida-strain female
<i>Maconellicoccus hirsutus</i>	Mealybug	SAMEA3699093	ERR1189167	9.8	160 Mb (GCA_900064465.1)	61	80	1.415	27.94	130	0	2	16	No	
<i>Operophtera brumata</i>	Winter moth	SAMN03121611	SRR1618545 SRR1618581 SRR1618582	22.2	640 Mb (GCA_001266575.1)	35	28	1.35	33.34	129	0	1	18	No	Female; adult; head and thorax; wild caught individual
<i>Pararge aegeria</i>	Speckled wood butterfly	SAMN02688782	SRR1190479	9.8			138	1.282	83.56	129	0	2	17	No	Whole adult, lab culture
<i>Pediaspis aceris</i> 1	Gall wasp	SAMEA3925672	ERR1355090	3.8			12	1.188	8.62	119	0	5	24	No	
<i>Pediaspis aceris</i> 2	Gall wasp	SAMEA3925673	ERR1355091	3.8			12	1.174	8.05	114	0	9	25	No	
<i>Polygonia c-album</i>	Comma butterfly	SAMN02688783	SRR1190476	6			61	1.463	22.09	129	0	1	18	No	Whole adult, lab cultured
<i>Pseudomyrmex</i> sp. PSW-54	Ant	SAMN03275520	SRR1742977	36.9	280 Mb (GCA_002006095.1 for congeners)	130	195	1.245	15.91	127	0	3	18	No	Adult female worker ant
<i>Rhagoletis pomonella</i>	Apple maggot fly	SAMN05388941 SRR3901027	SRR3900841	23.3	0.97 (C-value)	25	270	1.314	13.52	127	0	3	18	No	Single adult female fly

(continued on next page)

Table 4 (continued)

Host species/ ID	Description/ common name	BioSample accession number	SRA accession numbers	Total seq. data (Gb)	Host genome size (ref.)	Expected host coverage (x)	Wolbachia median cov. (x)	Wolbachia assembly size (Mbp)	Wolbachia assembly N50 (kb)	BUSCO comp.	BUSCO dup.	BUSCO frag.	BUSCO missing	Evidence of multiple infections?	Sample notes
<i>Trichogramma pretiosum</i>	Wasp	SAMN02439301	SRR1191749 SRR1191750 SRR1191751 SRR1191752 SRR1191753	68.6	190 Mb (GCA_000599845.3)	360	50	1,097	51.37	127	0	3	18	No	
<i>Acromyrmex echinatior</i>	Ant	SAMEA762107	ERR034187 ERR03416	13.2	300 Mb (GCA_000204515.1)	44	56	1,611	4.66	104	0	8	36	No	1 male
<i>Bembidion lapponicum</i>	Beetle	SAMN04276907	SRR2939026	8.5			11	1.151	1.79	70	0	13	65	No	Adult, whole body
<i>Biorhiza pallida</i> 3	Wasp	SAMEA2053314	ERR233313	4.1			4.2	0.645	0.7	3	0	8	137	No	
<i>Callosobruchus chinensis</i>	Bean weevil	SAMN02313283	SRR949786 SRR952345	32.1	0.75 (C-value)	44	340	2.894	4.06	78	6	12	52	Yes: bimodal coverage distribution; assembly size; BUSCO duplications	Male, head, thorax, feet
<i>Ceratina calcarata</i>	Carpenter bee	SAMN04210145	SRR2912519	16	200 Mb (GCA_001652005.1)	80	11	1.053	1.96	55	0	12	81	No	1 haploid male
Cynipini 1	Oak gall wasp	SAMEA1965365	ERR233303 ERR233304 ERR233305	9.4			20	1.216	6.57	96	0	13	39	No	
Cynipini 2	Oak gall wasp	SAMEA2053318	ERR233306	7.3			17	1.182	3.32	82	0	21	45	No	
<i>Dactylopius coccus</i>	Domestic cochineal	SAMN02725055	SRR1231828 SRR1231831 SRR1231832	6.2	21.1 Mb (estimate given in from NCBI BioSample entry)	290	110	2.563	5.95	47	80	2	19	Yes: assembly size; BUSCO duplications; bimodal coverage distribution	Bulk sample of 50 Oaxacan Mexican grana
<i>Drosophila melanogaster</i> 3	Fruit fly	SAMN05417645	SRR3931592	3.4	~175 Mb	19	6.5	1.171	6.14	102	0	15	31	No	Adult male whole body, wild caught from Africa
<i>Homalodisca vittripennis</i> 2	Glassy-winged sharpshooter (leafhopper)	SAMN02209957	SRR941998	39.5	1.45 Gb (GCA_000696855.2)	27	64	1.803	17.02	109	23	3	13	Maybe; BUSCO duplications	Lab-reared Florida-strain male
<i>Isocolus centaureae</i> 1	Gall wasp	SAMEA3930555	ERR1359249	3.4			7.9	0.998	2.19	63	0	16	69	No	

(continued on next page)

Table 4 (continued)

Host species/ ID	Description/ common name	BioSample accession number	SRA accession numbers	Total seq. data (Gb)	Host genome size (ref.)	Expected host coverage (x)	Wolbachia median cov. (x)	Wolbachia assembly size (Mbp)	Wolbachia assembly N50 (kb)	BUSCO comp.	BUSCO dup.	BUSCO frag.	BUSCO missing	Evidence of multiple infections?	Sample notes
<i>Isocolus centaureae</i> 2	Gall wasp	SAMEA3930556	ERR1359250	3.4		7.7	0.965	2.31	63	0	14	71	No		
<i>Megacopta cribraria</i>	Stink bug	SAMN02313994	SRR1145746	5.7		31	2.097	1.74	75	4	13	56	Yes; assembly size; BUSCO duplications; possible bimodal coverage distribution		
<i>Mycopsylla fici</i> 1	Fig psyllid	SAMN04226368	SRR2954433	0.9		9.9	1.171	2.32	76	0	6	66	No		
<i>Mycopsylla fici</i> 2	Fig psyllid	SAMN04226369	SRR2954467	0.8		12	1.238	2.68	77	0	14	57	No		
<i>Mycopsylla proxima</i>	Psyllid	SAMN04226370	SRR2954473	1.1		5.7	0.364	0.76	0	0	3	145	No		
<i>Rhagoletis zephyria</i>	Tephritid fly	SAMN04977950	SRR3670118 SRR3670117 SRR3670120	132.9	1.1 Gb (GCA_001687245.1)	120	1200	1.881	11.19	80	51	2	15	Yes; bimodal coverage distribution; assembly size; BUSCO duplications	Single adult female fly

our purposes this should be a reasonable approximation. We then estimated the expected sequencing depth of the host by dividing the total amount of sequencing data by the estimated host genome size. Although genome size data on some host species was lacking, large differences in sequencing depth between the host and endosymbiont support active *Wolbachia* infections in several species, including *Anopheles gambiae*, *Diabrotica virgifera*, *Diachasma alloeum*, *Diaphorina citri*, several *Drosophila* species, *Homalodisca vitripennis*, *Rhagoletis pomonella* and *R. zephyria*, *Trichogramma pretiosum*, *Callosobruchus chinensis*, *Ceratina calcarata*, and *Dactylopius coccus* (Table 4).

In a few cases, there was evidence of multiple infections in a single sample. This evidence included an unusual number of duplicated BUSCO reference genes in the assembly (e.g., *Homalodisca vitripennis* 1), the presence of multiple peaks in the coverage distribution histogram (e.g., *Callosobruchus chinensis*), assembly sizes much larger than previously sequenced *Wolbachia* genomes (e.g., *Dactylopius coccus*), or some combination of these (Table 4).

***Wolbachia* phylogeny**

All phylogenetic trees based on individual or concatenated datasets using the *ftsZ* and *groE* sequences show two distinct branches representing supergroups A and B (Fig. 1; Figs. S1–S2). The tree resulting from the concatenated dataset has the most robust bootstrap support for most clades. Positive control samples that were included in the phylogeny cluster with other control samples of the same known supergroup. Of the species where *Wolbachia* had been previously unidentified, according to all trees, the strains isolated from *D. alloeum* falls within supergroup A, while the *B. lapponicum*, *I. centaureae*, *G. Buenoi*, *D. nevermanni* and *D. oraia* isolates all fall within supergroup B (Figs. 1 and 2, Figs. S1–S2).

The phylogeny generated from whole-genome sequencing data (Fig. 2) was similar in overall topology to the trees based on *ftsZ* and *groE*, with two clear clades representing supergroups A and B, but with higher bootstrap support for most branches.

DISCUSSION

Observed low infection rates

While *Wolbachia* is estimated to infect between 20–76% of arthropod species (Werren, Windsor & Guo, 1995; Jeyaprakash & Hoy, 2000), in this set of data only 11.8% of species tested positive. Given the source, this low rate of infection can be hypothesized to be the result of five possible scenarios: (1) Underrepresentation in the amount of data available per host species. For example, only 43 out of the 288 (14.9%) species and subspecies tested had ≥ 10 samples available in the SRA that met the criteria of this study (Table S2). When >100 individuals are tested for *Wolbachia*, results are skewed towards finding a positive sample (Hilgenboecker et al., 2008). (2) Bias in the source of the samples. Sources vary between wild-caught individuals, lab stocks, and unreported sources. Since the phenotypic consequences of *Wolbachia* are well established, if uninfected individuals are needed for a study they may be selectively chosen (see Dorđević et al., 2017; Becking et al., 2017), or the researchers may even actively treat infections with antibiotics (Dobson & Rattanadechakul, 2001; Casiraghi et al., 2002; Koukou et al., 2006) or increased rearing temperature. In those




Figure 1 Concatenated phylogeny. Molecular phylogenetic analysis by maximum likelihood based on the concatenated dataset containing *ftsZ* and the *groE* operon (total of 1,381 nucleotide positions). Bold letters next to host species names indicate supergroup relationships of *Wolbachia* isolates identified in previous studies for each host. Asterisks indicate reference sequences (see also Table 2).

Full-size DOI: 10.7717/peerj.5486/fig-1




Figure 2 Whole-genome phylogeny. Maximum likelihood phylogeny based on whole-genome sequence data of *Wolbachia* isolates assembled here and previously sequenced reference *Wolbachia* genomes (indicated by samples with associated accession numbers), with a total of 133,744 nucleotide positions. Numbers by nodes indicate bootstrap support based on 200 replicates. Bold letters next to host species names indicate supergroup relationships of *Wolbachia* isolates identified in previous studies for each host. Isolates with accession numbers listed represent reference genome sequences from other studies.

[Full-size](#) DOI: 10.7717/peerj.5486/fig-2

cases the sequencing data will consequently test negative for *Wolbachia* using the methods employed here. (3) Tissue sampled. In some species infection has only been detectable in the gonads, indicating that infection density in somatic tissue may be variable or low ([Dobson et al., 1999](#)). For many samples in the SRA, specific tissue has not been indicated. (4) Bioinformatic removal of bacterial contaminants. Even if *Wolbachia* is sequenced with the host's DNA, the researcher may have eliminated these reads bioinformatically before depositing the reads as relatively standard practice in sequence processing ([Kunin et al., 2008](#); [Schmieder & Edwards, 2011](#); [Derks et al., 2015](#)). (5) False negatives. It is possible that some infections may have been missed due to the limited set of available reference genes; more divergent strains might not have been detected in these analyses.

Strain supergroup affiliation

For 13 of the species that tested positive, previous information was available about supergroup affiliations of *Wolbachia* strains that have been found to infect them (Table 3). Our results are mostly consistent with previously reported phylogenetic relationships. Previously, *C. chinensis*, *D. coccus*, and *D. simulans* have been found to be infected with A and/or B strains (Kondo, Shimada & Fukatsu, 1999; Kondo et al., 2002; Riegler et al., 2004; Ellegaard et al., 2013; Ramirez-Puebla et al., 2016). Here, evidence of both A and B supergroup strains was found in *D. simulans* (Fig. 1), though the whole-genome phylogeny was somewhat inconsistent here, suggesting possible recombination for some genes. Moreover, while the single-gene phylogenies suggested that the endosymbionts of these *C. chinensis* and *D. coccus* samples were members of supergroups A and B, respectively, the whole-genome assemblies for both of these endosymbionts contained strong evidence of dual infections, so we cannot rule out the presence of both A and B supergroup strains in these samples.

Wolbachia infection has also been documented prior to this study in *B. pallida*, *P. aegeria*, and *P. sp. PSW-54* but the supergroup relationships were not reported (Subandiyah et al., 2000; Rokas et al., 2001; Russell et al., 2012; Kautz, Rubin & Moreau, 2013; Baldini et al., 2014). Our concatenated results suggest that a supergroup A strain infects *B. pallida* and *P. sp. PSW-54*, while a B strain infects *P. aegeria*.

Two species showed a different supergroup strain than what has been previously reported—*D. spinosa* and *A. gambiae*. *D. spinosa* has previously been identified to harbor a supergroup A strain, but here we discovered an infection that clusters within supergroup B. It may be possible for *D. spinosa* to harbor both A and B strains since other species in the genus have been shown to have supergroup B infections (Plantard et al., 1999).

Particularly notable is our identification of a supergroup B strain in *A. gambiae*. Anopheline mosquitoes were once thought to lack infection by *Wolbachia* in nature (Kittayapong et al., 2000; Ricci et al., 2002; Rasgon & Scott, 2004), though they are capable of experimental infection in the lab (Hughes et al., 2011). However, there have been recent reports of natural infections in wild populations (Baldini et al., 2014; Gomes et al., 2017). In particular, a supergroup A strain was found to infect *A. gambiae* mosquitoes in Mali that reduces the transmission of the malaria parasite (Gomes et al., 2017). The strain identified here clearly belongs to supergroup B, and is related to strains infecting Hymenoptera and Lepidoptera, rather than fleas (Siphonaptera) like the previously identified supergroup A strain. In addition, other recent surveys have found evidence of diverse *Wolbachia* strains, including supergroup B strains, within the *A. gambiae* species complex (Ayala et al., 2018; Jeffries et al., 2018). Combined, these results suggest that the diversity of *Wolbachia* infections in *Anopheles* may be currently underappreciated. Importantly, we have good evidence that the sample here is an actual *Wolbachia* infection rather than an integrated piece of *Wolbachia* DNA in the host genome. First, we assembled a nearly complete *Wolbachia* genome from this dataset; more importantly, given that the *A. gambiae* genome is roughly 280 Mb (Holt, 2002), and this dataset contained roughly 9.1 Gb of raw sequence reads, we would expect to have roughly ~32× coverage of the host genome, but the

Wolbachia genome had only $\sim 9 \times$ coverage, suggesting that *Wolbachia* DNA was present at lower densities in this sample than the host DNA.

The assembled *ftsZ* and *groE* sequences from *C. calcarata*, *E. simulans*, *M. fici*, and *M. proxima* assemblies were too short to be included in our individual gene-based phylogenetic reconstruction; the infection density, and thus the sequencing coverage, for these species may have been too low to yield reliable assemblies for these genes. We were able to assign the *E. simulans* infection to supergroup B based on its draft genome sequence, but the supergroup relationships for the others are still unknown. Previously, *Wolbachia* sequence information has been isolated in *M. fici*, and *M. proxima* (Fromont et al., unpublished data; Table 3) but supergroup affiliation was not suggested. According to our literature search this is the first detection of *Wolbachia* in *C. calcarata* and *E. simulans*. The other six previously unidentified species infections were included in the phylogeny. *D. alloicum* clustered with known supergroup A infections while *B. lapponicum*, *D. oraia*, *D. nevermanni*, *G. buenoi*, *I. centaureae* isolate clustered within the supergroup B clade.

Finally, our phylogeny also offers some hints into possible mechanisms of horizontal transmission of *Wolbachia* infections. In particular, the strain identified here infecting *Diachasma alloicum* is closely related to the strain found in *Rhagoletis pomonella* (as well as *D. melanogaster* and *D. yakuba*). This is intriguing because *D. alloicum* is a parasitoid wasp that uses *R. pomonella* and *R. mendax* as its host (Maier, 1981; Stelinski, Pelz & Liburd, 2004), suggesting that this may represent a natural horizontal transfer of *Wolbachia* from one lineage to another; previous studies have found evidence of horizontal transmission between predators and prey or hosts and parasites (Heath et al., 1999; Le Clec'h et al., 2013). However, contamination by host material in parasitoid samples, or vice versa, could also explain this outcome, so this result should be interpreted cautiously until this path of transmission can be experimentally confirmed.

Multiple infections and integration of wolbachia into the host genome

Double (Perrot-Minnot, Guo & Werren, 1996; Narita, Nomura & Kageyama, 2007) and even triple *Wolbachia* infections (Rousset, Braig & O'Neill, 1999; Kondo et al., 2002) have been reported in arthropod populations and individuals, both naturally and through experimental injection. The initial screening methods presented here are not capable of identifying multiple infections because we only looked for a positive or negative test result and then used only the single longest contig for phylogenetic construction. In conventional PCR there is a tradeoff between specificity and sensitivity of primers; additionally no one primer is capable of identifying *Wolbachia* in all samples (Simões et al., 2011). PCR is useful in initial infection confirmation but sequencing is usually necessary to confirm group relationships. Techniques used to identify multiple infections currently include quantitative PCR with highly specific primers (Kondo et al., 2002; Narita, Nomura & Kageyama, 2007), cloning and sequencing (Jannongluk et al., 2002), and Southern hybridization (Perrot-Minnot, Guo & Werren, 1996).

We were able to identify evidence of possible multiple infections through genome assembly. In some cases, the assembly was approximately double the expected size, contained a large number of duplicated genes, or showed evidence of multiple peaks in a

coverage histogram, all of which are signs of infection by multiple, independent strains. Again, these results should be interpreted cautiously pending experimental validation. For instance, some of the multiply infected samples consisted of pooled DNA from multiple individuals (e.g., *Drosophila yakuba* and *Diabrotica vinifera*), so the “multiple” infection might simply result from different individuals in the sample harboring different endosymbiont strains. Nevertheless, these results show that high-throughput sequencing can be a powerful way to detect multiple infections, especially when a priori sequence information for designing strain-specific primers is unavailable.

A related issue is that *Wolbachia* DNA is frequently integrated into host genomes ([Vavre et al., 1999](#); [Leclercq et al., 2016](#)); in some cases, these insertions even consist of nearly whole *Wolbachia* genome sequences ([Dunning Hotopp et al., 2007](#)). This complicates our analyses because some of the identified “infections” could actually be *Wolbachia* DNA integrated into the host genome; in fact, horizontally transferred *Wolbachia* DNA has already been identified in four orders which are all represented by the positive results in this study, Coleoptera, Diptera, Hemiptera, and Hymenoptera ([Hotopp, 2011](#)). We were able to rule out horizontally transferred DNA in some, but not all, cases of positive samples, using sequencing depth information; if the sequencing depth of the assembled *Wolbachia* contigs differs from the sequencing depth of the host’s nuclear DNA, that suggests a true, active infection. True infections could also be validated experimentally when necessary, for example, using fluorescence in situ hybridization ([Hughes et al., 2011](#)). Either way, horizontally transferred *Wolbachia* DNA would still indicate that a species at least had a history of infection at some point in the past.

This work shows that it is often possible to assemble draft genomes of endosymbionts from host DNA, similar to previous studies in which *Wolbachia* genomes were assembled from sequencing host organisms ([Ghedin et al., 2004](#); [Salzberg et al., 2005](#); [Richardson et al., 2012](#); [Saha et al., 2012](#); [Campana, Robles García & Tuross, 2015](#); [Derks et al., 2015](#); [Lindsey et al., 2016b](#)), even when the endosymbiont was not the focus or original reason for performing the sequencing in the first place. Although they may be fragmented, these draft genomes can still provide valuable information about the phylogenetics and evolution of the endosymbiont. While *Wolbachia* is relatively well studied, there are many other endosymbionts that have received less attention, such as some *Spiroplasma*, *Cardinium*, *Arsenophonus*, and *Flavobacterium* species ([Duron et al., 2008](#)), and others await discovery. This study shows that extensive field sampling may not even be necessary to get a better understanding of the diversity of these endosymbionts; the sequencing data are probably already available in public databases. With the right reference databases and metagenomics software, there is a lot of potential to learn more about these endosymbionts just from already existing resources.

CONCLUSIONS AND RECOMMENDATIONS

Wolbachia is a well-known endosymbiont of many arthropod species and while standard *Wolbachia* diagnostic techniques utilize various *Wolbachia* primers to confirm infection via PCR ([Simões et al., 2011](#)) there are trade-offs that limit large scale surveys. Here, we

present a method to identify *Wolbachia* bioinformatically using publicly accessible host raw sequencing data. In eight arthropod species, *Wolbachia* was identified where infection has not previously been reported, and in 27 other arthropod species infection was confirmed. Isolates of *Wolbachia* from positive samples all clustered within either supergroups A or B, and for seven of the newly identified hosts we identified the supergroup of the strain. From these isolates we assembled draft *Wolbachia* genomes, which provided robustly supported phylogenetic information as well as information about potential HGT events or signs of multiple infection.

These results highlight the importance of depositing raw sequencing datasets to public archives like the NCBI SRA and the value that they have in studying endosymbionts. At the same time, we offer some suggestions for best practices when depositing sequence data into public archives to maximize its usefulness for future researchers ([Wilkinson et al., 2016](#); [Griffin et al., 2017](#)). First, we encourage everyone performing high-throughput sequencing to deposit their data into public databases like the NCBI SRA, where it can easily be searched and accessed, as opposed to depositing only in smaller, taxon-specific databases or personal/lab web sites. Second, data should be minimally filtered; while “contaminant” sequences like endosymbiont DNA may be a nuisance to those who generated the data, they may be of interest to others. Finally, all sequence data should be accompanied by as much metadata as possible. Without this information, interpreting results can be difficult. For example, many of the sequences we used in this study lacked detailed information about the source of the DNA in the associated BioSample entries (e.g., whether it came from a lab strain or wild-caught specimens, its geographic origin if field collected, whether it was from a single individual or a pooled sample, whether the specimen was male or female, whether it was a whole body or specific tissues, etc.). Including this information would have helped us better understand possible biases in the dataset, such as how well the results may reflect the frequency of infection in natural populations, or whether a sample might give a false negative result because *Wolbachia* is not present at high densities in the tissues sampled for DNA.

ACKNOWLEDGEMENTS

We thank P Brannock, S Borrelli, and the editor and three anonymous reviewers for their helpful feedback and suggestions on earlier versions this manuscript. We also thank V Buonaccorsi, C Walls, and GCAT-SEEKquence for computing support, as well as the National Center for Genome Analysis Support at Indiana University.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the National Science Foundation (NSF DEB-1453298 to Christopher Chandler, and ABI-1458641 to Indiana University). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

National Science Foundation: NSF DEB-1453298.

Indiana University: ABI-1458641.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Jane Pascar and Christopher H. Chandler conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The raw data and code are provided in the [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5486#supplemental-information>.

REFERENCES

- Abe Y, Miura K. 2002. Does *Wolbachia* induce unisexuality in oak gall wasps? (Hymenoptera: Cynipidae). *Annals of the Entomological Society of America* 95:583–586 DOI 10.1603/0013-8746(2002)095[0583:DWIUIO]2.0.CO;2.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215:403–410 DOI 10.1016/S0022-2836(05)80360-2.
- Augustinos AA, Santos-Garcia D, Dionyssopoulou E, Moreira M, Papapanagiotou A, Scarvelakis M, Doudoumis V, Ramos S, Aguiar AF, Borges PAV, Khadem M, Latorre A, Tsiamis G, Bourtzis K. 2011. Detection and characterization of *Wolbachia* infections in natural populations of Aphids: is the hidden diversity fully unraveled? *PLOS ONE* 6:e28695 DOI 10.1371/journal.pone.0028695.
- Ayala D, Akone-Ella O, Rahola N, Kengne P, Ngangue MF, Mezeme F, Makanga BK, Costantini C, Simard F, Prugnolle F, Roche B, Duron O, Paupy C. 2018. Natural *Wolbachia* infections are common in the major malaria vectors in Central Africa. *bioRxiv*. DOI 10.1101/343715.
- Baldini F, Segata N, Pompon J, Marcenac P, Robert Shaw W, Dabiré RK, Diabaté A, Levashina EA, Catteruccia F. 2014. Evidence of natural *Wolbachia* infections in field populations of *Anopheles gambiae*. *Nature Communications* 5:3985 DOI 10.1038/ncomms4985.

- Baldo L, Hotopp JCD, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, Hayashi C, Maiden MCJ, Tettelin H, Werren JH.** 2006. Multilocus sequence typing system for the endosymbiont Wolbachia pipiensis. *Applied and Environmental Microbiology* 72:7098–7110 DOI 10.1128/AEM.00731-06.
- Baldo L, Werren JH.** 2007. Revisiting Wolbachia supergroup typing based on WSP: spurious lineages and discordance with MLST. *Current Microbiology* 55:81–87 DOI 10.1007/s00284-007-0055-8.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotnik AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA.** 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* 19:455–477 DOI 10.1089/cmb.2012.0021.
- Bazzocchi C, Jamnongluk W, O'Neill SL, Anderson TJC, Genchi C, Bandi C.** 2000. wsp gene sequences from the Wolbachia of filarial nematodes. *Current Microbiology* 41:96–100 DOI 10.1007/s002840010100.
- Becking T, Giraud I, Raimond M, Moumen B, Chandler C, Cordaux R, Gilbert C.** 2017. Diversity and evolution of sex determination systems in terrestrial isopods. *Scientific Reports* 7:1084 DOI 10.1038/s41598-017-01195-4.
- Bertels F, Silander OK, Pachkov M, Rainey PB, Van Nimwegen E.** 2014. Automated reconstruction of whole-genome phylogenies from short-sequence reads. *Molecular Biology and Evolution* 31:1077–1088 DOI 10.1093/molbev/msu088.
- Bing X-L, Xia W-Q, Gui J-D, Yan G-H, Wang X-W, Liu S-S.** 2014. Diversity and evolution of the Wolbachia endosymbionts of Bemisia (Hemiptera: Aleyrodidae) whiteflies. *Ecology and Evolution* 4:2714–2737 DOI 10.1002/ece3.1126.
- Bouchon D, Rigaud T, Juchault P.** 1998. Evidence for widespread Wolbachia infection in isopod crustaceans: molecular identification and host feminization. *Proceedings of the Royal Society B: Biological Sciences* 265:1081–1090 DOI 10.1098/rspb.1998.0402.
- Bourtzis K, Nirgianaki A, Onyango P, Savakis C.** 1994. A prokaryotic dnaA sequence in Drosophila melanogaster Wolbachia infection and cytoplasmic incompatibility among laboratory strains. *Insect Molecular Biology* 3:131–142 DOI 10.1111/j.1365-2583.1994.tb00160.x.
- Braig HR, Zhou W, Dobson SL, O'Neill SL.** 1998. Cloning and characterization of a gene encoding the major surface protein of the bacterial endosymbiont Wolbachia pipiensis. *Journal of Bacteriology* 180:2373–2378.
- Braquart-Varnier C, Lachat M, Herbinière J, Johnson M, Caubet Y, Bouchon D, Sicard M.** 2008. Wolbachia mediate variation of host immunocompetence. *PLOS ONE* 3:e3286 DOI 10.1371/journal.pone.0003286.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.** 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:1 DOI 10.1186/1471-2105-10-421.
- Campana MG, Robles García NM, Tuross N.** 2015. America's red gold: multiple lineages of cultivated cochineal in Mexico. *Ecology and Evolution* 5:607–617 DOI 10.1002/ece3.1398.

- Casiraghi M, Bordenstein SR, Baldo L, Lo N, Beninati T, Wernegreen JJ, Werren JH, Bandi C.** 2005. Phylogeny of Wolbachia pipiensis based on gltA, groEL and ftsZ gene sequences: clustering of arthropod and nematode symbionts in the F supergroup, and evidence for further diversity in the Wolbachia tree. *Microbiology* 151:4015–4022 DOI [10.1099/mic.0.28313-0](https://doi.org/10.1099/mic.0.28313-0).
- Casiraghi M, McCall J, Simoncini L, Kramer L, Sacchi L, Genchi C, Werren J, Bandi C.** 2002. Tetracycline treatment and sex-ratio distortion: a role for Wolbachia in the moulting of filarial nematodes? *International Journal for Parasitology* 32:1457–1468 DOI [10.1016/S0020-7519\(02\)00158-3](https://doi.org/10.1016/S0020-7519(02)00158-3).
- Castresana J.** 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17:540–552 DOI [10.1093/oxfordjournals.molbev.a026334](https://doi.org/10.1093/oxfordjournals.molbev.a026334).
- Charlat S, Ballard JWO, Mercot H.** 2004. What maintains noncytoplasmic incompatibility inducing Wolbachia in their hosts: a case study from a natural *Drosophila yakuba* population. *Journal of Evolutionary Biology* 17(2):322–330.
- Chevreux B, Wetter T, Suhai S.** 1999. Genome sequence assembly using trace signals and additional sequence information. *Computer Science and Biology: Proceedings of the German Conference on Bioinformatics* 99:45–56.
- Clark TL, Meinke LJ, Skoda SR, Foster JE.** 2001. Occurrence of Wolbachia in selected diabroticite (Coleoptera: Chrysomelidae) beetles. *Annals of the Entomological Society of America* 94:877–885 DOI [10.1603/0013-8746\(2001\)094\[0877:OOWISD\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2001)094[0877:OOWISD]2.0.CO;2).
- Cordaux R, Bouchon D, Grève P.** 2011. The impact of endosymbionts on the evolution of host sex-determination mechanisms. *Trends in Genetics* 27:332–341 DOI [10.1016/j.tig.2011.05.002](https://doi.org/10.1016/j.tig.2011.05.002).
- Cordaux R, Michel-Salzat A, Bouchon D.** 2001. Wolbachia infection in crustaceans: novel hosts and potential routes for horizontal transmission. *Journal of Evolutionary Biology* 14:237–243 DOI [10.1046/j.1420-9101.2001.00279.x](https://doi.org/10.1046/j.1420-9101.2001.00279.x).
- Cordaux R, Pichon S, Ben Afia Hatira H, Doublet V, Grève P, Marcadé I, Braquart-Varnier C, Souty-Grosset C, Charfi-Cheikhrouha F, Bouchon D.** 2012. Widespread Wolbachia infection in terrestrial isopods and other crustaceans. *ZooKeys* 176:123–131 DOI [10.3897/zookeys.176.2284](https://doi.org/10.3897/zookeys.176.2284).
- Cordaux R, Pichon S, Ling A, Pérez P, Delaunay C, Vavre F, Bouchon D, Grève P.** 2008. Intense transpositional activity of insertion sequences in an ancient obligate endosymbiont. *Molecular Biology and Evolution* 25:1889–1896 DOI [10.1093/molbev/msn134](https://doi.org/10.1093/molbev/msn134).
- Darby AC, Armstrong SD, Bah GS, Kaur G, Hughes MA, Kay SM, Koldkjaer P, Rainbow L, Radford AD, Blaxter ML, Tanya VN, Trees AJ, Cordaux R, Wastling JM, Makepeace BL.** 2012. Analysis of gene expression from the Wolbachia genome of a filarial nematode supports both metabolic and defensive roles within the symbiosis. *Genome Research* 22:2467–2477 DOI [10.1101/gr.138420.112](https://doi.org/10.1101/gr.138420.112).
- Darriba D, Taboada GL, Doallo R, Posada D.** 2012. JModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9:772 DOI [10.1038/nmeth.2109](https://doi.org/10.1038/nmeth.2109).

- Derks MFL, Smit S, Salis L, Schijlen E, Bossers A, Mateman C, Pijl AS, De Ridder D, Groenen MAM, Visser ME, Megens H-J.** 2015. The genome of winter moth (*Operophtera brumata*) provides a genomic perspective on sexual dimorphism and phenology. *Genome Biology and Evolution* 7:2321–2332 DOI [10.1093/gbe/evv145](https://doi.org/10.1093/gbe/evv145).
- Dittmer J, Bouchon D.** 2018. Feminizing Wolbachia influence microbiota composition in the terrestrial isopod *Armadillidium vulgare*. *Scientific Reports* 8:6998 DOI [10.1038/s41598-018-25450-4](https://doi.org/10.1038/s41598-018-25450-4).
- Dorđević M, Stojković B, Savković U, Immonen E, Tucić N, Lazarević J, Arnqvist G.** 2017. Sex-specific mitonuclear epistasis and the evolution of mitochondrial bioenergetics, ageing, and life history in seed beetles. *Evolution* 71:274–288 DOI [10.1111/evol.13109](https://doi.org/10.1111/evol.13109).
- Dobson SL, Bourtzis K, Braig HR, Jones BF, Zhou W, Rousset F, O'Neill SL.** 1999. Wolbachia infections are distributed throughout insect somatic and germ line tissues. *Insect Biochemistry and Molecular Biology* 29:153–160 DOI [10.1016/S0965-1748\(98\)00119-2](https://doi.org/10.1016/S0965-1748(98)00119-2).
- Dobson SL, Rattanadechakul W.** 2001. A novel technique for removing Wolbachia infections from *Aedes albopictus* (Diptera: Culicidae). *Journal of Medical Entomology* 38:844–849 DOI [10.1603/0022-2585-38.6.844](https://doi.org/10.1603/0022-2585-38.6.844).
- Dunning Hotopp JC, Clark ME, Oliveira DCSG, Foster JM, Fischer P, Torres MCM, Giebel JD, Kumar N, Ishmael N, Wang S, Ingram J, Nene RV, Shepard J, Tomkins J, Richards S, Spiro DJ, Ghedin E, Slatko BE, Tettelin H, Werren JH.** 2007. Widespread lateral gene transfer from intracellular bacteria to multicellular Eukaryotes. *Science* 317:1753–1756 DOI [10.1126/science.1142490](https://doi.org/10.1126/science.1142490).
- Duron O, Bouchon D, Boutin S, Bellamy L, Zhou L, Engelstadter J, Hurst GD.** 2008. The diversity of reproductive parasites among arthropods: Wolbachia do not walk alone. *BMC Biology* 6:27 DOI [10.1186/1741-7007-6-27](https://doi.org/10.1186/1741-7007-6-27).
- Ellegaard KM, Klasson L, Näslund K, Bourtzis K, Andersson SGE.** 2013. Comparative genomics of Wolbachia and the bacterial species concept. *PLOS Genetics* 9:e1003381 DOI [10.1371/journal.pgen.1003381](https://doi.org/10.1371/journal.pgen.1003381).
- Frost CL, Fernández-Marín H, Smith JE, Hughes WOH.** 2010. Multiple gains and losses of Wolbachia symbionts across a tribe of fungus-growing ants. *Molecular Ecology* 19:4077–4085 DOI [10.1111/j.1365-294X.2010.04764.x](https://doi.org/10.1111/j.1365-294X.2010.04764.x).
- Gerth M, Gansauge M-T, Weigert A, Bleidorn C.** 2014. Phylogenomic analyses uncover origin and spread of the Wolbachia pandemic. *Nature Communications* 5:5117 DOI [10.1038/ncomms6117](https://doi.org/10.1038/ncomms6117).
- Gerth M, Hurst GDD.** 2017. Short reads from honey bee (*Apis sp.*) sequencing projects reflect microbial associate diversity. *PeerJ* 5:e3529 DOI [10.7717/peerj.3529](https://doi.org/10.7717/peerj.3529).
- Ghedin E, Wang S, Foster JM, Slatko BE.** 2004. First sequenced genome of a parasitic nematode. *Trends in Parasitology* 20:151–153 DOI [10.1016/j.pt.2004.01.011](https://doi.org/10.1016/j.pt.2004.01.011).
- Giordano R, Jackson JJ, Robertson HM.** 1997. The role of Wolbachia bacteria in reproductive incompatibilities and hybrid zones of *Diabrotica* beetles and *Gryllus* crickets. *Proceedings of the National Academy of Sciences of the United States of America* 94:11439–11444 DOI [10.1073/pnas.94.21.11439](https://doi.org/10.1073/pnas.94.21.11439).

- Glowska E, Dragun-Damian A, Dabert M, Gerth M.** 2015. New Wolbachia supergroups detected in quill mites (Acari: Syringophilidae). *Infection, Genetics and Evolution* **30**:140–146 DOI [10.1016/J.MEEGID.2014.12.019](https://doi.org/10.1016/J.MEEGID.2014.12.019).
- Gomes FM, Hixson BL, Tyner MDW, Ramirez JL, Canepa GE, Alvese Silva TL, Molina-Cruz A, Keita M, Kane F, Traoré B, Sogoba N, Barillas-Mury C.** 2017. Effect of naturally occurring Wolbachia in *Anopheles gambiae* s.l. mosquitoes from Mali on *Plasmodium falciparum* malaria transmission. *Proceedings of the National Academy of Sciences of the United States of America* **114**:12566–12571 DOI [10.1073/pnas.1716181114](https://doi.org/10.1073/pnas.1716181114).
- Griffin PC, Khadake J, LeMay KS, Lewis SE, Orchard S, Pask A, Pope B, Roessner U, Russell K, Seemann T, Treloar A, Tyagi S, Christiansen JH, Dayalan S, Gladman S, Hangartner SB, Hayden HL, Ho WWH, Keeble-Gagnère G, Korhonen PK, Neish P, Prestes PR, Richardson MF, Watson-Haigh NS, Wyres KL, Young ND, Schneider MV.** 2017. Best practice data life cycle approaches for the life sciences. *F1000Research* **6**:1618 DOI [10.12688/f1000research.12344.1](https://doi.org/10.12688/f1000research.12344.1).
- Gurevich A, Saveliev V, Vyahhi N, Tesler G.** 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* **29**:1072–1075 DOI [10.1093/bioinformatics/btt086](https://doi.org/10.1093/bioinformatics/btt086).
- Haegeman A, Vanholme B, Jacob J, Vandekerckhove TTM, Claeys M, Borgonie G, Gheysen G.** 2009. An endosymbiotic bacterium in a plant-parasitic nematode: member of a new Wolbachia supergroup. *International Journal for Parasitology* **39**:1045–1054 DOI [10.1016/J.IJPARA.2009.01.006](https://doi.org/10.1016/J.IJPARA.2009.01.006).
- Haine ER, Pickup NJ, Cook JM.** 2005. Horizontal transmission of Wolbachia in a *Drosophila* community. *Ecological Entomology* **30**:464–472 DOI [10.1111/j.0307-6946.2005.00715.x](https://doi.org/10.1111/j.0307-6946.2005.00715.x).
- Heath BD, Butcher RDJ, Whitfield WGF, Hubbard SF.** 1999. Horizontal transfer of Wolbachia between phylogenetically distant insect species by a naturally occurring mechanism. *Current Biology* **9**:313–316 DOI [10.1016/S0960-9822\(99\)80139-0](https://doi.org/10.1016/S0960-9822(99)80139-0).
- Hertig M.** 1936. The Rickettsia, Wolbachia pipiens (gen. et sp.n.) and associated inclusions of the mosquito, *culex pipiens*. *Parasitology* **28**(4):453–486 DOI [10.1017/S0031182000022666](https://doi.org/10.1017/S0031182000022666).
- Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH.** 2008. How many species are infected with Wolbachia?—a statistical analysis of current data. *FEMS Microbiology Letters* **281**:215–220 DOI [10.1111/j.1574-6968.2008.01110.x](https://doi.org/10.1111/j.1574-6968.2008.01110.x).
- Hoffmann AA, Montgomery BL, Popovici J, Iturbe-Ormaetxe I, Johnson PH, Muzzi F, Greenfield M, Durkan M, Leong YS, Dong Y, Cook H, Axford J, Callahan AG, Kenny N, Omodei C, McGraw EA, Ryan PA, Ritchie SA, Turelli M, O'Neill SL.** 2011. Successful establishment of Wolbachia in *Aedes* populations to suppress dengue transmission. *Nature* **476**:454–457 DOI [10.1038/nature10356](https://doi.org/10.1038/nature10356).
- Holden PR, Brookfield JFY, Jones P.** 1993. Cloning and characterization of an *ftsZ* homologue from a bacterial symbiont of *Drosophila melanogaster*. *Molecular & General Genetics* **240**:213–220 DOI [10.1007/BF00277059](https://doi.org/10.1007/BF00277059).
- Holt RA.** 2002. The genome sequence of the malaria mosquito *Anopheles gambiae*. *Science* **298**:129–149 DOI [10.1126/science.1076181](https://doi.org/10.1126/science.1076181).

- Hotopp JCD.** 2011. Horizontal gene transfer between bacteria and animals. *Trends in Genetics* **27**(4):157–163.
- Hughes GL, Koga R, Xue P, Fukatsu T, Rasgon JL.** 2011. Wolbachia infections are virulent and inhibit the human malaria parasite Plasmodium falciparum in Anopheles gambiae. *PLOS Pathogens* **7**:e1002043 DOI [10.1371/journal.ppat.1002043](https://doi.org/10.1371/journal.ppat.1002043).
- Hurst GDD, Jiggins FM.** 1999. Male-killing Wolbachia in two species of insect. *Proceedings of the Royal Society B: Biological Sciences* **266**:735–740 DOI [10.1098/rspb.1999.0698](https://doi.org/10.1098/rspb.1999.0698).
- Husnik F, McCutcheon JP.** 2016. Repeated replacement of an intrabacterial symbiont in the tripartite nested mealybug symbiosis. *Proceedings of the National Academy of Sciences of the United States of America* **113**:e5416–E5424 DOI [10.1073/pnas.1603910113](https://doi.org/10.1073/pnas.1603910113).
- Ioannidis P, Hotopp JCD, Sapountzis P, Siozios S, Tsiamis G, Bordenstein SR, Baldo L, Werren JH, Bourtzis K.** 2007. New criteria for selecting the origin of DNA replication in Wolbachia and closely related bacteria. *BMC Genomics* **8**(1):182.
- Jamnongluk W, Kittayapong P, Baimai V, O'Neill SL.** 2002. Wolbachia infections of Tephritid fruit flies: molecular evidence for five distinct strains in a single host species. *Current Microbiology* **45**:255–260 DOI [10.1007/s00284-002-3746-1](https://doi.org/10.1007/s00284-002-3746-1).
- Jeffries CL, Lawrence GG, Golovko G, Kristan M, Orsborne J, Spence K, Hurn E, Bandibabone J, Tantely LM, Raharimalala FN, Keita K, Camara D, Barry Y, Watsenga F, Manzambi EZ, Afrane YA, Mohammed AR, Abeku TA, Hegde S, Khanipov K, Pimenova M, Fofanova Y, Boyer S, Irish SR, Hughes GL, Walker T.** 2018. Novel Wolbachia strains in *Anopheles* malaria vectors from Sub-Saharan Africa. *bioRxiv* DOI [10.1101/338434](https://doi.org/10.1101/338434).
- Jeyaprakash A, Hoy MA.** 2000. Long PCR improves Wolbachia DNA amplification: wsp sequences found in 76% of sixty-three arthropod species. *Insect Molecular Biology* **9**:393–405 DOI [10.1046/j.1365-2583.2000.00203.x](https://doi.org/10.1046/j.1365-2583.2000.00203.x).
- Jiggins FM, Bentley JK, Majerus MEN, Hurst GDD.** 2002. Recent changes in phenotype and patterns of host specialization in Wolbachia bacteria. *Molecular Ecology* **11**:1275–1283 DOI [10.1046/j.1365-294X.2002.01532.x](https://doi.org/10.1046/j.1365-294X.2002.01532.x).
- Kageyama D, Ohno M, Sasaki T, Yoshida A, Konagaya T, Jouraku A, Kuwazaki S, Kanamori H, Katayose Y, Narita S, Miyata M, Riegler M, Sahara K.** 2017. Femi-nizing Wolbachia endosymbiont disrupts maternal sex chromosome inheritance in a butterfly species. *Evolution Letters* **1**:232–244 DOI [10.1101/115386](https://doi.org/10.1101/115386).
- Katoh K, Standley DM.** 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**:772–780 DOI [10.1093/molbev/mst010](https://doi.org/10.1093/molbev/mst010).
- Kautz S, Rubin BER, Moreau CS.** 2013. Bacterial infections across the ants: frequency and prevalence of Wolbachia, Spiroplasma, and Asaia. *Psyche: A Journal of Entomology* **2013**:1–11 DOI [10.1155/2013/936341](https://doi.org/10.1155/2013/936341).
- Kikuchi Y, Fukatsu T.** 2003. Diversity of Wolbachia endosymbionts in heteropteran bugs. *Applied and Environmental Microbiology* **69**:6082–6090 DOI [10.1128/AEM.69.10.6082-6090.2003](https://doi.org/10.1128/AEM.69.10.6082-6090.2003).

- Kittayapong P, Baisley KJ, Baimai V, O'Neill SL.** 2000. Distribution and diversity of Wolbachia infections in Southeast Asian mosquitoes (Diptera: Culicidae). *Journal of Medical Entomology* **37**:340–345
DOI [10.1603/0022-2585\(2000\)037\[0340:DADOWI\]2.0.CO;2](https://doi.org/10.1603/0022-2585(2000)037[0340:DADOWI]2.0.CO;2).
- Kodandaramaiah U, Weingartner E, Janz N, Dalén L, Nylin S.** 2011. Population structure in relation to host-plant ecology and Wolbachia infestation in the comma butterfly. *Journal of Evolutionary Biology* **24**:2173–2185
DOI [10.1111/j.1420-9101.2011.02352.x](https://doi.org/10.1111/j.1420-9101.2011.02352.x).
- Kondo N, Ijichi N, Shimada M, Fukatsu T.** 2002. Prevailing triple infection with Wolbachia in *Callosobruchus chinensis* (Coleoptera: Bruchidae). *Molecular Ecology* **11**:167–180 DOI [10.1046/j.0962-1083.2001.01432.x](https://doi.org/10.1046/j.0962-1083.2001.01432.x).
- Kondo N, Shimada M, Fukatsu T.** 1999. High prevalence of Wolbachia in the Azuki Bean Beetle *Callosobruchus chinensis* (Coleoptera, Bruchidae). *Zoological Science* **16**:955–962 DOI [10.2108/zsj.16.955](https://doi.org/10.2108/zsj.16.955).
- Koukou K, Pavlikaki H, Kilias G, Werren JH, Bourtzis K, Alahiotis SN.** 2006. Influence of antibiotic treatment and Wolbachia curing on sexual isolation among *Drosophila melanogaster* cage populations. *Evolution* **60**:87–96
DOI [10.1111/j.0014-3820.2006.tb01084.x](https://doi.org/10.1111/j.0014-3820.2006.tb01084.x).
- Kraaijeveld K, Franco P, De Knijff P, Stouthamer R, Van Alphen JJ.** 2011. Clonal genetic variation in a Wolbachia-infected asexual wasp: horizontal transmission or historical sex? *Molecular Ecology* **20**(17):3644–3652
DOI [10.1111/j.1365-294X.2011.05150.x](https://doi.org/10.1111/j.1365-294X.2011.05150.x).
- Kumar S, Stecher G, Tamura K.** 2016. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* **33**:1870–1874
DOI [10.1093/molbev/msw054](https://doi.org/10.1093/molbev/msw054).
- Kunin V, Copeland A, Lapidus A, Mavromatis K, Hugenholtz P.** 2008. A Bioinformatician's guide to metagenomics. *Microbiology and Molecular Biology Reviews* **72**:557–578 DOI [10.1128/MMBR.00009-08](https://doi.org/10.1128/MMBR.00009-08).
- Kyei-Poku GK, Giladi M, Coghlin P, Mokady O, Zchori-Fein E, Floate KD.** 2006. Wolbachia in wasps parasitic on filth flies with emphasis on *Spalangia cameroni*. *Entomologia Experimentalis et Applicata* **121**:123–135
DOI [10.1111/j.1570-8703.2006.00469.x](https://doi.org/10.1111/j.1570-8703.2006.00469.x).
- Le Clec'h W, Chevalier FD, Genty L, Bertaux J, Bouchon D, Sicard M.** 2013. Cannibalism and predation as paths for horizontal passage of Wolbachia between terrestrial isopods. *PLOS ONE* **8**(4):e60232 DOI [10.1371/journal.pone.0060232](https://doi.org/10.1371/journal.pone.0060232).
- Leclercq S, Thézé J, Chebbi MA, Giraud I, Moumen B, Ernenwein L, Grève P, Gilbert C, Cordaux R.** 2016. Birth of a W sex chromosome by horizontal transfer of Wolbachia bacterial symbiont genome. *Proceedings of the National Academy of Sciences of the United States of America* **113**:15036–15041 DOI [10.1073/pnas.1608979113](https://doi.org/10.1073/pnas.1608979113).
- Li H.** 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. ArXiv preprint. [arXiv:1303.3997](https://arxiv.org/abs/1303.3997).
- Lindsey AR, Bordenstein SR, Newton IL, Rasgon JL.** 2016a. Wolbachia pipiensis should not be split into multiple species: a response to Ramírez-Puebla et al., species in

Wolbachia? proposal for the designation of ‘Candidatus Wolbachia bourtzisii’, ‘Candidatus Wolbachia onchocercicola’, ‘Candidatus Wolbachia blax. *Systematic and Applied Microbiology* **39**:220–222 DOI [10.1016/j.syapm.2016.03.001](https://doi.org/10.1016/j.syapm.2016.03.001).

Lindsey ARI, Werren JH, Richards S, Stouthamer R. 2016b. Comparative genomics of a parthenogenesis-inducing wolbachia symbiont. *G3: Genes-Genomes-Genetics* **6**:2113–2123 DOI [10.1534/g3.116.028449](https://doi.org/10.1534/g3.116.028449).

Lo N, Casiraghi M, Salati E, Bazzocchi C, Bandi C. 2002. How many Wolbachia supergroups exist? *Molecular Biology and Evolution* **19**:341–346 DOI [10.1093/oxfordjournals.molbev.a004087](https://doi.org/10.1093/oxfordjournals.molbev.a004087).

Lo N, Paraskevopoulos C, Bourtzis K, O'Neill SL, Werren JH, Bordenstein SR, Bandi C. 2007. Taxonomic status of the intracellular bacterium Wolbachia pipiensis. *International Journal of Systematic and Evolutionary Microbiology* **57**:654–657 DOI [10.1099/IJS.0.64515-0](https://doi.org/10.1099/IJS.0.64515-0).

Maier CT. 1981. Parasitoids emerging from Puparia of *Rhagoletis Pomonella* (Diptera: Tephritidae) infesting hawthorn and apple in Connecticut. *The Canadian Entomologist* **113**:867–870 DOI [10.4039/Ent113867-9](https://doi.org/10.4039/Ent113867-9).

Masui S, Sasaki T, Ishikawa H. 1997. groE-homologous operon of Wolbachia, an intracellular symbiont of arthropods: a new approach for their phylogeny. *Zoological Science* **14**:701–706 DOI [10.2108/zsj.14.701](https://doi.org/10.2108/zsj.14.701).

Mateos M, Castrezana SJ, Nankivell BJ, Estes AM, Markow TA, Moran NA. 2006. Heritable endosymbionts of Drosophila. *Genetics* **174**:363–376 DOI [10.1534/genetics.106.058818](https://doi.org/10.1534/genetics.106.058818).

Mock A, Tajovsky K, Žurovcová M, Jarošová A, Kocourek P, Gruber J, Angyal D, Spelda J. 2016. Hungarosoma bokori Verhoeff, 1928 (Diplopoda: Chordeumatida): new insights into its taxonomy, systematics, molecular genetics, biogeography and ecology. *Zootaxa* **4178**(2):234–256 DOI [10.11646/zootaxa.4178.2.4](https://doi.org/10.11646/zootaxa.4178.2.4).

Narita S, Nomura M, Kageyama D. 2007. Naturally occurring single and double infection with Wolbachia strains in the butterfly *Eurema hecabe*: transmission efficiencies and population density dynamics of each Wolbachia strain. *FEMS Microbiology Ecology* **61**:235–245 DOI [10.1111/j.1574-6941.2007.00333.x](https://doi.org/10.1111/j.1574-6941.2007.00333.x).

Ødegaard F. 2000. How many species of arthropods? Erwin’s estimate revised. *Biological Journal of the Linnean Society* **71**:583–597 DOI [10.1006/BIJL.2000.0468](https://doi.org/10.1006/BIJL.2000.0468).

O’Neill SL, Giordano R, Colbert AM, Karr TL, Robertson HM. 1992. 16S rRNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic incompatibility in insects. *Proceedings of the National Academy of Sciences of the United States of America* **89**:2699–2702 DOI [10.1073/pnas.89.7.2699](https://doi.org/10.1073/pnas.89.7.2699).

Osborne SE, Leong YS, O’Neill SL, Johnson KN. 2009. Variation in antiviral protection mediated by different Wolbachia strains in *Drosophila simulans*. *PLOS Pathogens* **5**:e1000656 DOI [10.1371/journal.ppat.1000656](https://doi.org/10.1371/journal.ppat.1000656).

Paraskevopoulos C, Bordenstein SR, Wernegreen JJ, Werren JH, Bourtzis K. 2006. Toward a Wolbachia multilocus sequence typing system: discrimination of Wolbachia strains present in *Drosophila* species. *Current Microbiology* **53**:388–395 DOI [10.1007/s00284-006-0054-1](https://doi.org/10.1007/s00284-006-0054-1).

- Perrot-Minnot MJ, Guo LR, Werren JH. 1996.** Single and double infections with Wolbachia in the parasitic wasp *Nasonia vitripennis*: effects on compatibility. *Genetics* **143**:961–972.
- Pintureau B, Chaudier S, Lassablière F, Charles H, Grenier S. 2000.** Addition of wsp sequences to the Wolbachia phylogenetic tree and stability of the classification. *Journal of Molecular Evolution* **51**:374–377 DOI [10.1007/s002390010099](https://doi.org/10.1007/s002390010099).
- Plantard O, Rasplus JY, Mondor G, Le Clainche I, Solignac M. 1999.** Distribution and phylogeny of Wolbachia inducing thelytoky in Rhoditini and Aylacini (Hymenoptera: Cynipidae). *Insect Molecular Biology* **8**:185–191 DOI [10.1046/j.1365-2583.1999.820185.x](https://doi.org/10.1046/j.1365-2583.1999.820185.x).
- Quinlan AR, Hall IM. 2010.** BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**:841–842 DOI [10.1093/bioinformatics/btq033](https://doi.org/10.1093/bioinformatics/btq033).
- Ramirez-Puebla ST, Ormeno-Orrillo E, Vera-Ponce de Leon A, Lozano L, Sanchez A, Rosenblueth M, Martinez-Romero E. 2016.** Genomes of *Candidatus Wolbachia Bourtzisii* wDacA and *Candidatus Wolbachia Pipiensis* wDacB from the Cochineal Insect *Dactylopius coccus* (Hemiptera: Dactylopiidae). *G3: Genes-Genomes-Genetics* **6**:3343–3349 DOI [10.1534/g3.116.031237](https://doi.org/10.1534/g3.116.031237).
- Rasgon JL, Scott TW. 2004.** An initial survey for *Wolbachia* (Rickettsiales: Rickettsiaceae) infections in selected California Mosquitoes (Diptera: Culicidae): Table 1. *Journal of Medical Entomology* **41**:255–257 DOI [10.1603/0022-2585-41.2.255](https://doi.org/10.1603/0022-2585-41.2.255).
- Raychoudhury R, Baldo L, Oliveira DCSG, Werren JH. 2009.** Modes of acquisition of Wolbachia: horizontal transfer, hybrid introgression, and codivergence in the *Nasonia* species complex. *Evolution* **63**:165–183 DOI [10.1111/j.1558-5646.2008.00533.x](https://doi.org/10.1111/j.1558-5646.2008.00533.x).
- Ricci I, Cancrini G, Gabrielli S, D'amelio S, Favia G. 2002.** Searching for *Wolbachia* (Rickettsiales: Rickettsiaceae) in Mosquitoes (Diptera: Culicidae): large polymerase chain reaction survey and new identifications. *Journal of Medical Entomology* **39**:562–567 DOI [10.1603/0022-2585-39.4.562](https://doi.org/10.1603/0022-2585-39.4.562).
- Richardson MF, Weinert LA, Welch JJ, Linheiro RS, Magwire MM, Jiggins FM, Bergman CM. 2012.** Population genomics of the Wolbachia endosymbiont in *Drosophila melanogaster*. *PLOS Genetics* **8**(12):e1003129 DOI [10.1371/journal.pgen.1003129](https://doi.org/10.1371/journal.pgen.1003129).
- Riegler M, Charlat S, Stauffer C, Merçot H. 2004.** Wolbachia transfer from *Rhagoletis cerasi* to *Drosophila simulans*: investigating the outcomes of host-symbiont coevolution. *Applied and Environmental Microbiology* **70**:273–279 DOI [10.1128/AEM.70.1.273-279.2004](https://doi.org/10.1128/AEM.70.1.273-279.2004).
- Rogers EE, Backus EA. 2014.** Anterior foregut microbiota of the glassy-winged sharpshooter explored using deep 16S rRNA gene sequencing from individual insects. *PLOS ONE* **9**(9):e106215 DOI [10.1371/journal.pone.0106215](https://doi.org/10.1371/journal.pone.0106215).
- Rokas A, Atkinson RJ, Brown GS, West SA, Stone GN. 2001.** Understanding patterns of genetic diversity in the oak gallwasp *Biorhiza pallida*: demographic history or a Wolbachia selective sweep? *Heredity* **87**:294–304 DOI [10.1046/j.1365-2540.2001.00872.x](https://doi.org/10.1046/j.1365-2540.2001.00872.x).

- Rokas A, Atkinson RJ, Nieves-Aldrey JL, West SA, Stone GN.** 2002. The incidence and diversity of Wolbachia in gallwasps (Hymenoptera; Cynipidae) on oak. *Molecular Ecology* **11**:1815–1829 DOI [10.1046/j.1365-294X.2002.01556.x](https://doi.org/10.1046/j.1365-294X.2002.01556.x).
- Ros VID, Fleming VM, Feil EJ, Breeuwer JAJ.** 2009. How diverse is the genus Wolbachia? Multiple-gene sequencing reveals a putatively new Wolbachia supergroup recovered from spider mites (Acarina: Tetranychidae). *Applied and Environmental Microbiology* **75**:1036–1043 DOI [10.1128/AEM.01109-08](https://doi.org/10.1128/AEM.01109-08).
- Ros VID, Fleming VM, Feil EJ, Breeuwer JAJ.** 2012. Diversity and recombination in Wolbachia and Cardinium from Bryobia spider mites. *BMC Microbiology* **12**(1):S13 DOI [10.1186/1471-2180-12-S1-S13](https://doi.org/10.1186/1471-2180-12-S1-S13).
- Rousset F, Braig HR, O'Neill SL.** 1999. A stable triple Wolbachia infection in Drosophila with nearly additive incompatibility effects. *Heredity* **82**(Pt. 6):620–627 DOI [10.1046/j.1365-2540.1999.00501.x](https://doi.org/10.1046/j.1365-2540.1999.00501.x).
- Rowley SM, Raven RJ, McGraw EA.** 2004. Wolbachia pipiensis in Australian Spiders. *Current Microbiology* **49**(3):208–214 DOI [10.1007/s00284-004-4346-z](https://doi.org/10.1007/s00284-004-4346-z).
- Russell JA, Funaro CF, Giraldo YM, Goldman-Huertas B, Suh D, Kronauer DJC, Moreau CS, Pierce NE.** 2012. A veritable menagerie of heritable bacteria from ants, butterflies, and beyond: broad molecular surveys and a systematic review. *PLOS ONE* **7**:e51027 DOI [10.1371/journal.pone.0051027](https://doi.org/10.1371/journal.pone.0051027).
- Saha S, Hunter WB, Reese J, Morgan JK, Marutani-Hert M, Huang H, Lindeberg M.** 2012. Survey of endosymbionts in the *Diaphorina citri* metagenome and assembly of a *Wolbachia* wDi draft genome. *PLOS ONE* **7**:e50067 DOI [10.1371/journal.pone.0050067](https://doi.org/10.1371/journal.pone.0050067).
- Salzberg SL, Dunning Hotopp JC, Delcher AL, Pop M, Smith DR, Eisen MB, Nelson WC.** 2005. Serendipitous discovery of Wolbachia genomes in multiple Drosophila species. *Genome Biology* **6**(3):R23 DOI [10.1186/gb-2005-6-3-r23](https://doi.org/10.1186/gb-2005-6-3-r23).
- Saridaki A, Bourtzis K.** 2010. Wolbachia: more than just a bug in insects' genitals. *Current Opinion in Microbiology* **13**:67–72 DOI [10.1016/j.mib.2009.11.005](https://doi.org/10.1016/j.mib.2009.11.005).
- Schilthuizen M, Honda J, Stouthamer R.** 1998. Parthenogenesis-inducing Wolbachia in *Trichogramma kaykai* (hymenoptera: Trichogrammatidae) originates from a single infection. *Annals of the Entomological Society of America* **91**:410–414 DOI [10.1093/aesa/91.4.410](https://doi.org/10.1093/aesa/91.4.410).
- Schilthuizen M, Stouthamer R.** 1998. Distribution of Wolbachia among the guild associated with the parthenogenetic gall wasp *Diplolepis rosae*. *Heredity* **81**:270–274 DOI [10.1038/sj.hdy.6883850](https://doi.org/10.1038/sj.hdy.6883850).
- Schmieder R, Edwards R.** 2011. Fast identification and removal of sequence contamination from genomic and metagenomic datasets. *PLOS ONE* **6**:e17288 DOI [10.1371/journal.pone.0017288](https://doi.org/10.1371/journal.pone.0017288).
- Schuler H, Arthofer W, Riegler M, Bertheau C, Krumböck S, Köppler K, Vogt H, Teixeira LAF, Stauffer C.** 2011. Multiple Wolbachia infections in *Rhagoletis pomonella*. *Entomologia Experimentalis et Applicata* **139**:138–144 DOI [10.1111/j.1570-7458.2011.01115.x](https://doi.org/10.1111/j.1570-7458.2011.01115.x).

- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015.** BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**:3210–3212 DOI [10.1093/bioinformatics/btv351](https://doi.org/10.1093/bioinformatics/btv351).
- Simões PM, Mialdea G, Reiss D, Sagot M-F, Charlat S. 2011.** Wolbachia detection: an assessment of standard PCR Protocols. *Molecular Ecology Resources* **11**:567–572 DOI [10.1111/j.1755-0998.2010.02955.x](https://doi.org/10.1111/j.1755-0998.2010.02955.x).
- Stamatakis A. 2014.** RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**:1312–1313 DOI [10.1093/bioinformatics/btu033](https://doi.org/10.1093/bioinformatics/btu033).
- Stelinski LL, Pelz KS, Liburd OE. 2004.** Field observations quantifying attraction of the Parasitic Wasp, *Diachasma Alloeum* (Hymenoptera: Braconidae) to Blueberry fruit infested by the Blueberry Maggot Fly, *Rhagoletis mendax* (Diptera: Tephritidae). *Florida Entomologist* **87**:124–129 DOI [10.1653/0015-4040\(2004\)087\[0124:FOQAOT\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2004)087[0124:FOQAOT]2.0.CO;2).
- Stouthamer R, Luck RF, Hamilton WD. 1990.** Antibiotics cause parthenogenetic Trichogramma (Hymenoptera/Trichogrammatidae) to revert to sex. *Proceedings of the National Academy of Sciences of the United States of America* **87**:2424–2427 DOI [10.1073/pnas.87.7.2424](https://doi.org/10.1073/pnas.87.7.2424).
- Subandiyah S, Nikoh N, Tsuyumu S, Somowiyarjo S, Fukatsu T. 2000.** Complex endosymbiotic microbiota of the citrus psyllid *Diaphorina citri* (Homoptera: Psylloidea). *Zoological Science* **17**:983–989 DOI [10.2108/zsj.17.983](https://doi.org/10.2108/zsj.17.983).
- Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. 2015.** Sambamba: fast processing of NGS alignment formats. *Bioinformatics* **31**:2032–2034 DOI [10.1093/bioinformatics/btv098](https://doi.org/10.1093/bioinformatics/btv098).
- Teixeira L, Ferreira Á, Ashburner M. 2008.** The bacterial symbiont Wolbachia induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLOS Biology* **6**:e1000002 DOI [10.1371/journal.pbio.1000002](https://doi.org/10.1371/journal.pbio.1000002).
- Tsai K-H, Lien J-C, Huang C-G, Wu W-J, Chen W-J. 2004.** Molecular (sub) grouping of endosymbiont Wolbachia infection among mosquitoes of Taiwan. *Journal of Medical Entomology* **41**:677–683 DOI [10.1603/0022-2585-41.4.677](https://doi.org/10.1603/0022-2585-41.4.677).
- Van Borm S, Wenseleers T, Billen J, Boomsma J. 2003.** Cloning and sequencing of wsp encoding gene fragments reveals a diversity of co-infecting Wolbachia strains in Acromyrmex leafcutter ants. *Molecular Phylogenetics and Evolution* **26**:102–109 DOI [10.1016/S1055-7903\(02\)00298-1](https://doi.org/10.1016/S1055-7903(02)00298-1).
- Vavre F, Fleury F, Lepetit D, Fouillet P, Boulétreau M. 1999.** Phylogenetic evidence for horizontal transmission of Wolbachia in host-parasitoid associations. *Molecular Biology and Evolution* **16**:1711–1723 DOI [10.1093/oxfordjournals.molbev.a026084](https://doi.org/10.1093/oxfordjournals.molbev.a026084).
- Verne S, Johnson M, Bouchon D, Grandjean F. 2007.** Evidence for recombination between feminizing Wolbachia in the isopod genus *Armadillidium*. *Gene* **397**:58–66 DOI [10.1016/j.gene.2007.04.006](https://doi.org/10.1016/j.gene.2007.04.006).
- Walker T, Moreira LA. 2011.** Can Wolbachia be used to control malaria? *Memórias do Instituto Oswaldo Cruz* **106**:212–217 DOI [10.1590/S0074-02762011000900026](https://doi.org/10.1590/S0074-02762011000900026).

- Wang Y, Chandler C.** 2016. Candidate pathogenicity islands in the genome of *Candidatus Rickettsiella isopodorum*, an intracellular bacterium infecting terrestrial isopod crustaceans. *PeerJ* 4:e2806 DOI 10.7717/peerj.2806.
- Weeks AR, Turelli M, Harcombe WR, Reynolds KT, Hoffmann AA.** 2007. From parasite to mutualist: rapid evolution of Wolbachia in natural populations of *Drosophila*. *PLOS Biology* 5:e114 DOI 10.1371/journal.pbio.0050114.
- Werren JH.** 1997a. Wolbachia run amok. *Proceedings of the National Academy of Sciences of the United States of America* 94:11154–11155 DOI 10.1073/pnas.94.21.11154.
- Werren JH.** 1997b. Biology of Wolbachia. *Annual Review of Entomology* 42:587–609 DOI 10.1146/annurev.ento.42.1.587.
- Werren JH, Bartos JD.** 2001. Recombination in Wolbachia. *Current Biology* 11:431–435 DOI 10.1016/S0960-9822(01)00101-4.
- Werren JH, Windsor DM.** 2000. Wolbachia infection frequencies in insects: evidence of a global equilibrium? *Proceedings of the Royal Society B: Biological Sciences* 267:1277–1285 DOI 10.1098/rspb.2000.1139.
- Werren JH, Windsor D, Guo L.** 1995. Distribution of Wolbachia among Neotropical Arthropods. *Proceedings of the Royal Society B: Biological Sciences* 262:197–204 DOI 10.1098/rspb.1995.0196.
- Werren JH, Zhang W, Guo LR.** 1995. Evolution and phylogeny of Wolbachia: reproductive parasites of arthropods. *Proceedings of the Royal Society B: Biological Sciences* 261:55–63 DOI 10.1098/rspb.1995.0117.
- Westermann AJ, Gorski SA, Vogel J.** 2012. Dual RNA-seq of pathogen and host. *Nature Reviews Microbiology* 10:618–630 DOI 10.1038/nrmicro2852.
- Wilkinson MD, Dumontier M, Aalbersberg IJ, Appleton G, Axton M, Baak A, Blomberg N, Boiten J-W, Da Silva Santos LB, Bourne PE, Bouwman J, Brookes AJ, Clark T, Crosas M, Dillo I, Dumon O, Edmunds S, Evelo CT, Finkers R, Gonzalez-Beltran A, Gray AJG, Groth P, Goble C, Grethe JS, Heringa J, 't Hoen PAC, Hooft R, Kuhn T, Kok R, Kok J, Lusher SJ, Martone ME, Mons A, Packer AL, Persson B, Rocca-Serra P, Roos M, Van Schaik R, Sansone S-A, Schultes E, Sengstag T, Slater T, Strawn G, Swertz MA, Thompson M, Van der Lei J, Van Mulligen E, Velterop J, Waagmeester A, Wittenburg P, Wolstencroft K, Zhao J, Mons B.** 2016. The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data* 3:160018 DOI 10.1038/sdata.2016.18.
- Wiwatwanaratanaabutr I, Kittayapong P, Caubet Y, Bouchon D.** 2009. Molecular phylogeny of wolbachia strains in arthropod hosts based on groe-homologous gene sequences. *Zoological Science* 26:171–177 DOI 10.2108/zsj.26.171.
- Woolfit M, Iturbe-Ormaetxe I, Brownlie JC, Walker T, Riegler M, Seleznev A, Popovici J, Rancés E, Wee BA, Pavlides J, Sullivan MJ, Beatson SA, Lane A, Sidhu M, McMeniman CJ, McGraw EA, O'Neill SL.** 2013. Genomic evolution of the pathogenic Wolbachia strain, wMelPop. *Genome Biology and Evolution* 5(11):2189–2204.
- Zabalou S, Charlat S, Nirgianaki A, Lachaise D, Merçot H, Bourtzis K.** 2004. Natural Wolbachia infections in the *Drosophila yakuba* species complex do not induce

- cytoplasmic incompatibility but fully rescue the wRi modification. *Genetics* **167**(2):827–834.
- Zerbino DR, Birney E. 2008.** Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Research* **18**:821–829 DOI [10.1101/gr.074492.107](https://doi.org/10.1101/gr.074492.107).
- Zha X, Zhang W, Zhou C, Zhang L, Xiang Z, Xia Q. 2014.** Detection and characterization of Wolbachia infection in silkworm. *Genetics and Molecular Biology* **37**:573–580 DOI [10.1590/S1415-47572014000400014](https://doi.org/10.1590/S1415-47572014000400014).
- Zhou W, Rousset F, O’Neil S. 1998.** Phylogeny and PCR-based classification of Wolbachia strains using wsp gene sequences. *Proceedings of the Royal Society B: Biological Sciences* **265**:509–515 DOI [10.1098/rspb.1998.0324](https://doi.org/10.1098/rspb.1998.0324).