



Association of burnout with doctor–patient relationship and common stressors among postgraduate trainees and house officers in Lahore—a cross-sectional study

Waqas Ahmad^{*}, Huma Ashraf^{*}, Afnan Talat^{*}, Aleena Ahmad Khan^{*}, Ammad Anwar Baig^{*}, Iqra Zia^{*}, Zohak Sarfraz^{*}, Hifsa Sajid^{*}, Marium Tahir^{*}, Usman Sadiq^{*} and Hira Imtiaz^{*}

CMH Lahore Medical College, Lahore, Pakistan

^{*}These authors contributed equally to this work.

ABSTRACT

Introduction. Burnout is defined as a prolonged state of physical and psychological exhaustion. Doctors, due to the demanding nature of their job, are susceptible to facing burnout, which has far reaching implications on their productivity and motivation. It affects the quality of care they provide to patients, thus eroding the doctor–patient relationship which embodies patient centeredness and autonomy. The study aims at addressing the stressors leading to burnout and its effect on the doctor–patient relationship.

Methods. A descriptive, cross-sectional study design with convenience (non-probability) sampling technique was employed in six major hospitals of Lahore, Pakistan. A total of 600 doctors were approached for the study which included house officers or “HOs” (recent graduates doing their 1 year long internship) and post-graduate trainees or “PGRs” (residents for 4–5 years in their specialties). Burnout was measured using the Copenhagen Burnout Inventor (CBI) while attitudes towards the doctor–patient relationship was measured using the Patient Practitioner Orientation Scale (PPOS), which measures two components of the relationship: power sharing and patient caring. Pearson correlation and linear regression analysis were used to analyze the data via SPSS v.21.

Results. A total of 515 doctors consented to take part in the study (response rate 85.83%). The final sample consisted of 487 doctors. The burnout score was not associated with the total and caring domain scores of PPOS ($P > 0.05$). However, it was associated with the power sharing sub-scale of PPOS. Multiple linear regression analysis yielded a significant model, by virtue of which CBI scores were positively associated with factors such as female gender, feeling of burn out, scoring high on sharing domain of PPOS and a lack of personal control while CBI scores were negatively associated with private medical college education, having a significant other, accommodation away from home and a sense of never ending competition. Burnout levels varied significantly between house officers and post graduate trainees. Twenty-three percent of the participants (mostly house officers) had high/very high burnout levels on the

Submitted 24 February 2018
Accepted 6 August 2018
Published 10 September 2018

Corresponding author
Huma Ashraf,
humaashraff@gmail.com

Academic editor
Antonio Palazón-Bru

Additional Information and
Declarations can be found on
page 9

DOI 10.7717/peerj.5519

 Copyright
2018 Ahmad et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

CBI (Kristenson's burnout scoring). Both groups showed significant differences with respect to working hours, smoking status and income.

Conclusion. Although burnout showed no significant association with total and caring domain scores of PPOS (scale used to assess doctor–patient relationship), it showed a significant association with the power sharing domain of PPOS suggesting some impact on the overall delivery of patient care. Thus, it necessitates the monitoring of stressors in order to provide an atmosphere where patient autonomy can be practiced.

Subjects Psychiatry and Psychology, Public Health

Keywords Burnout, Distress, Doctor patient relationship, Lahore, Stressors, PPOS, CBI, Patient care, Doctors

INTRODUCTION

Stress, in moderation, is a driving force for humans to thrive but when present in excess leads, among other things, to burnout—“a state of prolonged physical and psychological exhaustion.” (*Dyrbye et al., 2014*). Burnout has become an integral part of the lives of medical students (*Dyrbye et al., 2014*) and practicing health professionals (*Dyrbye et al., 2006*).

Burnout is a well-documented phenomenon amongst students of medicine worldwide. A study done in Minnesota on medical students found a high (45%) prevalence of burnout. (*Dyrbye et al., 2006*). Similarly, another study found burnout rates as high as 49.5% among Greek medical students (*Msaouel et al., 2010*). In 2014, we conducted a study at two medical colleges of Pakistan and discovered that 47% of students had high levels of burnout. In addition, 30% of the students with high levels were not aware of being burnt out (*Muzafar et al., 2015*). Medical students, due to the uncertainties of the future, rigorous system of evaluation and never-ending need to keep up with the latest trends in the evolving medical field whilst maintaining a social life, inevitably tend to burn out (*Dyrbye et al., 2014*). It is hypothesised that this will continue to affect the work of students as they progress through medical training as house officers (an MBBS intern working under the supervision of an attending physician) and as postgraduate trainee (resident) doctors.

In a study conducted in Kerala, India, patient related burnout was reported to be the highest among the medical and surgical interns (68.62%) in contrast to non-medical/non-surgical residents. The study also showed a positive correlation between the number of years of residency and the degree of burnout (*Ratnakaran, Prabhakaran & Karunakaran, 2016*). Moreover, risk factors like working more than 8 hours per day, high workload, and a lack of self-care were linked to burnout (*Rizo-Baeza et al., 2018*). Both in the work and personal sphere, people expect equality of gain for their investment. A higher level of depersonalization, emotional exhaustion and a decreased sense of personal accomplishment was reported in student nurses who felt that they were committing more to their hospital and patients than what they were receiving back (*Schaufeli, Dierendonck & Gorp, 1996*). In addition to work, family stressors like being a single parent can also play a role in causing burnout (*Rizo-Baeza et al., 2018*). In contrast, an empowering leadership leads to

more engaged workers and hence causes a decrease in burnout (*Laschinger et al., 2015*). Healthcare workers with support from their workplace (e.g., from supervisors) reported a decrease in emotional exhaustion, which can ultimately decrease burnout (*Constable & Russell, 1986*).

Studies have found that many medical trainees experience burnout (*Dyrbye et al., 2014*) which may be problematic as it might erode a healthy doctor patient relationship; this relationship embodies trust and empathy (*Ha & Longnecker, 2010*). The hallmark of an accurate diagnosis and the effectiveness of the treatment administered is attributed to the quality of the doctor–patient relationship (*Hellin, 2002*). A patient-centred or egalitarian relationship gives the patient complete freedom over the choice and approach of his treatment, thus bridging the gap between the doctor and patient with lesser incidence of litigations and complaints (*Beckman et al., 1994*). On the other hand, a doctor-centred or paternalistic relationship limits the exchange of information and takes away the patient’s control over his treatment. This type of relationship is believed to result from cynicism and the lack of empathy that are linked to burnout (*Paro et al., 2014*).

A number of studies have been conducted to find the association between burnout and the doctor–patient relationship. A study conducted by *Ratanawongsa et al. (2008)* showed no association between burnout and differences in the doctor’s behaviour towards their patients. However, burnout did seem to have a positive correlation with the patient’s communication with the physician; patients demonstrated more rapport-building during interactions with physicians who had high burn out (*Ratanawongsa et al., 2008*). In contrast, a meta-analysis concluded that burnout is attributed to a poor doctor-relationship (*Sablik, Samborska-Sablik & Drożdż, 2013*). This notion, that burnout causes difficulty in establishing a mutual doctor–patient relationship, still remains questionable.

It is for these reasons and the lack of adequate research and reliable data, particularly from Pakistan (one of the most populated countries), that necessitated us to evaluate the relationship between burnout and the doctor–patient relationship. In lieu of these observations we aim: (1) to assess the level of clinically significant burnout among the sample of post-graduate trainees and house officers; (2) to analyse various personal, professional and organisational factors contributing to a physician’s burnout; (3) to draw a comparison between HOs and PGRs for burnout and its risk factors, and; (4) to explore a possible association of burnout with the doctor–patient relationship and assess attitude of doctors towards the relationship.

METHODOLOGY

Study sample

A descriptive, cross-sectional study design with convenience (non-probability) sampling technique was employed. In Pakistan after completion of basic medical education (MBBS) doctors embark on 1 year long training post of “House Officer” or “HO” after which they are legally allowed to practice medicine in Pakistan. If a doctor chooses to pursue specialty training they do so by becoming “Post Graduate Residents” or “PGRs” for 4–5 years in their respective specialties. 600 doctors were approached for the study out of which 515

consented to take part (response rate was 85.83%). Twenty-eight respondents were excluded due to lack of basic information e.g., age, gender, etc., along with missing more than three and two responses in PPOS (Patient Practitioner Orientation Scale) and CBI (Copenhagen Burnout Inventory), respectively. The final sample size thus consisted of 487 doctors. Data collection was done entirely in Lahore city among various hospitals, both public and private, between August 2017 and December 2017. A series of demographic questions were included in the questionnaire to record some basic information.. A comparison between the postgraduate trainees and house officers was made in this research.

Instruments

Copenhagen Burnout Inventory

Copenhagen Burnout Inventory (CBI) was employed which measures three aspects of burnout; Personal, Work related and Patient related. Only personal sub-domain of burnout was assessed in this study with a six-item scale. Cronbach's alpha is 0.80. All burnout items were intermixed. The responses have the following categories: Always, Often, Sometimes, Seldom, Never/almost never. Each corresponds to a score of 100, 75, 50, 25, and 0. The total score on the scale is the average of the scores on the items. Kristensen's classification was also used which divides the scores by setting up cut off points such as: nil, low, moderate, high and very high burnout, where: nil=score equal to lowest 25% of distribution of scores, low=score of next higher 25% of distribution of scores, moderate=score of the next 12.5% of distribution of scores, high=score of the next 12.5% of distribution of scores, and very high=score of the highest 25% of distribution of scores ([Winwood & Winefield, 2004](#); [Kristensen et al., 2005](#)).

Patient Practitioner Orientation Scale

The attitudes of participants towards doctor-patient relationship were assessed by using a reliable instrument called Patient Practitioner Orientation Scale (PPOS) ([Krupat et al., 2000](#)). The PPOS contains 18-items and measures the subject's leaning towards a doctor-centred or a patient-centred care. Cronbach's alpha is 0.83. Each item has six possible responses ranging from 1 (strongly agree) to 6 (strongly disagree). PPOS also contains two subscales which measure two sub-domains of doctor-patient relationship: Sharing and Caring. Sharing refers to an individual's belief that a patient should have the right to be an equal part of decision making process along with the physician. Caring refers to a doctor's belief that a patient should be treated as a whole rather than just as a disease. Both sub-scales have nine items each. All the scores are reported as mean of the total score ranging from 1 (doctor centred) to 6 (patient centred).

Stressors' scale

Stressful events were measured on a 30 item questionnaire which has a five point likert scale ranging from none of the time to all of the time. The headings are classified as: lifestyle, psychological, work and personal stressors.

Statistical analysis

SPSS Inc., (Chicago, IL, USA) version 21 software was used for analysis. Descriptive statistics are shown for age, gender, level of training, etc. ([Table 1](#)). Independent-sample

Table 1 Demographics of the participants ($N = 487$).

Parameter	<i>n</i> (%)
Mean age \pm SD	26.39 \pm 2.97
Gender	
Male	216(44.4)
Female	271(55.6)
Level of training	
HO	282(58.0)
PGR	205(42.0)
Living status	
With family	243(49.9)
Hostel	244(50.1)
Relationship status	
Single	289(59.3)
Married	114(23.4)
In a relationship	27(5.5)
Engaged	56(11.5)
Divorced	1(0.2)

t-test was used for the comparison of means. Five groups of responses for stressful events (None of the time, Rarely, Some of the time, Often and All of the time) were regrouped into two groups for the simplicity of analysis. None of the time, rarely and some of the time were grouped together, while often and all of the time were grouped together. Multiple Linear Regression analysis was run to analyse association of scores on CBI (dependent variable) with demographic characteristics of respondents, age, income, stressors and PPOS as predictor variables. Histogram and P-P plots were visualized to assess the assumption of normality of data. Durbin Watson test, Cook's distance, case wise diagnostics and values of variance inflation factor (VIF), tolerance (TOL) were run to ensure all assumptions of multiple regression analysis were met. Categorical variables were re-coded as dummy variables with values 0 and 1. The variables with more than 1 option e.g., relationship status were regrouped into two logical groups (having a significant other and having no significant other) for a robust regression model (final sample, $N = 487$).

Ethical statement

Ethical approval (reference number 40 ERC/CMHLMC) from Ethics Review Committee of CMH Lahore Medical College and Institute of Dentistry was obtained. Informed written consent was sought from each participant and their personal details were kept confidential.

RESULTS

Demographics

Demographic characteristics of the participants are shown in [Table 1](#). According to [Table 1](#), the sample at hand is quite young and represents both genders almost equally. More HOs than PGRs took part in the study, which simply reflects their abundance over PGRs.

Table 2 Significant determinants of burnout (regression analysis) ($N = 487$).

Model	Unstandardized coefficients		p -value	95.0% Confidence interval for B	
	B	Std. error		Lower bound	Upper bound
(Constant)	31.298	12.050	.010	7.616	54.981
Participant's Gender (female)	6.035	1.815	<0.001	2.468	9.602
Graduated from a Private Institute	-7.335	2.438	.003	-12.125	-2.544
Relationship status?	-3.420	1.679	.042	-6.721	-.120
Burn-out is a state of prolonged physical and psychological exhaustion. Do you feel burnt out?	11.794	1.709	.001	8.435	15.153
SharingPPOS (mean)	2.510	1.235	.043	.083	4.938
CaringPPOS (mean)	-1.407	1.465	.338	-4.287	1.474
Accommodation away from home.	-5.007	1.938	.010	-8.815	-1.198
Lack of personal control over what you do.	5.160	2.280	.024	.679	9.642
Sense of never ending competition.	-6.896	1.828	<0.001	-10.488	-3.304

Notes.

$R = 0.637$, $R_2 = 0.405$, $R_3 = 0.405$, Durbin Watson = 1.846, Dependent Variable: CBI Score.

Risk factors

Mean PPOS score of the entire sample was 3.63 ± 0.45 . Subscale scores for caring and sharing domain were 3.92 ± 0.55 and 3.34 ± 0.67 respectively. CBI and total PPOS score along with caring domain had no significant linear correlation. Sharing domain, however, was significantly correlated with CBI scores in the multiple linear regression model (Table 2). Total PPOS, sharing and caring domain yielded no significant multiple regression models with demographics and stressful events.

Mean CBI scores were 51.18 ± 20.29 . CBI scores yielded a significant regression model with the demographics and stressful events (Table 2). High scores on CBI were positively associated with female gender, perception of being burnt out, scoring high on sharing domain of PPOS and lack of personal control over what one does. Similarly graduating from a private medical college (Table 2), having a significant other (Table 2), accommodation away from home (Table 2) and sense of never ending competition (Table 2) were negatively correlated with CBI scores. All other predictor variables did not contribute significantly towards predicting CBI scores. Of the participants stating self-perception of burnout, only 36% had high/very high burnout levels.

Comparison

A comparison between HO and PGRs was made in Table 3 where they are compared by Pearson chi-square and Independent Sample t -test.

DISCUSSION

In our study, the first hypothesis that house officers (HOs) and post graduate trainees (PGRs) would have a high burnout rate was supported by 29.8% of HOs and 16.1% of PGRs reporting high/very high burnout. A number of studies reported similar results (Al-Dubai et al., 2013; Shanafelt et al., 2010; Ratanawongsa et al., 2008; Shanafelt et al.,

Table 3 A comparison of house officers and post graduate residents.

Variable (total $n = 487$)	House officer ($n = 282$) $n(\%)$	Post graduate resident ($n = 205$) $n(\%)$	p -value
No income or \leq USD 312/month ^b	93.6	20.5	<0.001
Income \geq USD 312/Month ^b	6.4	79.5	<0.001
Graduated from govt. institute ^b	62.4	81.5	<0.001
Graduated from private institute ^b	37.6	18.5	<0.001
Smoke ^b	7.8	17.1	<0.001
Burn-out is a state of prolonged physical and psychological	62.4	45.4	<0.001
High/Very High Burnout (According to Kristensen's Classification of CBI) ^b	29.8	16.1	<0.001
Wouldn't change their profession if had the option ^b	65.2	56.6	0.033
Working hours/week mean (\pm SD) ^a	65.09 \pm 16.01	58.15 \pm 14.69	<0.001

Notes.^aIndependent sample t -test.^bPearson Chi-Square.

2015; Guthrie et al., 1998; Dyrbye et al., 2010; Marjani et al., 2008; Dyrbye et al., 2014; Ratnakaran, Prabhakaran & Karunakaran, 2016).

This observation can be explained by the emotionally depleting nature of the job, long working hours (Khamisa, Peltzer & Oldenburg, 2013), patient overload (Khamisa, Peltzer & Oldenburg, 2013) and challenging work environment. These factors may account for the significant difference observed between burnout scores of public and private hospitals (Table 2, $p = 0.003$). Moreover, our study showed that a high percentage of doctors with burnout reported feeling burnt out, showing self-awareness of their state. (Table 2, $p < 0.001$). Self-awareness is essential in order to seek help (Rickwood et al., 2005). This may prove to have a role in prevention of burnout. It was also found that females have a higher tendency for being burnt out as compared to males (Table 2, $p < 0.001$).

Another interesting finding was the negative relationship between the sense of never ending competition and burnout (Table 2, $p < 0.001$). Medical schools are known for their low acceptance rates and high academic merit. In addition to these, the type A personality of medical students and a continuous need for academic excellence can make medicine competitive. This finding indicates a positive attitude of trainees towards competition which is considered to be conducive for learning. Friendly competition has been linked to stronger motivation and improved performance (Burguillo, 2010); motivation is a key component in preventing burnout (Al-Dubai et al., 2013).

Moreover, social support which was reported as an important preventive factor of burnout (Chou, Li & Hu, 2014), was supported by our study, which found reduced levels of burnout among doctors having a significant other (Table 2, $p = 0.042$). In contrast, accommodation away from home was reported to have a negative association with burnout (Table 2, $p = 0.01$). This can be explained with improved communication, better transport and hostel facilities which may compensate for homesickness. Accommodation at hostels increases social interaction and support offered by peers experiencing similar problems,

which may help in reducing burnout. Previously, lack of peer support has been linked with increased levels of burnout (*Khamisa, Peltzer & Oldenburg, 2013*).

During the study a comparison was drawn between burnout rates of HOs and PGRs. It was observed that more HOs had a “high/ very high burnout” than PGRs (*Table 3*, $p < 0.001$). This was comparable to an Indian study which found high rates of personal burnout among HOs than PGRs (*Ratnakaran, Prabhakaran & Karunakaran, 2016*). The high level of burnout can be attributed to one major common factor: stressful nature of early medical training. HOs work longer hours with lower income than PGRs (*Table 3*, $p < 0.001$). Greater number of working hours lead to poor work-life balance leaving less personal time (*Shanafelt et al., 2015*) and increases patient contact which causes emotional strain for the doctor (*Sablik, Samborska-Sablik & Drożdż, 2013*). As time for recreation is important for regeneration of emotional energy, working hours need to be addressed with a decrease in number of hours to 48 hours as suggested by European Working Time Directive (EWTD). Low income implies more financial strain, an important stressor that affects quality of life, and lack of reward. The resulting fatigue caused by it may also increase stress.

However, HOs were found to be more self-aware, as they had a higher rate of self-reported feeling of burnout than PGRs and still opted not to change their profession showing more resilience. Healthy coping mechanisms are linked to resilience (*Hamill, 2003*) which may be the reason for lower percentage of HOs (*Table 3*, $p < 0.001$) smoking (7.8%) compared to PGRs trainees (17.1%). Smoking, an unhealthy strategy, is a popular choice to cope with stress among Pakistani medical students (*Shaikh et al., 2004*).

Doctors showed a tendency towards doctor centered relationship. This is comparable to a previous research where MBBS students showed a similar attitude towards doctor–patient relationship (*Ahmad et al., 2015*). There is lack of educational interventions like communication skills’ trainings aimed at improving doctor–patient relationship, which may be the reason for similarity of the attitude. Although burnout showed no significant association with total score and caring domain of PPOS, it showed significant association with power sharing domain of PPOS (*Table 2*, $p = 0.083$). Sharing implies equal autonomy of the patient in making decisions, which requires time and effort on part of the physician to patients’ concerns and their choices, resulting in exhaustion and inefficacy. Similarly, lack of control over what one does also showed to have a significant association with high burnout (*Table 2*, $p = 0.024$) in this study and a previous study (*Richter et al., 2014*). Given that sense of autonomy and efficacy are important determinants of intrinsic motivation (*Ryan & Deci, 2000*), it can be inferred that more power sharing results in lack of intrinsic motivation thus leading to burnout. Another study indicated doctor–patient interaction as an important stressor of burnout (*Sablik, Samborska-Sablik & Drożdż, 2013*). It explained that the helpers have to over-extend themselves to the point where they become emotionally saturated which onsets emotional exhaustion. When feeling exhausted, it is difficult to gain a sense of accomplishment, resulting in inefficacy (*Bresó, Salanova & Schaufeli, 2007*) that eventually leads to burnout.

CONCLUSION

Despite high rates of burnout and its impact on patient care, little research and preventive measures have been taken against it. Thus, it is necessary to comprehend as well as establish the causal relation between burnout and doctor-patient relationship. Moreover, to improve care delivery, it is essential to regulate the stressors that decrease the efficacy of doctors as exhibited by this study such as; long working hours, low income, emotional depletion, lack of autonomy with an evident preponderance of organizational over personal factors. However, it does not imply association of burnout with doctor-patient relationship, it only signifies relation to power sharing domain of doctor-patient relationship.

LIMITATIONS

(1) A cross-sectional study design rather than a case-control was used to study association of burnout and doctor-patient relationship. (2) Self-assessment questionnaires were employed to study burnout and doctor-patient relationship. It would have been more reliable to study behaviour of doctors or assess doctor-patient relationship through patient feedback. (3) It is suggested for future research to study other factors that may have an impact on doctor-patient relationship such an examination intervention (*Hur et al., 2014*). (4) A convenience sampling technique was employed but it is suggested for future research to use a random sampling technique.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The authors received no funding for this work.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Waqas Ahmad conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Huma Ashraf conceived and designed the experiments, performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Afnan Talat, Aleena Ahmad Khan, Iqra Zia, Zohak Sarfraz, Hifsa Sajid, Marium Tahir, Usman Sadiq and Hira Imtiaz conceived and designed the experiments, performed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Ammad Anwar Baig conceived and designed the experiments, performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

CMH Lahore Medical and Institute of Dentistry, Ethical Review Committee granted Ethical approval to carry out this study within its facilities (Reference number: 40 ERC/CMHLMC).

Data Availability

The following information was supplied regarding data availability:

Raw data includes: “Raw data.spss” which has forms that were coded and entered in SPSS.v5)and “Table 2 Raw result (regression analysis)” which has both significant and insignificant determinants of burnout

The raw data are provided in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5519#supplemental-information>.

REFERENCES

- Ahmad W, Krupat E, Asma Y, Fatima N-E, Attique R, Mahmood U, Waqas A. 2015.** Attitudes of medical students in Lahore, Pakistan towards the doctor–patient relationship. *PeerJ* 3:e1050 DOI 10.7717/peerj.1050.
- Al-Dubai SAR, Ganasegeran K, Perianayagam W, Rampal KG. 2013.** Emotional burnout, perceived sources of job stress, professional fulfillment, and engagement among medical residents in Malaysia. *The Scientific World Journal* 2013:Article 137620 DOI 10.1155/2013/137620.
- Beckman HB, Markakis KM, Suchman AL, Frankel RM. 1994.** The doctor–patient relationship and malpractice. Lessons from plaintiff depositions. *Archives of Internal Medicine* 154:1365–1370 DOI 10.1001/archinte.154.12.1365.
- Bresó E, Salanova M, Schaufeli WB. 2007.** In search of the third dimension of burnout: efficacy or inefficacy? *Applied Psychology* 56:460–478 DOI 10.1111/j.1464-0597.2007.00290.x.
- Burguillo JC. 2010.** Using game theory and competition-based learning to stimulate student motivation and performance. *Computers and Education* 55:566–575 DOI 10.1016/j.compedu.2010.02.018.
- Chou L-P, Li C-Y, Hu SC. 2014.** Job stress and burnout in hospital employees: comparisons of different medical professions in a regional hospital in Taiwan. *BMJ Open* 4:e004185 DOI 10.1136/bmjopen-2013-004185.
- Constable JF, Russell DW. 1986.** The effect of social support and the work environment upon burnout among nurses. *Journal of Human Stress* 12:20–26 DOI 10.1080/0097840X.1986.9936762.

- Dyrbye LN, Thomas MR, Huntington JL, Lawson KL, Novotny PJ, Sloan JA, Shanafelt TD. 2006. Personal life events and medical student burnout: a multicenter study. *Academic Medicine* 81:374–384 DOI 10.1097/00001888-200604000-00010.
- Dyrbye LN, Thomas MR, Power DV, Durning S, Moutier C, Massie FS, Harper W, Eacker A, Szydlo DW, Sloan JA, Shanafelt TD. 2010. Burnout and serious thoughts of dropping out of medical school: a multi-institutional study. *Academic Medicine* 85:94–102 DOI 10.1097/ACM.0b013e3181c46.
- Dyrbye LN, West CP, Satele D, Boone S, Tan L, Sloan J, Shanafelt TD. 2014. Burnout among US medical students, residents, and early career physicians relative to the general US population. *Academic Medicine* 89:443–451 DOI 10.1097/ACM.000000000000134.
- Guthrie E, Black D, Bagalkote H, Shaw C, Campbell M, Creed F. 1998. Psychological stress and burnout in medical students: a five-year prospective longitudinal study. *Journal of the Royal Society of Medicine* 91:237–243.
- Ha JF, Longnecker NP. 2010. doctor–patient communication: a review. *The Ochsner Journal* 10:38–43 DOI 10.3946/kjme.2014.26.2.99.
- Hamill SK. 2003. Resilience and self-efficacy: the importance of efficacy beliefs and coping mechanisms in resilient adolescents. *Colgate University Journal of the Sciences* 35:115–146.
- Hellin T. 2002. The physician patient relationship: recent developments and changes. *Haemophilia* 8:450–454.
- Hur Y, Kim S, Park JH, Cho A-R, Choi CJ. 2014. Changes in medical students' patient-centeredness attitudes by implementation of clinical performance examination. *Korean Journal of Medical Education* 26:99–106.
- Khamisa N, Peltzer K, Oldenburg B. 2013. Burnout in relation to specific contributing factors and health outcomes among nurses: a systematic review. *International Journal of Environmental Research and Public Health* 10:2214–2240 DOI 10.3390/ijerph10062214.
- Kristensen TS, Borritz M, Villadsen E, Christensen KB. 2005. The Copenhagen burnout inventory: a new tool for the assessment of burnout. *Work and Stress* 19:192–207 DOI 10.1080/02678370500297720.
- Krupat E, Rosenkranz SL, Yeager CM, Barnard K, Putnam SM, Inui TS. 2000. The practice orientations of physicians and patients: the effect of doctor–patient congruence on satisfaction. *Patient Education and Counseling* 39:49–59 DOI 10.1016/S0738-3991(99)00090-7.
- Laschinger HKS, Borgogni L, Consiglio C, Read E. 2015. The effects of authentic leadership, six areas of worklife, and occupational coping self-efficacy on new graduate nurses' burnout and mental health: a cross-sectional study. *International Journal of Nursing Studies* 52:1080–1089 DOI 10.1016/j.ijnurstu.2015.03.002.
- Marjani A, Gharavi AM, Jahanshahi M, Vahidirad A, Alizadeh F. 2008. Stress among medical students of Gorgan (South East of Caspian Sea), Iran. *Kathmandu University Medical Journal* 6:421–425 DOI 10.3126/kumj.v6i3.1726.

- Msaouel P, Keramaris NC, Tasoulis A, Kolokythas D, Syrmos N, Pararas N, Thireos E, Lionis C. 2010. Burnout and training satisfaction of medical residents in Greece: will the European work time directive make a difference? *Human Resources for Health* 8:16 DOI 10.1186/1478-4491-8-16.
- Muzafar Y, Khan HH, Ashraf H, Hussain W, Sajid H, Tahir M, Rehman A, Sohail A, Waqas A, Ahmad W. 2015. Burnout and its associated factors in medical students of Lahore, Pakistan. *Cureus* 7:e390 DOI 10.7759/cureus.390.
- Paro HBMS, Silveira PSP, Perotta B, Gannam S, Enns SC, Giaxa RRB, Bonito RF, Martins MA, Tempiski PZ. 2014. Empathy among medical students: is there a relation with quality of life and burnout? *PLOS ONE* 9:e94133 DOI 10.1371/journal.pone.0094133.
- Ratanawongsa N, Roter D, Beach MC, Laird SL, Larson SM, Carson KA, Cooper LA. 2008. Physician burnout and patient-physician communication during primary care encounters. *Journal of General Internal Medicine* 23:1581–1588 DOI 10.1007/s11606-008-0702-1.
- Ratnakaran B, Prabhakaran A, Karunakaran V. 2016. Prevalence of burnout and its correlates among residents in a tertiary medical center in Kerala, India: a cross-sectional study. *Journal of Postgraduate Medicine* 62:157–161 DOI 10.4103/0022-3859.184274.
- Richter A, Kostova P, Harth V, Wegner R. 2014. Children, care, career—a cross-sectional study on the risk of burnout among German hospital physicians at different career stages. *Journal of Occupational Medicine and Toxicology* 9(1):41 DOI 10.1186/s12995-014-0041-6.
- Rickwood D, Deane FP, Wilson CJ, Ciarrochi JV. 2005. Young people's help-seeking for mental health problems. *Australian e-Journal for the Advancement of Mental Health* 4:218–251 DOI 10.5172/jamh.4.3.218.
- Rizo-Baeza M, Mendiola-Infante SV, Sepehri A, Palazón-Bru A, Gil-Guillén VF, Cortés-Castell E. 2018. Burnout syndrome in nurses working in palliative care units: an analysis of associated factors. *Journal of Nursing Management* 26:19–25 DOI 10.1111/jonm.12506.
- Ryan RM, Deci EL. 2000. Intrinsic and extrinsic motivations: classic definitions and new directions. *Contemporary Educational Psychology* 25:54–67 DOI 10.1006/ceps.1999.1020.
- Sablik Z, Samborska-Sablik A, Drożdż J. 2013. Systematic review/meta-analysis Universality of physicians' burnout syndrome as a result of experiencing difficulty in relationship with patients. *Archives of Medical Science* 3:398–403 DOI 10.5114/aoms.2012.28658.
- Schaufeli WB, Dierendonck D Van, Gorp K Van. 1996. Burnout and reciprocity: towards a dual-level social exchange model. *Work and Stress* 10:225–237 DOI 10.1080/02678379608256802.
- Shaikh BT, Kahloon A, Kazmi M, Khalid H, Nawaz K, Khan NA, Khan S. 2004. Students, stress and coping strategies: a case of Pakistani Medical School. *Education for Health* 17:346–353 DOI 10.1080/13576280400002585.

- Shanafelt TD, Balch CM, Bechamps G, Russell T, Dyrbye L, Satele D, Collicott P, Novotny PJ, Sloan J, Freischlag J. 2010.** Burnout and medical errors among American surgeons. *Annals of Surgery* **251**:995–1000
[DOI 10.1097/SLA.0b013e3181bfdab3](https://doi.org/10.1097/SLA.0b013e3181bfdab3).
- Shanafelt TD, Hasan O, Dyrbye LN, Sinsky C, Satele D, Sloan J, West CP. 2015.** Changes in burnout and satisfaction with work-life balance in physicians and the general US working population between 2011 and 2014. *Mayo Clinic Proceedings* **90**:1600–1613 [DOI 10.1016/j.mayocp.2015.08.023](https://doi.org/10.1016/j.mayocp.2015.08.023).
- Winwood PC, Winefield AH. 2004.** Comparing two measures of burnout among dentists in Australia. *International Journal of Stress Management* **11**:282–289
[DOI 10.1037/1072-5245.11.3.282](https://doi.org/10.1037/1072-5245.11.3.282).