

## IMAGES IN CLINICAL RADIOLOGY

# A Symptomatic Killian-Jamieson Diverticulum Detected by Ultrasonography


M. Navez\*, P. Bosschaert\* and J. C. Degols†

Killian-Jamieson diverticulum (KJD) is a rare entity resulting from the mucosal herniation through a lateral area of weakness known as the Killian-Jamieson space below the cricopharyngeal muscle. Ultrasonographic diagnosis is exceptional. Moreover, symptoms are found in only eleven percent of patients. KJD in its symptomatic form must be correctly identified because its therapeutic management is in some points different of Zenker's diverticulum.


**Keywords:** ultrasonography; hypopharyngeal; lateral cervical; Killian-Jamieson diverticulum; Zenker's diverticulum

A 73-year-old woman was suffering of mild dysphagia with occasional regurgitations and depressible swelling during swallowing on the left side of her neck which appeared two years earlier. Ultrasonographic examination showed a hypoechoic mass surrounded by a digestive wall measuring 45mm located just behind the left thyroid lobe (**Fig. A**). Moving bright

hyperechoic foci caused by air bubbles were clearly observed. Barium swallow pharyngoesophagegraphy highlighted a large protruding sac arising from the left anterolateral pharyngoesophageal junction retaining contrast medium (**Fig. B**). No concomitant



**Figure A:** Ultrasonographic axial section showing KJD behind the thyroid left lobe.



**Figure B:** Radiological right anterolateral view after barium administration.

\* Department of Radiology, Clinique St-Pierre, Ottignies-LLN, Belgium  
 Marie\_navez@hotmail.com, Pierre.bosschaert@skynet.be


† Department of Otorhinolaryngology, Clinique St-Pierre, Ottignies-LLN, Belgium  
 Jean-christophe.degols@cspo.be

functional abnormalities were found. It was diagnosed as a Killian-Jamieson diverticulum (KJD) based on its specific lateral projection (**Fig. C**). Computed tomography on procubitus position with oral contrast ingestion was performed demonstrating the diverticulum environment and the absence of complications such as inflammation or fistula.

### Comment

KJD is rare entity related to the more common Zenker's diverticulum (ZD) with its classic posterior development. It results from the mucosal herniation through a lateral

area of weakness known as the Killian-Jamieson space below the cricopharyngeal muscle [1]. This space was first described by G. Killian in 1907 and confirmed by EB. Jamieson in 1934. Seventy-five percent are unilateral on the left side and twenty-five percent are bilateral. Elderly patients are more often affected by this condition. This diverticulum can mimic some thyroid nodules because it is located in the vicinity of the thyroid gland. Symptoms are found in eleven percent of patients. There is no case described of aspiration pneumonia provoked by this abnormality (Rubesin 2001). However, this complication is reported in twelve percent of ZDs. The singular anatomical disposition of the two types of diverticula presumably accounts for the explanation of this difference. Symptomatic KJD may be surgically treated. Although cricopharyngeal myotomy in addition to diverticulotomy is a mandatory component of the ZD treatment, cricopharyngeal myotomy is not necessary for KJD treatment because of the unproved dysfunction of this muscle. We must note here the importance with respect to the directly adjacent recurrent laryngeal nerve entry point into the larynx during the surgical procedure. Endoscopic treatment of this kind of diverticulum has not been found in literature to date and seems inadvisable. Diverticulopexy was only proposed in one case. In conclusion, KJD in its symptomatic form must be correctly identified because its therapeutic management is in some points different of ZD.



**Figure C:** Anatomical drawing.

### Competing Interests

The authors declare that they have no competing interests.

### Reference

1. Ekberg, O and Nylander, G. Lateral Diverticula from the Pharyngoesophageal Junction Area. *Radiology*. 1983; 146: 117–22. DOI: <http://dx.doi.org/10.1148/radiology.146.1.6401363>. PMID: 6401363.

**How to cite this article:** Navez, M, Bosschaert, P and Degols, J C 2015 A Symptomatic Killian-Jamieson Diverticulum Detected by Ultrasonography. *Journal of the Belgian Society of Radiology*, 99(2), pp. 74–75, DOI: <http://dx.doi.org/10.5334/jbr-btr.877>

**Published:** 30 December 2015

**Copyright:** © 2015 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported License (CC-BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/3.0/>.

**] *Journal of the Belgian Society of Radiology* is a peer-reviewed open access journal published by Ubiquity Press.**

**OPEN ACCESS** 