

Research Article

Nest description, new parasitoid associations and geographical range of *Trypoxyylon (Trypoxyylon) florale* Richards (Hymenoptera: Crabronidae)

Descripción de nido, nuevas asociaciones con parásitóides y rango geográfico de *Trypoxyylon (Trypoxyylon) florale* Richards (Hymenoptera: Crabronidae)

Nelson Wanderley Perioto¹ and Rogéria Inês Rosa Lara¹

¹ Laboratório de Sistemática e Bioecologia de Parasitoídeos e Predadores, Ribeirão Preto-SP, Brasil. E-mail: nperioto@biologico.br

ZooBank: urn:lsid:zoobank.org:pub:DD15AC10-C818-48F7-B9ED-CD02898AC332

Abstract. In this work the *Trypoxyylon (Trypoxyylon) florale* Richards, 1934 (Hymenoptera: Crabronidae) nest is described for the first time and the geographic ranges of *Trypoxyylon (Trypoxyylon) florale* and *Ceyxia latilabra* Andrade & Tavares, 2009 (Hymenoptera: Chalcididae) are extended about 500 km NE from the type locality and 2,800 km SE from Manaus (AM) and about 2,200 km S from Lago Verde (MA), respectively, the previously known records. Additionally are provided news data of associations between *Trypoxyylon (Trypoxyylon) florale* and *Ceyxia latilabra* and *Caenochrysis armata* (Mocsáry, 1889) (Hymenoptera: Chrysidae) based on the examination of exemplars collected at “Sertão do Cacau”, in São Sebastião municipality, São Paulo State, Brazil.

Key words: *Caenochrysis*, *Ceyxia*, Neotropical, Nesting, Parasitoids.

Resumen. En este trabajo se describe por primera vez el nido de *Trypoxyylon (Trypoxyylon) florale* Richards, 1934 (Hymenoptera: Crabronidae) y se amplían los rangos geográficos de *Trypoxyylon (Trypoxyylon) florale* y de *Ceyxia latilabra* Andrade & Tavares, 2009 (Hymenoptera: Chalcididae) a cerca de 500 km NE desde la localidad tipo, y cerca de 2.800 km SE desde Manaus (AM) y cerca de 2.200 km S desde Lago Verde (MA), respectivamente, los registros previamente conocidos. Además, se proporcionan nuevos datos de asociaciones entre *Trypoxyylon (Trypoxyylon) florale*, *Ceyxia latilabra* y *Caenochrysis armata* (Mocsáry, 1889) (Hymenoptera: Chrysidae) basadas en el examen de ejemplares recogidos en “Sertão do Cacau”, en el municipio de São Sebastião, São Paulo Estado, Brasil.

Palabras clave: *Caenochrysis*, *Ceyxia*, Neotropical, Ninho, Parasitoides.

Introduction

Trypoxyylon Latreille, 1796 is a diverse and widespread genus of solitary wasps of the family Crabronidae (Hymenoptera). Among Crabronidae, *Trypoxyylon* is the second largest genus in species number (Pulawski 2016) and comprises 630 species of spider-hunting wasps (Hanson and Menke 2006). These wasps build predominantly mud nests (Amarante 2005, 2006) on vegetation, crevices, and buildings or in pre-existing cavities and another wasps abandoned nests (Richards 1934; Hanson and Menke 2006). The genus is usually split into *Trypoxyylon* and *Trypargilum* subgenera; the first widespread and, the second,

Received 22 June 2018 / Accepted 1 August 2018 / Published online 17 August 2018

Responsible Editor: José Mondaca E.



Este es un artículo de acceso abierto distribuido bajo los términos de la licencia Creative Commons License (CC BY NC 4.0)

restricted to the New World (Richards 1934; Hanson and Menke 2006), with circa of 70 and 90 species in Neotropical Region, respectively (Amarante 2002, 2005). There are 37 species of *T. (Trypoxylon)* and same number of *T. (Trypargilum)* recorded in 23 states and Federal District of Brazil (Amarante 2002).

Ceyxia Girault, 1911 (Hymenoptera: Chalcididae), with twenty-seven valid species, has mainly Neotropical distribution, occurring from southern United States to southern Brazil and northern Argentina (Andrade and Tavares 2009). Meanwhile in Brazil there are 24 recorded species, at 18 states and Federal District (Andrade and Tavares 2009).

The most of 50 valid species of *Caenochrysis* Kimsey & Bohart, 1981 (Hymenoptera: Chrysidae) (Hymenoptera Online 2016) also has Neotropical distribution with six species occurring in North America (Kimsey 2006a) and 25 recorded species in Brazil (Lucena and Zanella 2015).

Several genera of parasitoids of three orders and different families of insects are associated with species of *Trypoxylon*; among the Hymenoptera, the Chalcidoidea: *Ceyxia* (Chalcididae), *Melittobia* Westwood, 1848 (Eulophidae), *Aximopsis* Ashmead, 1904 and *Eurytoma* Illiger, 1807 (Eurytomidae), *Monodontomerus* Westwood, 1833 and *Torymus* Dalman, 1820 (Torymidae), *Dibrachys* Förster, 1856 and *Epistenia* Westwood, 1832 (Pteromalidae) (Molumby 1996; Fox *et al.* 2012; Yu *et al.* 2012; Coudrain *et al.* 2013; Noyes 2016); the Ichneumonoidea: *Bathythrix* Förster, 1869, *Dolichomitus* Smith, 1877, *Ephialtes* Gravenhorst, 1829, *Hoplocryptus* Thomson, 1873, *Nematopodius* Gravenhorst, 1829, *Perithous* Holmgren, 1859, *Poemenia* Holmgren, 1859, *Townesia* Ozols, 1962, *Eurycryptus* Cameron, 1901, *Messataporus* Cushman 1929 and *Photocryptus* Viereck, 1913 (Ichneumonidae) (Aguiar and Santos 2009; Yu *et al.* 2012); the Evanioidea: *Gasteruption* Latreille, 1796 (Gasteruptiidae) (Pagliano and Scaramozzino 2000; Coudrain *et al.* 2013) and the Chrysididae: *Caenochrysis*, *Chrysis* Linnaeus, 1761 and *Trichrysis* Lichtenstein, 1876 (Chrysididae) (Kimsey 2006a; Coudrain *et al.* 2013; Bogusch *et al.* 2015); among the Diptera: *Antrax* Scopoli, 1763 (Bombyliidae) and an unidentified species of Sarcophagidae (Molumby 1996; Santoni *et al.* 2009; Fox *et al.* 2012) and among the Coleoptera: *Megatoma* Herbst, 1792 and *Trogoderma* Dejean, 1821 (Dermestidae) and *Trichodes* Herbst, 1792 (Cleridae) (Coudrain *et al.* 2013).

The objectives of this paper is to describe the *Trypoxylon (Trypoxylon) florale* Richards, 1934 nest, reporting its new associations with hymenopterous parasitoids and providing new distribution records to *T. (Trypoxylon) florale* and to *Ceyxia latilabra* Andrade & Tavares, 2009.

Material and Methods

A nest of *T. (Trypoxylon) florale*, a species of the *tridentata* group, was collected in January 2016 from a wall of a vacation home ($23^{\circ}46'03.53''S / 45^{\circ}37'59.48''W$) located in the “Sertão do Cacau”, in São Sebastião, State of São Paulo, Brazil (Fig. 1). The locality has an Af climate (Koeppen system), with an average temperature of $23.3^{\circ}C$ and 1,638 mm average annual rainfall (Alvares *et al.* 2013; ClimateData.Org 2016). The nest was carefully removed from the wall with the aid of a pocket knife and transported to the Laboratório de Sistemática e Bioecologia de Parasitoides e Predadores (LRRP) of the Instituto Biológico, Ribeirão Preto, São Paulo State, Brazil and kept in a clear plastic cage (12 cm in diameter x 7 cm height), at room temperature until the emergence of the adults. The collections were done under SISBIO license#16473-1.

The color images were obtained with a digital camera (DFC 295, Leica, Germany) attached to a stereomicroscope (M205C APO, Leica, Singapore). Serial images from different layers were combined with Helicon Focus software (version 5.3) and figures were prepared using Adobe Photoshop software (version 11.0).

A female and a male specimens of *T. (Trypoxylon) florale*, a female specimen of *C. latilabra* and a female specimen of *Caenochrysis armata* (Mocsáry, 1889) were deposited in




Figure 1. Map of Brazil showing State of São Paulo; the black dot indicate the place where the nest of *Trypoxylon (Trypoxylon) florale* Richards, 1934 was collected.

the Coleção Entomológica do Laboratório de Sistemática e Bioecologia de Parasitoides e Predadores (LRRP) do Instituto Biológico (Ribeirão Preto, SP, Brazil, Dr. N. W. Perioto, curator); a female specimen of *C. latilabra* in the Coleção Entomológica do Departamento de Ciências Biológicas, Universidade Federal do Espírito Santo (UFES) (Vitória, ES, Brazil, Dr. Marcelo Teixeira Tavares, curator) and a female specimen of *T. (Trypoxylon) florale* in the Coleção Entomológica Pe. Jesus Santiago Moure (DZUP), Universidade Federal do Paraná (Curitiba, PR, Brazil, Dr. Gabriel Augusto Rodrigues de Melo, curator).

Results and Discussion

The nest is composed of mud, with about 2 cm in diameter and with 13-15 cells (0.8 cm long and 0.3 cm wide) (Fig. 2). From the nest emerged two females and a male specimens of *T. (Trypoxylon) florale* (Fig. 3), two females specimens of *C. latilabra* (Fig. 4) and a female specimen of *C. armata* (Fig. 5).

Almost nothing is known about the biology and habits of *T. (Trypoxylon) florale*. The species was described by Richards (1934) and until this article was known from a single specimen, the female holotype, collected in Blumenau, Santa Catarina State, in 1885 and deposited in the Entomological Collection of the Natural History Museum of Vienna (NHMW) (Richards 1934). Thus this is the first documented record of *T. (Trypoxylon) florale* for São Paulo State and this finding extends its distribution range about 500 km NE from the type locality.


The association between *Ceyxia* and *Trypoxylon* species is well documented in the literature: *C. flaviscapus* Girault, 1911 is associated to *T. nitidum* F. Smith, 1856; *C. laticlipeata* Andrade & Tavares, 2009 to *T. fabricator* F. Smith, 1873; *C. perparva* Andrade & Tavares, 2009 to *Trypoxylon maidli bodkini* Richards, 1934; *C. villosa* (Olivier, 1790) to *T. albitarse* Fabricius, 1804 and *T. mexicanum* Saussure, 1867 (Burks 1960; Boucek 1992; Andrade and Tavares 2009). For the first time *C. latilabra* is associated as parasitoid of *T. (Trypoxylon) florale*. It has known distribution for Brazil (Amazonas and Maranhão States) (Andrade and Tavares

2009). Consequently, this finding extends the distribution range about 2,800 km SE from Manaus (AM) and about 2,200 km S from Lago Verde (MA), the previously known records.

There are references to species of *Brachymeria* Weswood, 1829 (Hymenoptera: Chalcididae) associated species of *Trypoxyton* (Peruquetti 2001; González and Matthews 2005; Yu *et al.* 2012; Noyes 2016). It is plausible that such organisms may be, in fact, species of *Ceyxia* given that Andrade and Tavares (2009), when studying the Neotropical species of Brachymeriini, grouped in *Ceyxia* those that show the presence of escrobo-malar carina, jaws typically elongated and tapered at the apex and are associated with wasps that inhabit nests of mud.

Caenochrysis armata is associated with *T. (Trypoxyton) florale* for the first time, but the exact association between them is not entirely clear. In general, Chrysidinae has been found as ecto or cleptoparasitoides (or both) of solitary Hymenoptera who build nests (Kimsey 2006b). Two species of *Caenochrysis* have been reported in association with species of *Trypoxyton*: *C. devensor* (Bohart, 1966) was reared from nests of *T. (Trypargilum) tridentatum* Packard, 1867 (Parker and Bohart 1966) and *C. doriae* (Gribodo, 1874) from nests of *T. (Trypoxyton) bidentatum* W. Fox, 1891 (Parker and Bohart 1966), *T. (Trypoxyton) frigidum* F. Smith, 1856; *T. (Trypargilum) collinum rubrocinctum* (Packard, 1867) (Krombein 1967) and *T. (Trypargilum) tridentatum* (Bohart and Kimsey 1982).

Finally, this study is searching increased the knowledge about the geographical distribution and associations among three species of Hymenoptera. More studies are essential because it's great our ignorance about the geographical distribution of the vast majority of the species of Hymenoptera and such scarcity makes activities such as collecting of specimens in the field and publication of new records essential to the establishment of bases for future biogeographical and ecological studies.



Figures 2-5. 2. Nest of *Trypoxyton (Trypoxyton) florale* Richards, 1934. (A) Ventral view. (B) Dorsal view. Scale bar: 1 cm. 3. *Trypoxyton (Trypoxyton) florale* Richards, 1934, female. Habitus, lateral view. 4. *Ceyxia latilabra* Andrade & Tavares, 2009, female. Habitus, lateral view. 5. *Caenochrysis armata* (Mocsáry, 1889), female. Habitus, lateral view.

Acknowledgments

We thank Dr. Marcelo Teixeira Tavares (Universidade Federal do Espírito Santo), MSc. Daercio Adam de Araújo Lucena (Universidade de São Paulo, campus Ribeirão Preto) and MSc. Bruno Bueno da Rosa (Universidade Federal do Paraná) for identification of the wasps. We also thank the Instituto Nacional de Ciência e Tecnologia dos Hymenoptera Parasitoides (CNPq/Fapesp/Capes) for the financial support.

Literature Cited

- Aguiar, A.P. and Santos, B.F. (2009)** Biological synopsis of *Photocryptus* Viereck (Hymenoptera, Ichneumonidae, Cryptinae), with eight new host records and fourteen new geographic records. *Zootaxa*, 2148: 49-45.
- Alvares, C.A., Stape, J.L., Sentelhas, P.C., de Moraes Gonçalves, J.L. and Sparovek, G. (2013)** Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift*, 22(6): 711-728.
- Amarante, S.T.P. (2002)** A synonymic catalog of the Neotropical Crabronidae and Sphecidae (Hymenoptera: Apoidea). *Arquivos de Zoologia*, 37(1): 1-139.
- Amarante, S.T.P. (2005)** Addendum and corrections to a synonymic catalog of the Neotropical Crabronidae and Sphecidae. *Papéis Avulsos de Zoologia*, 45: 1-18.
- Amarante, S.T.P. (2006)** Familia Crabronidae. In: Fernandez, F. and Sharkey, M.J., eds. *Introducción a los Hymenoptera de la Región Neotropical*. Bogotá: Sociedad Colombiana de Entomología y Universidad Nacional de Colombia, pp. 457-469.
- Andrade, T.V. and Tavare, M.T. (2009)** Revisão de *Ceyxia* Girault, stat. rev. (Hymenoptera, Chalcididae, Brachymeriini). *Revista Brasileira de Entomologia*, 53(4): 511-548.
- Bogusch, P., Astapenkova, Á.A. and Heneberg, P. (2015)** Larvae and nests of six aculeate Hymenoptera (Hymenoptera: Aculeata) nesting in reed galls induced by *Lipara* spp. (Diptera: Chloropidae) with a review of species recorded. *PLoS ONE*, 10(6): e0130802. Available from: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0130802>
- Bohart, R.M. and Kimsey, L.S. (1982)** A synopsis of the Chrysidiidae in America north of Mexico. *Memoirs of the American Entomological Institute*, 33: 1-266.
- Boucek, Z. (1992)** The New World genera of Chalcididae. *Memoirs of the American Entomological Institute*, 53: 49-118.
- Burks, B.D. (1960)** A revision of the genus *Brachymeria* Westwood in America north of Mexico (Hymenoptera: Chalcididae). *Transactions of the American Entomological Society*, 86: 225-273.
- ClimateData.Org. (2016)** Clima: São Sebastião. Viewed 20 July 2016. Available from: <http://pt.climate-data.org/location/34837/>
- Coudrain, V., Herzog, F. and Entling, M.H (2013)** Effects of habitat fragmentation on abundance, larval food and parasitism of a spider-hunting wasp. *PLoS One*, 8(3): e59286. Available from: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0059286>
- Fox, E.G.P., Solis, D.R., Rossi, M.L., Feitosa, R.M. and Brescovit, A.D. (2012)** The larval morphology and nest habits of *Trypoxylon (Trypargilum) rogenhoferi* Kohl 1884 (Insecta: Hymenoptera: Crabronidae). *Zootaxa*, 3251: 47-56.
- González, J.M. and Matthews, R.W. (2005)** An annotated bibliography of *Melittobia* (Hymenoptera: Eulophidae). *Caribbean Journal of Science*, 8: 1-48.
- Hanson, P.E. and Menke, A.S. (2006)** Las avispas apoideas: Ampulicidae, Sphecidae, Crabronidae. In: Hanson, P.E. and Gauld, I.D., eds. *Hymenoptera de la Región Neotropical*. Issue 77 of Memoirs of the American Entomological Institute. Gainesville: The American Entomological Institute, pp. 694-733.

- Hymenoptera Online - HOL (2016)** Viewed. 20 July 2016. Available from: <http://hol.osu.edu/search.html?limit=50&name=Caenochrysis>
- Kimsey, L.S. (2006a)** *California cuckoo wasps in the family Chrysidae (Hymenoptera)*. Issue 125 of University of California Press. Berkeley: University of California Press. 311 p. Available from: <http://escholarship.org/uc/item/3j6466jw#page-5>
- Kimsey, L.S. (2006b)** Familia Chrysidae. In: Hanson, P.E. and Gauld, I.D., eds. *Hymenoptera de la Región Neotropical*. Issue 77 of Memoirs of the American Entomological Institute. Gainesville: The American Entomological Institute, pp. 544-550.
- Krombein, K.V. (1967)** *Wasps and bees: life histories, nests, and associates*. Washington DC: Smithsonian Institute Press. 570 pp.
- Linsenmaier, W. (1984)** Das subgenus *Trichrysis* Lichtenstein in Nord und Südamerika. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 57: 194-224.
- Lucena, D.A.A. and Zanella, F.C.V. (2015)** Viewed 20 July 2016. Chrysidae Linnaeus, 1761. In: *Catálogo Taxonômico da Fauna do Brasil*. Available from: http://fauna.jbrj.gov.br/fauna/listaBrasil/ConsultaPublicaUC/BemVindoConsultaPublicaConsultar.do?iinvalidatePageControlCounter=14&lingua=&jsonRank=&rankTaxon=32767&nomeTaxon=&autor=&nomeVernaculo=&nomeCompleto=Chrysidae+Linnaeus%2C+1761&formaVida=QUALQUER&substrato=QUALQUER&oocorrencia=OCORRE®iao=QUALQUER&estado=QUALQUER&endemismo=TODOS&origem=TODOS&mostrarAte=ESPECIE&opcoesBusca=NOME_ACEITO
- Molumby, A. (1996)** Why make daughters larger? Maternal sex-allocation and sex-dependent selection for body size in a mass-provisioning wasp, *Trypoxyton politum*. *Behavioral Ecology*, 8(3): 279-287.
- Noyes, J.S. (2016)** Viewed 20 July 2016. *Universal Chalcidoidea Database*. World Wide Web electronic publication. Available from: <http://www.nhm.ac.uk/chalcidooids>
- Pagliano, G. and Scaramozzino, P.L. (2000)** Gasteruptiidae italiani (Hymenoptera, Evanioidea). *Bollettino Museo Regionale di Scienze Naturali Torino*, 17(1): 5-38.
- Parker, F.D. and Bohart, R.M. (1966)** Host-parasite association in some twig-nesting Hymenoptera from western North America. *Pan-Pacific Entomologist*, 42: 91-98.
- Peruquetti, R.C. (2001)** Comportamento de oviposição e tempo desenvolvimento de *Brachymeria villosa* (Oliver) (Hymenoptera, Chalcididae). *Revista Brasileira de Zoologia*, 18(3): 695-697.
- Pulawski, W. (2016)** Viewed 20 July 2016. Catalog of Sphecidae: number of species. Available from: http://researcharchive.calacademy.org/research/entomology/entomology_resources/hymenoptera/sphecidae/number_of_species.pdf
- Richards, O.W. (1934)** The American species of the genus *Trypoxyton* (Hymenopt., Sphecoidea). *Transactions of the Royal Entomological Society of London*, 82: 173-962.
- Santoni, M.M., Brescovit, A.D. and del Lama, M.A. (2009)** Ocupação diferencial do habitat por vespas do gênero *Trypoxyton* (*Trypargilum*) Latreille (Hymenoptera, Crabronidae). *Revista Brasileira de Entomologia*, 53(1): 107-114. Available from: <http://dx.doi.org/10.1590/S0085-56262009000100024>
- Yu, D.S., van Achterberg, C. and Horstmann, K. (2012)** Viewed 20 July 2016. *Taxapad Ichneumonoidea*. Available from: <http://www.taxapad.com>