

TECHNOLOGICAL AND ECONOMIC DEVELOPMENT OF ECONOMY

ISSN 2029-4913 / eISSN 2029-4921

2016 Volume 22(3): 453–469
doi:10.3846/20294913.2016.1171262

Intuitionistic fuzzy theory and its application in economy, technology and management

MULTI-CRITERIA DECISION-MAKING METHOD BASED ON INTUITIONISTIC TRAPEZOIDAL FUZZY PRIORITISED OWA OPERATOR

Peide LIU^a, Ye LI^{b,c}, Jurgita ANTUCHEVIČIENĖ^d^a*School of Management Science and Engineering, Shandong University of Finance and Economics,
Jinan Shandong 250014, China*^b*School of Economic and Management, Beijing Jiaotong University, Beijing 100044, China*^c*Beijing Engineering Research Center of Green Building Technology, Beijing 100070, China*^d*Department of Construction Technology and Management,
Vilnius Gediminas Technical University, LT-10223 Vilnius, Lithuania*

Received 23 June 2015; accepted 8 December 2015

Abstract. In the real decision-making, there are many multiple attribute decision-making (MADM) problems, in which there exists the prioritised relationship among decision-making attributes. In this paper, with respect to the prioritised multi-criteria decision-making problems under intuitionistic trapezoidal fuzzy information, a new decision-making method on the basis of the intuitionistic trapezoidal fuzzy prioritised ordered weighted aggregation operator has been proposed. Firstly, the definitions, operational rules and characteristics of intuitionistic trapezoidal fuzzy numbers and POWA operator have been introduced. Then, intuitionistic trapezoidal fuzzy prioritised ordered weighted aggregation (ITFPOWA) operator has been defined as well as the computational method of associated weight, and some properties have been studied and proved. Furthermore, based on the ITFPOWA operator, an approach to the multi-criteria decision-making with intuitionistic trapezoidal fuzzy numbers has been established. Finally, an illustrative example has been given to prove the evaluation procedures of the developed approach and to demonstrate its practicality and validity.

Keywords: MADM, prioritised, intuitionistic trapezoidal fuzzy numbers, POWA, ITFPOWA.

JEL Classification: C44, C60.

Introduction

There are many multi-attribute decision-making (MADM) problems, which received considerable attention in the past few years. Since Zadeh (1965) proposed the theory of fuzzy set (FS), the research on fuzzy MADM (FMADM) problems has become a hot focus

Corresponding author Jurgita Antuchevičienė
E-mail: jurgita.antucheviciene@vgtu.lt

(Merigó *et al.* 2015). Based on the fuzzy set, Atanassov (1986) presented the intuitionistic fuzzy set (IFS) by adding a new non-membership function, and then Atanassov and Gargov (1989), Atanassov (1994) proposed the interval-valued intuitionistic fuzzy set (IVIFS), which is a generalisation of the FS and IFS. The basic feature of the IVIFS is that the values of its membership function and non-membership function take the form of interval numbers rather than crisp numbers. On the basis of the presented theory, a great amount of extensions have been made (Yu, Shi 2015). Shu *et al.* (2006) presented the definition of intuitionistic triangular fuzzy number, and constructed an algorithm of the intuitionistic fuzzy fault-tree analysis. Zhang and Liu (2010) defined the triangular intuitionistic fuzzy numbers by extending the membership degree and the non-membership degree to the triangular fuzzy numbers, and proposed the weighted arithmetic averaging operator and the weighted geometric average operator. Furthermore, based on these operators, a method of multiple attribute group decision-making (MAGDM) with triangular intuitionistic fuzzy information has been established. New arithmetic operations and logic operators for triangular intuitionistic fuzzy numbers and a new method based on evidential reasoning were presented by Wang *et al.* (2013a, 2013b). The hesitant FMADM problem was investigated and some prioritised aggregation operators for aggregating hesitant fuzzy information were proposed, namely, hesitant fuzzy prioritised weighted average (HFPWA) operator and hesitant fuzzy prioritised weighted geometric (HFPWG) operator (Wei 2012). Some prioritised aggregation operators for aggregating triangular fuzzy information have been developed by Zhao *et al.* (2013). Fuzzy number intuitionistic fuzzy prioritised weighted average (FNIFPWA) operator and fuzzy number intuitionistic fuzzy prioritised weighted geometric (FNIFPWG) operator were proposed by Lin *et al.* (2013). The method with intuitionistic interval fuzzy information was applied to group decision-making (Wang *et al.* 2014).

At present, there are some studies on the intuitionistic trapezoidal fuzzy numbers mainly including the following: Wang (2008) proposed the conception of the intuitionistic trapezoidal fuzzy number and interval intuitionistic trapezoidal fuzzy number. Wang and Zhang (2008) proposed a programming method for the MADM problems with intuitionistic trapezoidal fuzzy number and incomplete weight information. Wan and Dong (2010) defined a new ranking method by using the coordinates of gravity centre about intuitionistic trapezoidal fuzzy number, and proposed the ordered weighted aggregation operator and hybrid aggregation operator to solve the MADM problems with intuitionistic trapezoidal fuzzy numbers. Wei (2010) proposed the intuitionistic trapezoidal fuzzy ordered weighted averaging (ITFOWA) operator and intuitionistic trapezoidal fuzzy hybrid aggregation (ITFHA) operator and pointed out a method with respect to MAGDM problems, where the attribute values are intuitionistic trapezoidal fuzzy information. The weighted geometric aggregation operator of trapezoidal fuzzy numbers for group decision-making was presented by Liu and Jin (2012a). Some new aggregation operators including interval intuitionistic trapezoidal fuzzy ordered weighted geometric (IITFOWG) operator and interval intuitionistic trapezoidal fuzzy hybrid geometric (IITFHG) operator were proposed, and some desirable properties of these operators were studied by Wei *et al.* (2012). Zhang *et al.* (2013) proposed a grey relational projection method for the MADM problems with the intuitionistic trapezoidal fuzzy number and the unknown weight information. Liu and Yu

(2013) proposed the method of calculating density weighted vector and developed some density aggregation operators based on interval numbers and intuitionistic trapezoidal fuzzy numbers. Then, a multiple attribute decision-making method for the MADM problems with the intuitionistic trapezoidal fuzzy numbers was presented. Liu, D. P. and Liu, Y. (2014) proposed an intuitionistic trapezoidal fuzzy power generalised weighted average (ITFPGWA) operator, and propounded an approach to deal with group decision-making problems under intuitionistic trapezoidal fuzzy information based on the ITFPGWA operator. Chen and Liu (2014) proposed the intuitionistic trapezoidal fuzzy generalised Heronian OWA operator, and applied it to the MADM problems where the evaluation of information is depicted by intuitionistic trapezoidal fuzzy information. Heronian mean operators were applied for group decision-making (Liu *et al.* 2014). Uncertain linguistic variables and uncertain linguistic operators were analysed (Liu, Jin 2012b, 2012c; Liu, Yu 2014; Liu, Wang 2014).

The above researches were mainly on the MADM problems, in which attributes have been independent of each other and had no precedence relationship. However, in real decision-making, there exists a kind of MADM problems with the precedence relationship between the attributes. Yager (2004) studied this kind of the decision-making problems and emphasised that the importance of attributes with a lower priority has been determined on the basis of the extent, to which alternatives should satisfy the attributes with higher priority. Then, Yager (2008) proposed a prioritised aggregated operator and further constructed the prior “anding” operator and the prior “oring” operator in 2008. Based on prioritised aggregated operator, Yager (2009) proposed the prioritised ordered weighted average (OWA) operator for the prioritised decision-making problems, in which there exists a relationship between the criteria. Zeng *et al.* (2014) applied the weighted average operator to business decision-making. Yan *et al.* (2011) used the Hamacher parameterised t-norms to induce the priority weight for each priority level and proposed a benchmark-based approach to induce the priority weight for each priority level. Then, target-oriented decision analysis has been utilised to obtain the benchmark achievement for fuzzy requirements. Some of Hamacher aggregation operators were applied to group decision-making (Liu 2014). Some new aggregation operators based on the Choquet integral and Einstein operations and considering the interactions phenomena among the decision-making criteria or their ordered positions were proposed by Xu *et al.* (2014). The intuitionistic fuzzy group decision-making problem, in which all the experts use the intuitionistic fuzzy preference relations (IFPRs) to express their preferences was analysed by Liao *et al.* (2015).

Xu *et al.* (2011) defined intuitionistic fuzzy prioritised ordered weighted average (IFPOWA) operator and established the corresponding method on the basis of IFPOWA operator. Guo *et al.* (2011) developed the prioritised ordered weighted C-OWA (POWC-OWA) operator and prioritised ordered weighted C-OWG (POWC-OWG) operator, and analysed some of their characteristics. Based on POWC-OWA operator, a new method was constructed for solving the MADM problems, in which the attribute values are interval numbers.

Based on the aforementioned analysis, aiming at the MADM problems, in which there exists a priority relationship between the criteria and the criteria values are intuitionistic

trapezoidal fuzzy numbers, we integrate intuitionistic trapezoidal fuzzy numbers and prioritised aggregated operator, and propose the intuitionistic trapezoidal fuzzy prioritised ordered weighted average (ITFPOWA) operator. Then, we propose an MADM method on the basis of the ITFPOWA operator. In the end, an illustrative example has been given to prove the effectiveness and feasibility of the method.

In order to do so, the remainder of this paper is organised as follows: in Section 1, we have briefly reviewed some basic concepts, operational rules, comparison method and the expected value of the intuitionistic trapezoidal fuzzy numbers, and introduced the definition of the POWA operator and the relevant weights. In Section 2, we have proposed an intuitionistic trapezoidal fuzzy prioritised ordered weighted aggregation (ITFPOWA) operator, and discussed some properties. Section 3 gives a method for the multi-criteria decision-making with intuitionistic trapezoidal fuzzy numbers based on the ITFPOWA operator. In Section 4, we have given an example to show the decision-making steps. The last Section ends this paper with some conclusions.

1. Preliminaries

1.1. Intuitionistic trapezoidal fuzzy numbers

Definition 1. Let \tilde{a} be an intuitionistic trapezoidal fuzzy number, if its membership function is defined as (Wang, Zhang 2008):

$$u_{\tilde{a}(x)} = \begin{cases} \frac{x-a}{b-a} u_{\tilde{a}}, & a \leq x < b; \\ u_{\tilde{a}}, & b \leq x \leq c; \\ \frac{d-x}{d-c} u_{\tilde{a}}, & c < x \leq d; \\ 0 & \end{cases} \tag{1}$$

and its non-membership function is defined as

$$v_{\tilde{a}(x)} = \begin{cases} \frac{b-x+v_{\tilde{a}}(x-a_1)}{b-a_1}, & a_1 \leq x < b; \\ v_{\tilde{a}}, & b \leq x \leq c; \\ \frac{x-\epsilon-v_{\tilde{a}}(d_1-x)}{d_1-c}, & c < x \leq d_1; \\ 0 & \end{cases} \tag{2}$$

where $0 \leq \mu_{\tilde{a}} \leq 1, 0 \leq v_{\tilde{a}} \leq 1$ and $0 \leq \mu_{\tilde{a}} + v_{\tilde{a}} \leq 1; a, a_1, b, c, d, d_1 \in R$. The intuitionistic trapezoidal fuzzy number is denoted as $\tilde{a} = \langle ([a, b, c, d]; \mu_{\tilde{a}}), ([a_1, b, c, d_1]; v_{\tilde{a}}) \rangle$. Comparing with trapezoidal fuzzy numbers, the intuitionistic trapezoidal fuzzy numbers have another parameter: non-membership function, which is used to depict the degree of which the decision makers think that the element does not belong to $[a_1, b, c, d_1]$. When $\mu_{\tilde{a}} = 1, v_{\tilde{a}} = 0$,

\tilde{a} is reduced to a trapezoidal fuzzy number. When $b = c$, the intuitionistic trapezoidal fuzzy numbers become intuitionistic triangular fuzzy numbers. Generally, there is $[a, b, c, d] = [a_1, b, c, d_1]$ in intuitionistic trapezoidal fuzzy number \tilde{a} , and then we can express it as $\tilde{a} = \langle [a, b, c, d]; \mu_{\tilde{a}}, \nu_{\tilde{a}} \rangle$. $\pi_{\tilde{a}} = 1 - u_{\tilde{a}} - v_{\tilde{a}}$ indicates the hesitated degree of \tilde{a} . The smaller $\pi_{\tilde{a}}$ is, the more certain the fuzzy number is. If $d \geq c \geq b \geq a \geq 0$, we can call \tilde{a} as positive intuitionistic trapezoidal fuzzy numbers.

Comparing with the intuitionistic fuzzy numbers, intuitionistic trapezoidal fuzzy numbers added to trapezoidal fuzzy numbers $[a, b, c, d]$ can express different dimensional decision-making information more exactly.

Definition 2 (Wang, Zhang 2008). Let $\tilde{a}_1 = \langle [a_1, b_1, c_1, d_1]; u_{\tilde{a}_1}, v_{\tilde{a}_1} \rangle$ and $\tilde{a}_2 = \langle [a_2, b_2, c_2, d_2]; u_{\tilde{a}_2}, v_{\tilde{a}_2} \rangle$ be two positive intuitionistic trapezoidal fuzzy numbers, $\lambda \geq 0$, then the operational laws between \tilde{a}_1 and \tilde{a}_2 can be defined as follow:

$$\tilde{a}_1 + \tilde{a}_2 = \langle [a_1 + a_2, b_1 + b_2, c_1 + c_2, d_1 + d_2]; u_{\tilde{a}_1} + u_{\tilde{a}_2} - u_{\tilde{a}_1} u_{\tilde{a}_2}, v_{\tilde{a}_1} v_{\tilde{a}_2} \rangle; \tag{3}$$

$$\tilde{a}_1 \tilde{a}_2 = \langle [a_1 a_2, b_1 b_2, c_1 c_2, d_1 d_2]; u_{\tilde{a}_1} u_{\tilde{a}_2}, v_{\tilde{a}_1} v_{\tilde{a}_2} - v_{\tilde{a}_1} v_{\tilde{a}_2} \rangle; \tag{4}$$

$$\lambda \tilde{a}_1 = \langle [\lambda a_1, \lambda b_1, \lambda c_1, \lambda d_1]; 1 - (1 - u_{\tilde{a}_1})^\lambda, v_{\tilde{a}_1}^\lambda \rangle; \tag{5}$$

$$\tilde{a}_1^\lambda = \langle [a_1^\lambda, b_1^\lambda, c_1^\lambda, d_1^\lambda]; u_{\tilde{a}_1}^\lambda, 1 - (1 - v_{\tilde{a}_1})^\lambda \rangle. \tag{6}$$

Theorem 1. Let $\tilde{a}_1 = \langle [a_1, b_1, c_1, d_1]; \mu_{\tilde{a}_1}, \nu_{\tilde{a}_1} \rangle$ and $\tilde{a}_2 = \langle [a_2, b_2, c_2, d_2]; \mu_{\tilde{a}_2}, \nu_{\tilde{a}_2} \rangle$ be two positive intuitionistic trapezoidal fuzzy numbers, then there are the following properties about the calculation rules between \tilde{a}_1 and \tilde{a}_2 :

$$\tilde{a}_1 + \tilde{a}_2 = \tilde{a}_2 + \tilde{a}_1; \tag{7}$$

$$\tilde{a}_1 \tilde{a}_2 = \tilde{a}_2 \tilde{a}_1; \tag{8}$$

$$\lambda(\tilde{a}_1 + \tilde{a}_2) = \lambda \tilde{a}_1 + \lambda \tilde{a}_2, \lambda \geq 0; \tag{9}$$

$$\lambda_1 \tilde{a}_1 + \lambda_2 \tilde{a}_1 = (\lambda_1 + \lambda_2) \tilde{a}_1, \lambda_1, \lambda_2 \geq 0; \tag{10}$$

$$\tilde{a}_1^{\lambda_1} \tilde{a}_1^{\lambda_2} = (\tilde{a}_1)^{\lambda_1 + \lambda_2}, \lambda_1, \lambda_2 \geq 0; \tag{11}$$

$$\tilde{a}_1^{\lambda_1} \tilde{a}_2^{\lambda_1} = (\tilde{a}_1 \tilde{a}_2)^{\lambda_1}, \lambda_1 \geq 0. \tag{12}$$

It is easy to prove Theorem 1; therefore, it is omitted here.

Definition 3. Let $\tilde{a}_1 = \langle [a_1, b_1, c_1, d_1]; u_{\tilde{a}_1}, v_{\tilde{a}_1} \rangle$ and $\tilde{a}_2 = \langle [a_2, b_2, c_2, d_2]; u_{\tilde{a}_2}, v_{\tilde{a}_2} \rangle$ be two positive intuitionistic trapezoidal fuzzy numbers, then the possibility degree of intuitionistic trapezoidal fuzzy numbers $\tilde{a}_1 \geq \tilde{a}_2$ is:

$$p(\tilde{a}_1 \geq \tilde{a}_2) = (1 + \frac{(\eta_{\tilde{a}_1} a_1 - \eta_{\tilde{a}_2} a_2) + (\eta_{\tilde{a}_1} b_1 - \eta_{\tilde{a}_2} b_2) + (\eta_{\tilde{a}_1} c_1 - \eta_{\tilde{a}_2} c_2) + (\eta_{\tilde{a}_1} d_1 - \eta_{\tilde{a}_2} d_2)}{|\eta_{\tilde{a}_1} a_1 - \eta_{\tilde{a}_2} a_2| + |\eta_{\tilde{a}_1} b_1 - \eta_{\tilde{a}_2} b_2| + |\eta_{\tilde{a}_1} c_1 - \eta_{\tilde{a}_2} c_2| + |\eta_{\tilde{a}_1} d_1 - \eta_{\tilde{a}_2} d_2|}) / 2; \tag{13}$$

where $\eta_{\tilde{a}_1} = 1 + u_{\tilde{a}_1} - v_{\tilde{a}_1}$, $\eta_{\tilde{a}_2} = 1 + u_{\tilde{a}_2} - v_{\tilde{a}_2}$.

Suppose $p(\tilde{a}_1 \geq \tilde{a}_2) = 0.5$, when $\tilde{a}_1 = \tilde{a}_2$, i.e. $a_1 = a_2, b_1 = b_2, c_1 = c_2, d_1 = d_2, u_{\tilde{a}_1} = u_{\tilde{a}_2}, v_{\tilde{a}_1} = v_{\tilde{a}_2}$.

For a set of intuitionistic trapezoidal fuzzy numbers $\tilde{a}_i = \langle [a_i, b_i, c_i, d_i]; u_{\tilde{a}_i}, v_{\tilde{a}_i} \rangle$, ($i = 1, 2, \dots, n$), the possibility degree $p_{ij} = p(\tilde{a}_i > \tilde{a}_j)$, ($i = 1, 2, \dots, n; j = 1, 2, \dots, m$) can be obtained by the pair-wise comparison, then the possibility degree matrix $P = (p_{ij \times n})$ was built. Let $\lambda_i = \sum_{j=1}^n p_{ij}$, we can get the ranking vector $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$. The bigger λ_i is, the bigger the intuitionistic trapezoidal fuzzy number is.

Definition 4. For the positive intuitionistic trapezoidal fuzzy number $\tilde{a} = \langle [a, b, c, d]; \mu_{\tilde{a}}, \nu_{\tilde{a}} \rangle$, its expected value is shown as follows:

$$I(\tilde{a}) = \frac{1}{8} \times [(a + b + c + d) \times (1 + \mu_{\tilde{a}} - \nu_{\tilde{a}})], \tag{14}$$

where $(a + b + c + d) / 4$ is the expectation of trapezoidal fuzzy number $[a, b, c, d]$, $(\mu_{\tilde{a}} + 1 - \nu_{\tilde{a}}) / 2$ is the expectation of intuitionistic fuzzy number $(\mu_{\tilde{a}}, \nu_{\tilde{a}})$.

1.2. Prioritised ordered weighted aggregated operator (POWA)

1.2.1. Definition of a POWA operator

Definition 5 (Yager 2009). Suppose that we have a collection of criteria partitioned into p distinct categories H_1, H_2, \dots, H_p such that $H_i = \{a_{i1}, a_{i2}, \dots, a_{in_i}\}$. Here a_{ij} are the criteria in the category H_i . We assume a prioritisation between these categories $H_1 > H_2 > \dots > H_p$.

The criteria in the class H_i have a higher priority than those in H_k if $i < k$. The total set of criteria is $C = \bigcup_{i=1}^p H_i$. We assume $n = \sum_{i=1}^p n_i$ is the total number of criteria. If $f_p : [0, 1]^n \rightarrow [0, 1]$, then

$$f_p(a_1, a_2, \dots, a_n) = \sum_{i=1}^p w_i f_i, \tag{15}$$

where $f_i = \sum_{j=1}^{n_i} \omega_{ij} a_{ij}$ is the aggregated value in the category H_i by weighted average operator, $\omega_i = (\omega_{i1}, \omega_{i2}, \dots, \omega_{in_i})^T$ is the attribute weight vector of a_{ij} ($j = 1, 2, \dots, n_i$).

Here, $0 \leq \omega_{ij} \leq 1, \sum_{i=1}^p \sum_{j=1}^{n_i} \omega_{ij} = 1$. Suppose $\tau = (\tau_1, \tau_2, \dots, \tau_p)$ is the position weight of f_i ($i = 1, 2, \dots, p$), calculated in the non-priority conditions, such that $0 \leq \tau_i \leq 1, \sum_{i=1}^p \tau_i = 1$ and $w = (w_1, w_2, \dots, w_p)^T$ is the weight vector associated with the category H_1, H_2, \dots, H_p , $w_i \in [0, 1], i = 1, 2, \dots, p, \sum_{i=1}^p w_i = 1$, then function f_p is called as POWA operator.

1.2.2. Position weight

About the position weight $\tau = (\tau_1, \tau_2, \dots, \tau_p)$, it can be determined according to the actual needs, or according to the following methods.

Firstly, we need to set the value $\alpha \in [0, 1]$ as expected attitude characteristics of the weighted vector. So, we calculate the position weighted vector $\tau = (\tau_1, \tau_2, \dots, \tau_p)$ used by the following math formula (O’Hagan 1990):

$$\begin{aligned}
 & \text{Max } - \sum_{j=1}^p \tau_j \ln(\tau_j) \\
 & \text{S/t } \sum_{j=1}^p \tau_j \frac{p-j}{p-1} = \alpha \\
 & \sum_{j=1}^p \tau_j = 1 \\
 & 0 \leq \tau_j \leq 1.
 \end{aligned}
 \tag{16}$$

In addition, the position weight vector τ can be determined by combinatorial numbers, as follows (Wang, Xu 2008):

$$\tau_{i+1} = \frac{C_{p-1}^i}{2^{p-1}}, \quad i = 0, 1, \dots, p-1.
 \tag{17}$$

1.2.3. Weight vector associated with the category H_1, H_2, \dots, H_p (Yager 2004, 2008)

Suppose T_i is the measurement of the relative importance in every prioritised level. Let $T_1 = 1, V_i = f_i$, then

$$T_i = \prod_{k=1}^{i-1} V_k = T_{i-1} V_{k-1}.
 \tag{18}$$

Let r_i be a normalised priority-based weight, and $r_{\sigma(i)}$ be the corresponding prioritised weight of $f_{\sigma(i)}$, $f_{\sigma(i)}$ is the i th largest element of $f_j, i = 1, 2, \dots, p$, and let $R_i = \sum_{i=1}^p r_{\sigma(i)}$ We have:

$$r_i = \frac{T_i}{\sum_{i=1}^p T_i}, \quad i = 1, 2, \dots, p.
 \tag{19}$$

Choose the BUM function:

$$h(z) = \sum_{i=1}^{j-1} \tau_i + \tau_j (pz - (j-1)), \quad (j-1)/p \leq z \leq j/p.
 \tag{20}$$

Let the weight vector associated with the category (H_1, H_2, \dots, H_p) be $w = (w_1, w_2, \dots, w_p)^T$ and $h(R_0) = 0$. Then,

$$w_i = h(R_{j+1}) - h(R_j).
 \tag{21}$$

2. Intuitionistic trapezoidal fuzzy prioritised ordered weighted aggregation (ITFPOWA) operator

2.1. Definition of ITFPOWA operator

Supposed there is a set of attributes $C = (C_{11}, C_{12}, \dots, C_{1n_1}, C_{21}, C_{22}, \dots, C_{2n_2}, C_{p1}, C_{p2}, \dots, C_{pn_p})$ and H_i is the corresponding prioritised degree, and meets $H_1 > H_2 > \dots > H_p$ so that $H_i = \{\tilde{a}_{i1}, \tilde{a}_{i2}, \dots, \tilde{a}_{in_i}\}$, \tilde{a}_{ij} are the criteria values in the category H_i under the attribute C_{ij} , and $\tilde{a}_{ij} = \langle [a_{ij}^a, a_{ij}^b, a_{ij}^c, a_{ij}^d]; u_{ij}, v_{ij} \rangle, 0 \leq a_{ij}^a, a_{ij}^b, a_{ij}^c, a_{ij}^d \leq 1, 0 \leq u_{ij} \leq 1; 0 \leq v_{ij} \leq 1, 0 \leq u_{ij} + v_{ij} \leq 1,$

$i = 1, 2, \dots, p, j = 1, 2, \dots, n_i$, where $n = \sum_{i=1}^p n_i$. $\omega_i = (\omega_{i1}, \omega_{i2}, \dots, \omega_{in_i})^T$ is the attribute weight of $C_{ij}, i = 1, 2, \dots, p, j = 1, 2, \dots, n_i, 0 \leq \omega_{ij} \leq 1, \sum_{i=1}^p \sum_{j=1}^{n_i} \omega_{ij} = 1$. $\tilde{f} = (\tilde{f}_1, \tilde{f}_2, \dots, \tilde{f}_p)$ is the aggregated vector in the category $H_i (i = 1, 2, \dots, p)$ by intuitionistic trapezoidal fuzzy weighted average operator, where

$$\begin{aligned} \tilde{f}_i = & \langle [f_i^a, f_i^b, f_i^c, f_i^d], fu_i, fv_i \rangle = \sum_{j=1}^{n_i} \omega_{ij} \tilde{a}_{ij} = \\ & \langle [\sum_{j=1}^{n_i} \omega_{ij} a_{ij}^a, \sum_{j=1}^{n_i} \omega_{ij} a_{ij}^b, \sum_{j=1}^{n_i} \omega_{ij} a_{ij}^c, \sum_{j=1}^{n_i} \omega_{ij} a_{ij}^d]; 1 - \prod_{j=1}^{n_i} (1 - u_{ij})^{\omega_{ij}}, \prod_{j=1}^{n_i} v_{ij}^{\omega_{ij}} \rangle. \end{aligned} \tag{22}$$

Definition 6. If $\tilde{F} : [0, 1]^n \rightarrow [0, 1]$, make:

$$\tilde{F} = \sum_{i=1}^p w_i \tilde{f}_i = \left\langle \left[\sum_{i=1}^p w_i f_i^a, \sum_{i=1}^p w_i f_i^b, \sum_{i=1}^p w_i f_i^c, \sum_{i=1}^p w_i f_i^d \right]; 1 - \prod_{i=1}^p (1 - fu_i)^{w_i}, \prod_{i=1}^p fv_i^{w_i} \right\rangle, \tag{23}$$

where $w = (w_1, w_2, \dots, w_p)^T$ is the associated weighted vector to the function \tilde{F} , $0 \leq w_i \leq 1, \sum_{i=1}^p w_i = 1$; then, we call the function \tilde{F} an intuitionistic trapezoidal fuzzy prioritised ordered weighted aggregated operator, referred to as the ITFPOWA operator.

2.2. Associated weighted vector

Suppose T_i is the measurement of the relative importance of intuitionistic trapezoidal fuzzy numbers in every prioritised level, let $T_1 = 1, V_i$ is the expected value of the intuitionistic trapezoidal fuzzy number \tilde{f}_i . Then,

$$T_i = \prod_{k=1}^{i-1} V_k = T_{i-1} V_{k-1}. \tag{24}$$

Let r_i be the normalised prioritised weight, $r_{\sigma(i)}$ – the corresponding prioritised weight of $\tilde{f}_{\sigma(i)}, \tilde{f}_{\sigma(i)}$ – the i th largest element of $(\tilde{f}_1, \tilde{f}_2, \dots, \tilde{f}_p)$, and let $R_i = \sum_{i=1}^p r_{\sigma(i)}$. We have:

$$r_i = \frac{T_i}{\sum_{i=1}^p T_i}, i = 1, 2, \dots, p. \tag{25}$$

Choose the BUM function:

$$h(z) = \sum_{i=1}^{j-1} \tau_i + \tau_j (pz - (j-1)) (j-1) / p \leq z \leq j / p. \tag{26}$$

Let the weight vector associated with the category H_1, H_2, \dots, H_p be $w = (w_1, w_2, \dots, w_p)^T$ and $h(R_0) = 0$. Then,

$$w_i = h(R_{j+1}) - h(R_j). \tag{27}$$

2.3. Characteristic of the ITFPOWA operator

Theorem 2. Let $\tilde{a}_i = \langle [a_i^a, a_i^b, a_i^c, a_i^d]; u_{\tilde{a}_i}, v_{\tilde{a}_i} \rangle (i = 1, 2, \dots, n)$ be a set of intuitionistic trapezoidal fuzzy numbers, then the integrated value by ITFPOWA operator is still an intuitionistic trapezoidal fuzzy number, and

$$\tilde{F} = \langle [\sum_{i=1}^n w_i a_{\sigma(i)}^a, \sum_{i=1}^n w_i a_{\sigma(i)}^b, \sum_{i=1}^n w_i a_{\sigma(i)}^c, \sum_{i=1}^n w_i a_{\sigma(i)}^d]; 1 - \prod_{i=1}^n (1 - u_{\sigma(i)})^{w_i}, \prod_{i=1}^n v_{\sigma(i)}^{w_i} \rangle .$$

Proof. We can use the mathematical induction method to prove this theorem. When $n = 2$,

$$\tilde{F} = w_1 \tilde{a}_{\sigma(1)} + w_2 \tilde{a}_{\sigma(2)} , \tag{28}$$

where $\tilde{a}_{\sigma(1)} = \max(\tilde{a}_1, \tilde{a}_2)$ and $\tilde{a}_{\sigma(2)} = \min(\tilde{a}_1, \tilde{a}_2)$ are both intuitionistic trapezoidal fuzzy numbers, $w_1, w_2 \in [0, 1]$ are both real numbers. From basic operations of intuitionistic trapezoidal fuzzy numbers, we can know, $\tilde{F}(\tilde{a}_1, \tilde{a}_2)$ is intuitionistic trapezoidal fuzzy number, and

$$\begin{aligned} \tilde{F} &= w_1 \tilde{a}_{\sigma(1)} + w_2 \tilde{a}_{\sigma(2)} = \\ &\langle [w_1 a_{\sigma(1)}^a + w_2 a_{\sigma(2)}^a, w_1 a_{\sigma(1)}^b + w_2 a_{\sigma(2)}^b, w_1 a_{\sigma(1)}^c + w_2 a_{\sigma(2)}^c, w_1 a_{\sigma(1)}^d + w_2 a_{\sigma(2)}^d], \\ &u_{\sigma(1)} + u_{\sigma(2)} - u_{\sigma(1)} u_{\sigma(2)}, v_{\sigma(1)} v_{\sigma(2)} \rangle . \end{aligned} \tag{29}$$

Supposed $n = k$, theorem 2 is right, that is

$$\begin{aligned} \tilde{F} &= \sum_{i=1}^k w_i \tilde{a}_{\sigma(i)} = \\ &\langle [\sum_{i=1}^k w_i a_{\sigma(i)}^a, \sum_{i=1}^k w_i a_{\sigma(i)}^b, \sum_{i=1}^k w_i a_{\sigma(i)}^c, \sum_{i=1}^k w_i a_{\sigma(i)}^d]; 1 - \prod_{i=1}^k (1 - u_{\sigma(i)})^{w_i}, \prod_{i=1}^k v_{\sigma(i)}^{w_i} \rangle . \end{aligned} \tag{30}$$

Then when $n = k + 1$, we can get

$$\begin{aligned} \tilde{F} &= \sum_{i=1}^k w_i \tilde{a}_{\sigma(i)} + w_{k+1} \tilde{a}_{k+1} = \\ &\langle [\sum_{i=1}^{k+1} w_i a_{\sigma(i)}^a, \sum_{i=1}^{k+1} w_i a_{\sigma(i)}^b, \sum_{i=1}^{k+1} w_i a_{\sigma(i)}^c, \sum_{i=1}^{k+1} w_i a_{\sigma(i)}^d]; 1 - \prod_{i=1}^{k+1} (1 - u_{\sigma(i)})^{w_i}, \prod_{i=1}^{k+1} v_{\sigma(i)}^{w_i} \rangle . \end{aligned} \tag{31}$$

So, when $n = k + 1$, theorem 2 is still right.

According to the principle of mathematical induction method, we can state that Theorem 2 holds.

Theorem 3 (Idempotency). Let $\tilde{a}_i = \langle [a_i, b_i, c_i, d_i]; u_{\tilde{a}_i}, v_{\tilde{a}_i} \rangle (i = 1, 2, \dots, n)$ be a set of intuitionistic trapezoidal fuzzy numbers, if the value of all intuitionistic trapezoidal fuzzy numbers is equal, that is $\tilde{a}_i = \langle [a_i, b_i, c_i, d_i]; u_{\tilde{a}_i}, v_{\tilde{a}_i} \rangle = \tilde{a} = \langle [a, b, c, d]; u_{\tilde{a}}, v_{\tilde{a}} \rangle$, then we have

$$\tilde{F} = \tilde{a} = \langle [a, b, c, d]; u_{\tilde{a}}, v_{\tilde{a}} \rangle . \tag{32}$$

Proof:

$$\begin{aligned}
 \tilde{F} &= \sum_{i=1}^n w_i \tilde{a}_{\sigma(i)} = \\
 &< [\sum_{i=1}^n w_i a_{\sigma(i)}, \sum_{i=1}^n w_i b_{\sigma(i)}, \sum_{i=1}^n w_i c_{\sigma(i)}, \sum_{i=1}^n w_i d_{\sigma(i)}]; 1 - \prod_{i=1}^n (1 - u_{\sigma(i)})^{w_i}, \prod_{i=1}^n v_{\sigma(i)}^{w_i} > = \\
 &< [\sum_{i=1}^n w_i a, \sum_{i=1}^n w_i b, \sum_{i=1}^n w_i c, \sum_{i=1}^n w_i d]; 1 - \prod_{i=1}^n (1 - u)^{w_i}, \prod_{i=1}^n v^{w_i} >, \tag{33}
 \end{aligned}$$

because

$$\begin{aligned}
 \sum_{i=1}^n w_i a &= a, \sum_{i=1}^n w_i b = b, \sum_{i=1}^n w_i c = c, \sum_{i=1}^n w_i d = d; \\
 1 - \prod_{i=1}^n (1 - u)^{w_i} &= 1 - (1 - u)^{\sum_{i=1}^n w_i} = 1 - (1 - u) = u, \\
 \prod_{i=1}^n v^{w_i} &= v^{\sum_{i=1}^n w_i} = v. \tag{34}
 \end{aligned}$$

So, $\tilde{F} = \tilde{a} = \langle [a, b, c, d]; u_{\tilde{a}}, v_{\tilde{a}} \rangle$.

Theorem 4 (Boundedness). Let $\tilde{a}_i = \langle [a_i, b_i, c_i, d_i]; u_{\tilde{a}_i}, v_{\tilde{a}_i} \rangle (i = 1, 2, \dots, n)$ be a set of intuitionistic trapezoidal fuzzy numbers, and $\tilde{a}^- = \min_i \langle [a_i, b_i, c_i, d_i]; u_{\tilde{a}_i}, v_{\tilde{a}_i} \rangle$, $\tilde{a}^+ = \max_i \langle [a_i, b_i, c_i, d_i]; u_{\tilde{a}_i}, v_{\tilde{a}_i} \rangle$, then $\tilde{a}^- \leq \tilde{F} \leq \tilde{a}^+$.

Proof:

Because $\tilde{a}_{\sigma(i)}$ is the i th largest element in $\tilde{a}_i (i = 1, 2, \dots, n)$, for any i , there is $\min(\tilde{a}_{\sigma(i)}) \leq \tilde{a}_{\sigma(i)} \leq \max(\tilde{a}_{\sigma(i)})$. Then, there is :

$$\sum_{i=1}^n w_i \tilde{a}_{\sigma(i)} \geq \sum_{i=1}^n w_i \min(\tilde{a}_{\sigma(i)}) = \min(\tilde{a}_{\sigma(i)}), \sum_{i=1}^n w_i \tilde{a}_{\sigma(i)} \leq \sum_{i=1}^n w_i \max(\tilde{a}_{\sigma(i)}) = \max(\tilde{a}_{\sigma(i)}).$$

So, we can get $\min(\tilde{a}_{\sigma(i)}) \leq \sum_{i=1}^n w_i \tilde{a}_{\sigma(i)} \leq \max(\tilde{a}_{\sigma(i)})$, i.e., $\tilde{a}^- \leq \tilde{F} \leq \tilde{a}^+$.

3. Multi-attribute decision-making method based on the ITFPOWA operator

Let $A = \{A_1, A_2, \dots, A_m\}$ be a discrete set of alternatives, and $C = (C_{11}, C_{12}, \dots, C_{1n_1}, C_{21}, C_{22}, \dots, C_{2n_2}, C_{p1}, C_{p2}, \dots, C_{pn_p})$ be the set of attributes, H_k — the corresponding prioritised degree and meets $H_1 > H_2 > \dots > H_p$ so that $H_k = \{\tilde{a}_{k1}^i, \tilde{a}_{k2}^i, \dots, \tilde{a}_{kn_k}^i\}$, \tilde{a}_{kj}^i — the criteria values in the category H_k under the attribute C_{kj} with respect to the alternative A_p , and $\tilde{a}_{kj}^i = \langle [a_{kj}^i, b_{kj}^i, b_{kj}^i, d_{kj}^i]; u_{kj}^i, v_{kj}^i \rangle, 0 \leq a_{kj}^i, b_{kj}^i, b_{kj}^i, d_{kj}^i \leq 1, 0 \leq u_{kj}^i \leq 1; 0 \leq v_{kj}^i \leq 1, 0 \leq u_{kj}^i + v_{kj}^i \leq 1, i = 1, 2, \dots, m, k = 1, 2, \dots, p, j = 1, 2, \dots, n_k$, where $n = \sum_{k=1}^p n_k$.

$\omega = (\omega_{11}, \omega_{12}, \dots, \omega_{1n_1}, \omega_{21}, \omega_{22}, \dots, \omega_{2n_2}, \omega_{p1}, \omega_{p2}, \dots, \omega_{pn_p})^T$ is the attribute weight vector of $C_{kj}, k = 1, 2, \dots, p, j = 1, 2, \dots, n_k, 0 \leq \omega_{kj} \leq 1, \sum_{k=1}^p \sum_{j=1}^{n_k} \omega_{kj} = 1$. $\tilde{f}^i = (\tilde{f}_1^i, \tilde{f}_2^i, \dots, \tilde{f}_p^i)$ is the aggregat-

ed vector in the category H_k ($k = 1, 2, \dots, p$) with respect to the alternative A_i by the intuitionistic trapezoidal fuzzy weighted average operator, $\tau = (\tau_1, \tau_2, \dots, \tau_p)$ is the position weight of $\tilde{f}^i = (\tilde{f}_1^i, \tilde{f}_2^i, \dots, \tilde{f}_p^i)$ calculated in the non-priority conditions, so that $0 \leq \tau_k \leq 1, \sum_{k=1}^p \tau_k = 1$, and $w_i = (w_{i1}, w_{i2}, \dots, w_{ip})^T$ is the weight vector associated with the category H_1, H_2, \dots, H_p with respect to the alternative $A_p, w_{ik} \in [0, 1], i = 1, 2, \dots, m, k = 1, 2, \dots, p, \sum_{k=1}^p w_{ik} = 1$. Then, the ranking of alternatives is required.

Next, we apply the ITFPOWA operator to MADM problems based on the intuitionistic trapezoidal fuzzy information.

Step 1. Get the aggregated information \tilde{f}_k^i in the same category H_k with respect to the alternative A_i .

We can use the intuitionistic trapezoidal fuzzy weighted average operator to integrate the information $(\tilde{a}_{k1}^i, \tilde{a}_{k2}^i, \dots, \tilde{a}_{kn_k}^i)$ in the same category H_k , and get

$$\tilde{f}_k^i = \sum_{j=1}^{n_k} \omega_{kj} \tilde{a}_{kj}^i. \tag{35}$$

Step 2. Calculate the position weight $\tau = (\tau_1, \tau_2, \dots, \tau_p)$ in the non-priority conditions by Formulas (16) or (17).

Step 3. Calculate associated weighted vector $w_i = (w_{i1}, w_{i2}, \dots, w_{ip})^T$ according to Formulas (24–27).

Step 4. Calculate the integrated value \tilde{f}^i by the ITFPOWA operator.

We can get

$$\tilde{f}^i = \sum_{k=1}^p w_{ik} \tilde{f}_k^i. \tag{36}$$

Step 5. Rank the alternatives A_1, A_2, \dots, A_m by the matrix of possibility degree.

Step 6. Finished.

4. Illustrative example

4.1. Problem formulation

Consider the recruitment of human resource managers.

There are 4 recruiters (evaluation objects), $A = (A_1, A_2, A_3, A_4)$, and 8 evaluation indicators, $C = (C_{11}, C_{12}, C_{13}, C_{21}, C_{31}, C_{41}, C_{42}, C_{51})$.

H_k is the corresponding prioritised level, and has prioritised relationship $H_1 > H_2 > H_3 > H_4 > H_5$. H_1 (Ability level): Learning ability C_{11} , Management innovation ability C_{12} , Communication and coordination ability C_{13} ; H_2 : Professional quality C_{21} ; H_3 : Moral character C_{31} ; H_4 (Knowledge level): Professional knowledge C_{41} , Laws and regulations knowledge C_{42} ; H_5 : State of health C_{51} .

$\omega_k = (\omega_{k1}, \omega_{k2}, \dots, \omega_{kn_k})^T$ is the attribute weight vector, $0 \leq \omega_{kj} \leq 1, \sum_{k=1}^p \sum_{j=1}^{n_k} \omega_{kj} = 1$. $\omega_1 = (0.1, 0.1, 0.1)^T, \omega_2 = 0.2, \omega_3 = 0.2, \omega_4 = (0.1, 0.1)^T, \omega_5 = 0.1$.

\tilde{a}_{kj}^i is the criteria value in the category H_k under the attribute C_{kj} with respect to the alternative A_p and $\tilde{a}_{kj}^i = \langle [a_{kj}^i, b_{kj}^i, b_{kj}^i, d_{kj}^i]; u_{kj}^i, v_{kj}^i \rangle$, $0 \leq a_{kj}^i, b_{kj}^i, b_{kj}^i, d_{kj}^i \leq 1$, $0 \leq u_{kj}^i \leq 1; 0 \leq v_{kj}^i \leq 1$, $0 \leq u_{kj}^i + v_{kj}^i \leq 1$, $i = 1, 2, 3, 4, k = 1, 2, 3, 4, 5, j = 1, 2, \dots, n_k$.

Decision-making information is shown in Table 1. $\tilde{f}^i = (\tilde{f}_1^i, \tilde{f}_2^i, \dots, \tilde{f}_p^i)$ is the aggregated vector in category H_k , ($k = 1, 2, 3, 4, 5$) with respect to the alternative A_i by the intuitionistic trapezoidal fuzzy weighted average (ITFWA) operator, $\tau = (\tau_1, \tau_2, \dots, \tau_p)$ is the position weight of $\tilde{f}^i = (\tilde{f}_1^i, \tilde{f}_2^i, \dots, \tilde{f}_p^i)$, calculated in the non-priority conditions, here $\tau = (0.3, 0.2, 0.1, 0.2, 0.2)^T$, and $w_i = (w_{i1}, w_{i2}, w_{i3}, w_{i4}, w_{i5})^T$ is the weight vector associated with the category H_1, H_2, H_3, H_4, H_5 with respect to the alternative A_p , $w_{ik} \in [0, 1], i = 1, 2, 3, 4, k = 1, 2, 3, 4, 5$. $\sum_{k=1}^5 w_{ik} = 1$. The aim is to select the most suitable candidate.

Table 1. Decision-making matrix $\tilde{A} = (\tilde{a}_{ij})_{m \times n}$

	C_{11}	C_{12}	C_{13}
A_1	$\langle [0.3, 0.4, 0.5, 0.6]; 0.5, 0.3 \rangle$	$\langle [0.7, 0.8, 0.9, 1.0]; 0.8, 0.2 \rangle$	$\langle [0.4, 0.5, 0.6, 0.7]; 0.5, 0.3 \rangle$
A_2	$\langle [0.4, 0.5, 0.6, 0.7]; 0.5, 0.2 \rangle$	$\langle [0.5, 0.6, 0.6, 0.7]; 0.5, 0.4 \rangle$	$\langle [0.7, 0.8, 0.9, 1.0]; 0.5, 0.4 \rangle$
A_3	$\langle [0.5, 0.6, 0.6, 0.7]; 0.7, 0.2 \rangle$	$\langle [0.4, 0.5, 0.6, 0.7]; 0.3, 0.2 \rangle$	$\langle [0.5, 0.6, 0.6, 0.7]; 0.6, 0.2 \rangle$
A_4	$\langle [0.6, 0.7, 0.8, 0.9]; 0.8, 0.1 \rangle$	$\langle [0.5, 0.6, 0.7, 0.8]; 0.6, 0.2 \rangle$	$\langle [0.4, 0.5, 0.6, 0.7]; 0.9, 0.1 \rangle$

Table 1. Decision-making matrix $\tilde{A} = (\tilde{a}_{ij})_{m \times n}$ (Continued 2)

	C_{21}	C_{31}	C_{41}
A_1	$\langle [0.7, 0.8, 0.9, 1.0]; 0.8, 0.2 \rangle$	$\langle [0.7, 0.8, 0.9, 1.0]; 0.8, 0.2 \rangle$	$\langle [0.7, 0.8, 0.9, 1.0]; 0.8, 0.2 \rangle$
A_2	$\langle [0.6, 0.7, 0.8, 0.9]; 0.7, 0.2 \rangle$	$\langle [0.5, 0.6, 0.6, 0.7]; 0.5, 0.4 \rangle$	$\langle [0.6, 0.7, 0.8, 0.9]; 0.7, 0.2 \rangle$
A_3	$\langle [0.3, 0.4, 0.5, 0.6]; 0.7, 0.2 \rangle$	$\langle [0.4, 0.5, 0.6, 0.7]; 0.3, 0.2 \rangle$	$\langle [0.3, 0.4, 0.5, 0.6]; 0.7, 0.2 \rangle$
A_4	$\langle [0.4, 0.5, 0.6, 0.7]; 0.5, 0.3 \rangle$	$\langle [0.5, 0.6, 0.7, 0.8]; 0.6, 0.2 \rangle$	$\langle [0.4, 0.5, 0.6, 0.7]; 0.5, 0.3 \rangle$

Table 1. Decision-making matrix $\tilde{A} = (\tilde{a}_{ij})_{m \times n}$ (Continued 1)

	C_{42}	C_{51}
A_1	$\langle [0.3, 0.4, 0.5, 0.6]; 0.5, 0.3 \rangle$	$\langle [0.7, 0.8, 0.9, 1.0]; 0.8, 0.2 \rangle$
A_2	$\langle [0.4, 0.5, 0.6, 0.7]; 0.5, 0.2 \rangle$	$\langle [0.5, 0.6, 0.6, 0.7]; 0.5, 0.4 \rangle$
A_3	$\langle [0.5, 0.6, 0.6, 0.7]; 0.7, 0.2 \rangle$	$\langle [0.4, 0.5, 0.6, 0.7]; 0.3, 0.2 \rangle$
A_4	$\langle [0.6, 0.7, 0.8, 0.9]; 0.8, 0.1 \rangle$	$\langle [0.5, 0.6, 0.7, 0.8]; 0.6, 0.2 \rangle$

4.2. Multi-attribute decision-making method based on the ITFPOWA operator

Step 1. Integrate the criteria values in 5 class criteria. Then, get the decision-making matrix $\tilde{f}^i = (\tilde{f}_1^i, \tilde{f}_2^i, \dots, \tilde{f}_p^i)$, respectively:

$$\tilde{f} = \begin{bmatrix} \langle [0.14, 0.17, 0.20, 0.23]; 0.26, 0.67 \rangle & \langle [0.14, 0.16, 0.18, 0.20]; 0.28, 0.72 \rangle & \langle [0.14, 0.16, 0.18, 0.20]; 0.28, 0.72 \rangle \\ \langle [0.16, 0.19, 0.21, 0.24]; 0.19, 0.71 \rangle & \langle [0.12, 0.14, 0.16, 0.18]; 0.21, 0.72 \rangle & \langle [0.10, 0.12, 0.12, 0.14]; 0.13, 0.83 \rangle \\ \langle [0.14, 0.17, 0.18, 0.21]; 0.22, 0.62 \rangle & \langle [0.06, 0.08, 0.10, 0.12]; 0.21, 0.72 \rangle & \langle [0.08, 0.10, 0.12, 0.14]; 0.07, 0.72 \rangle \\ \langle [0.15, 0.18, 0.21, 0.24]; 0.38, 0.54 \rangle & \langle [0.08, 0.10, 0.12, 0.14]; 0.13, 0.79 \rangle & \langle [0.10, 0.12, 0.14, 0.16]; 0.17, 0.72 \rangle \end{bmatrix}$$

$$\left. \begin{aligned} &<[0.10,0.12,0.14,0.16];0.21,0.75 > <[0.07,0.08,0.09,0.10];0.15,0.85 > \\ &<[0.10,0.12,0.14,0.16];0.17,0.72 > <[0.05,0.06,0.06,0.07];0.07,0.91 > \\ &<[0.08,0.10,0.11,0.13];0.21,0.72 > <[0.04,0.05,0.06,0.07];0.04,0.85 > \\ &<[0.10,0.12,0.14,0.16];0.21,0.70 > <[0.05,0.06,0.07,0.08];0.09,0.85 > \end{aligned} \right\}.$$

Step 2. Use the ITFPOWA operator to calculate the integrated values \tilde{f}^i for every alternative, respectively.

V is a matrix of expectations, where V_{ik} are expectations of \tilde{f}_k^i . We can get

$$V = \begin{bmatrix} 0.055 & 0.047 & 0.047 & 0.029 & 0.013 \\ 0.048 & 0.037 & 0.018 & 0.029 & 0.005 \\ 0.053 & 0.022 & 0.019 & 0.026 & 0.005 \\ 0.082 & 0.019 & 0.029 & 0.033 & 0.008 \end{bmatrix}.$$

Calculate the associated weighted vector $w_i = (w_{i1}, w_{i2}, \dots, w_{ip})^T$.

Take A_1 for an example. According to Formulas (24–27) we can get

$$\begin{aligned} T_1 &= 1, T_2 = 0.05455, T_3 = 0.00255, T_4 = 0.000119, T_5 = 0.000003; \\ r &= (0.94587, 0.051596, 0.002414, 0.000113, 0.000003); \\ R &= (0.945874, 0.99747, 0.99988, 0.99997, 1); \\ w_1 &= (0.94587, 0.051596, 0.002414, 0.000113, 0.000003). \end{aligned}$$

Calculate the integrated values \tilde{f}^i :

$$\tilde{f}^1 = ITFPOWA(A_1) = <[0.14, 0.17, 0.20, 0.23]; 0.26, 0.67 >.$$

Similarly, we can get

$$\begin{aligned} \tilde{f}^2 &= ITFPOWA(A_2) = <[0.16, 0.19, 0.21, 0.24]; 0.19, 0.71 >; \\ \tilde{f}^3 &= ITFPOWA(A_3) = <[0.14, 0.17, 0.18, 0.21]; 0.22, 0.62 >; \\ \tilde{f}^4 &= ITFPOWA(A_4) = <[0.14, 0.16, 0.17, 0.20]; 0.37, 0.62 >. \end{aligned}$$

Step 3. Use the matrix of possibility degree to rank all alternatives.

According to Formula (13), we can build the matrix of the possibility degree:

$$P = \begin{bmatrix} 0.5 & 1 & 0.864 & 0 \\ 0 & 0.5 & 0 & 0 \\ 0.136 & 1 & 0.5 & 0 \\ 1 & 1 & 1 & 0.5 \end{bmatrix}.$$

Then, we can get the sorting vector $\lambda = (2.364, 0.500, 1.636, 3.500)$.

As $\lambda_4 > \lambda_1 > \lambda_3 > \lambda_2$, so $A_4 \succ A_1 \succ A_3 \succ A_2$, we can conclude that A_4 is the best candidate.

In order to verify the validity of the proposed method, we used the weighted geometric aggregation operator as suggested by Liu and Jin (2012a) to solve this decision-making

problem, and we obtained the ranking result $A_4 \succ A_1 \succ A_3 \succ A_2$. Obviously, the same ranking results have been produced by two methods. However, the newly proposed decision-making method can consider the prioritised relationship among the attributes, and in the method proposed by Liu and Jin (2012a), the attribute values were considered to have an equal status.

Conclusions

In real decision-making, there are many multiple attribute decision-making (MADM) problems, in which there exists the prioritised relationship among decision-making attributes. At the same time, the intuitionistic trapezoidal fuzzy numbers make it easier to express the fuzzy information. So, the research on the prioritised fuzzy MADM method based on the intuitionistic trapezoidal fuzzy numbers has broad application prospects. In addition, the traditional POWA operator is generally suitable for aggregating the information taking the form of numerical values, and yet it fails in adapting to intuitionistic trapezoidal fuzzy numbers. Therefore, in this paper, with respect to MADM problems whose attributes have the prioritised relations and the values of attributes take the form of intuitionistic trapezoidal fuzzy numbers, we combined POWA operator with the intuitionistic trapezoidal fuzzy numbers, and have developed an intuitionistic trapezoidal fuzzy prioritised ordered weighted average (ITFPOWA) operator. Some properties of the operator have also been analysed. Furthermore, based on the above operator, we have proposed an approach to decision-making problems, in which the attribute values were intuitionistic trapezoidal fuzzy numbers. The prominent characteristic of the developed approach is that they can take the prioritised relationship between attributes into account. In the end, an illustrative example has been given to show the steps of the proposed method and to indicate its practicality and validity.

Because intuitionistic trapezoidal fuzzy numbers are more suitable to be used for depicting uncertain or fuzzy information, it has been widely used in real decision-making, such as science and technology project review, blind paper review and so on. At the same time, as the aforementioned method can consider the prioritised relationship among attributes, it is more scientific to do decision-making based on them. In the future, we will keep on studying the extension and applications of the established operators to other domains.

Acknowledgments

This paper has been supported by the National Natural Science Foundation of China (Nos. 71471172 and 71271124), the Special Funds of Taishan Scholars Project of Shandong Province, National Soft Science Project of China (2014GXQ4D192), the Humanities and Social Sciences Research Project of Ministry of Education of China (No. 13YJC630104), Shandong Provincial Social Science Planning Project (No. 15BGLJ06), and the Natural Science Foundation of Shandong Province (No. ZR2011FM036).

References

- Atanassov, K. T. 1986. Intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 20(1): 87–96. [http://dx.doi.org/10.1016/S0165-0114\(86\)80034-3](http://dx.doi.org/10.1016/S0165-0114(86)80034-3)
- Atanassov, K. T. 1994. Operators over interval-valued intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 64(2): 159–174. [http://dx.doi.org/10.1016/0165-0114\(94\)90331-X](http://dx.doi.org/10.1016/0165-0114(94)90331-X)
- Atanassov, K. T.; Gargov, G. 1989. Interval-valued intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 31(3): 343–349. [http://dx.doi.org/10.1016/0165-0114\(89\)90205-4](http://dx.doi.org/10.1016/0165-0114(89)90205-4)
- Chen, Y. B.; Liu, P. D. 2014. Multi-attribute decision-making approach based on intuitionistic trapezoidal fuzzy general Heronian OWA operator, *Journal of Intelligent & Fuzzy Systems* 27(3): 1381–1392.
- Guo, T. T.; Zhang, S. H.; Wei, C. P.; Wang, X. J. 2011. Method based on POWC-OWA operator for interval number multi-attribute information aggregation, *Journal of Qufu Normal University* 37(1): 19–23.
- Liao, H. C.; Xu, Z. S.; Zeng, X. J.; Merigó, J. M. 2015. Framework of group decision making with intuitionistic fuzzy preference information, *IEEE Transactions on Fuzzy Systems* 23(4): 1211–1227. <http://dx.doi.org/10.1109/TFUZZ.2014.2348013>
- Lin, R.; Zhao, X.; Wei, G. W. 2013. Fuzzy number intuitionistic fuzzy prioritized operators and their application to multiple attribute decision making, *Journal of Intelligent and Fuzzy Systems* 24(4): 879–888.
- Liu, P. D. 2014. Some Hamacher aggregation operators based on the interval-valued intuitionistic fuzzy numbers and their application to group decision making, *IEEE Transactions on Fuzzy Systems* 22(1): 83–97. <http://dx.doi.org/10.1109/TFUZZ.2013.2248736>
- Liu, P. D.; Jin, F. 2012a. A multi-attribute group decision-making method based on weighted geometric aggregation operators of interval-valued trapezoidal fuzzy numbers, *Applied Mathematical Modelling* 36(6): 2498–2509. <http://dx.doi.org/10.1016/j.apm.2011.09.006>
- Liu, P. D.; Jin, F. 2012b. Methods for aggregating intuitionistic uncertain linguistic variables and their application to group decision making, *Information Sciences* 205: 58–71. <http://dx.doi.org/10.1016/j.ins.2012.04.014>
- Liu, P. D.; Jin, F. 2012c. The trapezoid fuzzy linguistic Bonferroni mean operators and their application to multiple attribute decision making, *Scientia Iranica* 19(6): 1947–1959. <http://dx.doi.org/10.1016/j.scient.2012.06.028>
- Liu, P. D.; Liu, Y. 2014. An approach to multiple attribute group decision making based on intuitionistic trapezoidal fuzzy power generalized aggregation operator, *International Journal of Computational Intelligence Systems* 7(2): 291–304. <http://dx.doi.org/10.1080/18756891.2013.862357>
- Liu, P. D.; Liu, Z. M.; Zhang, X. 2014. Some intuitionistic uncertain linguistic Heronian mean operators and their application to group decision making, *Applied Mathematics and Computation* 230: 570–586. <http://dx.doi.org/10.1016/j.amc.2013.12.133>
- Liu, P. D.; Wang, Y. M. 2014. Multiple attribute group decision making methods based on intuitionistic linguistic power generalized aggregation operators, *Applied Soft Computing* 17: 90–104. <http://dx.doi.org/10.1016/j.asoc.2013.12.010>
- Liu, P. D.; Yu, X. C. 2013. Density aggregation operators based on the intuitionistic trapezoidal fuzzy numbers for multiple attribute decision making, *Technological and Economic Development of Economy* 19(supplement 1): S454–S470. <http://dx.doi.org/10.3846/20294913.2013.881436>
- Liu, P. D.; Yu, X. C. 2014. 2-dimension uncertain linguistic power generalized weighted aggregation operator and its application for multiple attribute group decision making, *Knowledge-Based Systems* 57(1): 69–80. <http://dx.doi.org/10.1016/j.knsys.2013.12.009>
- Merigó, J. M.; Gil-Lafuente, A. M.; Yager, R. R. 2015. An overview of fuzzy research with bibliometric indicators, *Applied Soft Computing* 27: 420–433. <http://dx.doi.org/10.1016/j.asoc.2014.10.035>

- O'Hagan, M. 1990. Using maximum entropy-ordered weighted average to construct a fuzzy neuron, *Proceedings of the 24th Annual IEEE Asilomar Conference on Signals, Systems and Computers*, 1990, Pacific Grove, CA, 618–623.
- Shu, M. H.; Cheng, C. H.; Chang, J. R. 2006. Using intuitionistic fuzzy sets for fault-tree analysis on printed circuit board assembly, *Microelectronics Reliability* 46(12): 2139–2148.
<http://dx.doi.org/10.1016/j.microrel.2006.01.007>
- Wan, S. P.; Dong, J. Y. 2010. Method of intuitionistic trapezoidal fuzzy number for multi-attribute group decision, *Control and Decision* 25(5): 773–776.
- Wang, J. Q. 2008. Overview on fuzzy multi-criteria decision-making approach, *Control and Decision* 23(6): 601–606.
- Wang, J. Q.; Han, Z. Q.; Zhang, H. Y. 2014. Multi-criteria group decision-making method based on intuitionistic interval fuzzy information, *Group Decision and Negotiation* 23(4): 715–733.
<http://dx.doi.org/10.1007/s10726-012-9316-4>
- Wang, J. Q.; Nie, R. R.; Zhang, H. Y.; Chen, X. H. 2013a. New operators on triangular intuitionistic fuzzy numbers and their applications in system fault analysis, *Information Sciences* 251: 79–95.
<http://dx.doi.org/10.1016/j.ins.2013.06.033>
- Wang, J. Q.; Nie, R. R.; Zhang, H. Y.; Chen, X. H. 2013b. Intuitionistic fuzzy multi-criteria decision-making method based on evidential reasoning, *Applied Soft Computing* 13(4): 1823–1831.
<http://dx.doi.org/10.1016/j.asoc.2012.12.019>
- Wang, J. Q.; Zhang, Z. 2008. Programming multi-criteria decision-making method based on intuitionistic fuzzy number with incomplete certain information, *Control and Decision* 23(10): 1145–1148.
- Wang, Y.; Xu, Z. S. 2008. A new method of giving OWA weights, *Mathematics in Practice and Theory* 38(3): 51–61.
- Wei, G. W. 2012. Hesitant Fuzzy prioritized operators and their application to multiple attribute group decision making, *Knowledge-Based Systems* 31: 176–182.
<http://dx.doi.org/10.1016/j.knsys.2012.03.011>
- Wei, G. W. 2010. Some arithmetic aggregation operators with intuitionistic trapezoidal fuzzy numbers and their application to group decision making, *Journal of Computers* 5(3): 345–351.
<http://dx.doi.org/10.4304/jcp.5.3.345-351>
- Wei, G. W.; Zhao, X.; Wang, H. 2012. An approach to multiple attribute group decision making with interval intuitionistic trapezoidal fuzzy information, *Technological and Economic Development of Economy* 18(2): 317–330. <http://dx.doi.org/10.3846/20294913.2012.676995>
- Xu, Y. J.; Sun, T.; Li, D. F. 2011. Intuitionistic trapezoidal fuzzy prioritized OWA operator and the application in the multi-criteria decision-making, *Control and Decision* 26(1): 129–132.
- Xu, Y. J.; Wang, H. M.; Merigó, J. M. 2014. Intuitionistic Einstein fuzzy Choquet integral operators for multiple attribute decision making, *Technological and Economic Development of Economy* 20(2): 227–253. <http://dx.doi.org/10.3846/20294913.2014.913273>
- Yager, R. R. 2004. Modeling prioritized multicriteria decision making, *IEEE Transactions on Systems, Man, and Cybernetics, Part B: Cybernetics* 34(6): 2396–2404.
<http://dx.doi.org/10.1109/TSMCB.2004.837348>
- Yager, R. R. 2008. Prioritized aggregated operators, *International Journal of Approximate Reasoning* 48(1): 263–274. <http://dx.doi.org/10.1016/j.ijar.2007.08.009>
- Yager, R. R. 2009. Prioritized OWA aggregated, *Fuzzy Optimization Decision Making* 10(8): 245–262.
<http://dx.doi.org/10.1007/s10700-009-9063-4>
- Yan, H. B.; Huynh, V. N.; Nakamori, Y.; Murai, T. 2011. On prioritized weighted aggregation in multi-criteria decision making, *Expert Systems with Applications* 38(1): 812–823.
<http://dx.doi.org/10.1016/j.eswa.2010.07.039>

- Yu, D.; Shi, S. 2015. Researching the development of Atanassov intuitionistic fuzzy set: Using a citation network analysis, *Applied Soft Computing* 32: 189–198. <http://dx.doi.org/10.1016/j.asoc.2015.03.027>
- Zadeh, I. A. 1965. Fuzzy sets, *Information and Control* 8(3): 338–353. [http://dx.doi.org/10.1016/S0019-9958\(65\)90241-X](http://dx.doi.org/10.1016/S0019-9958(65)90241-X)
- Zeng, S. Z.; Wang, Q. F.; Merigó, J. M.; Pan, T. J. 2014. Induced intuitionistic fuzzy ordered weighted averaging: Weighted average operator and its application to business decision-making, *Computer Science & Information Systems* 11(4): 839–857.
- Zhang, X.; Jin, F.; Liu, P. D. 2013. A grey relational projection method of multi-attribute decision making based on intuitionistic trapezoidal fuzzy number with weight information unknown, *Applied Mathematical Modelling* 37(5): 3467–3477. <http://dx.doi.org/10.1016/j.apm.2012.08.012>
- Zhang, X.; Liu, P. D. 2010. Method for aggregating triangular intuitionistic fuzzy information and its application to decision making, *Technological and Economic Development of Economy* 16(2): 280–290. <http://dx.doi.org/10.3846/tede.2010.18>
- Zhao, X.; Lin, R.; Wei, G. W. 2013. Fuzzy prioritized operators and their application to multiple attribute group decision making, *Applied Mathematical Modelling* 37(7): 4759–4770. <http://dx.doi.org/10.1016/j.apm.2012.09.048>

Peide LIU is Doctor and Professor at the School of Management Science and Engineering at Shandong University of Finance and Economics, and assistant director of the Enterprise's Electronic-commerce Engineering Research Center of Shandong. He has authored or co-authored more than 110 publications. His research interests include aggregation operators, fuzzy logic, fuzzy decision-making, and their applications.

Ye LI has Master's Degree received from the University of International Relations in International politics in 2008. Now she is General Manager of Beijing Engineering Research Center of Green Building Technology, and is studying for her doctorate in engineering and project management at Beijing Jiaotong University. She has authored or co-authored 6 publications. Research interests: construction supply chain management, construction technology and organisation, construction energy saving and construction engineering management, multiple criteria decision-making.

Jurgita ANTUVHEVIČIENĖ is Doctor and Professor at the Department of Construction Technology and Management, Vilnius Gediminas Technical University, Vilnius, Lithuania. She is a member of EURO Working Groups Multicriteria Decision Aiding and OR in Sustainable Development and Civil Engineering, Editorial Board member of three international research journals. Her research interests include sustainable development, construction business management and investment, multiple criteria analysis, decision-making theories and decision-support systems.