

# A new life for an old pump: V-ATPase and neurotransmitter release

Stefano Vavassori and Andreas Mayer

Département de Biochimie, Université de Lausanne, 1066 Epalinges, Switzerland

Neurons fire by releasing neurotransmitters via fusion of synaptic vesicles with the plasma membrane. Fusion can be evoked by an incoming signal from a preceding neuron or can occur spontaneously. Synaptic vesicle fusion requires the formation of trans complexes between SNAREs as well as  $\text{Ca}^{2+}$  ions. Wang et al. (2014, *J. Cell Biol.* <http://dx.doi.org/jcb.201312109>) now find that the  $\text{Ca}^{2+}$ -binding protein Calmodulin promotes spontaneous release and SNARE complex formation via its interaction with the  $V_0$  sector of the V-ATPase.

Evoked release of synaptic vesicles occurs after an action potential (or series of action potentials), whereas spontaneous release occurs in the absence of a presynaptic action potential (Kochubey et al., 2011). Both forms of release require the formation of trans complexes between SNARE (“soluble NSF attachment receptor”) proteins, one in the vesicle and the other anchored in the plasma membrane. SNARE proteins exist in two cognate forms: R-SNAREs (e.g., Synaptobrevin, which is located in the synaptic vesicles) and Q-SNAREs, which can contain up to three subunits (e.g., Syntaxin1A and SNAP-25, located in the plasma membrane). Both forms of neurotransmitter release also require  $\text{Ca}^{2+}$  ions. Synaptotagmins act as a  $\text{Ca}^{2+}$  sensor for evoked release. Little is known about how the  $\text{Ca}^{2+}$  dependence of spontaneous release is generated.

The V-ATPase is a proton pump consisting of two sub-complexes that can reversibly dissociate from each other: The membrane-integral  $V_0$  sector and the peripheral  $V_1$  sector, which carries the ATPase activity.  $V_0$  contains a cylinder of proteolipids (consisting of highly homologous isoforms of subunit c); the large, membrane-integral subunit a; and the peripheral subunit d and subunit e (Fig. 1 A). V-ATPase provides the electrochemical potential that contributes to loading of the secretory vesicles with neurotransmitters. V-ATPase-dependent luminal acidification can influence protein trafficking, e.g., between endolysosomal compartments (Wada et al., 2008; Huotari and Helenius, 2011). In addition, *in vivo* evidence points to a physical role of  $V_0$  in exocytosis and membrane fusion, which is independent of proton pumping. Observations that have uncovered a function for  $V_0$  in exocytosis and membrane fusion stem from regulated secretion in both *Drosophila melanogaster* (Hiesinger et al., 2005) and

mammalian cells (Sun-Wada et al., 2006; Di Giovanni et al., 2010), secretion of multivesicular bodies in *Caenorhabditis elegans* (Liégeois et al., 2006), phagosome-lysosome fusion in the zebrafish *Danio rerio* (Peri and Nüsslein-Volhard, 2008), and vacuole fusion in yeast (Bayer et al., 2003; Strasser et al., 2011). A recent study challenged a physical role for  $V_0$  in vacuole fusion *in vivo* (Coonrod et al., 2013); however, the main assay in that study suffered from a conceptual flaw, i.e., it could not measure vacuole fusion but only biosynthetic transport of the indicator proteins to the vacuoles. At the same time, another study provided a compelling *in vivo* demonstration that a lack of vesicle acidification impairs exocytosis, but that it does so by reducing the pool of free  $V_0$  sectors and thus impeding their physical function in the fusion process (Poëa-Guyon et al., 2013).


$V_0$  subunits interact with Calmodulin (Peters et al., 2001; Zhang et al., 2008) and with Q- and R-SNAREs (Galli et al., 1996; Peters et al., 2001; Takeda et al., 2008; Di Giovanni et al., 2010). In this issue, Wang et al. provide compelling evidence that Calmodulin regulates SNARE complex assembly via  $V_0$ . They used the  $V_0$  subunit a from *Drosophila* (v100) with point mutations in its Calmodulin binding site to selectively disrupt the interaction of Calmodulin and v100. This allele (v100<sup>WFI</sup>) rescues most defects resulting from the loss of v100 (Hiesinger et al., 2005; Williamson et al., 2010), notably endolysosomal acidification and endolysosomal protein sorting. Thus, v100<sup>WFI</sup> retains its functionality as part of the V-ATPase proton pump. v100<sup>WFI</sup> rescues evoked neurotransmitter release but it diminishes spontaneous transmitter release by >90%. An extensive biochemical characterization revealed that v100 disrupts the assembly of Q-SNARE complexes by competitively binding to Syntaxin1A and SNAP-25.  $\text{Ca}^{2+}$ -Calmodulin can disrupt the competitive interactions of v100 with the SNAREs, permit the Q-SNAREs to form a complex, and also incorporate the R-SNARE VAMP and catalyze fusion. These results indicate that v100, when associated with Calmodulin, can serve as a regulator of SNARE complex formation (Fig. 1 B).

This finding adds a novel aspect to the role of  $V_0$  in exocytosis. However, other known effects and interactions of  $V_0$  subunits illustrate that regulating SNARE complex formation cannot be the only way in which  $V_0$  influences membrane fusion and

© 2014 Vavassori and Mayer This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

Correspondence to Andreas Mayer: [Andreas.Mayer@unil.ch](mailto:Andreas.Mayer@unil.ch)

**Figure 1. The V-ATPase  $V_0$  sector affects trans-SNARE pairing and lipid mixing via its subunits a and c.** (A) The V-ATPase is composed of a peripheral sector  $V_1$  (blue, 10 subunits) and a membrane integral sector  $V_0$  (red and yellow).  $V_0$  contains a cylinder of proteolipids (subunits c, yellow), the proton-conducting subunit a, and subunits d and e (red).  $V_1$  and  $V_0$  can dissociate from each other in a regulated and reversible fashion. This equilibrium can be influenced by SNAREs. B and C show two roles of  $V_0$  in membrane fusion, which are not mutually exclusive. The emphasis is on subunit a or subunits c, respectively. (B) Regulation of SNARE complex formation by subunit a. Binding of subunit a to  $\text{Ca}^{2+}$ -Calmodulin (CaM, green) alleviates the block of SNARE complex assembly caused by the interaction of subunit a with the Q-SNARE. (C) Model of how proteolipids (yellow) might enhance the capacity of SNAREs to stimulate lipid mixing. Stimulated by their interaction with trans-SNARE complexes, the proteolipids undergo a conformational change, which might expose hydrophobic surfaces between the proteolipid subunits and facilitate lipid reorientation and membrane fusion (Strasser et al., 2011).  $V_0$  subunits e and d are not shown because there are no data implicating them in fusion.  $V_0$  subunit a is not shown for the sake of clarity.



exocytosis. If  $V_0$  subunit a were only a negative regulator of SNARE complex formation, as suggested by the results of Wang et al. (2014), its deletion should leave fusion intact or even stimulate it. The opposite is the case, which suggests that  $V_0$  also serves to promote fusion. This is also evident from the effects of mutations in other  $V_0$  subunits, such as the central ring of proteolipids (subunit c), which interfere with fusion. Proteolipid rings can adopt at least two conformations (Clare et al., 2006) and they can form  $\text{Ca}^{2+}$ -inducible pores in the membrane that are permeable to hydrophilic molecules (Morel, 2003). Single amino acid substitutions in proteolipid transmembrane domains and proteolipid fusion proteins impede lipid mixing, but, unlike  $v100^{\text{WFI}}$ , they permit the formation of normal levels of trans-SNARE complexes (Strasser et al., 2011). These effects could only be explained by a conformational change in  $V_0$ . Therefore, the current working model postulates that  $V_0$  proteolipids can exist in at least two conformations: one conducive to fusion and one supporting V-ATPase assembly and proton pumping (Strasser et al., 2011; Poëa-Guyon et al., 2013).

According to this model, the molecular interactions of  $V_0$  with SNAREs (Peters et al., 2001; Di Giovanni et al., 2010) might influence  $V_0$  conformation and/or compete with the attachment of  $V_1$  to  $V_0$  (Fig. 1 A). The recruitment of  $V_0$  into the  $V_1$ - $V_0$  V-ATPase holoenzyme can inhibit exocytosis by depleting the pool of

free  $V_0$  sectors that are necessary to support fusion (Poëa-Guyon et al., 2013). Experimental observations from yeast support a role for SNAREs in regulating this pool of free  $V_0$ : Deletion of the vacuolar R-SNARE NYV1 (a Synaptobrevin homologue in yeast) increases  $V_0$ - $V_1$  association, and NYV1 overexpression reduces it (Strasser et al., 2011). Furthermore, structural data suggest that there is substantial space between the subunits of the proteolipid cylinder that might be invaded by lipids (Clare et al., 2006). These data gave rise to a hypothesis explaining how  $V_0$  might promote membrane fusion. In this hypothesis, it is assumed that SNAREs destabilize the  $V_0$ - $V_1$  interaction and favor a  $V_0$  conformation that supports fusion by allowing lipids to invade the space between subunits of the proteolipid cylinder, facilitate their reorientation, and thereby promote the merger of the membrane leaflets (Fig. 1 C). The role of  $v100$  in SNARE complex assembly that Wang et al. (2014) describe is not mutually exclusive with a role of  $V_0$  in lipid reorientation (Fig. 1, B and C). It adds an interesting new aspect of  $V_0$  function and suggests that  $V_0$  is intimately linked with the membrane fusion apparatus. This illustrates that much remains to be discovered before we will fully understand the mechanisms by which  $V_0$  subunits regulate and promote the different stages of the fusion process.

Submitted: 10 March 2014  
Accepted: 25 March 2014

## References

- Bayer, M.J., C. Reese, S. Buhler, C. Peters, and A. Mayer. 2003. Vacuole membrane fusion: V<sub>0</sub> functions after trans-SNARE pairing and is coupled to the Ca<sup>2+</sup>-releasing channel. *J. Cell Biol.* 162:211–222. <http://dx.doi.org/10.1083/jcb.200212004>
- Clare, D.K., E.V. Orlova, M.A. Finbow, M.A. Harrison, J.B.C. Findlay, and H.R. Saibil. 2006. An expanded and flexible form of the vacuolar ATPase membrane sector. *Structure.* 14:1149–1156. <http://dx.doi.org/10.1016/j.str.2006.05.014>
- Coonrod, E.M., L.A. Graham, L.N. Carpp, T.M. Carr, L. Stirrat, K. Bowers, N.J. Bryant, and T.H. Stevens. 2013. Homotypic vacuole fusion in yeast requires organelle acidification and not the V-ATPase membrane domain. *Dev. Cell.* 27:462–468. <http://dx.doi.org/10.1016/j.devcel.2013.10.014>
- Di Giovanni, J., S. Boudkazi, S. Mochida, A. Bialowas, N. Samari, C. Lévêque, F. Youssouf, A. Brechet, C. Iborra, Y. Maulet, et al. 2010. V-ATPase membrane sector associates with synaptobrevin to modulate neurotransmitter release. *Neuron.* 67:268–279. <http://dx.doi.org/10.1016/j.neuron.2010.06.024>
- Galli, T., P.S. McPherson, and P. De Camilli. 1996. The V<sub>0</sub> sector of the V-ATPase, synaptobrevin, and synaptophysin are associated on synaptic vesicles in a Triton X-100-resistant, freeze-thawing sensitive, complex. *J. Biol. Chem.* 271:2193–2198. <http://dx.doi.org/10.1074/jbc.271.4.2193>
- Hiesinger, P.R., A. Fayyazuddin, S.Q. Mehta, T. Rosenmund, K.L. Schulze, R.G. Zhai, P. Verstreken, Y. Cao, Y. Zhou, J. Kunz, and H.J. Bellen. 2005. The v-ATPase V0 subunit a1 is required for a late step in synaptic vesicle exocytosis in *Drosophila*. *Cell.* 121:607–620. <http://dx.doi.org/10.1016/j.cell.2005.03.012>
- Huotari, J., and A. Helenius. 2011. Endosome maturation. *EMBO J.* 30:3481–3500. <http://dx.doi.org/10.1038/emboj.2011.286>
- Kochubey, O., X. Lou, and R. Schneggenburger. 2011. Regulation of transmitter release by Ca<sup>2+</sup> and synaptotagmin: insights from a large CNS synapse. *Trends Neurosci.* 34:237–346. <http://dx.doi.org/10.1016/j.tins.2011.02.006>
- Liégeois, S., A. Benedetto, J.-M. Garnier, Y. Schwab, and M. Labouesse. 2006. The V0-ATPase mediates apical secretion of exosomes containing Hedgehog-related proteins in *Caenorhabditis elegans*. *J. Cell Biol.* 173:949–961. <http://dx.doi.org/10.1083/jcb.200511072>
- Morel, N. 2003. Neurotransmitter release: the dark side of the vacuolar-H+ATPase. *Biol. Cell.* 95:453–457. [http://dx.doi.org/10.1016/S0248-4900\(03\)00075-3](http://dx.doi.org/10.1016/S0248-4900(03)00075-3)
- Peri, F., and C. Nüsslein-Volhard. 2008. Live imaging of neuronal degradation by microglia reveals a role for v0-ATPase a1 in phagosomal fusion in vivo. *Cell.* 133:916–927. <http://dx.doi.org/10.1016/j.cell.2008.04.037>
- Peters, C., M.J. Bayer, S. Bühler, J.S. Andersen, M. Mann, and A. Mayer. 2001. Trans-complex formation by proteolipid channels in the terminal phase of membrane fusion. *Nature.* 409:581–588. <http://dx.doi.org/10.1038/35054500>
- Poëa-Guyon, S., M.R. Ammar, M. Erard, M. Amar, A.W. Moreau, P. Fossier, V. Gleize, N. Vitale, and N. Morel. 2013. The V-ATPase membrane domain is a sensor of granular pH that controls the exocytotic machinery. *J. Cell Biol.* 203:283–298. <http://dx.doi.org/10.1083/jcb.201303104>
- Strasser, B., J. Iwaszkiewicz, O. Michielin, and A. Mayer. 2011. The V-ATPase proteolipid cylinder promotes the lipid-mixing stage of SNARE-dependent fusion of yeast vacuoles. *EMBO J.* 30:4126–4141. <http://dx.doi.org/10.1038/emboj.2011.335>
- Sun-Wada, G.-H., T. Toyomura, Y. Murata, A. Yamamoto, M. Futai, and Y. Wada. 2006. The a3 isoform of V-ATPase regulates insulin secretion from pancreatic β-cells. *J. Cell Sci.* 119:4531–4540. <http://dx.doi.org/10.1242/jcs.03234>
- Takeda, K., M. Cabrera, J. Rohde, D. Bausch, O.N. Jensen, and C. Ungermann. 2008. The vacuolar V1/V0-ATPase is involved in the release of the HOPS subunit Vps41 from vacuoles, vacuole fragmentation and fusion. *FEBS Lett.* 582:1558–1563. <http://dx.doi.org/10.1016/j.febslet.2008.03.055>
- Wada, Y., G.-H. Sun-Wada, H. Tabata, and N. Kawamura. 2008. Vacuolar-type proton ATPase as regulator of membrane dynamics in multicellular organisms. *J. Bioenerg. Biomembr.* 40:53–57. <http://dx.doi.org/10.1007/s10863-008-9128-z>
- Wang, D., D. Epstein, O. Khalaf, S. Srinivasan, W.R. Williamson, A. Fayyazuddin, F.A. Quijcho, and P.R. Hiesinger. 2014. Ca<sup>2+</sup>-Calmodulin regulates SNARE assembly and spontaneous neurotransmitter release via v-ATPase subunit V0a1. *J. Cell Biol.* 205:21–31. <http://dx.doi.org/10.1083/jcb.201312109>
- Williamson, W.R., D. Wang, A.S. Haberman, and P.R. Hiesinger. 2010. A dual function of V0-ATPase a1 provides an endolysosomal degradation mechanism in *Drosophila melanogaster* photoreceptors. *J. Cell Biol.* 189:885–899. <http://dx.doi.org/10.1083/jcb.201003062>
- Zhang, W., D. Wang, E. Volk, H.J. Bellen, P.R. Hiesinger, and F.A. Quijcho. 2008. V-ATPase V0 sector subunit a1 in neurons is a target of calmodulin. *J. Biol. Chem.* 283:294–300. <http://dx.doi.org/10.1074/jbc.M708058200>