

RESEARCH ARTICLE

Open Access

Phospholipase A₂ inhibitors protect against prion and A β mediated synapse degeneration

Clive Bate*¹, Mourad Tayebi¹ and Alun Williams^{1,2}

Abstract

Background: An early event in the neuropathology of prion and Alzheimer's diseases is the loss of synapses and a corresponding reduction in the level of synaptophysin, a pre-synaptic membrane protein essential for neurotransmission. The molecular mechanisms involved in synapse degeneration in these diseases are poorly understood. In this study the process of synapse degeneration was investigated by measuring the synaptophysin content of cultured neurones incubated with the prion derived peptide (PrP^{Sc}) or with A β ₁₋₄₂, a peptide thought to trigger pathogenesis in Alzheimer's disease. A pharmacological approach was used to screen cell signalling pathways involved in synapse degeneration.

Results: Pre-treatment with phospholipase A₂ inhibitors (AACOCF₃, MAFP and aristolochic acids) protected against synapse degeneration in cultured cortical and hippocampal neurones incubated with PrP^{Sc} or A β ₁₋₄₂. Synapse degeneration was also observed following the addition of a specific phospholipase A₂ activating peptide (PLAP) and the addition of PrP^{Sc} or A β ₁₋₄₂ activated cytoplasmic phospholipase A₂ within synapses. Activation of phospholipase A₂ is the first step in the generation of platelet-activating factor (PAF) and PAF receptor antagonists (ginkgolide B, Hexa-PAF and CV6029) protected against synapse degeneration induced by PrP^{Sc}, A β ₁₋₄₂ and PLAP. PAF facilitated the production of prostaglandin E₂, which also caused synapse degeneration and pre-treatment with the prostanoid E receptor antagonist AH13205 protected against PrP^{Sc}, A β ₁₋₄₂ and PAF induced synapse degeneration.

Conclusions: Our results are consistent with the hypothesis that PrP^{Sc} and A β ₁₋₄₂ trigger abnormal activation of cytoplasmic phospholipase A₂ resident within synapses, resulting in elevated levels of PAF and prostaglandin E₂ that cause synapse degeneration. Inhibitors of this pathway that can cross the blood brain barrier may protect against the synapse degeneration seen during Alzheimer's or prion diseases.

Background

In the transmissible spongiform encephalopathies, otherwise known as the prion diseases, changes in synaptic function and a reduction in synaptophysin levels within the brain occur at a time before any gross neuronal loss is observed [1-3]. These synaptic alterations are associated with the accumulation of a differentially folded, and protease-resistant isoform (PrP^{Sc}), of the host encoded cellular prion protein (PrP^C) [4]. The formation of PrP^{Sc} is accompanied by a decreased expression of proteins

involved in exocytosis and neurotransmission, such as synaptophysin, SNAP-25 and synapsins in the brains of scrapie-infected mice [2,5] and in humans affected with Creutzfeldt-Jakob disease (CJD) [6].

The molecular mechanisms that underlie synapse degeneration in prion diseases are not understood. Such processes have been examined by incubating cultured neurones with PrP^{Sc} or specific prion-derived peptides. A major PrP fragment spanning amino acid residues 81-82 to 144-153 was isolated from the brains of patients with the hereditary prion disease Gerstmann-Sträussler-Scheinker disease [7]. Synthetic peptides containing amino acid residues 82 to 146 (PrP^{Sc}-146) had similar structural and biochemical properties to PrP^{Sc} suggesting that this

* Correspondence: cbate@rvc.ac.uk¹ Department of Pathology and Infectious Diseases, Royal Veterinary College, Hawkshead Lane, North Mymms, Herts, AL9 7TA, UK

Full list of author information is available at the end of the article

fragment was the neurotoxic species generated in prion diseases. This hypothesis was strengthened by observations that both partially purified PrP^{Sc} preparations and PrP82-146 caused synapse degeneration in cortical and hippocampal neurones [8]. The effect of PrP82-146 on synapses in neuronal cultures was measured using an enzyme linked immunoassay (ELISA) to quantify the amount of synaptophysin [9]. Synaptophysin is a pre-synaptic membrane protein essential for neurotransmitter release and the recycling of synaptic vesicles and hence neurotransmission [10-13]. The amount of synaptophysin has been used to access synaptic density in the brain [14,15] and in cultured neurones [8]. Although immunocytochemistry is commonly used to examine synapse density this method is susceptible to errors in counting and field selection. The use of an ELISA overcame such problems by measuring synaptic density throughout neuronal cultures.

Synaptic failure is also thought to contribute to the neuropathogenesis of Alzheimer's disease (AD) [16] and the loss of synaptic proteins is the best correlate of dementia in AD [14,17-20]. The amyloid hypothesis of AD pathogenesis maintains that the primary event is the production of neurotoxic amyloid- β (A β) peptides following the proteolytic cleavage of the amyloid precursor protein into different fragments [21,22]. These fragments include A β_{1-42} which is widely regarded as the main pathogenic species in AD. Recent studies showed the importance of small soluble oligomers of A β or A β derived diffusible ligands in neurotoxicity [23,24]. In this study we sought to determine whether PrP82-146 and A β induced synapse degeneration was mediated through specific cell signalling pathways. We report that PrP82-146 and A β_{1-42} induced synapse degeneration was prevented by pharmacological inhibition of PLA₂ and that both PrP82-146 and A β_{1-42} peptides increased activation of cytoplasmic phospholipase A₂ (cPLA₂) within synapses suggesting that activation of this enzyme triggers synapse degeneration. This hypothesis was supported by the observation that the synapse degeneration was also observed following the addition of a specific PLA₂ activating peptide (PLAP). Activation of PLA₂ is the first step in the production of bioactive prostaglandins and platelet-activating factor (PAF), specific antagonists of which also reduced PrP82-146 and A β_{1-42} induced synapse degeneration.

Results

PLA₂ inhibitors protected against PrP82-146 induced synapse degeneration

The addition of the prion derived peptide PrP82-146 reduced the synaptophysin content of cortical neurones indicative of a loss of synapses. This effect was a consequence of the specific amino acid sequence of PrP82-146

as a control peptide (PrP82-146scram) did not affect synapses [8]. The synaptophysin content was reduced by greater than 80% by 1 μ M PrP82-146 (Figure 1A) which did not affect neuronal survival as measured by thiazyl blue tetrazolium (98% neuronal survival \pm 6 compared with 100% \pm 6, n = 9, P = 0.53). Immunoblot analysis showed that PrP82-146 caused a dose-related reduction in the amounts of synaptophysin in cell extracts without affecting the amounts of β -actin (Figure 1B).

To determine if the degeneration of synapses resulted from activation of specific cell signalling pathways cortical neurones were pre-treated with inhibitors of some common cell signalling pathways and incubated with PrP82-146. Pre-treatment with the PLA₂ inhibitors (1 μ M AACOCF₃, 1 μ M MAFP or 5 μ g/ml aristolochic acids) protected against the PrP82-146 mediated decrease in synaptophysin (Figure 1A). The concentration of PrP82-146 required to reduce the synaptophysin content of vehicle treated cortical neurones by 50% (EC₅₀) was approximately 60 nM. By comparison, the EC₅₀ of PrP82-146 for cortical neurones treated with 1 μ M AACOCF₃ or 1 μ M MAFP was greater than 10 μ M, and in neurones




Figure 1 PLA₂ inhibitors protected cortical neurones against PrP82-146 induced synapse degeneration. (A) The synaptophysin content of cortical neurones pre-treated for with a vehicle control (■), 1 μ M AACOCF₃ (○), 1 μ M MAFP (●), 5 μ g/ml aristolochic acid (□), 10 μ M U73122 (▲) or 10 μ M ethyl-18-OCH₃ (Δ) and subsequently incubated with varying concentrations of PrP82-146 for 24 hours. Values shown are the mean average amount of synaptophysin \pm standard deviation (SD), n = 15. (B) Immunoblots showing the amount of synaptophysin and β -actin in cell extracts from neurones incubated with PrP82-146 (1.25 - 0.05 μ M).

pre-treated with 5 µg/ml aristolochic acids the EC_{50} was 5 µM. In contrast, pre-treatment with phospholipase C inhibitors (10 µM U73122 or ethyl-18-OCH₃) did not affect the PrP82-146 induced loss of synaptophysin from cortical neurones. The synaptophysin content of neurones was not significantly affected by treatment with 5 µM AACOCF₃ (100 units synaptophysin ± 4 compared to 101 ± 4, n = 12, P = 0.5), 5 µM MAFP (100 ± 4 v 102 ± 6, n = 12, P = 0.4) or 5 µg/ml aristolochic acids (100 ± 4 compared to 98 ± 4, n = 12, P = 0.2) alone showing that these drugs did not stimulate synaptogenesis, nor did they damage synapses.

PLA₂ inhibitors protected hippocampal neurones against PrP82-146 induced synapse degeneration

To determine if the effect of PLA₂ inhibitors was selective for cortical neurones, their effect on hippocampal neurones was also examined. PrP82-146 reduced the amount of synaptophysin in hippocampal neurones; the EC_{50} of PrP82-146 in hippocampal neurones was 10 nM and while the addition of 200 nM PrP82-146 reduced the synaptophysin content to 20% of control cultures, it did not alter cell survival as measured by thiazyl blue tetrazolium (97% cell survival ± 7 compared with 100% ± 9, n = 9, P = 0.7). Pre-treatment with 1 µM AACOCF₃ or 1 µM MAFP prevented the PrP82-146 mediated decrease in the synaptophysin content of hippocampal neurones and in treated neurones the EC_{50} was increased to greater than 1 µM (Figure 2).




Figure 2 PLA₂ inhibitors protected hippocampal neurones against PrP82-146 induced synapse degeneration. The synaptophysin content of cultured hippocampal neurones pre-treated with a vehicle control (■), 1 µM AACOCF₃ (○) or 1 µM MAFP (●) and incubated with varying concentrations of PrP82-146 for 24 hours. Values shown are the mean average amount of synaptophysin ± SD, n = 9.

PLA₂ inhibitors protected against synapse degeneration induced by Aβ₁₋₄₂

The synapse degeneration that occurs in AD is thought to be caused by Aβ₁₋₄₂ [25,26]. This process can be modelled *in vitro* and the addition of Aβ₁₋₄₂, but not Aβ₄₂₋₁, reduced the synaptophysin content of cultured cortical and hippocampal neurones [27]. Here we report that pre-treatment with 1 µM AACOCF₃ or 1 µM MAFP protected cortical neurones against the Aβ₁₋₄₂ induced reduction in synaptophysin content (Figure 3A). While the EC_{50} of Aβ₁₋₄₂ in vehicle treated cortical neurones was 50 nM, the EC_{50} of Aβ₁₋₄₂ for cortical neurones treated with 1 µM




Figure 3 PLA₂ inhibitors protected against Aβ₁₋₄₂ induced synapse degeneration. (A) The synaptophysin content of cortical neurones pre-treated with a vehicle control (■), 1 µM AACOCF₃ (○) or 1 µM MAFP (●) and incubated with varying concentrations of Aβ₁₋₄₂ for 24 hours. Values shown are the mean average amount of synaptophysin ± SD, n = 12. (B) The synaptophysin content of hippocampal neurones pre-treated with a vehicle control (■) or 1 µM AACOCF₃ (○) and incubated with varying concentrations of Aβ₁₋₄₂ for 24 hours. Values shown are the mean average amount of synaptophysin ± SD, n = 6.

AACOFC₃ was greater than 10 μM. PLA₂ inhibitors also protected hippocampal neurones against Aβ₁₋₄₂ induced synapse degeneration. Thus, the EC₅₀ of Aβ₁₋₄₂ for vehicle treated hippocampal neurones was 10 nM, while in hippocampal neurones pre-treated with 1 μM AACOCF₃ the EC₅₀ of Aβ₁₋₄₂ was increased to greater than 1 μM (Figure 3B).

Conditioned medium from 7PA2 cells (7PA2-CM), which contains naturally secreted Aβ oligomers [28], also reduced the synaptophysin content of cortical neurones, whereas conditioned medium from non-transfected Chinese hamster ovary cells (CHO-CM), which did not contain Aβ oligomers, had no affect (Figure 4A). Pre-treatment of cortical neurones with either 1 μM

AACOFC₃ or 1 μM MAFP prevented the 7PA2-CM mediated decrease in synaptophysin (Figure 4B).

PLA₂ inhibitors did not affect the accumulation of PrP82-146 in synapses

Prior studies showed that PrP82-146 accumulated within synaptosomes isolated from cortical neurones prior to synaptic degeneration [8]. That observation raised the possibility that the protective effect of PLA₂ inhibitors might be because they reduced the binding/accumulation of PrP82-146 within synapses. Firstly, we showed that the amount of synaptophysin was not significantly different between synaptosomes isolated from cortical neurones and neurones treated 1 μM AACOCF₃ or 1 μM MAFP; indicating that these drugs did not affect synapse formation. Next cortical neurones were pre-treated with 1 μM AACOCF₃ before the addition of 100 nM PrP82-146. The amount of PrP82-146 found within synaptosomes isolated from vehicle and AACOCF₃ treated neurones were not significantly different (Figure 5). Similar results were obtained with synaptosomes from cortical neurones pre-treated with 1 μM MAFP indicating that the PLA₂ inhibitors did not prevent the binding or transport of PrP82-146 to synapses.




Figure 4 PLA₂ inhibitors protected against 7PA2-CM induced synapse degeneration. (A) The synaptophysin content of cortical neurones incubated for 24 hours with 7PA2-CM (□) or CHO-CM (■). Values shown are the mean average amount of synaptophysin ± SD, n = 12. (B) The synaptophysin content of cortical neurones incubated for pre-treated with a vehicle control (□), 1 μM AACOCF₃ (■) or 1 μM MAFP (striped bars) and incubated with doubling dilutions of 7PA2-CM for 24 hours. Values shown are the mean average amount of synaptophysin ± SD, n = 12.




Figure 5 PLA₂ inhibitors did not affect the accumulation of PrP82-146 in synapses. The amount of PrP82-146 detected in synaptosomes isolated from cortical neurones incubated with a vehicle control (■), 1 μM AACOCF₃ (□) or 1 μM MAFP (striped bars). Treated cortical neurones were incubated with 100 nM PrP82-146 for the time periods shown. Synaptosomes were isolated from neuronal cultures and the amount of PrP82-146 was measured by ELISA. Values shown are the mean average amount of PrP82-146 (nM) ± SD, n = 9.

PLA₂ inhibitors protected against synapse degeneration induced by PLAP

Next we sought to determine whether synapse degeneration resulted from activation of an endogenous PLA₂. The addition of PLAP, a peptide that activated endogenous cPLA₂ [29,30], had similar effects as PrP82-146 and Aβ₁₋₄₂ on synapses; it reduced the synaptophysin content of cortical neurones in a dose dependent manner. The addition of 1 μM PLAP reduced the synaptophysin content of cultured neurones by approximately 80% without affecting the survival of neurones as measured by thiazyl blue tetrazolium (96% neuronal survival ± 5 compared with 100% ± 8, n = 9, P = 0.7). Pre-treatment with 1 μM AACOCF₃ or 1 μM MAFP protected cortical neurones against PLAP induced synapse degeneration; in vehicle treated neurones the EC₅₀ of PLAP was 100 nM, while in neurones pre-treated with 1 μM AACOCF₃ or 1 μM MAFP the addition of 1 μM PLAP did not reduce the synaptophysin content by more than 20% (Figure 6).

cPLA₂ is enriched in synaptosomes

To investigate the hypothesis that synapse degeneration resulted from activation of an endogenous cPLA₂, the amount of cPLA₂ protein in synaptosomes was measured. The cPLA₂ protein was enriched in synaptosomes isolated from cortical neurones compared to whole cell membrane extracts (100 units cPLA₂/mg protein ± 31 compared to 6 units/mg ± 1, n = 12, P < 0.01). There were

no significant differences in the amount of total cPLA₂ protein in synaptosomes isolated from vehicle treated cortical neurones and those treated for 1 hour with 200 nM PrP82-146, 200 nM PrP82-146scrambled, 1 μM PLAP, 100 nM Aβ₁₋₄₂ or 100 nM Aβ₄₂₋₁ (Table 1). Measurements were made after 1 hour, before any synapse degeneration was observed. Next specific antibodies (to cPLA₂ phosphorylated at serine 505) were used to determine the amount of activated cPLA₂ within synaptosomes. The amount of activated cPLA₂ within synaptosomes was significantly increased following the addition of 200 nM PrP82-146, 1 μM PLAP or 100 nM Aβ₁₋₄₂ but not after the addition of PrP82-146scrambled or Aβ₄₂₋₁ (Table 1). Moreover, there was a significant inverse correlation between the amount of activated cPLA₂ in synaptosomes after 1 hour and the amount of synaptophysin in neurones after 24 hours following the addition of different amounts of PrP82-146 (range 1.25 to .005 μM), Pearson's coefficient = -0.76, P < 0.01 (Figure 7).

PAF antagonists protected against PrP82-146 induced synapse degeneration

The activation of PLA₂ is the first step in the production of PAF [31] that has been shown to cause synapse degeneration *in vitro* [27]. The addition of PAF receptor antagonists (Hexa-PAF, CV6209 or ginkgolide B), in the range 0.1-10 μM, did not affect the amount of synaptophysin in cortical neurones. However, pre-treatment with 2 μM Hexa-PAF, 2 μM CV6209 or 1 μM ginkgolide B provided protection against PrP82-146 induced synapse degeneration (Figure 8). The EC₅₀ of PrP82-146 in vehicle treated cortical neurones was 50 nM while the EC₅₀ of PrP82-146 for neurones treated with 2 μM Hexa-PAF was 2 μM, for neurones treated with 2 μM CV6209 the EC₅₀ was 5 μM and for neurones pre-treated with 1 μM ginkgolide B the EC₅₀ was 20 μM. Pre-treatment with Hexa-PAF, ginkgolide or CV6209 also prevented the reduction in the synaptophysin content of cortical neurones incubated with 100 nM Aβ₁₋₄₂ or 1 μM PLAP (Table 2). In a similar manner, the amount of synaptophysin in hippocampal neurones incubated with 50 nM PrP82-146 was significantly increased by pre-treatment with 1 μM ginkgolide B (97 units synaptophysin ± 9 compared to 18 ± 7, n = 6, P < 0.01), 2 μM Hexa-PAF (94 ± 11 compared to 18 ± 7, n = 6, P < 0.01) or 2 μM CV6209 (94 ± 11 compared to 18 ± 7, n = 6, P < 0.01).

Prostaglandin E₂ induced synapse degeneration

Prior studies showed that PAF facilitates the production of prostaglandins [32] suggesting that one or more of the prostaglandins produced in response to PrP82-146, Aβ₁₋




Figure 6 PLA₂ inhibitors protected against PLAP induced synapse degeneration. The synaptophysin content of cortical neurones pre-treated with a vehicle control (■), 1 μM AACOCF₃ (○) or 1 μM MAFP (●) and incubated with varying concentrations of PLAP for 24 hours. Values shown are the mean average amount of synaptophysin ± SD, n = 12.


Table 1: PrP82-146 and A β ₁₋₄₂ activated cPLA₂ within synapses.

Treatment	Total cPLA ₂ protein (units/mg protein)	Activated cPLA ₂ (units/mg protein)
None (units)	100 ± 31	100 ± 19
200 nM PrP82-146	106 ± 18	773 ± 51 *
200 nM PrP82-146scrambled	95 ± 22	111 ± 34
1 μ M PLAP	119 ± 30	928 ± 84 *
100 nM A β ₁₋₄₂	109 ± 21	655 ± 62 *
100 nM A β ₄₂₋₁	92 ± 27	121 ± 34

Primary cortical neurones were incubated with PrP, A β or PLAP peptides as shown for 1 hour. Synaptosomes were subsequently isolated from treated neurones and the amount of cPLA₂ protein, and the amount of activated cPLA₂ (phosphorylated at serine 505) were measured by ELISA. Values shown are the mean average amount of cPLA₂ protein or activated cPLA₂ (units/mg protein) \pm SD, n = 12. * = amount of activated cPLA₂ significantly greater than in control synaptosomes.

₄₂ or PLAP are responsible for synapse degeneration. This hypothesis was tested by treating cortical neurones with individual prostaglandins. We report that the addition of prostaglandin E₂, but not other prostaglandins reduced the amount of synaptophysin in cortical neurones (Figure 9). Prostaglandin E₂, acts via specific prostanoid E receptors [33] and pre-treatment with the prostanoid E recep-

tor antagonist AH13205, but not the prostanoid D receptor antagonist BWA868C, prevented the loss of synaptophysin in cortical neurones incubated with PrP82-146, A β ₁₋₄₂, PLAP, PAF or prostaglandin E₂ (Table 3). Taken together, these results show that the effects of PrP82-146 or A β ₁₋₄₂ on synapses were ultimately mediated through prostanoid E receptors.



Discussion

The addition of PrP82-146 reduced the synaptophysin content of cortical neurones indicative of a loss of synapses. This occurred at concentrations of peptide that did not affect neuronal survival [8], results consistent with *in vivo* observations that the loss of synapses in prion diseases occurs before any significant neuronal death is seen [1,2]. PrP82-146 also damaged synapses in hippocampal neuronal cultures. The hippocampus is involved in memory formation and the loss of synapses in this area is consistent with the cognitive deficits that occur during the early stages of scrapie infections [3,34].

A pharmacological approach was used to examine the molecular mechanisms involved in prion induced synapse degeneration. Critically, pre-treatment with selective PLA₂ inhibitors reduced PrP82-146 induced synapse degeneration, whereas pre-treatment with inhibitors of some other common cell signalling pathways including phospholipase C had no affect. PrP^{Sc} is detected within synapses in CJD patients and experimental scrapie infections [2,35,36] and PrP82-146 was found within synaptosomes isolated from cultured neurones [8]. We found that pre-treatment with PLA₂ inhibitors did not alter the

Table 2: PAF receptor antagonists reduced synapse degeneration induced by A β ₁₋₄₂ or PLAP.

Synaptophysin (units/10 ⁶ cells)				
Treatment	Vehicle	Hexa-PAF	Ginkgolide B	CV6209
Vehicle	100 ± 6	98 ± 7	102 ± 5	96 ± 8
100 nM A β ₁₋₄₂	32 ± 6	94 ± 6	98 ± 9	89 ± 8
1 μ M PLAP	27 ± 7	95 ± 8	94 ± 8	82 ± 7

Primary cortical neurones were pre-treated with a vehicle control, or with PAF receptor antagonists (2 μ M Hexa-PAF, 1 μ M ginkgolide or 2 μ M CV6209) for 1 hour prior to incubation with A β ₁₋₄₂ or PLAP as shown for 24 hours. Values shown are the mean average amount of synaptophysin (units/10⁶ cells) \pm SD, n = 12.

amount of PrP82-146 found within synaptosomes, indicating the protective effect of these drugs was not due to the reduced binding of PrP82-146 to synaptic receptors, or a reduction in the trafficking of PrP82-146 to synapses.

Next, the possibility that PrP82-146 might activate cPLA₂ resident within synapses was examined. Previous studies reported that PLA₂ was found at synapse and regulated synaptic vesicle recycling [37,38] and we observed that cPLA₂ was highly enriched in synaptosomes. These observations imply that cPLA₂ is part of the normal function of synapses and complete inhibition of cPLA₂ may not be desirable as it may interfere with neurotransmission. However, the addition of PrP82-146 increased the

amount of activated cPLA₂ within synaptosomes nearly 8 fold suggesting that PrP82-146 induced activation of cPLA₂ resident within synapses was responsible for synapse degeneration. This hypothesis was supported by our observations that there was a significant inverse correlation between the amount of activated cPLA₂ and synaptic density and that synapse degeneration was also observed after the addition of PLAP. In addition, the effect of both PLAP and PrP82-146 on synapses was greatly reduced by pre-treatment with the cPLA₂ inhibitors AACOCF₃ and MAFP. Collectively, these results strongly suggesting that activation of an endogenous cPLA₂ is a pivotal event in PrP82-146 mediated synapse degeneration.

The activation of PLA₂ leads to the formation of a number of bioactive factors including prostaglandins, leucotrienes, docosanoids and PAF [31,39]. PAF receptors are present at synaptic endings [40] and PAF facilitates neurotransmission [41]. However, higher concentrations of PAF are implicated in the neurotoxicity of glutamate, human immunodeficiency virus infection, prion diseases and AD [39,42]. Reports that synaptic activity becomes excitotoxic in neurones exposed to elevated levels of PAF [43] may explain the role of PAF in these diseases. Here we showed that synapse degeneration induced by PrP82-146 was reduced by pre-treatment with PAF receptor antagonists, ginkgolide B, Hexa-PAF and CV6209. These PAF receptor antagonists also reduced synapse degeneration induced by PLAP suggesting that the effect of PrP82-146 and PLAP were a secondary effect mediated by activation of an endogenous cPLA₂ and the production of PAF.

PAF affects the production of prostaglandin E₂ [32] which is found in close association with PrP^{Sc} deposition and neuronal degeneration in scrapie [44]. In addition, the levels of prostaglandin E₂ are raised in the cerebrospinal fluid of patients with CJD [45]. While prostaglandin




Figure 8 PAF receptor antagonists protected against PrP82-146 induced synapse degeneration. The synaptophysin content of cortical neurones pre-treated with a vehicle control (■), 1 μ M ginkgolide B (○), 2 μ M Hexa-PAF (●) or 2 μ M CV6209 (□) and subsequently incubated with varying concentrations of PrP82-146 for 24 hours. Values shown are the mean average amount of synaptophysin \pm SD, n = 12.

Table 3: A prostanoid E receptor antagonist protected against synapse degeneration

Treatment	Synaptophysin (units)		
	Vehicle control	Prostanoid E receptor antagonist AH13205	Prostanoid D receptor antagonist BWA868C
Vehicle control	100 ± 6	99 ± 7	99 ± 3
200 nM PrP82-146	22 ± 5	87 ± 8	30 ± 7
100 nM Aβ ₁₋₄₂	32 ± 6	93 ± 5	34 ± 6
1 μM PLAP	27 ± 7	91 ± 7	38 ± 6
10 nM PAF	31 ± 4	85 ± 9	32 ± 5
10 nM Prostaglandin E ₂	30 ± 5	97 ± 3	33 ± 6

Primary cortical neurons were pre-treated with a vehicle control, a prostanoid E receptor antagonist (100 nM AH13205) or a prostanoid D receptor antagonist (100 nM BWA868C) for 1 hour prior to incubation with PrP82-146, Aβ₁₋₄₂, PLAP, PAF or prostaglandin E₂ as shown for 24 hours. Values shown are the mean average amount of synaptophysin ± SD, n = 9.

E₂ may modify hippocampal synaptic transmission via a pre-synaptic prostanoid E receptors [46], higher concentrations cause synapse degeneration [27]. As shown in Figure 9 the addition of prostaglandin E₂, but not other prostaglandins, caused synapse degeneration. The effects of prostaglandin E₂ are mediated by specific membrane

bound prostanoid E receptors [47] and synapse degeneration induced by PrP82-146 was reduced by pre-treatment with the prostanoid E receptor antagonist AH13205 suggesting that its effect was mediated through the prostanoid E receptor.

The results presented here may also be pertinent for other diseases in which synapse degeneration is common. For example, the neurotoxicity of Aβ₁₋₄₂, widely believed to be the main toxin responsible for neurodegeneration in AD [23,48-50], was reduced following knockdown on cPLA₂ [51,52]. In this study we showed that Aβ₁₋₄₂ strongly activated cPLA₂ within synapses and that pre-treatment with PLA₂ inhibitors protected against the synapse degeneration induced by Aβ₁₋₄₂ and naturally secreted Aβ oligomers (7PA2-CM). This observation is consistent with reports that inhibition of cPLA₂ protected against cognitive impairment in a mouse model of AD [53]. A recent study indicated that PrPC acts as a receptor for Aβ₁₋₄₂ oligomers [54]. PrPC also acts as a receptor for PrP peptides [55] suggesting that both PrP and Aβ₁₋₄₂ activate PLA₂ through their interaction with PrPC. We note that oligomers, rather than monomers, of PrP or Aβ₁₋₄₂ are neurotoxic [56-58]. Since oligomers of PrP and Aβ₁₋₄₂, but not monomers, have the capacity to cross-link PrPC, our results support the hypothesis that it is the cross-linkage of PrPC by PrP or Aβ₁₋₄₂ oligomers that activated cPLA₂ leading to the production of PAF and prostaglandins. We propose that it is the persistent activation of cPLA₂ by PrP82-146 or Aβ₁₋₄₂ that results in high concen-




Figure 9 Prostaglandin E₂ caused synapse degeneration. The synaptophysin content of cortical neurons treated with varying concentrations of prostaglandins E₂ (●) D₂ (○), F_{2a} (□), I₂ (■) or 15d-J₂ (▲) for 24 hours. Values shown are the mean average amount of synaptophysin ± SD, n = 9.

trations of PAF and prostaglandin E₂ that cause synapse degeneration [27,43].

Conclusion

Drugs that protect the synapse provide a rational strategy to treat many neurodegenerative diseases. In this study we showed that PLA₂ inhibitors, PAF receptor antagonists and a prostanoid E receptor antagonist all protected against the synapse degeneration induced by PrP82-146 or Aβ₁₋₄₂ *in vitro*. These peptides did not have a direct effect on synapses; rather synapse degeneration was caused after PrP82-146 or Aβ₁₋₄₂ induced activation of an endogenous cPLA₂ and the production of PAF and prostaglandin E₂. Such results suggest that specific cPLA₂ inhibitors or PAF receptor antagonists that are able to cross the blood-brain barrier should be considered for further testing in animal models of prion and Alzheimer's diseases.

Methods

Primary neuronal cultures

Primary cortical neurones were prepared from the brains of mouse embryos (day 15.5). Neuronal precursors were plated at 2×10^5 cells/well in 48 well plates in Ham's F12 containing 5% foetal calf serum (FCS) for 2 hours. Cultures were shaken (600 r.p.m for 5 minutes) and non-adherent cells removed by 3 washes in phosphate buffered saline (PBS). Neurones were grown in neurobasal medium (NBM) containing B27 components (PAA) for 7 days. Immunohistochemistry showed that the cells were greater than 97% neurofilament positive. Less than 3% stained for GFAP (astrocytes) or for F4/80 (microglial cells). Hippocampal neurones were prepared from the brains of adult mice as described [59]. Briefly, hippocampi were dissected and triturated in Ham's F12 containing 5% FCS, 0.35% glucose, 0.025% trypsin, and 0.1% type IV collagenase. After 30 minutes at 37°C, the cells were triturated again and the cell suspension was passed through a 100 μm cell strainer. Cells were collected by centrifugation, washed twice in Ham's F12 containing 5% FCS and plated at 2×10^5 cells/well in 48 well plates for 24 hours. Cultures were shaken (600 r.p.m for 5 minutes) to remove non-adherent cells, washed twice with PBS and neurones were cultured in NBM containing B27 components and 10 ng/ml glial-derived neurotrophic factor (Sigma) for 10 days. Neurones were incubated with test compounds for 3 hours before the addition of peptides and the amount of synaptophysin in treated neurones was measured 24 hours later. The survival of neurones was determined 5 days later using 25 μM thiazyl blue tetrazolium for 3 hours; neuronal survival was reported as a percentage of controls (vehicle treated neurones).

Cell extracts

Neurones were homogenised in a buffer containing 150 mM NaCl, 10 mM Tris-HCl, 10 mM EDTA, 0.5% Nonidet P-40, 0.5% sodium deoxycholate and mixed protease inhibitors (AEBSE, Aprotinin, Leupeptin, Bestain, Pepstatin A and E-46 (Sigma)) and a phosphatase inhibitor cocktail including PP1, PP2A, microcystin LR, cantharidin and p-bromotetramisole (Sigma) at 10⁶ cells/ml. Nuclei and cell debris was removed by low speed centrifugation (300 × g for 5 minutes). For immunoblots, cells were mixed 1:1 with Laemmli buffer containing β-mercaptoethanol, boiled for 5 minutes and run on a 12% polyacrylamide gel. Proteins were transferred onto a Hybond-P PVDF membrane (Amersham Biotech) by semi-dry blotting. Membranes were blocked using 10% milk powder; synaptophysin was detected using a mouse monoclonal antibody (mAb) anti-synaptophysin SY38 (Abcam) and β-actin was detected by incubation with a mouse mAb (clone AC-74, Sigma). These were visualised using a combination of biotinylated rabbit anti-mouse IgG (Dako), extravidin-peroxidase and an enhanced chemiluminescence kit (Invitrogen).

Synaptophysin ELISA

Synaptophysin levels in cell extracts were measured by ELISA as described [9,27]. A mouse monoclonal anti-synaptophysin MAB368 (Millipore) was used as a capture antibody and bound synaptophysin was detected using rabbit polyclonal anti-synaptophysin (Abcam) followed by a biotinylated anti-rabbit IgG, extravidin-alkaline phosphatase and 1 mg/ml 4-nitrophenol phosphate (Sigma). Absorbance was measured on a microplate reader at 405 nm and synaptophysin content was calculated from a standard curve. Samples were expressed as "units synaptophysin" where 100 units was defined as the amount of synaptophysin in 10⁶ vehicle treated neurones. A standard curve was generated from this sample using sequential log 2 dilutions (range 100 to 1.56 units).

Synaptosome preparations

Cortical neurones were pre-treated with drugs for 3 hours prior to the addition of 100 nM PrP82-146. After different time points (10, 30, 60 or 90 minutes) neurones were washed 3 times to remove unbound peptide and synaptosomes prepared on a discontinuous Percoll gradient as described [60]. Briefly, 10⁶ cortical neurones were homogenized at 4°C in 1 ml of SED solution (0.32 M sucrose, 50 mM Tris-HCl, 1 mM EDTA, and 1 mM dithiothreitol, pH 7.4 and mixed protease/phosphatase inhibitors (as above)). The preparation was centrifuged at 1000 × g for 10 minutes. The supernatant was transferred to a 4-step gradient of 3, 7, 15, and 23% Percoll in SED solution and centrifuged at 16,000 × g for 30 minutes at

4°C. The synaptosome fraction was collected from the interface of the 15% and 23% Percoll steps, washed twice (16,000 × *g* for 30 minutes at 4°C) and suspended in extraction buffer (150 mM NaCl, 10 mM Tris-HCl, 10 mM EDTA, 0.2% SDS and mixed protease/phosphatase inhibitors).

PrP82-146 ELISA

The amount of PrP82-146 in synaptosomes was determined by ELISA. Nunc Maxisorb Immunoplates were coated with 0.5 µg/ml of mouse mAb 3F4 (reactive with residues 109-112 of human PrP (Abcam)) this mAb does not bind to murine PrP [61]. Samples were applied and PrP82-146 was detected with 0.5 µg/ml biotinylated IC5M35 (D-gen, <http://www.d-gen.co.uk>), followed by streptavidin-alkaline phosphatase (Dako) and 1 mg/ml 4-nitrophenyl phosphate (Sigma). Absorbance was measured on a microplate reader at 405 nm and the amount of PrP82-146 in cell extracts was calculated by reference to a standard curve of PrP82-146.

cPLA₂ ELISA

The amount of cPLA₂ in synaptosomes was measured by ELISA [62]. Nunc Maxisorb Immunoplates were coated with 0.5 µg/ml of mouse mAb anti-cPLA₂, clone CH-7 (Upstate) and blocked with 10% FCS. Samples were incubated for 1 hour and the amount of cPLA₂ was detected using a goat polyclonal anti-cPLA₂ (Santa-Cruz Biotech). Bound antibodies were detected with biotinylated anti-goat IgG, extravidin-alkaline phosphatase and 1 mg/ml 4-nitrophenyl phosphate. Absorbance was measured at 405 nm and the amount of cPLA₂ was calculated from a standard curve using nonlinear regression. The amount of cPLA₂ protein was expressed in units (100 units = amount of cPLA₂ in synaptosomes isolated from 10⁶ untreated primary cortical neurones). The activation of cPLA₂ is accompanied by phosphorylation of the 505 serine residue which creates a unique epitope. To measure the amount of activated cPLA₂, an ELISA using a mAb (anti-cPLA₂, clone CH-7) combined with rabbit polyclonal anti-phospho-cPLA₂ (Cell Signalling Technology) was used. Bound antibodies were detected with biotinylated anti-rabbit IgG (Dako), extravidin-alkaline phosphatase and 1 mg/ml 4-nitrophenyl phosphate. Samples were expressed as "units activated cPLA₂" where 100 units were defined as the amount of activated cPLA₂ in synaptosomes derived from 10⁶ untreated primary cortical neurones.

Drugs

U73122 and Ethyl-18-OCH₃ were obtained from Biomol. Arachidonyl trifluoromethyl ketone (AACOCF₃), aristo-

lochic acids, ginkgolide B, 1-O-Hexadecyl-2-acetyl-*sn*-glycerol-3-phospho-(N,N,N-trimethyl)-hexanolamine (Hexa-PAF), methyl arachidonyl fluorophosphonate (MAFP), prostaglandin G₂, AH13205 and BWA868C were obtained from Sigma. Prostaglandins 15d-J₂, E₂, D₂, F₂α, H₂ and I₂ were obtained from Novabiochem. Stock solutions were dissolved in ethanol or di-methyl sulphoxide (DMSO) and diluted in medium to obtain final working concentrations. Vehicle controls consisted of equivalent dilutions of ethanol or DMSO in NBM.

Peptides

Peptides containing amino acids 82 to 146 of the human PrP protein (PrP82-146) corresponding to a PrP fragment found in certain prion-infected human brains [7], a control peptide (PrP82-146scrambled) were synthesised by solid-phase chemistry and purified by reverse-phase high performance liquid chromatography. A synthetic peptide containing the amino acid residues 1 to 42 of Aβ (Aβ₁₋₄₂) of the Aβ protein and a control peptide consisting of the same amino acids in reverse order (Aβ₄₂₋₁) were obtained from Bachem. Aβ peptides were first dissolved in hexafluoroisopropanol, lyophilised and subsequently solubilised and stored in DMSO. PLA₂ Activating Peptide (PLAP) was purchased from Bachem. Stock solutions of peptides were thawed on the day of use and sonicated before dilution in culture medium and addition to cells.

CHO cells stably transfected with a cDNA encoding APP₇₅₁ containing the Val717Phe familial AD mutation (referred to as 7PA2 cells) were cultured in DMEM with 10% FCS [28]. Conditioned medium (CM) from these cells contains stable Aβ oligomers (7PA2-CM). Conditioned medium from non-transfected CHO cells (CHO-CM) was used as controls.

Statistical Methods

Differences between treatment groups were assessed using two sample t tests or analysis of variance techniques. In all tests statistical significance was set at the 1% level.

List of abbreviations

(AD): Alzheimer's disease; (Aβ): amyloid-β; (AACOCF₃): Arachidonyl trifluoromethyl ketone; (CHO): Chinese hamster ovary; (CM): conditioned medium; (CJD): Creutzfeldt-Jakob disease; (DMSO): di-methyl sulphoxide; (ELISA): Enzyme linked immunoassay; (Hexa-PAF): 1-O-Hexadecyl-2-acetyl-*sn*-glycerol-3-phospho-(N,N,N-trimethyl)-hexanolamine; (MAFP): methyl arachidonyl fluorophosphonate; (NBM): neurobasal medium; (PBS): phosphate buffered saline; (PLA₂): phospholipase A₂; (cPLA₂): cytoplasmic PLA₂; (PLAP): PLA₂ Activating

Peptide; (PAF): platelet-activating factor; (PrP): prion protein.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CB and MT were responsible for the conception, planning and performance of experiments and for writing the manuscript. AW contributed to the planning of experiments, interpretation of results and the writing of the manuscript. All authors approved the final manuscript.

Acknowledgements

This work was supported by the European Commission FP6 "Neuroprior" Network of Excellence. We thank Professor Mario Salmons (Milan) for supplying PrP82-146 and PrP82-146scrambled peptides.

Author Details

¹Department of Pathology and Infectious Diseases, Royal Veterinary College, Hawkshead Lane, North Mymms, Herts, AL9 7TA, UK and ²Department of Veterinary Medicine, University of Cambridge, Madingley Road, Cambridge, CB3 0ES, UK

Received: 27 January 2010 Accepted: 8 April 2010
Published: 8 April 2010

References

- Cunningham C, Deacon R, Wells H, Boche D, Waters S, Diniz CP, Scott H, Rawlins JN, Perry VH: **Synaptic changes characterize early behavioural signs in the ME7 model of murine prion disease.** *Eur J Neurosci* 2003, **17**(10):2147-2155.
- Jeffrey M, Halliday WG, Bell J, Johnston AR, MacLeod NK, Ingham C, Sayers AR, Brown DA, Fraser JR: **Synapse loss associated with abnormal PrP precedes neuronal degeneration in the scrapie-infected murine hippocampus.** *Neuropathol Appl Neurobiol* 2000, **26**(1):41-54.
- Johnston AR, Fraser JR, Jeffrey M, MacLeod N: **Synaptic Plasticity in the CA1 Area of the Hippocampus of Scrapie-Infected Mice.** *Neurobiology of Disease* 1998, **5**(3):188-195.
- Prusiner SB: **Prions.** *Proc Natl Acad Sci USA* 1998, **95**(23):13363-13383.
- Siso S, Puig B, Varea R, Vidal E, Acin C, Prinz M, Montrasio F, Badiola J, Aguzzi A, Pumarola M, et al.: **Abnormal synaptic protein expression and cell death in murine scrapie.** *Acta Neuropathol* 2002, **103**(6):615-626.
- Ferrer I, Puig B, Blanco R, Martí E: **Prion protein deposition and abnormal synaptic protein expression in the cerebellum in Creutzfeldt-Jakob disease.** *Neuroscience* 2000, **97**(4):715-726.
- Salmons M, Morbin M, Massignan T, Colombo L, Mazzoleni G, Capobianco R, Diomedea L, Thaler F, Mollica L, Musco G, et al.: **Structural properties of Gerstmann-Strausler-Scheinker disease amyloid protein.** *J Biol Chem* 2003, **278**(48):48146-48153.
- Bate C, Tayebi M, Salmons M, Diomedea L, Williams A: **Polyunsaturated fatty acids protect against prion-mediated synapse damage in vitro.** *Neurotox Res* 2010, **17**(3):203-214.
- Lipton AM, Cullum CM, Satumtira S, Sontag E, Hynan LS, White CL, Bigio EH: **Contribution of asymmetric synapse loss to lateralizing clinical deficits in frontotemporal dementias.** *Arch Neurol* 2001, **58**(8):1233-1239.
- Elferink LA, Scheller RH: **Synaptic vesicle proteins and regulated exocytosis.** *J Cell Sci* 1993, **17**:75-79.
- Weimer RM, Jorgensen EM: **Controversies in synaptic vesicle exocytosis.** *J Cell Sci* 2003, **116**(18):3661-3666.
- Wiedenmann B, Franke WW: **Identification and localization of synaptophysin, an integral membrane glycoprotein of Mr 38,000 characteristic of presynaptic vesicles.** *Cell* 1985, **41**(3):1017-1028.
- Daly C, Sugimori M, Moreira JE, Ziff EB, Llinas R: **Synaptophysin regulates clathrin-independent endocytosis of synaptic vesicles.** *Proc Natl Acad Sci USA* 2000, **97**(11):6120-6125.
- Reddy PH, Mani G, Park BS, Jacques J, Murdoch G, Whetsell W Jr, Kaye J, Manczak M: **Differential loss of synaptic proteins in Alzheimer's disease: implications for synaptic dysfunction.** *J Alzheimers Dis* 2005, **7**(2):103-117.
- Counts SE, Nadeem M, Lad SP, Wu J, Mufson EJ: **Differential expression of synaptic proteins in the frontal and temporal cortex of elderly subjects with mild cognitive impairment.** *J Neuropathol Exp Neurol* 2006, **65**(6):592-601.
- Selkoe DJ: **Alzheimer's Disease Is a Synaptic Failure.** *Science* 2002, **298**(5594):789-791.
- Heinonen O, Soininen H, Sorvari H, Kosunen O, Paljarvi L, Koivisto E, Riekkinen Sr: **Loss of synaptophysin-like immunoreactivity in the hippocampal formation is an early phenomenon in Alzheimer's disease.** *Neuroscience* 1995, **64**(2):375-384.
- Terry RD, Masliah E, Salmon DP, Butters N, DeTeresa R, Hill R, Hansen LA, Katzman R: **Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment.** *Ann Neurol* 1991, **30**(4):572-580.
- DeKosky ST, Scheff SW: **Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity.** *Ann Neurol* 1990, **27**(5):457-464.
- Hamos JE, DeGennaro LJ, Drachman DA: **Synaptic loss in Alzheimer's disease and other dementias.** *Neurology* 1989, **39**(3):355-361.
- Vassar R, Citron M: **A β -Generating Enzymes: Recent Advances in β and γ -Secretase Research.** *Neuron* 2000, **27**(3):419-422.
- Hardy J, Selkoe DJ: **The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics.** *Science* 2002, **297**(5580):353-356.
- Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, et al.: **Diffusible, nonfibrillar ligands derived from A β ₁₋₄₂ are potent central nervous system neurotoxins.** *Proc Natl Acad Sci USA* 1998, **95**(11):6448-6453.
- Chromy BA, Nowak RJ, Lambert MP, Viola KL, Chang L, Velasco PT, Jones BW, Fernandez SJ, Lacor PN, Horowitz P, et al.: **Self-assembly of A β ₁₋₄₂ into globular neurotoxins.** *Biochemistry* 2003, **42**(44):12749-12760.
- Takahashi RH, Almeida CG, Kearney PF, Yu F, Lin MT, Milner TA, Gouras GK: **Oligomerization of Alzheimer's β -Amyloid within Processes and Synapses of Cultured Neurons and Brain.** *J Neurosci* 2004, **24**(14):3592-3599.
- Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, Brett FM, Farrell MA, Rowan MJ, Lemere CA, et al.: **Amyloid- β protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory.** *Nat Med* 2008, **14**(8):837-842.
- Bate C, Tayebi M, Williams A: **Ginkgolides protect against amyloid- β ₁₋₄₂-mediated synapse damage in vitro.** *Mol Neurodegen* 2008, **3**(1):1.
- Podlisny MB, Ostaszewski BL, Squazzo SL, Koo EH, Rydel RE, Teplow DB, Selkoe DJ: **Aggregation of secreted amyloid β -protein into sodium dodecyl sulfate-stable oligomers in cell culture.** *J Biol Chem* 1995, **270**(16):9564-9570.
- Pilane CM, LaBelle EF: **cPLA₂ activator peptide, PLAP, increases arachidonic acid release and apoptosis of vascular smooth muscle cells.** *J Cell Physiol* 2004, **198**(1):48-52.
- Sattar AA, Haque R: **Cytosolic PLA₂ in zymogen granule fusion and amylase release: inhibition of GTP-induced fusion by arachidonyl trifluoromethyl ketone points to cPLA₂ in G-protein-mediated secretory vesicle fusion.** *J Biochem* 2007, **141**(1):77-84.
- Francescangeli E, Domanska-Janik K, Goracci G: **Relative contribution of the de novo and remodelling pathways to the synthesis of platelet-activating factor in brain areas and during ischemia.** *J Lipid Mediat Cell Signal* 1996, **14**(1-3):89-98.
- Teixeira MM, al Rashed S, Rossi AG, Hellewell PG: **Characterization of the prostanoid receptors mediating inhibition of PAF-induced aggregation of guinea-pig eosinophils.** *Br J Pharmacol* 1997, **121**(1):77-82.
- Sugimoto Y, Narumiya S: **Prostaglandin E Receptors.** *J Biol Chem* 2007, **282**(16):11613-11617.
- Mallucci GR, White MD, Farmer M, Dickinson A, Khatun H, Powell AD, Brandner S, Jefferys JG, Collinge J: **Targeting cellular prion protein reverses early cognitive deficits and neurophysiological dysfunction in prion-infected mice.** *Neuron* 2007, **53**(3):325-335.
- Kovacs GG, Preusser M, Strohschneider M, Budka H: **Subcellular Localization of Disease-Associated Prion Protein in the Human Brain.** *Am J Pathol* 2005, **166**(1):287-294.
- Kitamoto T, Shin RW, Doh-Ura K, Tomokane N, Miyazono M, Muramoto T, Tateishi J: **Abnormal isoform of prion proteins accumulates in the**

- synaptic structures of the central nervous system in patients with Creutzfeldt-Jakob disease. .
37. Moskowitz N, Schook W, Puszkin S: **Regulation of endogenous calcium-dependent synaptic membrane phospholipase A₂.** *Brain Res* 1984, **290**(2):273-279.
 38. Moskowitz N, Schook W, Puszkin S: **Interaction of brain synaptic vesicles induced by endogenous Ca²⁺-dependent phospholipase A₂.** *Science* 1982, **216**(4543):305-307.
 39. Sun GY, Xu J, Jensen MD, Simonyi A: **Phospholipase A₂ in the central nervous system: implications for neurodegenerative diseases.** *J Lipid Res* 2004, **45**(2):205-213.
 40. Marcheselli VL, Rossowska MJ, Domingo MT, Braquet P, Bazan NG: **Distinct platelet-activating factor binding sites in synaptic endings and in intracellular membranes of rat cerebral cortex.** *J Biol Chem* 1990, **265**(16):9140-9145.
 41. Clark GD, Happel LT, Zorumski CF, Bazan NG: **Enhancement of hippocampal excitatory synaptic transmission by platelet-activating factor.** *Neuron* 1992, **9**(6):1211-1216.
 42. Farooqui AA, Horrocks LA: **Phospholipase A₂-Generated Lipid Mediators in the Brain: The Good, the Bad, and the Ugly.** *The Neuroscientist* 2006, **12**(3):245-260.
 43. Bellizzi MJ, Lu SM, Masliah E, Gelbard HA: **Synaptic activity becomes excitotoxic in neurons exposed to elevated levels of platelet-activating factor.** *J Clin Invest* 2005, **115**(11):3185-3192.
 44. Williams A, Van Dam AM, Ritchie D, Eikelenboom P, Fraser H: **Immunocytochemical appearance of cytokines, prostaglandin E₂ and lipocortin-1 in the CNS during the incubation period of murine scrapie correlates with progressive PrP accumulations.** *Brain Res* 1997, **754**(1-2):171-180.
 45. Minghetti L, Cardone F, Greco A, Puopolo M, Levi G, Green AJ, Knight R, Pocchiari M: **Increased CSF levels of prostaglandin E(2) in variant Creutzfeldt-Jakob disease.** *Neurology* 2002, **58**(1):127-129.
 46. Sang N, Zhang J, Marcheselli V, Bazan NG, Chen C: **Postsynaptically Synthesized Prostaglandin E2 (PGE2) Modulates Hippocampal Synaptic Transmission via a Presynaptic PGE2 EP2 Receptor.** *J Neuroscience* 2005, **25**(43):9858-9870.
 47. Wilson RJ, Rhodes SA, Wood RL, Shield VJ, Noel LS, Gray DW, Giles H: **Functional pharmacology of human prostanoid EP2 and EP4 receptors.** *Eur J Pharmacol* 2004, **501**(1-3):49-58.
 48. Lacor PN, Buniel MC, Furlow PW, Sanz Clemente A, Velasco PT, Wood M, Viola KL, Klein WL: **Aβ Oligomer-Induced Aberrations in Synapse Composition, Shape, and Density Provide a Molecular Basis for Loss of Connectivity in Alzheimer's Disease.** *J Neurosci* 2007, **27**(4):796-807.
 49. Shankar GM, Bloodgood BL, Townsend M, Walsh DM, Selkoe DJ, Sabatini BL: **Natural Oligomers of the Alzheimer Amyloid-β Protein Induce Reversible Synapse Loss by Modulating an NMDA-Type Glutamate Receptor-Dependent Signaling Pathway.** *J Neurosci* 2007, **27**(11):2866-2875.
 50. Haass C, Selkoe DJ: **Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid β-peptide.** *Nat Rev Mol Cell Biol* 2007, **8**(2):101-112.
 51. Kriem B, Sponne I, Fifre A, Malaplate-Armand C, Lozac'h-Pillot K, Koziel V, Yen-Potin FT, Bihain B, Oster T, Olivier JL, et al.: **Cytosolic phospholipase A₂ mediates neuronal apoptosis induced by soluble oligomers of the amyloid-β peptide.** *The FASEB Journal* 2004.
 52. Malaplate-Armand C, Florent-Bechard S, Youssef I, Koziel V, Sponne I, Kriem B, Leininger-Muller B, Olivier J-L, Oster T, Pillot T: **Soluble oligomers of amyloid-β peptide induce neuronal apoptosis by activating a cPLA₂-dependent sphingomyelinase-ceramide pathway.** *Neurobiol Dis* 2006, **23**(1):178-189.
 53. Sanchez-Mejia RO, Newman JW, Toh S, Yu G-Q, Zhou Y, Halabisky B, Cisse M, Scearce-Levie K, Cheng IH, Gan L, et al.: **Phospholipase A₂ reduction ameliorates cognitive deficits in a mouse model of Alzheimer's disease.** *Nat Neurosci* 2008, **11**(11):1311-1318.
 54. Lauren J, Gimbel DA, Nygaard HB, Gilbert JW, Strittmatter SM: **Cellular prion protein mediates impairment of synaptic plasticity by amyloid-β oligomers.** *Nature* 2009, **457**(7233):1128-1132.
 55. Lee KS, Linden R, Prado MAM, Brentani RR, Martins VR: **Towards cellular receptors for prions.** *Reviews in Medical Virology* 2003, **13**(6):399-408.
 56. Dahlgren KN, Manelli AM, Stine WB Jr, Baker LK, Krafft GA, LaDu MJ: **Oligomeric and Fibrillar Species of Amyloid-β Peptides Differentially Affect Neuronal Viability.** *J Biol Chem* 2002, **277**(35):32046-32053.
 57. Simoneau S, Rezaei H, Sales N, Kaiser-Schulz G, Lefebvre-Roque M, Vidal C, Fournier JG, Comte J, Wopfner F, Grosclaude J, et al.: **In vitro and in vivo neurotoxicity of prion protein oligomers.** *PLoS Pathog* 2007, **3**(8):e125.
 58. Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ: **Naturally secreted oligomers of amyloid β protein potently inhibit hippocampal long-term potentiation in vivo.** *Nature* 2002, **416**(6880):535-539.
 59. Brewer GJ: **Isolation and culture of adult rat hippocampal neurons.** *J Neurosci Meth* 1997, **71**(2):143-155.
 60. Thais ME, Carqueja CL, Santos TG, Silva RV, Stroeh E, Machado RS, Wahlheim DO, Bianchin MM, Sakamoto AC, Brentani RR, et al.: **Synaptosomal glutamate release and uptake in mice lacking the cellular prion protein.** *Brain Res* 2006, **1075**(1):13-19.
 61. Kascsak RJ, Rubenstein R, Merz PA, Tonna-DeMasi M, Fersko R, Carp RI, Wisniewski HM, Diring H: **Mouse polyclonal and monoclonal antibody to scrapie-associated fibril proteins.** *J Virol* 1987, **61**(12):3688-3693.
 62. Bate C, Tayebi M, Williams A: **Sequestration of free cholesterol in cell membranes by prions correlates with cytoplasmic phospholipase A₂ activation.** *BMC Biol* 2008, **6**(1):8.

doi: 10.1186/1750-1326-5-13

Cite this article as: Bate et al., Phospholipase A2 inhibitors protect against prion and Aβ mediated synapse degeneration *Molecular Neurodegeneration* 2010, **5**:13

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

